Deceptive combined effects of short allele dominance and stuttering: an example

with *Ixodes scapularis*, the main vector of Lyme disease in the U.S.A.

Thierry De Meeûs[1,*], Cynthia T. Chan[2,3], John M. Ludwig[2,4], Jean I. Tsao[5], Jaymin Patel[2,6],

Jigar Bhagatwala[2,7], and Lorenza Beati[2].

1. Intertryp, IRD, Cirad, Univ Montpellier, Montpellier, France.

2. The U.S. National Tick Collection, Institute for Coastal Plain Science, Georgia Southern

University, Statesboro, GA, USA.

3. College of Agricultural & Environmental Sciences, University of Georgia, Athens, GA,

USA.

4. Department of Microbiology, University of Georgia, Athens, GA, USA

5. Department of Fisheries and Wildlife, Michigan State University, East Lansing, MI, USA

6. Division of Hospital Medicine, College of Medicine, University of Florida, Gainesville, FL,

USA.

7. Medical College of Georgia, Augusta University, Augusta, GA, USA.

* Corresponding author: thierry.demeeus@ird.fr

Keywords: Microsatellite loci, short allele dominance, stuttering, heterozygote deficit,

linkage disequilibrium, curing microsatellite data.

Running title: Short allele dominance and stuttering

Abstract

Null alleles, short allele dominance (SAD), and stuttering increase the perceived

relative inbreeding of individuals and subpopulations as measured by Wright's F_{IS} and F_{ST}.

Such amplifying problems are usually caused by inaccurate primer design (if developed

from a different species or a distant population), poor DNA quality, low DNA concentration,

or a combination of some or all these sources of inaccuracy. When combined, these

issues can increase the correlation between polymorphism at concerned loci and,

consequently, of linkage disequilibrium (LD) between those. In this note, we studied an

original microsatellite data set generated by analyzing nine loci in *Ixodes scapularis* ticks

from the eastern U.S.A. To detect null alleles and SAD we used correlation methods and

variation measures. To detect stuttering, we evaluated heterozygote deficit between alleles

displaying a single repeat difference. We demonstrated that an important proportion of loci

affected by amplification problems (one with null alleles, two with SAD and three with

stuttering) lead to highly significant heterozygote deficits (F_{IS}=0.12, *p*-value<0.0001). This

occurred together with a prohibitive proportion (31%) of pairs of loci in significant LD and a

significant variation in the measure of population subdivision across loci (Wright's F_{ST}).

This suggested a strong Wahlund effect and/or homogenizing selection at several loci. By

finding small peaks corresponding to previously disregarded larger alleles in some

homozygous profiles for loci with SAD and by pooling alleles close in size for loci with

stuttering, we generated an amended dataset. Except for one locus with null alleles, the

analyses of the corrected dataset revealed a significant excess of heterozygotes (F_{IS}=-

0.058) as expected in dioecious and strongly subdivided populations, with a more

reasonable proportion (17%) of pairs of loci characterized by significant LD. Strong

subdivision was also confirmed by the standardized F_{ST}' corrected for null alleles

(F_{ST}'=0.28).

Introduction

Null alleles, short allele dominance (SAD) and stuttering are frequent consequences of poor PCR amplifications, in particular for microsatellite markers. Amplification problems usually arise when primers are designed by using DNA of a different species or a distant population, when DNA is degraded or at too low of a concentration (Chapuis and Estoup, 2007), or any combination of these listed causes.

Null alleles occur when a mutation on the flanking sequence of the targeted locus affects the hybridization of the corresponding primer, resulting in amplification failure. Heterozygous individuals with one null allele falsely appear to be homozygous, while homozygous individuals for null alleles are considered to be missing data.

SAD, also called long allele dropout (Van Oosterhout et al., 2004), known from minisatellite markers, was also discovered to occur in microsatellite loci (Wattier et al., 1998). In heterozygous samples, longer alleles are less successfully amplified than shorter alleles through competition for Taq polymerase. This can lead to misinterpreting heterozygous individuals as homozygous for the shortest allele (De Meeûs et al., 2007).

Stuttering is the result of inaccurate PCR amplification through Taq slippage of a specific DNA strand. This generates several PCR products that differ from each other by one repeat and can cause difficulties when discriminating between fake and true homozygotes, such as heterozygous individuals for dinucleotide microsatellite allele sequences with a single repeat difference.

Allelic dropout is akin to SAD, but occurs randomly to any allele irrespective of its size.

The consequence of these issues is a homozygous excess when compared to the expected Castle-Weinberg proportions (Castle, 1903; Weinberg, 1908) measured by Wright's F_{IS} (Wright, 1965). These problems, like all others associated with amplification, are locus specific (Van Oosterhout et al., 2004; De Meeûs et al., 2007; De Meeûs, 2018) and thus lead to locus specific variation (namely, an increase) of F_{IS}. A less well known, though well documented (Chapuis and Estoup, 2007; Séré et al., 2017; Manangwa et al., 2019) effect of such amplification problems consists of an increase of Wright's F_{ST} (Wright, 1965) that is commonly used to measure the degree of genetic differentiation between subpopulations.

While an analytical cure exists for null alleles (Chapuis and Estoup, 2007; Séré et al., 2017), such remediation is unavailable for SAD and stuttering. To the best of our knowledge, the impact of amplification problems on linkage disequilibrium (LD) between locus pairs has yet to be investigated. When combined, the effect of the occurrence of null

alleles, SAD, and stuttering may artificially generate a positive correlation between allele
occurrences at affected loci and then increase the perceived LD between them.
In this note, we utilize an original data set generated through the analysis of nine
microsatellite loci in *Ixodes scapularis*, sampled across the eastern U.S. to show that the
combined effect of SAD, stuttering, and null alleles can induce an increase in the number
of locus pairs in significant LD. We then propose and test an efficient way to amend such
data.
96
Material and Methods
Sampling and DNA extraction
Larvae, nymphs, and adults of *I. scapularis* were sampled haphazardly from
different sites across the eastern U.S. on different occasions, extending from November
2001 to May 2014, by means of dragging and flagging the vegetation (Figure 1 and Table
1) (Rulison et al., 2013).
Immatures, particularly the larval offspring belonging to a single female, tend to
cluster on vegetation and can thus constitute different lineages within the same subsample
(Kempf et al., 2011). Consequently, to avoid possible Wahlund effects, where a
heterozygote deficit results from the admixture of individuals from genetically distant
subpopulations (e.g. see (De Meeûs, 2018)), we removed immature stages from the
present study. The remaining 387 adult ticks were subdivided into cohorts, with each
cohort comprising samples collected across two consecutive years in the fall, the following
winter and spring across the tick distribution range. This subdivision was based on
observations showing that Northeastern adults active in Fall can undergo winter
quiescence and resume activity in spring (Yuval and Spielman, 1990).
Many publications have emphasized the importance of mitochondrial clades in
different populations of *I. scapularis* across the U.S. (Norris et al., 1996; Qiu et al., 2002;
Sakamoto et al., 2014). Thus, to account for the mitochondrial clade representation and to
(again) avoid possible Wahlund effects, all ticks were assigned clades by phylogenetic
analysis of their 12S rDNA gene sequences. We identified 6 main clades in our dataset,
the previously identified American clade was subdivided in two lineages (AMI and AMII),
and the so-called southern clade was subdivided in 4 lineages (SOI, SOII, SOIII and SOIV)
(Table 1).
In conclusion, the combination of Site-Clade-Cohort data defined 45 subsamples
within the 387 individual adult ticks. Some subsamples included a small number of

individuals (1-4). While such subsamples were expected to exert a negligible weight on our
analyses, they were not eliminated.
Procedures for all DNA extractions followed published modified protocols (Beati and
Keirans, 2001) with a DNeasy Blood & Tissue Kit (Qiagen, Valencia, CA).
127
Selection and characterization of microsatellite markers
Thirteen of the first batches of genome sequences of *I. scapularis* that were
accessioned by VectorBase (www.vectorbase.org; Giraldo-Calderón *et al.* 2015) in
GenBank (AC205653.1, AC205652.1, AC205650.1, AC205647.1, AC205646.1,
AC205643.1, AC205642.1, AC205641.1, AC205638.1, AC205635.1, AC205634.1,
AC205632.1, AC205630.1) were used to manually detect motifs with at least 6 repeats of
AG, AT, CA, TA, TG, CT, GC, ACG, GTT, TTA, CAC, GAT, and AAAC. Primer pairs were
selected in the flanking regions by using Oligo v.5 (Molecular Biology Insights, Colorado
Springs, CO). DNA, extracted and pooled from six ticks from Connecticut, was used to test
whether the selected primer sets successfully amplified fragments of the expected size.
PCRs were performed using the 5-Prime Master PCR kit (5-Prime, Gaithersburg, MD) and
a single touch-down amplification protocol consisting of 5 min. denaturation at 93°C; 5
cycles: 20 sec. denaturation at 93°C, 20 sec. annealing at 55°C-1.5°C/cycle, 30 sec.
elongation at 72°C; 30 cycles: 20 sec. denaturation at 93°C, 25 sec. annealing at 47°C, 30
sec. elongation at 70°C; final extension at 70°C for 5 min. Amplicons were run on 4% E-
gels (Life Technologies Co., Carlsbad, CA). The risk of having selected primers within
repeated portions of the genome had to be considered due to the fact that large repeated
genomic fragments are known to occur abundantly in *I. scapularis* (Gulia-Nuss et al.,
2016). In order to confirm that the primers were amplifying the targeted loci, the amplicon
of one randomly chosen tick was cloned with a TOPO-TA PCR cloning kit (Life
Technologies Co, Carlsbad, CA) for each locus. Five cloned colonies were picked
randomly for each tick and the insert amplified and sequenced (DNA Analysis Facility on
Science Hill, Yale University). Finally, as microsatellite primers are known to amplify
sometimes more than one closely related species, the same set of primers was tested on
DNA samples of *Ixodes ricinus*, *Ixodes pacificus*, and *Ixodes persulcatus* (LB, personal
collection), all taxa belonging to the *I. ricinus* complex of ticks (Keirans et al., 1999).
The primer pairs that yielded amplicons of the expected size were then used to
individually amplify a subset of 67 DNA samples from ticks collected by flagging or
dragging in Alabama (10), Georgia (15), Connecticut (16), Massachusetts (14), New York
(2), Pennsylvania (2), and South Carolina (8) (Table 1). For these amplifications, forward

primers were labeled with fluorescent dyes (Applied Biosystems, Thermo Fisher Scientific,

CA) as listed in Table 2. The amplicons were sent to the DNA Analysis Facility on Science

Hill (Yale University, New Haven, CT) for genotyping. The allele peaks were scored using

GeneMarker (SoftGenetics, State College, PA). All data were recorded in an Excel

spreadsheet for further ease of conversion.

Figure 1: Sampling sites for *Ixodes scapularis* from the eastern U.S.A. (State codes as in
Table 1).

Table 1: State, site, GPS coordinates (decimal degrees), 12S clade membership, date of
sampling, corresponding cohort membership and size (*N*) of *Ixodes scapularis* adult
subsamples from the eastern U.S.A.

State	Site	latitude	longitude	Clade	Date	Cohort	N
	AL1	33.24	-86.13	AMI	2011 Jan	C7	21
	AL2	32.95	-87.14	AMI	2012 Dec	C9	3
Alabama	AL1	33.24	-86.13	AMII	2011 Jan	C7	18
	AL2	32.95	-87.14	AMII	2012 Dec	C9	2
	AL1	33.24	-86.13	SOI	2011 Jan	C7	6
	AL1	33.24	-86.13	SOII	2011 Jan	C7	6
Connecticut	CT1	41.35	-72.76	AMI	2001 Nov	C1	3
	CT1	41.35	-72.76	AMI	2003 Jun	C2	21
	FL2	30.06	-81.37	AMI	2011 Jan	C7	2
	FL2	30.06	-81.37	AMII	2011 Jan	C7	3
Florida	FL2	30.06	-81.37	SOI	2011 Jan	C7	1
	FL2	30.06	-81.37	SOII	2011 Jan	C7	4
	FL1	30.65	-84.21	SOII	2012 Dec	C9	4
	GA1	32.45	-81.78	AMI	2009 Dec	C5	11
Georgia	GA1	32.45	-81.78	SOI	2009 Dec	C5	5
	GA1	32.45	-81.78	SOII	2009 Dec	C5	17
	GA1	32.45	-81.78	SOIII	2009 Dec	C5	1
Iowa	IA1	42.67	-91.59	AMI	2007 May	C3	3
	LA1	30.94	-91.46	AMI	2012 Dec	C9	3
Louisiana	LA1	30.94	-91.46	AMII	2012 Dec	C9	2
	LA1	30.94	-91.46	SOIII	2012 Dec	C9	1
	LA1	30.94	-91.46	SOIV	2012 Dec	C9	2
Massachusetts	MA1	41.71	-69.92	AMI	2010 May	C5	6
Maryland	MD1	39.29	-76.88	AMI	2010 Oct	C7	11
	NC1	33.91	-78.39	AMI	2009 Jan	C4	22
North-Carolina	NC1	33.91	-78.39	SOI	2009 Jan	C4	4
	NC1	33.91	-78.39	SOIII	2009 Jan	C4	2
New-York	NY1	40.76	-72.83	AMI	2009 Oct	C5	10
	PA2	41.06	-79.48	AMI	2014 May	C10	33
Pennsylvania	PA1	40.67	-75.96	AMI	2010 Oct	C7	5

	SC1	33.33	-81.66	AMI	2011 Apr	C7	12
	SC5	34.29	-79.87	AMI	2012 Dec	C9	3
	SC1	33.33	-81.66	AMII	2011 Apr	C7	9
South-Carolina	SC1	33.33	-81.66	SOI	2011 Apr	C7	7
	SC1	33.33	-81.66	SOII	2011 Apr	C7	13
	SC5	34.29	-79.87	SOII	2012 Dec	C9	2
	SC1	33.33	-81.66	SOIII	2011 Apr	C7	1
Tennessee	TN1	35.37	-86.07	AMI	2010 Dec	C7	23
	TX2	30.24	-100.72	AMI	2012 Dec	C9	11
Texas	TX1	31.80	-96.23	SOI	2012 Dec	C9	6
	TX2	30.24	-100.72	SOIV	2012 Dec	C9	14
Virginia	VA1	38.29	-78.29	AMI	2010 Oct	C7	7
	VA2	37.32	-80.73	AMI	2012 Dec	C9	4
Wisconsin	WI1	43.95	-90.70	AMI	2011 Oct	C8	24
	WI2	44.04	-90.65	AMI	2012 Oct	C9	19

Table 1: List of primer sets used or developed for this study. PAC = *I. pacificus*, PER = *I. persulcatus*, RIC = *I. ricinus.* Approximate size range and number of alleles correspond to 67 individuals used for the initial tests.

Locus name	Repeat array	Primer name	Primer sequences (5'→3')	Reference or GenBank accession number	Dye	Approx. Size range (bp)	No of alleles	Cross-species amplification
IS1	(AG)10	Amy1-IsAG25a	AAATGTCCGAACAGCCTTAT	Fagerberg *et al* (2001)	6 FAM	93-193	17	PAC/PER/RIC
		Amy2-IsAG25b	GCCCTTGAGTCTACCCACTA					
IS3	(GTT)$_5$	bac1d_a	GCAGATCTCTTGGGCTAG	AC205653	VIC	76-100	7	none
		bac1d_b	AAGCTAAGGCGTTCGTTG					
IS4	(AT)21	bac1m_a	TGTCGGTTTGATGCCAA	AC205653	VIC	88-126	17	PAC/PER/RIC
		bac1m_b	GGCTCCATTCACCAGTC					
IS5	(CA)9	bac3dh_a	TGCCTGTGACGAAACCA	AC205650	NED	62-140	17	none
		bac3dh_b	TCTCCCAAGAGATCTAGGTA					
IS6	(TA)10	bac1j_a	TCTCCCAAGAGATCTAGGTA	AC205653	VIC	100-186	13	PER
		bac1j_b	ATCTGTTCAGTGGGCACA					
IS7	(TA)11	bac1k_a	GGGACTGGACACACGA	AC205653	VIC	48-170	26	none
		bac1k_b	CTAGGTGGCGCAAGTC					
IS8	(CA)14	bac3s_a	CGTTTCAAAGTCGGAGA	AC205650	PET	96-194	11	PER
		bac3s_b	GATGTGAGGGCGTGGT					
IS9	(AAAC)5	bac4cef_a	CGCCTTTTGTCCCAACC	AC205647	6 FAM	85-125	12	PER
		bac4cef_b	GACTAACAGCATTGGAGCA					
IS10	(TTA)9	bac5cf_a	TCCCCCAACAAGATTGATG	AC205646	6 FAM	77-137	15	none
		bac5cf_b	GAGACGACGTAGATTCTTG					
IS11	(TTA)$_6$	bac5g_a	GCTTTAGCGGGCTGGT	AC205646	PET	81-165	12	PER
		bac5g_b	TACGTGAATACGTCCTTGG					
IS12	(TA)43	bac6a_a	GCAAGCTTCGCTATTCTC	AC205643	6 FAM	111-229	26	none
		bac6a_b	CAGTAATTTCGCATCGGTT					
IS13	(TA)22	bac6c_a	TAGGTACAAGAAAACGTGCT	AC205643	NED	37-91	17	none
		bac6c_b	CAAGGTAATTGTTCTCGTCA					
IS14	(TA)5	bac6d_a	CCTTGCCTTACATGGTT	AC205643	HEX	57-105	13	PAC/PER/RIC

		bac6d_b	CGTACCAAACCAAAGCAAG					
IS15	$(AT)_8$	bac6e_a	TATTGTAACCGACGCTAGG					
		bac6e_b	GACAATCTCTACGCAAATCC	AC205643	NED	79-125	18	none
IS16	$(CA)_8$	bac6f_a	CCCCCAAACACGCACA					
		bac6f_b	TTGCTTCATGCAGGGAAC	AC205643	VIC	80-106	12	RIC
IS17	$(CA)_6$	bac7e_a	CCAGCATTTAACCCTCAAG					
		bac7e_b	TAGTGGGGTATGGCACTG	AC205642	HEX	139-197	12	PER/RIC
IS18	$(TG)_6$	bac8a_a	GTAGGTACCCTAAGAAGGAT		6			
		bac8a_b	TTGAGGAAGCAGAATGTAGG	AC205641	FAM	75-195	16	PER/RIC
IS19	$(CT)_7$	bac9a_a	AGAACCAGTTCAGCATTCC					
		bac9a_b	GAACATTTTCACGTGTTGC	AC205638	PET	94-166	6	PER
IS20	$(GC)9$	bac11a_a	CGCTCCCTTCGAAGTTC					
		bac11a_b	GAGAAGACAGTTTCCATCG	This study	HEX	76-106	13	PAC/PER/RIC
IS21	$(ACG)6$	bac11c_a	CGAATCGCGCACACTAG					
		bac11c_b	GCTGTGTTGCTGGTCAC	This study	NED	109-251	14	PAC/PER
IR27	$(AC)9$		ATACCCGTAGAACGAGAG	Delaye *et al*	6			
			GTTTTTCAAGATTTCCGCC	(2008)	FAM			RIC

Genotyping

Nine microsatellite loci (IR27, IS1, IS3, IS11, IS15, IS16, IS17, IS18, and IS19) were
used for genotyping at the continental scale (Table 2). Of these, IR27 (Delaye et al., 1998)
and IS1 (Fagerberg et al., 2001), were drawn from previously published studies. The loci
were amplified and genotyped using the procedures described above, although PCR
conditions had to be slightly optimized for markers IS11 and IS15 (touchdown annealing
temperature decreased from 58°C to 50°C) and IR27 (touchdown annealing temperature
decreased from 56°C to 53°C) (Table 2).

Population genetics analyses

The raw data set was coded and converted into all required formats using Create
(Coombs et al., 2008).

To test for LD, we used the *G*-based test first described by Goudet et al. (Goudet et
al., 1996), adapted for contingency tables of locus pairs with 10000 reshuffling of
genotypes (or 15000 when needed). The *G* statistics obtained for each subsample were
then summed over all subsamples to get a single statistic and hence, a single test across
subsamples. This procedure was shown to be the most powerful (De Meeûs et al., 2009)
and was implemented within Fstat 2.9.4 (Goudet, 2003) an updated version of the original
1995 Fstat software (Goudet, 1995). There are as many tests as locus pairs and these
tests are correlated (one locus is used as many times as there is any other locus). To take
into account this repetition of correlated tests, we used Benjamini and Yekutieli (BY) false
discovery rate procedure (Benjamini and Yekutieli, 2001) with R version 3.5.1 (R-Core-
Team, 2018). We also undertook a Fisher exact test with Rcmdr version 2.3-1 (Fox, 2005;
Fox, 2007) when examining the eight tests in which each locus was involved in, in order to
see if some pairs of loci were found in significant LD more often than by chance.

For a hierarchy with three levels (individuals, subsamples, and total sample), three
F-statistics can be defined (Wright, 1965). F_{IS} measures inbreeding of individuals relative
to inbreeding of subsamples or relative deviation of genotypic proportions from local
random mating proportions. F_{ST} measures inbreeding of subsamples relative to total
inbreeding or relative inbreeding due to the subdivision of the total population into several
isolated subpopulations. F_{IT} measures inbreeding of individuals relative to total inbreeding.
Under the null hypothesis (panmixia and no subdivision), all these statistics are expected
to be null. Otherwise, F_{IS} and F_{IT} can vary from -1 (one heterozygote class) to +1 (all
individuals homozygous) and F_{ST} from 0 (all subsamples share similar allele frequencies)

to +1 (all subsamples fixed for one or the other allele). These statistics were estimated with

Weir and Cockerham's unbiased estimators (Weir and Cockerham, 1984) with Fstat.

In dioecious species (like ticks), heterozygote excess occurs over all loci (e.g. (De

Meeûs et al. 2007)) and is proportional to subpopulation size (N_e=-1/(2×F_{IS})-F_{IS}/(1+F_{IS}))

(Balloux, 2004). Therefore, the finding of homozygous excesses really represents a strong

deviation from random mating expectations. Technical problems, like null alleles,

stuttering, SAD or allele dropouts unevenly affects some loci, producing a positive F_{IS} with

an important variation across loci with significant outliers (De Meeûs 2018). Significant

departure from 0 of these F-statistics was tested with 10000 randomizations of alleles

between individuals within subsample (deviation from local random mating test) or of

individuals between subsamples within the total sample (population subdivision test). For

F_{IS}, the statistic used was f (Weir and Cockerham's F_{IS} estimator). To test for subdivision,

we used the G-based test (Goudet et al. 1996) over all loci, which is the most powerful

procedure when combining tests across loci (De Meeûs et al. 2009).

To compute 95% confidence intervals (95%CI) of F-statistics, we used the standard

error of F_{IS} (StrdErrFIS) and F_{ST} (StrdErrFST) computed by jackknifing over populations, or

5000 bootstraps over loci as described elsewhere (De Meeûs et al. 2007). For jackknives,

the number of usable subsamples was restricted to subsamples with at least 5 ticks (23

subsamples) (e.g. (De Meeûs, 2012) p 73).

In case of significant homozygote excess and LD we tried to discriminate

demographic from technical causes with the determination key proposed by De Meeûs (De

Meeûs 2018). Null alleles better explain the data if the StrdErrFIS becomes at least twice

as high as StrdErrFST; F_{IS} and F_{ST} are positively correlated; and a positive correlation

links F_{IS} and the number of missing data (putative null homozygotes). The significance of

correlations was tested with a unilateral (ρ>0) Spearman's rank correlation test with R. The

presence of null alleles was also verified with MicroChecker v 2.2.3 (Van Oosterhout et al.

2004) and null allele frequencies estimated with Bookfield's second method (Brookfield

1996). The adjustment between observed and expected numbers of missing data was

tested with a unilateral exact binomial test in R with the alternative hypothesis being "there

is not enough missing data as expected if heterozygote deficits were entirely explained by

null alleles under panmixia". MicroChecker also detects stuttering and SAD. Stuttering is

detected when MicroChecker reveals an observed proportion of heterozygous individuals

for alleles with one repeat difference significantly smaller than the expected value. The

occurrence of SAD was also checked with an unilateral (ρ<0) Spearman's rank correlation

test between allele size and F_{IT} as proposed by Manangwa et al. (Manangwa et al., 2019).

This test is more powerful than with F_{IS} as was proposed earlier (Wattier et al., 1998; De

Meeûs et al., 2004). If previous tests are not significant and if StrdErrFIS > StrdErrFST,

then a Wahlund effect better explains the data (De Meeûs, 2018), this is especially true in

instances of significant LD (Manangwa et al., 2019). In these cases, a positive correlation

between the number of times a locus is found in significant LD (NLD) and its total genetic

diversity as measured by Nei's H_T (Nei and Chesser, 1983) (Spearman's correlation above

0.1) suggests an admixture of individuals from several subpopulations of the same species

but with an important degree of genetic differentiation between admixed subpopulations

(i.e. number of immigrants N_em=2). If the correlation is negative and the proportion of

significant LDs is above 40%, an admixture of different strongly divergent entities (e.g.

species) better explains the data (Manangwa et al., 2019). We thus undertook a bilateral

Spearman's test.

When diagnosed, there is no analytical remedy for SAD or stuttering as there is for

null alleles (Chapuis and Estoup, 2007; Séré et al., 2017). Nevertheless, SAD can be

cured by going back to the chromatograms of homozygous individuals and trying to find a

micro-peak (see the Results and discussion section), with a larger size, ignored in the first

reading. If enough profiles can be corrected this might salvage the incriminated locus.

Stuttering can be addressed by pooling alleles close in size. However, this procedure

requires that none of the pooled allele groups is constituted of rare alleles only. Indeed,

pooling rare alleles, usually found in heterozygosity with a more frequent allele, will tend to

artificially generate heterozygous excesses. In order to avoid this consequence, each

pooled group should contain at least one frequent allele (e.g. with $p≥0.05$).

In dioecious small populations, a heterozygote excess is expected. However, loci

with null alleles may display heterozygote deficits. In such situations a bilateral test (F_{IS} is

not different from 0) is needed and obtained as $p_{bilateral}=p_{mini}+1-p_{maxi}$, where p_{mini} is the

minimum unilateral p-value (for heterozygote deficit or excess) and p_{max} is the maximum

one.

Finally, a more accurate estimate of F_{ST} can be made for datasets with null alleles

after recoding missing genotypes as homozygous for allele 999 (null allele) with the ENA

method as implemented in FreeNA (Chapuis and Estoup, 2007). This estimate can then

be corrected for polymorphism with the formula $F_{ST}'=F_{ST}/(1-H_S)$ (Hedrick, 1999).

Results and discussion

Primer selection and characterization of loci

The inspection of the GenBank genomic sequences revealed the presence of 67

short tandem repeated motifs. The program Oligo v.5 did not find suitable primers for 17 of

them. Of the remaining 50 primer pairs, 22 amplified the pooled DNA sample and the sizes

of the amplicons were approximately as expected. The cloned amplified inserts all

contained the expected microsatellite repeats and flanking regions. The 22 primer sets

consistently amplified DNA from the 67 *I. scapularis* ticks and some of them also amplified

DNA of the other *Ixodes* species (Table 2).

Raw data analyses

All data are available in the supplementary File S1.

In order to maximize precision, LD tests were implemented over 15000

permutations. There was a very important proportion of locus pairs in significant LD

(~31%), with a negative correlation between NLD and H_T (ρ=-0.10, *p*-value=0.7974). None

of these tests remained significant after BY correction, although one pair displayed a very

low *p*-value (0.06). No single locus was found more often in significant LD than the others

(*p*-value=0.1626).

There was a highly significant heterozygote deficit (F_{IS}=0.115, *p*-value<0.0001), with

a considerable variation across loci (Figure 2).

299	Figure 2: F_{IS} values for each locus and averaged across those (All) of *Ixodes scapularis*
300	from the eastern U.S.A. with 95% jackknife confidence intervals over subsamples
301	(for each locus) and bootstraps over loci (All). Results of tests for panmixia, short
302	allele dominance and number of subsamples with stuttering for each locus are also
303	indicated.

Locus	IR27	IS1	IS3	IS11	IS15	IS16	IS17	IS18	IS19	All
p-value for F_{IS} not > 0	0.3825	0.0001	0.9992	0.0001	0.0001	0.1872	0.0004	0.9999	0.1038	0.0001
p-value for short allele dominance	0.2895	0.0008	0.6128	0.7303	0.0059	0.6377	0.9642	0.5078	0.3962	
Number of subsamples with stuttering	0	0	0	0	3	3	1	0	2	

StdrdErrFIS (0.07) was almost four times higher than StrdErrFST (0.019); the correlation between F_{IS} and F_{ST} was negative (ρ=-0.55, *p*-value=0.9397) and the correlation between F_{IS} and the number of blanks was positive but not significant (ρ=0.18, *p*-value=0.3218). These results suggested that locus-specific effects were involved. Nevertheless, null alleles only weakly explained the observed pattern. The substantial proportion of significant LDs suggested the existence of a strong Wahlund effect. Small subsamples due to the partitioning of the data into cohorts and 12S clades was expected to considerably lower the power of these tests, which may also explain why none of them remained significant after BY correction, which is a rather stringent procedure.

Subdivision was substantial (F_{ST}=0.098, 95% CI [0.063,0.133], *p*-value<0.0001), but highly variable across loci (Figure 3). Some loci (IS1, IS15 and IS17) displayed very low values (Figure 3). Interestingly, loci IS15 and IS17 were found four and two times in significant LD, respectively. Homogenizing selection (overdominance, balanced selection) might have produced the pattern observed at loci IS15 and IS17.

Figure 3: F_{ST} values for each locus and averaged across those (All) of *Ixodes scapularis*
from the eastern U.S.A. with 95% jackknife confidence intervals over subsamples
(for each locus) and bootstrap over loci (All). Results of tests for significant
subdivision (*p*-value) are also indicated.

Two loci displayed highly significant SAD (Figure 2) (loci IS1 and IS15) and
stuttering was diagnosed for four loci (IS15, IS16, IS17 and IS19).

Cured data set

To prevent withdrawing five (out of nine) loci, due to stuttering, SAD and/or possible
selection, we went back to the data. We first scanned the chromatograms for previously
ignored micro-peaks in homozygous individuals at loci displaying SAD (IS1 and IS15)
(Figure 4). SAD in IS15 might well have explained why we also found stuttering at this
locus (see below). We then tried to pool alleles close in size as described above for loci
IS16, IS17 and IS19. For IS16, alleles 88, 90 and 92 were recoded as 94; and allele 96 as
98. For locus IS17, alleles 170 and 172 were recoded as 174; alleles 178, 180, 182 and
184 were recoded as 186; alleles 188, 190 and 192 were recoded as 194; and alleles 196
and 198 as 200. Finally, for locus IS19, allele 91 was recoded as 93; alleles 97, 99, 101
and 103 as105; and alleles 107 and 109 as 111. The obtained amended dataset was
called "Cured dataset" (Supplementary file S1).

| 343 | Figure 4: Examples of an initially dismissed micro-peak that produced SAD at locus IS15 |
| 344 | and of stuttering at locus IS19 |

| 345 | |
| 346 | |

With the cured dataset, the proportion of locus pairs in significant LD dropped to
17% and the smallest p-value after BY correction was 0.1052. In particular, for loci IS15
and IS17, significant LD occurred only once for the former and never for the latter. The
correlation between NLD and H_T was much higher (ρ=0.27, p-value=0.4742). A
heterozygote deficit was no longer observed (F_{IS}=0, bilateral p-value=0.951) (Figure 5).
StrdErrFIS=0.063 was three times higher than StrdErrFST=0.021. There was no
correlation between F_{IS} and F_{ST} (ρ=-0.28, p-value=0.78). The correlation between number
of missing genotypes and F_{IS} was positive though marginally not significant (ρ=0.517, p-
value=0.0809). MicroChecker diagnosed null alleles in 13 subsamples for locus IS11,
which correspondingly displayed the highest F_{IS}=0.391, and in one subsample for IS19
(F_{IS}=-0.08). Null alleles probably explained these results, even if many missing data
probably did not correspond to null homozygotes. If we remove IS11 from the data, global
F_{IS} became significantly negative as expected for a small dioecious species (F_{IS}=-0.058, p-
value=0.0002). As for subdivision, F_{ST} remained almost unaffected, even with the ENA
method (F_{ST}=0.0964 in 95%CI=[0.0606..0.1338], p-value<0.0001). With H_S=0.66, the
standardized value was F_{ST}'=0.28.

364	Figure 5: F_{IS} values for each locus and averaged across those (All) of *Ixodes scapularis*
365	cured data from the eastern U.S.A. with 95% jackknife confidence intervals over
366	subsamples (for each locus) and bootstraps over loci (All). Results of tests for
367	panmixia for each locus are also indicated.

| 368 | |
| 369 | |

Conclusion

Combinations of amplification errors manifesting as null alleles, SAD or stuttering

lead not only to heterozygote deficits but also to an overall increase in LD. For this, the

different cures proposed are: in chromatograms, hunting for small and hard-to-detect

peaks in homozygous individuals to correct for SAD and pooling alleles with close sizes to

correct for stuttering. The obvious influence these problems can have would have led to

the unnecessary withdrawal of the flawed loci. The proposed amendments, together with

null allele's management with the ENA algorithm (Chapuis and Estoup, 2007), resulted in

analyses which revealed heterozygote excesses that would be expected in small dioecious

subpopulations, a more reasonable proportion of significant LD tests, and consequently

more accurate estimates of the degree of population subdivision.

In this case, the relatively important global LD across loci is probably due to small

effective sizes of the *I. scapularis* subpopulations, as confirmed by the important measure

of subdivision between subsamples ($F_{ST} \approx 0.3$).

The correlation between mitochondrial clade allocation and genetic structure of *I.*

scapularis is not within the scope of this work and will be treated in detail in a further study

that will also include immature stages.

Acknowledgements

This work was funded by NSF Grant # EF0914390 to L.B. and EEID EF-0914476 to
J.T. We thank members of the Lyme Gradient Consortium and many individuals who
provided ticks. We are also grateful to Heather Walker, Jenny Dickson, Keely Duff, Laquita
Burton, Alysha Benn and Nina Griffin, and the undergraduate students who provided field
and laboratory assistance.

References

Balloux, F., 2004. Heterozygote excess in small populations and the heterozygote-excess
effective population size. Evolution 58, 1891-1900.

Beati, L., Keirans, J.E., 2001. Analysis of the systematic relationships among ticks of the
genera *Rhipicephalus* and *Boophilus* (Acari : Ixodidae) based on mitochondrial 12S
ribosomal DNA gene sequences and morphological characters. J. Parasitol. 87, 32-
48.

Benjamini, Y., Yekutieli, D., 2001. The control of the false discovery rate in multiple testing
under dependency. Ann. Stat. 29, 1165–1188.

Castle, W.E., 1903. The laws of heredity of Galton and Mendel, and some laws governing
race improvement by selection. Proc Am Acad Arts Sci 39, 223-242.

Chapuis, M.P., Estoup, A., 2007. Microsatellite null alleles and estimation of population
differentiation. Mol. Biol. Evol. 24, 621-631.

Coombs, J.A., Letcher, B.H., Nislow, K.H., 2008. CREATE: a software to create input files
from diploid genotypic data for 52 genetic software programs. Mol. Ecol. Res. 8,
578–580.

De Meeûs, T., 2012. Initiation à la génétique des populations naturelles: Applications aux
parasites et à leurs vecteurs. IRD Editions, Marseille.

De Meeûs, T., 2018. Revisiting F_{IS}, F_{ST}, Wahlund effects and null alleles. J. Hered. 109,
446-456.

De Meeûs, T., Guégan, J.F., Teriokhin, A.T., 2009. MultiTest V.1.2, a program to
binomially combine independent tests and performance comparison with other
related methods on proportional data. BMC Bioinformatics 10, 443.

De Meeûs, T., Humair, P.F., Grunau, C., Delaye, C., Renaud, F., 2004. Non-Mendelian
transmission of alleles at microsatellite loci: an example in *Ixodes ricinus*, the vector
of Lyme disease. Int. J. Parasitol. 34, 943-950.

De Meeûs, T., McCoy, K.D., Prugnolle, F., Chevillon, C., Durand, P., Hurtrez-Boussès, S.,
Renaud, F., 2007. Population genetics and molecular epidemiology or how to
"débusquer la bête". Infect. Genet. Evol. 7, 308-332.

Delaye, C., Aeschlimann, A., Renaud, F., Rosenthal, B., De Meeûs, T., 1998. Isolation and
characterization of microsatellite markers in the *Ixodes ricinus* complex (Acari :
Ixodidae). Mol. Ecol. 7, 360-361.

Fagerberg, A.J., Fulton, R.E., Black, W.C., 2001. Microsatellite loci are not abundant in all
arthropod genomes: analyses in the hard tick, *Ixodes scapularis* and the yellow
fever mosquito, *Aedes aegypti.* Insect Mol. Biol. 10, 225-236.

Fox, J., 2005. The R commander: a basic statistics graphical user interface to R. J. Stat.
Software 14, 1–42.

Fox, J., 2007. Extending the R commander by "plug in" packages. R News 7, 46–52.

Goudet, J., 1995. FSTAT (Version 1.2): A computer program to calculate F-statistics. J.
Hered. 86, 485-486.

Goudet, J., 2003. Fstat (ver. 2.9.4), a program to estimate and test population genetics
parameters. Available at http://www.t-de-meeus.fr/Programs/Fstat294.zip, Updated
from Goudet (1995).

Goudet, J., Raymond, M., De Meeûs, T., Rousset, F., 1996. Testing differentiation in
diploid populations. Genetics 144, 1933-1940.

Gulia-Nuss, M., Nuss, A.B., Meyer, J.M., Sonenshine, D.E., Roe, R.M., Waterhouse, R.M.,
Sattelle, D.B., de la Fuente, J., Ribeiro, J.M., Megy, K., Thimmapuram, J., Miller,
J.R., Walenz, B.P., Koren, S., Hostetler, J.B., Thiagarajan, M., Joardar, V.S.,
Hannick, L.I., Bidwell, S., Hammond, M.P., Young, S., Zeng, Q.D., Abrudan, J.L.,
Almeida, F.C., Ayllon, N., Bhide, K., Bissinger, B.W., Bonzon-Kulichenko, E.,
Buckingham, S.D., Caffrey, D.R., Caimano, M.J., Croset, V., Driscoll, T., Gilbert, D.,
Gillespie, J.J., Giraldo-Calderon, G.I., Grabowski, J.M., Jiang, D., Khalil, S.M.S.,
Kim, D., Kocan, K.M., Koci, J., Kuhn, R.J., Kurtti, T.J., Lees, K., Lang, E.G.,
Kennedy, R.C., Kwon, H., Perera, R., Qi, Y.M., Radolf, J.D., Sakamoto, J.M.,
Sanchez-Gracia, A., Severo, M.S., Silverman, N., Simo, L., Tojo, M., Tornador, C.,
Van Zee, J.P., Vazquez, J., Vieira, F.G., Villar, M., Wespiser, A.R., Yang, Y.L., Zhu,
J.W., Arensburger, P., Pietrantonio, P.V., Barker, S.C., Shao, R.F., Zdobnov, E.M.,
Hauser, F., Grimmelikhuijzen, C.J.P., Park, Y., Rozas, J., Benton, R., Pedra, J.H.F.,
Nelson, D.R., Unger, M.F., Tubio, J.M.C., Tu, Z.J., Robertson, H.M., Shumway, M.,
Sutton, G., Wortman, J.R., Lawson, D., Wikel, S.K., Nene, V.M., Fraser, C.M.,

Collins, F.H., Birren, B., Nelson, K.E., Caler, E., Hill, C.A., 2016. Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease. Nat Commun 7.

Hedrick, P.W., 1999. Perspective: Highly variable loci and their interpretation in evolution and conservation. Evolution 53, 313-318.

Keirans, J.E., Needham, G.R., Oliver, J.H., 1999. The *Ixodes* (*Ixodes*) *ricinus* complex worldwide. Diagnosis of the species in the complex, hosts and distribution, in: Needham, G.R., R., M., Horn, D.J., Welbourn, W.C. (Eds.), Acarology IX: Symposia. Ohio Biological Survey, Columbus, Ohio, pp. 341-347.

Kempf, F., De Meeûs, T., Vaumourin, E., Noel, V., Taragel'ová, V., Plantard, O., Heylen, D.J.A., Eraud, C., Chevillon, C., McCoy, K.D., 2011. Host races in *Ixodes ricinus*, the European vector of Lyme borreliosis. Infect. Genet. Evol. 11, 2043-2048.

Manangwa, O., De Meeûs, T., Grébaut, P., Ségard, A., Byamungu, M., Ravel, S., 2019. Detecting Wahlund effects together with amplification problems: cryptic species, null alleles and short allele dominance in Glossina pallidipes populations from Tanzania. Mol. Ecol. Res. 19, 757–772.

Nei, M., Chesser, R.K., 1983. Estimation of fixation indices and gene diversities. Ann. Hum. Genet. 47, 253-259.

Norris, D.E., Klompen, J.S.H., Keiransand, J.E., Black, I.W.C., 1996. Population genetics of *Ixodes scapularis* (Acari: Ixodidae) based on mitochondrial 16S and 12S genes. J. Med. Entomol. 33, 78–89.

Qiu, W.G., Dykhuizen, D.E., Acosta, M.S., Luft, B.J., 2002. Geographic uniformity of the Lyme disease spirochete (Borrelia burgdorferi) and its shared history with tick vector (*Ixodes scapularis*) in the Northeastern United States. Genetics 160, 833-849.

R-Core-Team, 2018. R: A Language and Environment for Statistical Computing, Version 3.5.0 (2018-04-23) ed. R Foundation for Statistical Computing, Vienna, Austria, http://www.R-project.org.

Rulison, E.L., Kuczaj, I., Pang, G., Hickling, G.J., Tsao, J.I., Ginsberg, H.S., 2013. Flagging versus dragging as sampling methods for nymphal *Ixodes scapularis* (Acari: Ixodidae). J. Vector Ecol. 38, 163-167.

Sakamoto, J.M., Goddard, J., Rasgon, J.L., 2014. Population and demographic structure of *Ixodes scapularis* Say in the eastern United States. PLoS One 9, e101389.

Séré, M., Thévenon, S., Belem, A.M.G., De Meeûs, T., 2017. Comparison of different genetic distances to test isolation by distance between populations. Heredity 119, 55-63.

Van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M., Shipley, P., 2004. MICRO-
CHECKER: software for identifying and correcting genotyping errors in
microsatellite data. Mol Ecol Notes 4, 535-538.
Wattier, R., Engel, C.R., Saumitou-Laprade, P., Valero, M., 1998. Short allele dominance
as a source of heterozygote deficiency at microsatellite loci: experimental evidence
at the dinucleotide locus Gv1CT in *Gracilaria gracilis* (Rhodophyta). Mol. Ecol. 7,
1569-1573.
Weinberg, W., 1908. Über den Nachweis der Verebung beim Menschen. Jahresheft des
Vereins fur Vaterlundische Naturkunde in Wurttemberg 64, 368-382.
Weir, B.S., Cockerham, C.C., 1984. Estimating F-statistics for the analysis of population
structure. Evolution 38, 1358-1370.
Wright, S., 1965. The interpretation of population structure by F-statistics with special
regard to system of mating. Evolution 19, 395-420.
Yuval, B., Spielman, A., 1990. Duration and Regulation of the Developmental Cycle of
Ixodes dammini (Acari: Ixodidae). J. Med. Entomol. 27, 196-201.
505
506
507