1	**Title:**	The fungal sesquiterpenoid pyrenophoric acid B uses the
2		plant ABA biosynthetic pathway to inhibit seed
3		germination.
4		
5	**Running title:**	Cross-kingdom activity of the fungal metabolite pyrenophoric acid
6		B on seed germination.
7		

8	**Authors:**	Jorge Lozano-Juste[a,*], Marco Masi[b], Alessio Cimmino[b], Suzette
9		Clement[c], Maria A. Fernández[a], Regina Antoni[a], Susan Meyer[c],
10		Pedro L. Rodriguez[a] and Antonio Evidente[b,*]
11		

12	**Affiliation:**	[a]*Instituto de Biologia Molecular y Celular de Plantas (IBMCP),*
13		*Consejo Superior de Investigaciones Científicas (CSIC),*
14		*Universidad Politécnica de Valencia (UPV), C/Ingeniero Fausto*
15		*Elio s/n Edificio 8E, 46022 Valencia, Spain*
16		[b]*Dipartimento di Scienze Chimiche, Università di Napoli Federico*
17		*II, Complesso Universitario Monte Sant'Angelo, Via Cintia 4,*
18		*80126 Napoli, Italy*
19		[c] *Shrub Sciences Laboratory, U.S. Forest Service Rocky Mountain*
20		*Research Station, 735 North 500 East, Provo, Utah 84606, United*
21		*States*
22		
23		

E-mail addresses: Marco Masi marco.masi@unina.it

ORCID: 0000-0003-0609-8902

Alessio Cimmino alessio.cimmino@unina.it

ORCID: 0000-0002-1551-4237

Suzette Clement sclement@fs.fed.us

Maria A. Fernández maferlo9@upvnet.upv.es

Regina Antoni regiantoni@gmail.com

Susan Meyer semeyer@xmission.com

Pedro L. Rodriguez prodriguez@ibmcp.upv.es

Antonio Evidente evidente@unina.it Phone: +39- 081-2539178

Corresponding author. ORCID: 0000-0001-9110-1656

Jorge Lozano-Juste lojujo@ibmcp.upv.es Phone: +34-698255910

Corresponding author. ORCID: 0000-0001-7034-566X

* Corresponding authors: lojujo@ibmcp.upv.es and evidente@unina.it

Date of submission: 22 April 2019

Number of figures and tables: 4 Figures + 1 supplementary figure (All the Figures in

color in print)

Word count: 3565 words

46	**Highlight:**	The fungus *Pyrenophora semeniperda* produces pyrenophoric acid
47		B, a small molecule that exploits the plant ABA biosynthetic
48		pathway to reduce seed germination, increasing its reproductive
49		success.

Abstract

Pyrenophoric acid (P-Acid), P-Acid B and P-Acid C are three phytotoxic

sesquiterpenoids produced by the ascomycete seed pathogen *Pyrenophora semeniperda,*

a fungus proposed as a mycoherbicide for biocontrol of cheatgrass, an extremely invasive

weed. When tested in cheatgrass bioassays these metabolites were able to delay seed

germination, with P-Acid B being the most active compound. Here, we have investigated

the cross-kingdom activity of P-Acid B and its mode of action and found that it activates

the ABA signaling pathway in order to inhibit seedling establishment. P-Acid B inhibits

seedling establishment in wild-type *Arabidopsis thaliana* while several mutants affected

in the early perception as well as in downstream ABA signaling components were

insensitive to the fungal compound. However, in spite of structural similarities between

ABA and P-Acid B, the latter is not able to activate the PYR/PYL family of ABA

receptors. Instead, we have found that P-Acid B uses the ABA biosynthesis pathway at

the level of alcohol dehydrogenase ABA2 to reduce seedling establishment. We propose

that the fungus *Pyrenophora semeniperda* manipulates plant ABA biosynthesis as a

strategy to reduce seed germination, increasing its ability to cause seed mortality and

thereby increase its fitness through higher reproductive success.

Keywords

ABA2, ABA biosynthesis, Abscisic acid, Cross-kingdom activity, Pyrenophoric acid,

Pyrenophoric acid B, Pyrenophoric acid C, *Pyrenophora semeniperda*, PYR/PYL, Seed

germination.

Introduction

The fungus *Pyrenophora semeniperda* (Brittlebank and Adams) Shoemaker, a naturally

occurring necrotrophic seed pathogen, has been proposed as a potential biocontrol agent

against cheatgrass (*Bromus tectorum*) (Meyer *et al.*, 2013). This invasive weed is

dramatically altering the semi-arid shrubland ecosystems in the western U.S.A.

increasing wildfire frequency and intensity (Germino *et al.*, 2016). Thus, the ability of *P.*

semeniperda to produce toxins *in vitro*, which could potentially be useful in biological

control, was investigated (Meyer *et al.*, 2015; Cimmino *et al.*, 2015). *P. semeniperda*

produces large amounts of cytochalasin B in solid cultures (Masi *et al.*, 2014*a*) while

spirocyclic γ-lactams, including spirostaphylotrichins A, C, D, R, V, W and triticone E,

are produced when the fungus is grown in potato dextrose broth (Masi *et al.*, 2014*b*) .

Furthermore, from the same fungal organic extract smaller quantities of cytochalasins A,

F and deoxaphomin were isolated together with a new phytotoxic sesquiterpenoid, named

pyrenophoric acid (P-Acid) (Fig. 1A) (Masi *et al.*, 2014*a,c*). Finally, these compounds

were produced in *B. tectorum* seed culture together with other two phytotoxic

sesquiterpenoids named pyrenophoric acids B and C (P-Acid B and P-Acid C,

respectively) (Fig. 1A) (Masi *et al.*, 2014*d*). P-Acid, P-Acid B and P-Acid C were able to

inhibit both root and coleoptile growth of *B. tectorum* (Masi *et al.*, 2014*c,d*). In particular,

P-Acid B was the most active sesquiterpenoid among the P-Acids studied (Masi *et al.*,

2014*d*). It was able to reduce germination at 7 days after sowing and produced

germination delay relative to the control treatment. P-Acid B was also more bioactive in

reducing coleoptile and root growth than the other sesquiterpenoids, P-Acid and P-Acid

C. Interestingly, the production of small molecules able to delay germination could

benefit *P. semeniperda*, as its fitness is increased on slow-germinating or dormant seeds.

This is because slow-germinating seeds cannot escape pathogen-caused mortality, thus

increasing resources available for pathogen growth and reproduction (Meyer *et al.*, 2015;

Cimmino *et al.*, 2015).

P-Acid, P-Acid B and P-Acid C (Fig. 1A) are sesquiterpenoids closely related to

abscisic acid (ABA) (Fig. 1A). ABA is a plant hormone involved in many physiological

processes in plants including seed development and germination, root growth and

abscission among others (Leung and Giraudat, 1998). It has also a central role in the

adaptation to environmental stresses like drought, high salinity and low temperatures

(Yamaguchi-Shinozaki and Shinozaki, 2006). However, ABA is not a plant-specific

metabolite. It is produced by different organisms including fungi, suggesting that it is a

ubiquitous and versatile small compound that can modulate a myriad of physiological

functions (Hartung, 2010; Takezawa *et al.*, 2011). In *Arabidopsis thaliana* (Arabidopsis),

ABA biosynthesis begins with the conversion of zeaxanthin to all-*trans*-violaxanthin by a

zeaxanthin epoxidase known as ABA1 (Marin et al., 1996, Audran et al., 2001). Next, in

the rate-limiting step of ABA biosynthesis, 9-*cis*-epoxycarotenoid dioxygenase (NCED)

enzymes break down C_{40} carotenoids to produce xanthoxin (Schwartz *et al.*, 1997; Iuchi

et al., 2000). Xanthoxin is further transformed into abscisic aldehyde by the Arabidopsis

short-chain alcohol dehydrogenase ABA2 (González-Guzmán *et al.*, 2002). Finally,

abscisic aldehyde is used as substrate by the aldehyde oxidase AAO3, in a reaction that

depends on the activity of the molybdenum cofactor sulfurase ABA3, to produce ABA (

Schwartz *et al.*, 1997; Seo *et al.*, 2000; González-Guzmán *et al.*, 2004). ABA is

perceived by the PYR/PYL/RCAR family of ABA receptors, comprised of 14 members

in the eudicot plants Arabidopsis and tomato and 12 and 9 members in the monocot

species rice and *Brachypodium distachyon,* respectively (Ma *et al.*, 2009; Park *et al.*,

2009; He *et al.*, 2014; Pri-Tal *et al.*, 2017). ABA-bound PYR/PYL/RCAR proteins

interact with Protein Phosphatases type 2C (PP2C) and inhibit their activity allowing the

downstream phosphorylation cascades to activate the ABA response, including inhibition

of seed germination and stomatal closure, among others.

Here we have studied the effect and the mode of action of the fungus-produced

pyrenophoric acids on plant seed germination. Among P-Acids, P-Acid B was able to

inhibit germination of Arabidopsis seeds. We have found that P-Acid B makes use of the

ABA biosynthesis pathway at the level of ABA2 to activate the ABA signaling and

inhibit germination, increasing the probability of host seed mortality and thereby

increasing pathogen fitness.

Materials and Methods

Chemicals

Pyrenophoric acid and pyrenophoric acids B and C (Fig. 1A) were isolated from the solid

B. tectorum (cheatgrass) seed culture of *P. semeniperda* (strain WRK10-22), as

previously reported (Masi *et al.*, 2014*c*,*d*)

Plant material and growth conditions

Arabidopsis thaliana seeds were surface sterilized with 30% bleach containing 0.02%

Tween-20 for 10 minutes followed by rinsing with sterile water 5 times. Seeds were sown

on solid or liquid Murashige and Skoog (MS) media and stratified 3 days at 4°C. For

seedling establishment experiments, seeds were sown on 24-well plates filled with 1 mL

of MS media + 1% agar supplemented with the different chemicals dissolved in DMSO.

Seeds were kept in a Sanyo incubator under long day conditions at temperatures 23°C

/21°C (day/night) for 3 days. Pictures were taken with a Leica DMS1000 macroscope and

seedling establishment (seedlings with open and green cotyledons) scored. For luciferase

assays, seeds were incubated for 7 days under long day conditions in liquid MS media.

The different mutants used in this work, *112458* (Gonzalez-Guzman *et al.*, 2012), *abi3-9*

(Nambara *et al.*, 2002), *abi4-11* (Nambara *et al.*, 2002), *aba1-101* (Barrero *et al.*, 2005),

aba2-1 (Léon-Kloosterziel *et al.*, 1996), *nced3,5* (Frey *et al.*, 2012) *aba2-11* (González-

Guzmán *et al.*, 2002), *aao3-2* (González-Guzmán *et al.*, 2004), *aba3-1* (Léon-

Kloosterziel *et al.*, 1996), and the transgenic lines *pMAPKKK18-LUC$^+$* (Vaidya *et al.*,

2017) and *35S-HAB1^{W385A}* (Dupeux *et al.*, 2011) have been described previously. To

generate the double *nced2nced3* mutant, homozygous plants from the T-DNA lines

SALK_090937C (*nced2*) and GK-129B08.01 (*nced3*) were crossed and the F2 selected

on 100 mM NaCl plates. PCR-verified plants using the primers FORNced2:

ATGGTTTCTCTTCTTACAATGCCG,   RVNced2:

TTCCGGTTAACCATACCAATCTC, NewpROK2: GCCGATTTCGGAACCACCATC,

FWNced3GABI: CCTAGTGTTCAGATCGCCGGA,   RVNced3GABI:

GAGATTCTCGTCAGACTCGTT,   GK-o8474:

ATAATAACGCTGCGGACATCTACATTT were selfed and the F3 seed was used for

the different experiments.

Luciferase imaging

pMAPKKK18-LUC$^+$ seedlings were grown in 24-well plates filled with 1 mL liquid MS

media for 7 days. MS media was changed with MS media supplemented with 100 µM D-

luciferin, potassium salt (GoldBio) and the different treatments; 25 µM ABA, 200 µM P-

Acid B or DMSO as control, and incubated for 6 hours. Luminescence was recorded with

a LAS-3000 imager (Fujifilm) equipped with a CCD camera using 2 minutes exposures.

8-bit images were transformed in rainbow false color and quantified using Fiji

(Schindelin *et al.*, 2012).

Protein purification

The different expression clones were described previously (Antoni *et al.*, 2012; Castillo

et al., 2015). Protein induction was performed by adding 1mM IPTG to 250mL cultures

of BL21(DE3) cells at OD=0.6-0.8. Cells were grown overnight at 16 ^{o}C and collected by

centrifugation. Cells were lysed in 50 mM tris-HCl (pH 7.6), 250 mM KCl, 10mM 2-ME,

10mM Imidazole, 0.1% Tween 20, 10% glycerol and freeze/thawed at -80 $^{\circ}$C twice. After

sonication, the protein extract was cleared by centrifugation and applied into 1mL Ni-

NTA agarose. Ni-NTA beads were washed 3 times with 10 column volumes of 50 mM

tris-HCl (pH 7.6), 250 mM KCl, 10mM 2-ME, 20% glycerol, 0.1% Tween 20 buffer

supplemented with 30mM imidazole. Proteins were eluted from the Ni-NTA beads with

the same buffer containing 250mM imidazole.

PP2C enzymatic assays

For PP2C assays 1µM dNHAB1 and 2 µM PYR/PYL were used (Santiago *et al.*, 2009).

Proteins were incubated with the different chemicals (DMSO, ABA or P-Acid B) for 10

minutes in a volume of 50 µL containing 10mM $MnCl_2$. Reactions were initiated by the

addition of 50 µL 50mM Tris pH=8.0, 50mM *p*-nitrophenyl phosphate (pNPP).

Dephosphorylation of pNPP was followed by monitoring the absorbance at 405 nm

during 20 minutes. The activity of dNHAB1 in the absence of any receptor was set to

100%.

Results

Effect of pyrenophoric acids on seed germination.

P-Acid, P-Acid B and C (Fig. 1A) were purified from an organic extract of *P.*

semeniperda (strain WRK10-22) grown in solid *B. tectorum* (cheatgrass) seed culture

following the procedure described previously (Masi *et al.*, 2014*c,d*) and identified by

comparing the physical and spectroscopic properties with those previously reported (Masi

et al., 2014*c,d*). To assess the effects of P-Acids *in planta*, Arabidopsis WT Col-0 seeds

were germinated in the presence of 100 µM of each P-Acid, 1 µM ABA or DMSO as a

control. P-Acid and P-Acid C did not have any effect on seed germination while P-Acid

B severely delayed germination and markedly inhibited seedling establishment (Fig. 1B).

At day 3 after sowing 100% of Col-0 seedlings established under control (DMSO), P-

Acid or P-Acid C conditions. However, the P-Acid B treatment reduced seedling

establishment to 10%, similar to a treatment with 1 µM ABA (Fig. 1C). Therefore, it

seems that among P-Acids, pyrenophoric acid B, produced by the fungus *P. semeniperda*,

has cross-kingdom activity, as it is able to inhibit plant seed germination.

Pyrenophoric acid B activates the ABA signalling pathway.

In order to gain further insight into the mode of action of P-Acid B we used several

mutants affected in the early and late steps of ABA perception and signalling. We used

the *112458* sextuple mutant impaired in PYR1, PYL1, PYL2, PYL4, PYL5 and PYL8

ABA receptors (Gonzalez-Guzman *et al.*, 2012); 35S:HAB1^{W385A}, a transgenic line

expressing an engineered mutant form of the PP2C HAB1 unable to interact with ABA

receptors (Dupeux *et al.*, 2011); and *abi3* and *abi4* that carry mutations in 2 transcription

factors central to the activation of ABA signaling (Nambara *et al.*, 2002). All mutants

tested were insensitive to P-Acid B application (Fig. 2A). While only 10% of Col-0

seedlings established in the presence of 100 µM P-Acid B, all the mutants reached nearly

100% seedling establishment 3 days after the treatment (Fig. 2B). These results indicate

that the pathogen-produced P-Acid B uses the plant ABA signaling pathway to inhibit

seed germination. To elucidate if P-Acid B could activate ABA-dependent gene

expression we used an ABA reporter line that carries the ABA inducible promoter

pMAPKKK18 (Okamoto *et al.*, 2013) fused to the firefly luciferase enzyme. This reporter

line is highly activated by ABA (Fig. 2C,D) (Vaidya *et al.*, 2017). P-Acid B was able to

activate the reporter line since *pMAPKKK18-LUC$^+$* seedlings treated with 200 µM P-

Acid B during 6 hours showed around 4.5 fold increase on LUC activity due to the

activation of the ABA responsive promoter *pMAPKKK18* (Fig. 2D). In conclusion, P-

Acid B is able to activate ABA signalling, including ABA-dependent gene expression, to

inhibit seed germination.

Pyrenophoric acid B does not activate ABA receptors

P-Acid B inhibits seed germination and the sextuple mutant *112458* is insensitive to the

application of P-Acid B (Fig. 2A,B). In addition, pyrenophoric acids are sesquiterpenoids

with structural similarities to ABA (Fig. 1A). We reasoned that P-Acid B could activate

the ABA signaling pathway by directly binding to ABA receptors. To investigate this

possibility we used PP2C phosphatase assays with different ABA receptors and the PP2C

phosphatase HAB1. The PYR/PYL/RCAR family of ABA receptors binds ABA within a

hydrophobic cavity that defines the ABA binding pocket (Santiago *et al.*, 2009; Melcher

et al., 2009). After ABA binding, allosteric changes induced in ABA receptors facilitates

their interaction with PP2Cs inhibiting phosphatase activity. Therefore, the binding of

small molecules to ABA receptors can be followed by *in vitro* PP2C phosphatase assays

as reported elsewhere (Park *et al.*, 2009; Okamoto *et al.*, 2013; Vaidya *et al.*, 2017). We

performed *in vitro* PP2C phosphatase assays using recombinant PYR1, PYL1, PYL2,

PYL4, PYL5, PYL6, PYL8 and PYL10. As expected, ABA bound to all receptors tested

producing the concomitant inhibition of HAB1 phosphatase activity by 80-90%.

However, P-Acid B was not able to inhibit HAB1 in the presence of any of the receptors

tested (Fig. 3), strongly suggesting that despite structural and functional similarities with

ABA, P-Acid B does not activate ABA receptors.

P-Acid B exploits the plant ABA biosynthetic pathway to inhibit seed germination

P-Acid B inhibits seed germination and ABA-insensitive mutants are also insensitive to

P-Acid B, suggesting that P-Acid B needs an intact ABA signaling pathway to inhibit

seed germination. To better understand P-Acid B activity we analyzed the effect of P-

Acid B treatment on the germination of mutants affected in different steps of the ABA

biosynthetic pathway. We used *aba1-101*, compromised in the early steps of ABA

biosynthesis (Barrero *et al.*, 2005), the double mutants *nced2nced3 (nced2,3)* and

nced3,5, impaired in the production of xanthoxin; *aba2-1* and *aba2-11*, loss of function

alleles of *ABA2* that cannot convert xanthoxin into ABA-aldehyde (González-Guzmán *et*

al., 2002; Cheng *et al.*, 2002) and lastly, *aao3-2* and *aba3-1*, blocked in the last step of

ABA biosynthesis where ABA-aldehyde is oxidized to ABA (González-Guzmán *et al.*,

2004; Schwartz *et al.*, 1997*b*) . As expected, while all the genotypes germinated and

established 3 days after sowing in control (DMSO) conditions, 1 µM ABA treatment

inhibited germination and seedling establishment in the wild-type Col-0 and also in all

the ABA deficient mutants tested (Fig. 4A,B; Fig. S1). As in previous experiments, P-

Acid B delayed germination and severely affected seedling establishment of Col-0 seeds.

Interestingly, *aba2-1*, *aba2-11*, *aao3-2 and aba3-1* mutants were resistant to P-Acid B

treatment indicating that ABA biosynthesis from ABA2 onwards is required for P-Acid B

bioactivity on seed germination. However, *nced2,3* which is impaired in the upstream

step of ABA biosynthesis that precedes ABA2 activity, was sensitive to P-Acid B, with a

clear reduction in seedling establishment in response to this treatment (Fig. 4A,B). This

result was confirmed using a different *NCED* double mutant; *nced3,5*, and also the

mutant *aba1-101*, impaired in the first step of ABA biosynthesis (Fig. S1). Thus, while P-

Acid B reduced seedling establishment of Col-0, *aba1-101*, *nced2,3* and *nced3,5* seeds to

below 10%, the compound was unable to produce any effect on the other ABA

biosynthetic mutants *aba2-1*, *aba2-11*, *aao3-2* and *aba3-1* (Fig. 4A,B; Fig. S1).

Therefore, P-Acid B can complement the ABA biosynthetic defect of *aba1-101*, *nced2,3*

and *nced3,5* but requires downstream enzymatic steps. These results suggest that P-Acid

B requires ABA2 activity to inhibit seed germination and establishment. Actually, since

P-Acid B is structurally related to xanthoxin and ABA2 is able to convert xanthoxin-

related compounds into abscisic aldehyde (González-Guzmán *et al.*, 2002; Nambara and

Marion-Poll, 2005), we hypothesized that P-Acid B could be used by ABA2 as a

substrate to produce ABA or an ABA-mimic, thus explaining the activation of the ABA

signaling pathway by P-Acid B and its negative effect on seed germination.

Discussion

The fungus *P. semeniperda* is a plant pathogen specialized in infecting seed tissues. Due

to its ability to infect cheatgrass, one of the most invasive plant species in the USA, it has

been lately considered as a biocontrol agent for invasive grass species. For this purpose,

P. semeniperda produces a series of toxins that kill the host cells, including the

mycotoxin cytochalasin B, which impedes the correct polymerization of actin filaments

leading to cell death (Scherlach *et al.*, 2010). However, cytochalasin B is not the only low

molecular weight compound produced by *P. semeniperda* that has a role in infection.

Recently, a series of metabolites was isolated and identified from *P. semeniperda* cultures

and among them, P-Acid, P-Acid B and P-Acid C were found (Masi *et al.*, 2014*c,d*). P-

Acids are bioactive compounds able to inhibit seed germination and coleoptile growth,

with P-Acid B the most active of these compounds (Masi *et al.*, 2014*d*). In this work we

have investigated the molecular mechanism underlying P-Acids mode of action. We have

found that P-Acid B can inhibit *Arabidopsis thaliana* seed germination through the ABA

signaling pathway. P-Acid B needs an intact ABA signaling pathway to inhibit seed

germination since mutants affected in the first steps of ABA perception as well as

mutants impaired in central transcription factors essential for the activation of ABA

responses were insensitive to P-Acid B application.

P-Acids are ABA-related compounds with structural similarities to the plant hormone. It

has been previously reported that ABA receptors can accommodate ABA-like molecules

within its hydrophobic pocket (Kepka *et al.*, 2011; Benson *et al.*, 2015). For example, the

ABA degradation by-product, phaseic acid (PA), is able to bind to certain ABA receptors

(Weng *et al.*, 2016). PA has submicromolar activity towards the ABA receptors PYL3,

PYL5 and PYL6 and a role for PA on water stress tolerance has been proposed (Weng *et*

al., 2016; Lozano-Juste and Cutler, 2016; Rodriguez, 2016). Other hormone receptors can

also accommodate hormone analogs or different forms of the hormone. GID1, the

gibberellic-acid receptor binds different gibberellins (Murase *et al.*, 2008; Shimada *et al.*,

2008) and the jasmonic acid (JA) receptor COI1 is activated not only by its plant derived

ligand, JA-Ile, but also by the JA analog coronatin, produced by biotrophic bacteria

(Sheard *et al.*, 2010). Therefore we reasoned that the ABA-related sequiterpenoid P-Acid

B could directly bind to ABA receptors. However, we could not find activation to any of

the ABA receptors studied indicating that P-Acid B, despite its structural similarities to

ABA, does not activate ABA receptors directly, suggesting a different mechanism of

action. An alternative hypothesis for P-Acid B bioactivity is acting as precursor of ABA,

feeding the biosynthetic pathway and promoting ABA accumulation. This hypothesis

proved to be correct since *aba3-1*, *aao3-2*, *aba2-1* and *aba2-11* mutants affected in ABA

biosynthesis could establish in the presence of P-Acid B while Col-0 plants could not.

Therefore, P-Acid B may require *de novo* synthesis of ABA to inhibit seedling

establishment. In order to pinpoint the entrance of P-Acid B into the ABA biosynthetic

pathway, we have compared sensitivity to P-Acid B of mutants impaired in different

steps of ABA biosynthesis. Thus, when we analyzed the effect of P-Acid B on the *aba1-*

101, *nced2,3* and *nced3,5* mutants, we found that these strains are not resistant to P-Acid

B, while *aba2*, *aao3* and *aba3* single mutants could germinate and establish in presence

of P-Acid B. These data suggest that P-Acid B enters the ABA biosynthetic pathway

downstream of NCED and requires ABA2, AAO3 and ABA3 for bioactivity. AAO3

encodes an aldehyde oxidase and might restore the carboxylic group of P-Acid B in case

it was reduced to aldehyde by the reductive ability of plant cells for exogenous

molecules. Indeed, the reduction of non-activated carboxylic groups in different

substrates has been described for at least nine species of cultured plant cells (Villa and

Molinari, 2008). ABA2 is a member of the short-chain alcohol dehydrogenase protein

family crucial for ABA biosynthesis (González-Guzmán *et al.*, 2002; Cheng *et al.*, 2002).

It catalyzes the conversion of xanthoxin into ABA-aldehyde that is used by the AAO3

protein to produce ABA. Accordingly, the levels of ABA in *aba2* mutants are severely

reduced (Léon-Kloosterziel *et al.*, 1996; González-Guzmán *et al.*, 2002), indicating that

this is a key step in the basal production of ABA in plants. The conversion of xanthoxin

into ABA-aldehyde by ABA2 involves not one but several catalytic steps. ABA2 is able

to oxidize the 4'-hydroxyl group of xanthoxin into a 4'-keto group but it also isomerizes

the 1',2',-epoxide into a 1'-hydroxy-2'ene group (González-Guzmán *et al.*, 2002). Due to

structural similarities between xanthoxin and P-Acid B we hypothesized that ABA2

could oxidize the 4'hydroxyl group and dehydrogenate C-2' and C-3' of P-Acid B

generating ABA. Actually, the enzyme ABA2 can use certain alcohol substrates different

from xanthoxin being able to oxidize 3,5,5′-trimethylcyclohexanol alcohol although with

higher K_m values (González-Guzmán *et al.*, 2002). Taken together we propose P-Acid B

as a fungal compound with cross-kingdom activity that might be able to use the plant

enzyme ABA2 to induce the synthesis of ABA and inhibit seed germination. A body of

evidence indicates that the fungus *P. semeniperda* can benefit from reducing seed

germination and seedling establishment through P-Acid B production. *P. semeniperda*

reproductive success is often associated with slow germinating or dormant seeds (Finch-

Boekweg *et al.*, 2015) while fast germinating seeds usually tolerate *P. semeniperda*

infection and develop into normal plants (Finch-Boekweg *et al.*, 2015). Additionally, the

distribution of *P. semeniperda* populations is correlated with environmental conditions

that reduce seed germination rates and therefore, *P. semeniperda* is found more

frequently in dry scenarios rather than humid ones. Dry conditions reduce seed

germination and induce postgermination growth arrest (Beckstead *et al.*, 2007, 2016;

Meyer *et al.*, 2015).

In summary, we propose that the fungus *P. semeniperda* produces pyrenophoric acid B to

inhibit seed germination through the activation of plant ABA biosynthesis, in order to

increase its fitness by increasing the probability of pathogen-caused seed mortality,

thereby increasing resources available for pathogen reproduction.

Supplementary data

Fig.S1. Pyrenophoric acid B (P-Acid B) requires ABA2 and onwards to inhibit seedling

establishment.

Acknowledgements

This research was funded in part by Grant JFSP-11-S-2-6 to S.M. from the Joint Fire

Sciences Program of the U.S. Departments of Agriculture and Interior and in part, to

M.M., by Programme STAR 2017 financially supported by UniNA and Compagnia di

San Paolo. J. L.-J. is funded by a Marie-Sklodowska Curie Reintegration Grant H2020-

MSCA-707477. Work in P. L. R. lab is supported by RTC-2017-6019-2 and BIO2017-

82503-R grants from Ministerio de Ciencia, Innovación y Universidades. M.A.F. is a

recipient of a FPU fellowship from MECD. A. Evidente is associated to the Istituto di

Chimica Biomolecolare del CNR, Pozzuoli, Italy. We would like to thank Ebe Merilo

(University of Tartu) for sharing *nced3,5* and *aba3-1* mutant seeds.

REFERENCES

Audran C, Borel C, Frey A, Sotta B, Meyer C, Simonneau T, Marion-Poll A. 1998. Expression studies of the zeaxanthin epoxidase gene in *Nicotiana plumbaginifolia*. Plant Physiology **118**, 1021-1028.

Antoni R, Gonzalez-Guzman M, Rodriguez L, Rodrigues A, Pizzio GA, Rodriguez PL. 2012. Selective inhibition of clade A phosphatases type 2C by PYR/PYL/RCAR abscisic acid receptors. Plant Physiology **158**, 970–980.

Barrero JM, Piqueras P, González-Guzmán M, Serrano R, Rodríguez PL, Ponce MR, Micol JL. 2005. A mutational analysis of the ABA1 gene of *Arabidopsis thaliana* highlights the involvement of ABA in vegetative development. Journal of Experimental Botany **56**, 2071–2083.

Beckstead J, Meyer SE, Ishizuka TS, McEvoy KM, Coleman CE. 2016. Lack of Host Specialization on Winter Annual Grasses in the Fungal Seed Bank Pathogen *Pyrenophora semeniperda*. PLOS ONE **11**, e0151058.

Beckstead J, Meyer SE, Molder CJ, Smith C. 2007. Germination Rate Determines Pathogen-caused Seed Mortality. Annals of Botany **99**, 907–914.

Benson CL, Kepka M, Wunschel C, Rajagopalan N, Nelson KM, Christmann A, Abrams SR, Grill E, Loewen MC. 2015. Abscisic acid analogs as chemical probes for dissection of abscisic acid responses in *Arabidopsis thaliana*. Phytochemistry **113**, 96–107.

Castillo M-C, Lozano-Juste J, González-Guzmán M, Rodriguez L, Rodriguez PL, León J. 2015. Inactivation of PYR/PYL/RCAR ABA receptors by tyrosine nitration may enable rapid inhibition of ABA signaling by nitric oxide in plants. Science Signaling **8**, ra89.

Cheng W-H, Endo A, Zhou L, *et al.* 2002. A Unique Short-Chain Dehydrogenase/Reductase in Arabidopsis Glucose Signaling and Abscisic Acid Biosynthesis and Functions. The Plant Cell **14**, 2723–2743.

Cimmino A, Masi M, Evidente M, Superchi S, Evidente A. 2015. Fungal phytotoxins with potential herbicidal activity: chemical and biological characterization. Natural Product Reports **32**, 1629–1653.

Dupeux F, Antoni R, Betz K, *et al.* 2011. Modulation of Abscisic Acid Signaling in Vivo by an Engineered Receptor-Insensitive Protein Phosphatase Type 2C Allele. Plant Physiology **156**, 106–116.

Finch-Boekweg H, Gardner JS, Allen PS, Geary B. 2015. Postdispersal Infection and Disease Development of *Pyrenophora semeniperda* in *Bromus tectorum* Seeds. Phytopathology **106**, 236–243.

Germino MJ, Chambers JC, Brown CS (Eds.). 2016. *Exotic Brome-Grasses in Arid and Semiarid Ecosystems of the Western US: Causes, Consequences, and Management Implications.* Springer International Publishing.

González-Guzmán M, Abia D, Salinas J, Serrano R, Rodríguez PL. 2004*a*. Two New Alleles of the abscisic aldehyde oxidase 3 Gene Reveal Its Role in Abscisic Acid Biosynthesis in Seeds. Plant Physiology **135**, 325–333.

González-Guzmán M, Abia D, Salinas J, Serrano R, Rodríguez PL. 2004*b*. Two New Alleles of the abscisic aldehyde oxidase 3 Gene Reveal Its Role in Abscisic Acid Biosynthesis in Seeds. Plant Physiology **135**, 325–333.

González-Guzmán M, Apostolova N, Bellés JM, Barrero JM, Piqueras P, Ponce MR, Micol JL, Serrano R, Rodríguez PL. 2002. The Short-Chain Alcohol Dehydrogenase ABA2 Catalyzes the Conversion of Xanthoxin to Abscisic Aldehyde. The Plant Cell **14**, 1833–1846.

Gonzalez-Guzman M, Pizzio GA, Antoni R, *et al.* 2012. Arabidopsis PYR/PYL/RCAR Receptors Play a Major Role in Quantitative Regulation of Stomatal Aperture and Transcriptional Response to Abscisic Acid. The Plant Cell **24**, 2483–2496.

Hartung W. 2010. The evolution of abscisic acid (ABA) and ABA function in lower plants, fungi and lichen. Functional Plant Biology **37**, 806–812.

He Y, Hao Q, Li W, Yan C, Yan N, Yin P. 2014. Identification and characterization of ABA receptors in *Oryza sativa*. PloS One **9**, e95246.

Iuchi S, Kobayashi M, Yamaguchi-Shinozaki K, Shinozaki K. 2000. A Stress-Inducible Gene for 9-cis-Epoxycarotenoid Dioxygenase Involved in Abscisic Acid Biosynthesis under Water Stress in Drought-Tolerant Cowpea. Plant Physiology **123**, 553–562.

Kepka M, Benson CL, Gonugunta VK, Nelson KM, Christmann A, Grill E, Abrams SR. 2011. Action of natural abscisic acid precursors and catabolites on abscisic acid receptor complexes. Plant Physiology **157**, 2108–2119.

Léon-Kloosterziel KM, Gil MA, Ruijs GJ, Jacobsen SE, Olszewski NE, Schwartz SH, Zeevaart JA, Koornneef M. 1996. Isolation and characterization of abscisic acid-deficient Arabidopsis mutants at two new loci. The Plant Journal: For Cell and Molecular Biology **10**, 655–661.

Leung J, Giraudat and J. 1998. Abscisic Acid Signal Transduction. Annual Review of Plant Physiology and Plant Molecular Biology **49**, 199–222.

Lozano-Juste J, Cutler SR. 2016. Hormone signalling: ABA has a breakdown. Nature Plants **2**, 16137.

Ma Y, Szostkiewicz I, Korte A, Moes D, Yang Y, Christmann A, Grill E. 2009. Regulators of PP2C Phosphatase Activity Function as Abscisic Acid Sensors. Science **324**, 1064–1068.

Marin E, Nussaume L, Quesada A, Gonneau M, Sotta B, Hugueney P, Frey A, Marion-Poll A. 1996. Molecular identification of zeaxanthin epoxidase of Nicotiana plumbaginifolia, a gene involved in abscisic acid biosynthesis and corresponding to the ABA locus of Arabidopsis thaliana. EMBO J, **15**, 2331-2342.

Masi M, Evidente A, Meyer S, Nicholson J, Muñoz A. 2014*a*. Effect of strain and cultural conditions on the production of cytochalasin B by the potential mycoherbicide *Pyrenophora semeniperda* (Pleosporaceae, Pleosporales). Biocontrol science and technology **24**, 53–64.

Masi M, Meyer S, Clement S, Andolfi A, Cimmino A, Evidente A. 2014b. Spirostaphylotrichin W, a spirocyclic γ-lactam isolated from liquid culture of *Pyrenophora semeniperda*, a potential mycoherbicide for cheatgrass (*Bromus tectorum*) biocontrol. Tetrahedron, *70*, 1497-1501..

Masi M, Meyer S, Cimmino A, Andolfi A, Evidente A. 2014*c*. Pyrenophoric acid, a phytotoxic sesquiterpenoid penta-2,4-dienoic acid produced by a potential mycoherbicide, *Pyrenophora semeniperda*. Journal of Natural Products **77**, 925–930.

Masi M, Meyer S, Cimmino A, Clement S, Black B, Evidente A. 2014*d*. Pyrenophoric Acids B and C, Two New Phytotoxic Sesquiterpenoids Produced by *Pyrenophora semeniperda*. Journal of Agricultural and Food Chemistry **62**, 10304–10311.

Melcher K, Ng L-M, Zhou XE, *et al.* 2009. A gate-latch-lock mechanism for hormone signalling by abscisic acid receptors. Nature **462**, 602–608.

Meyer SE, Clement S, Beckstead J. 2013. Annual brome control using a native fungal seed pathogen.

Meyer SE, Masi M, Clement S, Davis TL, Beckstead J. 2015. Mycelial growth rate and toxin production in the seed pathogen *Pyrenophora semeniperda*: resource trade-offs and temporally varying selection. Plant Pathology **64**, 1450–1460.

Murase K, Hirano Y, Sun T, Hakoshima T. 2008. Gibberellin-induced DELLA recognition by the gibberellin receptor GID1. Nature **456**, 459–463.

Nambara E, Marion-Poll A. 2005. Abscisic Acid Biosynthesis and Catabolism. Annual Review of Plant Biology **56**, 165–185.

Nambara E, Suzuki M, Abrams S, McCarty DR, Kamiya Y, McCourt P. 2002. A Screen for Genes That Function in Abscisic Acid Signaling in *Arabidopsis thaliana*. Genetics **161**, 1247–1255.

Okamoto M, Peterson FC, Defries A, Park S-Y, Endo A, Nambara E, Volkman BF, Cutler SR. 2013. Activation of dimeric ABA receptors elicits guard cell closure, ABA-regulated gene expression, and drought tolerance. Proceedings of the National Academy of Sciences **110**, 12132–12137.

Park S-Y, Fung P, Nishimura N, *et al.* 2009. Abscisic Acid Inhibits Type 2C Protein Phosphatases via the PYR/PYL Family of START Proteins. Science **324**, 1068–1071.

Pri-Tal O, Shaar-Moshe L, Wiseglass G, Peleg Z, Mosquna A. 2017. Non-redundant functions of the dimeric ABA receptor BdPYL1 in the grass Brachypodium. The Plant Journal: For Cell and Molecular Biology **92**, 774–786.

Rodriguez PL. 2016. Abscisic Acid Catabolism Generates Phaseic Acid, a Molecule Able to Activate a Subset of ABA Receptors. Molecular Plant **9**, 1448–1450.

Santiago J, Dupeux F, Round A, Antoni R, Park S-Y, Jamin M, Cutler SR, Rodriguez PL, Márquez JA. 2009. The abscisic acid receptor PYR1 in complex with abscisic acid. Nature **462**, 665–668.

Scherlach K, Boettger D, Remme N, Hertweck C. 2010. The chemistry and biology of cytochalasans. Natural Product Reports **27**, 869–886.

Schindelin J, Arganda-Carreras I, Frise E, *et al.* 2012. Fiji: an open-source platform for biological-image analysis. Nature Methods **9**, 676–682.

Schwartz SH, Tan BC, Gage DA, Zeevaart JAD, McCarty DR. 1997. Specific Oxidative Cleavage of Carotenoids by VP14 of Maize. Science **276**, 1872–1874.

Seo M, Peeters AJM, Koiwai H, Oritani T, Marion-Poll A, Zeevaart JAD, Koornneef M, Kamiya Y, Koshiba T. 2000. The Arabidopsis aldehyde oxidase 3 (AAO3) gene product catalyzes the final step in abscisic acid biosynthesis in leaves. Proceedings of the National Academy of Sciences **97**, 12908–12913.

Sheard LB, Tan X, Mao H, *et al.* 2010. Jasmonate perception by inositol-phosphate-potentiated COI1–JAZ co-receptor. Nature **468**, 400–405.

Shimada A, Ueguchi-Tanaka M, Nakatsu T, Nakajima M, Naoe Y, Ohmiya H, Kato H, Matsuoka M. 2008. Structural basis for gibberellin recognition by its receptor GID1. Nature **456**, 520–523.

Takezawa D, Komatsu K, Sakata Y. 2011. ABA in bryophytes: how a universal growth regulator in life became a plant hormone? Journal of Plant Research **124**, 437–453.

Vaidya AS, Peterson FC, Yarmolinsky D, *et al.* 2017. A rationally designed agonist defines subfamily IIIA ABA receptors as critical targets for manipulating transpiration. ACS Chemical Biology.

Villa R, Molinari F. 2008. Reduction of carbonylic and carboxylic groups by plant cell cultures. Journal of Natural Products **71**, 693–696.

Weng J-K, Ye M, Li B, Noel JP. 2016. Co-evolution of Hormone Metabolism and Signaling Networks Expands Plant Adaptive Plasticity. Cell **166**, 881–893.

Yamaguchi-Shinozaki K, Shinozaki K. 2006. Transcriptional Regulatory Networks in Cellular Responses and Tolerance to Dehydration and Cold Stresses. Annual Review of Plant Biology **57**, 781–803.

Figure legends

Figure 1. Pyrenophoric acid B inhibits seed germination.

(A) Structures of abscisic acid, ABA, pyrenophoric acids, P-Acid, P-Acid B and P-Acid C used in this work. (B) Pictures of Arabidopsis Col-0 seeds treated with DMSO, 1 μM ABA, and 100 μM of the different pyrenophoric acids, 3 days after shown. (C) Quantification of seedling establishment 3 days after shown Col-0 seeds on DMSO, 1 μM ABA, and 100 μM of the different pyrenophoric acids. Values represent mean ± SD of 90 seed used in three different experiments.

Figure 2. Pyrenophoric acid B (P-Acid B) activates the ABA signaling pathway to inhibit seed germination.

(A) Pictures of Arabidopsis seeds treated with 100 μM Pyrenophoric acid B or DMSO as control 3 days after sown. Col-0 seed was used along with mutants in the ABA signaling pathway. (B) Quantification of seedling establishment 3 days after sown seeds on DMSO or 100 μM of P-Acid B. Values represent mean ± SD of 90 seed used in three different experiments. (C) Luminescence detection (LUC) of seedlings of the ABA reporter line pMAPKKK18-LUC[+] treated with DMSO as control, 200 μM P-Acid B or 25 μM ABA for 6 hours. The image of the seedlings under bright light (BF) is also included. (D) Quantification of the pMAPKKK18-LUC[+] expression shown in C.

Figure 3. Pyrenophoric acid B (P-Acid B) does not activate ABA receptors.

PP2C phosphatase assay using HAB1 (1 μM) and different ABA receptors (2 μM) in the presence of DMSO as control, 10 μM ABA or 100 μM P-Acid B. HAB1 phosphatase activity was set to 100% in the absence of receptor and pNPP was used as a chromogenic substrate. Values represent mean ± SD (n=3).

Figure 4. Pyrenophoric acid B (P-Acid B) uses the ABA biosynthesis pathway to inhibit seed germination.

(A) Pictures of Arabidopsis seeds treated with 1 μM ABA, 100 μM P-Acid B or DMSO as control, 3 days after sown. Col-0 was used along with mutants in the ABA

biosynthesis pathway. (B) Quantification of seedling establishment 3 days after sown seeds on DMSO, 1 μM ABA or 100 μM P-Acid B. Values represent mean ± SD of 90 seed used in three different experiments.

Figure S1. Pyrenophoric acid B (P-Acid B) requires ABA2 and onwards to inhibit seedling establishment.

(A) Pictures of Arabidopsis seeds treated with 1 μM ABA, 100 μM P-Acid B or DMSO as control, 3 days after sown. (B) Quantification of seedling establishment 3 days after sown seeds on 100 μM P-Acid B. Values represent mean ± SD of 90 seed used in three different experiments.

A

	Col0	*112458*	HAB1^{W385A}	*abi3*	*abi4*
Control					
P-ACID B					

B

Seedling establishment (%)

□ Control
■ P-ACID B

Col0 112458 HAB1W385A abi3 abi4

C

	DMSO	P-ACID B	ABA
LUC			
BF			

D

MAP3K18-LUC expression (a.u.)

DMSO   P-ACID B   ABA