A survey of vaccine-induced measles IgG antibody titer and the verification of changes

in temporal differences of measles vaccination in young adults

Hiraku Sasaki[1*], Tomoko Fukunaga[2], Ai Asano[2], Yoshio Suzuki[3], Yuko Nakanishi[1],

Junzi Kondo[1], Hiroki Ishikawa[4], Nobuto Shibata[1, 2]

[1] Department of Health Science, [2] Section of Health Management, and [3] Department of

Sports Science, Faculty of Health and Sports Science, Juntendo University, 1-1

Hiraga-gakuendai, Inzai, Chiba 270–1695, Japan.

[4] Department of Microbiology and Immunology, Showa University School of Medicine,

1-5-8 Hatanodai, Tokyo 142–8555, Japan.

Correspondence: Dr. Hiraku Sasaki, Laboratory of Microbiology and Hygiene,

Department of Health Science, Faculty of Health and Sports Science, Juntendo

University 1-1 Hiraga-gakuendai, Inzai, Chiba 270-1695, Japan.

Abstract

In Japan, sporadic measles cases increased rapidly in 2019 compared to that in

past 6 years. To clarify the persistence of immunity against measles in young adult, this

study explored the prevalence of IgG antibody titer against measles virus in 18- to

24-year-old young adult participants. Measles-specific IgG antibody titers determined

by enzyme immunoassay in serum samples collected from 506 participants between 18

to 24 years were assessed through statistical analyses. Multiple regression analysis

revealed that the distribution of measles IgG antibody titers was significantly correlated

with medical history ($P < 0.05$), while there was no significant difference among the

number of vaccinations related to measles IgG titers. Further, measles IgG titers were

significantly different, which was determined by the temporal change that elapsed

period after last vaccination ($P < 0.05$). These results indicate that periodic vaccination

against measles is required for young and older adults to prevent even sporadic measles

infection.

Introduction

In Japan, more than 300 cases of measles were reported, which has increased

rapidly compared to that in the past 6 years [1]. Measles cases were reported in 21 of 47

prefectures, and in particular, a third part of measles case were found in Osaka

prefecture within 1st to 10th week in 2019 [1]. Taking measles vaccination twice in

lifetime induced acquired immunity against measles in more than 95% of people [2, 3].

For the measles vaccination program in Japan, a person born after April 2006 is defined

to have received 2 doses of measles vaccine as a periodic inoculation at 1-2 year and 5-7

years of age [4]. Before the Application of these regulations, there used to be only

primary vaccination after 12 months of age. Thereafter, there was a change in the

regulations of the Ministry of Health, Labor and Welfare recommending 2 doses of

measles vaccination for people born before April 2006 with secondary inoculation being

received within the duration of 5 years (2008–2012, equivalent to 13 and 18 years of

age) as special measures for periodic inoculation [4]. Within the 5 -years, the rate of

vaccination ranged from 77.3 to 94.3% [5]; however, recent vaccination rate exhibited

more than 90% of people receiving measles vaccines twice in Japan [6]. Except for the

people who received secondary vaccination at limited periods, almost all young adults

born after April 2006 were vaccinated twice before they were 7 years of age. A large

measles outbreak is thought to be physically impossible in later childhood due to the

implementation of the measles vaccine program; however, there was insufficient

knowledge about the existence of protective antibody titer against measles in young

adults who received secondary measles vaccine within the 5 years duration limit.

Currently, measles cases are based on age bracket. In brief, more than 50% of measles

cases were seen in 15–29-year-old individuals, while less than 10–14-year-old

constituted only 12% [1]. Almost all young adults around 20 years of age were

subjected to the special measures of periodic inoculation but all of them did not receive

the same manner of measles vaccination. In particular, temporal changes in protective

serum IgG antibody against measles virus may vary in spite of receiving secondary

vaccination lately compared to young adults. However, temporal changes in measles

IgG antibody titer in generation of young adults that was subjected to special measures

of measles vaccine are obscure. Thus, there is a need to elucidate the factors involved in

the risk of the infection as well as the ways of reducing the occurrence of the infection.

To clarify the persistence of immunity against measles in young adults, this

study explored the prevalence of IgG antibody titer against measles virus in 18–

24-year-old young adult participants, and determined the relationships between

temporal changes in the antibody.

Materials and Methods

Study design

The surveillance was carried out by obtaining samples of sera from 506 young

adults between 18 to 24 years of age, who were first-year students in Juntendo

University to assess the prevalence of specific IgG antibodies against measles virus. The

serum samples were obtained from January 2018 to April 2018. Simultaneously,

vaccine history was collected from each individual's maternity passbook. The studied

participants also filled out questionnaires aimed at obtaining information about the

medical history of vaccination against measles and natural measles infection. The

collected information was used for proper interpretation of the obtained results.

The study protocol was approved by the Ethics Committee of Faculty of Health

and Sports Science, Juntendo University (number 30-2) and informed consent was

obtained from all participants and their parents before the study.

Laboratory methods

85 To determine measles-specific serum IgG antibody titer, the enzyme

86 immunoassay (EIA) was performed in the laboratory of BML, INC. (Tokyo, Japan)

87 using a measles virus immunoglobulin test kit (the Measles IgG-EIA manufactured by

88 Denka Seiken., Co., Ltd, Tokyo, Japan). For the virus antigen, Toyoshima strain was

89 used.

Statistical analysis

To identify the factors that influence measles-specific IgG antibody titer, multiple

regression analysis was performed. To avoid multicollinearity, factors were preliminary

determined with bivariate analysis and were confirmed to have no correlation.

Differences in the IgG antibody titers between the participants that had a medical

history of measles and those without history were evaluated with the unpaired *t*-tests.

To compare differences in IgG antibody titers among the numbers of vaccination and

temporal characteristics, a one-way analysis of variance (one-way ANOVA) was

employed. When there were significant statistical differences, the data were further

analyzed using the Bonferroni *post hoc* test to determine the significance between the

groups. Differences were considered significant for P values of less than 0.05.

Results

Outline of survey participants

During the blood collection for determining antibody titers in this surveillance,

participants who were 18 years of age were 80% of total survey participants while the

remaining 20% of participants were $\geq$ 19 years old. All the subjects who were 19 years

old did not receive secondary inoculations. Approximately 80% of the participants

received secondary vaccination, and 10% received only primary vaccination (Table 1).

Approximately 4% of participants had a medical history of measles based on

self-assessment information.

Table 1 Outline of measles-vaccine induced antibody survey participants

Characteristic	Number	Percentage
Total	506	100
Age,Y		
Mean	18	-
Range	17-24	-
Sex		
Female	160	32
Male	346	68
Vaccination		
1	56	11
2	399	79
3	21	4
Unvaccinated	30	6
Medical history	22	4

Measles-IgG titer and temporal characterization

The mean value of measles-specific IgG antibody titer determined by EIA

exhibited 13.4 ±12.0 (Mean ± standard deviation) ranging from 0.8 to 128 (Table 2).

The period of the month when primary and secondary vaccines were received exhibited

23.6±32.5 and 152.2 ±25.4 month, respectively. The recommended age for primary

measles vaccination was to be from 12 to 90 months during 1995 to 2000, and therefore,

the duration for primary vaccination was extended. Almost all participants corresponded

to the special measure that was recommended for secondary vaccination at 13 years of

age during 2008 to 2013, indicating that more than 80% of participants received

secondary vaccination at the assigned period. Regarding all the vaccinated participants,

an average of 7 years passed from last vaccination.

Table 2 Measles-specific IgG antibody titer and temporal characteristics of participants.

Item	Mean	Range	Number
Measles-specificIgG titer	13.4	0.8 - 128	506
1st vaccination months	23.6	11- 263	476
2nd vaccination months	152.2	16 - 265	420
3rd vaccination months	156.0	53 - 238	21
Elapsed months after last vaccination	83.3	0 - 237	476

Divergence of measles-IgG antibody titers by medical history

To discern mode of measles IgG antibody titer's distribution, multiple regression

analysis was performed for the IgG antibody titers and temporal characteristics. It was

observed that almost all parameters related to temporal characteristics had

multicollinearity. Thus, multiple regression analysis was performed based on Table 1

listed items devoid of temporal information. Of these, the distribution of measles IgG

antibody titers was significantly correlated with medical history of measles ($P < 0.05$).

Then, the differences in measles IgG antibody titers between presence and absence of

medical history were compared by the unpaired t-tests (Figure 1). The IgG antibody

titers collected from participants who had medical history of measles (27.0 ± 31.8) were

significantly higher than titers from participants who had no history (12.8 ± 9.9, $P <$

0.05).

Figure 2 shows measles IgG antibody titers that were divided into each numbers

of measles vaccinated and unvaccinated participants. The measles IgG antibody titers

from the participants who had a medical history of measles were excluded from Figure

2 and statistical analysis since almost all participants who had a medical history had

161 been unvaccinated and exhibited higher IgG antibody titer than participants who had no

162 history. The results of one-way ANOVA showed that there was no significant

163 difference among the number of vaccinations.

165 Temporal changes in measles-IgG antibody titers

166   To verify the temporal changes in measles IgG antibody titers after the last

167 vaccination, the elapsed periods were divided into 5 periods including 3 month or less

168 (n = 9), 4–12 months (n = 8), 13–60 months (n = 31), 61–72 months (n = 279), and 73

169 months or more (n = 73). In measles vaccine program, almost all participants received

170 secondary vaccination at 13 years of age, and thus, approximately 70% of participants

171 were observed in the period 61–72 months. Excluding measles IgG antibody titers from

172 the participants who had a medical history and those who were unvaccinated, one-way

173 ANOVA showed that there were significant differences among the 5-periods (Figure 3,

174 $P < 0.05$). Although secondary period (4-12 month) contained a few dispersed measles

175 IgG antibody titer values, the mean values of measles IgG antibody titer decreased with

176 the prolonged period. The Bonferroni *post hoc* test showed that there were 4 significant

177 differences between each period ($P < 0.05$), indicating that measles IgG antibody titer

178 retains high value if elapsed period is short after last vaccination.

180 Discussion

To protect from measles virus infection, measles-specific serum IgG antibody

titer determined by the EIA method was required 12 or more, and in case of unsatisfied

titer, additional vaccination was recommended [7]. Although average measles IgG

antibody titer of total participants slightly exceeded 12, 56% of measles IgG antibody

titers from 284 participants showed less than 12. These results indicate that more than

half of the studied participants required additional dose to protect against measles.

According to WHO announcement, Japan was verified as having achieved measles

elimination that was defined as interruption of endemic measles virus transmission for

at least 36 months [8]. Nevertheless, measles cases from foreigners in Japan have

occurred sporadically [1, 9-12]. Based on our surveillance, one of the causes may be

non-persistent protection by measles IgG antibody titer in young adults. When measles

was endemic in Japan, approximately 200,000 people, mainly children, were infected

with measles virus in the year 2000 [13, 14]. Thereafter, an attempt to raise coverage of

measles vaccination and to enforce 2 doses vaccination rigorously was successfully

achieved, which led to the elimination-period [8]. Currently, protection against measles

infection has been achieved; "however, the generation with insufficient immunization

against measles virus is mainly the young adults suggesting that periodic monitoring of

measles epidemic and acquired immunity against measles virus in young adult is

required.

Further, this surveillance focused on the temporal status of measles IgG antibody

from last vaccination, although there was no significant difference in measles IgG

antibody titers after primary and secondary vaccination. In the United States,

vaccination schedule and patterns are similar to those in Japan, and measles was

declared eliminated with the absence of continuous measles transmission for a period

greater than 12 months in the United States in the year 2000 [15]. However, endemic

outbreaks of measles were reported to be yet to occur [16-18], and in rarity, a part of

those outbreaks was caused by unvaccinated population [19]. For primary vaccine

failure, CD46 and TLR8 variants were considered involved in the occurrence of measles

vaccine failure [20]. Although it will be possible to have such cases, epidemiological

studies have demonstrated the efficacy of measles vaccine. In brief, 95% of children

who received measles vaccine acquired immunity against measles virus, and further,

additional secondary vaccine led to more than 99% immunization in children [21-23].

These results revealed that epidemic outbreaks might be caused by the unvaccinated

population or large number of international travelers [3]. Further, according to the large

surveillance by healthcare workers, serum measles IgG antibody titers from adults less

than 29 years of age showed susceptibility to measles [24]. Single dose of measles

vaccine in adults was reported to have significantly increased serum IgG titers even in

the initial insufficient IgG titers [25]. Sporadic infection may occur in young and older

adults under unprotected conditions and with reduced vaccine-induced IgG antibody

titer due to the temporal changes after last vaccination and consequent susceptibility to

infection. These results also suggest that preventive vaccination against measles is

required for young and older adults to prevent even sporadic measles cases.

Acknowledgements

This work was partially supported by JSPS KAKENHI Grant Number 16K07095.

References

[1] National Institute of Infectious Disease, Infectious Disease Surveillance Center.

Infectious diseases weekly report Japan 2019;21:10.

https://www.niid.go.jp/niid/ja/idwr-dl/2019.html.

[2] Demicheli V, Rivetti A, Debalini MG, Di Pietrantonj C. Vaccines for measles,

mumps and rubella in children. Cochrane Database Syst Rev 2012;(2):CD004407.

[3] Moss WJ, Griffin DE. Measles. Lancet 2012;379(9811):153–64.

[4] Ministry of Health, Labor and Welfare. Measles,

https://www.mhlw.go.jp/seisakunitsuite/bunya/kenkou_iryou/kenkou/kekkaku-kans

enshou/measles_eng/index.html.

[5] Tanaka-Taya K. Progress towards the 2012 measles elimination goal in Japan.

Uirusu 2010;60(1):59–68.

[6] Ministry of Health, Labor and Welfare.

https://www.mhlw.go.jp/bunya/kenkou/kekkaku-kansenshou21/hashika.html.

[7]   Takayama N, Saika S, Ichinohe S. EIA-IgG antibody measles prevention level

estimated from measles neutralizing, particle agglutination and

hemagglutination-inhibition antibody titer. Kansenshogaku Zasshi 2009;83(5):519–

24.

[8]   World Health Organization. Disease outbreak news. Emergencies preparedness,

response. Measles – Japan,

https://www.who.int/csr/don/20-june-2018-measles-japan/en/;2018.

[9]   Komabayashi K, Seto J, Tanaka S, Suzuki Y, Ikeda T, Onuki N, Yamada K, Ahiko

T, Ishikawa H, Mizuta K. The largest measles outbreak, including 38 modified

measles and 22 typical measles cases in its elimination era in Yamagata, Japan in

2017. Jpn J Infect Dis 2018;71(6):413–8. doi: 10.7883/yoken.JJID.2018.083.

[10]Miyoshi M, Komagome R, Yamaguchi H, Ohnishi A, Kikuchi M, Ishida S, Nagano

H, Okano M. Detection of measles virus genotypes B3, D4, D5, D8, and H1 in the

surveillance system in Hokkaido, Japan, 2006-2015, the last decade toward the

elimination. Jpn J Infect Dis 2017;70(3):317–9. doi: 10.7883/yoken.JJID.2016.253.

[11]Nishiura H, Mizumoto K, Asai Y. Assessing the transmission dynamics of measles

in Japan in 2016. Epidemics 2017;20:67–72. doi: 10.1016/j.epidem.2017.03.005.

[12]Shimizu K, Kinoshita R, Yoshii K, Akhmetzhanov AR, Jung S, Lee H, Nishiura H.

An investigation of a measles outbreak in Japan and China, Taiwan, China,

March-May 2018. Western Pac Surveill Response J 2018;9(3):25–31. doi:

10.5365/wpsar.2018.9.2.005.

[13]Nakatani H, Sano T, Iuchi T. Development of vaccination policy in Japan: current

issues and policy directions. Jpn J Infect Dis 2002;55(4):101–11.

[14]Gomi H, Takahashi H. Why is measles still endemic in Japan? Lancet.

2004;364(9431):328–9.

[15]Papania MJ, Wallace GS, Rota PA, Icenogle JP, Fiebelkorn AP, Armstrong GL,

Reef SE, Redd SB, Abernathy ES, Barskey AE, Hao L, McLean HQ, Rota JS,

Bellini WJ, Seward JF. Elimination of endemic measles, rubella, and congenital

rubella syndrome from the Western hemisphere: the US experience. JAMA Pediatr

2014;168(2):148–55.

[16]Fiebelkorn AP, Redd SB, Gastañaduy PA, Clemmons N, Rota PA, Rota JS, Bellini

WJ, Wallace GS. A comparison of postelimination measles epidemiology in the

United States, 2009-2014 versus 2001-2008. J Pediatric Infect Dis Soc

2017;6(1):40–8.

[17]Ehresmann K, Banerjee E, Kenyon C, Christianson B, Griffith J, Heath J, Roddy

M. MDH IDEPC Measles Outbreak Team. Response to "It could have been much

worse: The Minnesota measles outbreak of 2017". Vaccine 2019;37(5):683–4. doi:

10.1016/j.vaccine.2018.11.031.

[18]Clemmons NS, Wallace GS, Patel M, Gastañaduy PA. Incidence of measles in the

United States, 2001-2015. JAMA 2017;318(13):1279–81. doi:

10.1001/jama.2017.9984.

[19]Nelson R. US measles outbreak concentrated among unvaccinated children. Lancet

Infect Dis 2019;19(3):248. doi: 10.1016/S1473-3099(19)30074-X.

[20]Clifford HD, Hayden CM, Khoo SK, Naniche D, Mandomando IM, Zhang G,

Richmond P, Le Souëf PN. Polymorphisms in key innate immune genes and their

effects on measles vaccine responses and vaccine failure in children from

Mozambique. Vaccine 2012;30(43):6180–5. doi: 10.1016/j.vaccine.2012.07.063.

[21]Moss WJ, Strebel P. Biological feasibility of measles eradication. J Infect Dis

2011;204 Suppl 1:S47–53. doi: 10.1093/infdis/jir065.

[22]Meissner HC, Strebel PM, Orenstein WA. Measles vaccines and the potential for

worldwide eradication of measles. Pediatrics 2004;114(4):1065–9.

[23]Moss WJ, Griffin DE. Measles. Lancet. 2012;379(9811):153–64. doi:

10.1016/S0140-6736(10)62352-5.

[24]Kumakura S, Shibata H, Onoda K, Nishimura N, Matsuda C, Hirose M.

Seroprevalence survey on measles, mumps, rubella and varicella antibodies in

healthcare workers in Japan: sex, age, occupational-related differences and vaccine

efficacy. Epidemiol Infect 2014;142(1):12–9.

[25]Ogawa T, Uchiyama-Nakamura F, Sugata-Tsubaki A, Yamada Y, Uno K, Kasahara

K, Maeda K, Konishi M, Mikasa K. Antibody response to live attenuated vaccines

in adults in Japan. Open Med (Wars) 2016;11(1):482–8. doi:

10.1515/med-2016-0085.

Figure legends

Figure 1. Comparison of measles-specific IgG antibody titers based on the medical

history of participants.

The plots of IgG antibody titer that were more than 60 were omitted from Figure 1. The

whiskers extended to data points that were less than 1.5 x IQR away from 1st/3rd

quartile.

Figure 2. Comparison of measles-specific IgG antibody titers between the number of

vaccinated and unvaccinated participants.

The plots of IgG titer that were more than 60 were omitted from Figure 2. The measles

IgG antibody titers from the participants who had a medical history were excluded from

Figure 2 and statistical analysis. The whiskers extended to data points that were less

than 1.5 x IQR away from 1st/3rd quartile. There were no significant differences

between number of vaccinated and unvaccinated participants based on one-way

ANOVA ($P \geq 0.05$).

Figure 3. Comparison of measles-specific IgG antibody titer among 5-periods of

elapsed months after last vaccination

The plots of IgG titer that were more than 60 were omitted from Figure 3. The measles

IgG antibody titers of both the participants who had a medical history and the

unvaccinated participants were excluded from Figure 3 and statistical analysis. The

whiskers extended to data points that were less than 1.5 x IQR away from 1st/3rd

quartile. One-way ANOVA showed that there were significant differences among the

5-periods ($P < 0.05$). An asterisk denotes a significant difference between periods with

connected line ($P < 0.05$).

Figure 1

Measles-specific IgG titer

Number of vaccination

Figure 2

Figure 3