Plant A20/AN1 proteins coordinate different immune responses including RNAi

pathway for antiviral immunity

Li Chang[a,b,c], Ho-Hsiung Chang[b], Yi-Shu Chiu[b], Jui-Che Chang[b], Duen-Wei Hsu[d], Yuh

Tzean[b], An-Po Cheng [b], Hsiang-Chia Lu[e], and Hsin-Hung Yeh[b,e#]

[a]Molecular and Biological Agricultural Sciences Program, Taiwan International

Graduate Program, National Chung-Hsing University, Academia Sinica, Taiwan

[b]Agricultural Biotechnology Research Center, Academia Sinica, Taipei, Taiwan

[c]Graduate Institute of Biotechnology, National Chung-Hsng University, Taichung,

Taiwan

[d]Department of Biotechnology, National Kaohsiung Normal University, Kaohsiung,

Taiwan.

[e]Department of Plant Pathology and Microbiology, National Taiwan University, Taipei,

Taiwan.

Running Heads: Plant SAPs coordinate RNAi-mediated antiviral immunity

#Address correspondence to Hsin-Hung Yeh, hyeh@sinica.edu.tw

ABSTRACT

Salicylic acid (SA)-mediated immunity plays important roles in combating virus in plants. Two plant stress associated protein (SAPs) containing dual A20/AN1 zinc-finger domain were found to play important roles in SA-mediated immunity; however, detailed mechanisms remain elusive. In this study, another orchid homolog gene of *Pha13*, *Pha21*, was analyzed. Pha21 confers antiviral immunity in both transgenic orchid and *Arabidopsis* overexpressing Pha21. Expression of *Pha21* is early-induced by SA treatment, and is involved in the expression of the orchid homolog of the master regulator *NPR1*. Pha21 but not Pha13 is involved in the expression of key RNAi-related genes, *Dicer-like nuclease 4* (*DCL4*) and *Argonaut 1* (*AGO1*) in orchids. The involvement of SAPs in expression of orchid *DCL4* and *AGO1* is not limited to orchid, as AtSAP5 also plays essential role in the expression of *Arabidopsis DCL4* and *AGO1*. However, unlike Pha13 and AtSAP5, Pha21 does not play positive role in the expression of orchid homolog gene of *RNA-dependent RNA polymerase 1* (*RdR1*), an important gene in RNAi pathway. Pha21 can be found localized in the nucleus, and confers self-E3 ligase and ubiquitin binding activities. Functional domain analysis revealed that both A20 and AN1 domains of Pha21 are required for decreasing virus accumulation, and the AN1 domain plays a more important role in the expression of orchid *DCL4*. Collectively, our data suggests SA regulated SAPs play important roles in antiviral immunity and is involved in delicate regulation of key genes in RNAi-mediated pathway.

IMPORTANCE

Salicylic acid (SA)-mediated antiviral immunity plays an important role to protect

plants from virus infection; however, the detailed mechanisms remain elusive. We

previously demonstrated that two plant A20/AN1 proteins, orchid Pha13 and

Arabidopsis AtSAP5, function similarly and serve as an important hub to regulate SA-

mediated antiviral immunity. In this study, we identified another orchid A20/AN1

protein, Pha21, which is involved in SA-mediated antiviral immunity. In contrast to

Pha13 and AtSAP5, Pha21 plays minor negative roles in the expression of *PhaRdR1*

(orchid homolog of *RNA-dependent RNA polymerase 1*). However, Pha21 and AtSAP5,

but not Pha13, are involved in the expression of important players in RNAi pathway,

Dicer-like nuclease 4 (*DCL4*) and *Argonaut 1* (*AGO1*), in orchid and *Arabidopsis*. Our

data demonstrates that plant A20/AN1 proteins are conserved players in SA-mediated

antiviral resistance among plants, and provide links between the A20/AN1 proteins and

the RNAi pathway.

INTRODUCTION

The plant hormone salicylic acid (SA) is important to trigger plant immunity

especially against biotrophic pathogens such as viruses (1-4). In facing pathogen

invasion, plants recognize conserved microbe-associated molecular patterns (MAMPs)

by pattern-recognition receptors and trigger pathogen (or pattern)-triggered immunity

(PTI) as the first line of defense (5, 6). Although PTI provides protection from invasion

by most pathogens, some have evolved, and can utilize different effectors to suppress

PTI for successful infection (7). Plants have evolved resistance (R) proteins capable of

detecting these effectors to trigger effecter-triggered immunity (ETI) for a second line

of plant defense (7). ETI is generally associated with programmed cell death leading to

the formation of necrotic lesions, which is also known as hypersensitive response (HR)

to prevent the further spreading of pathogen infection (1).

Current evidence suggests that the PTI is important to limit virus infection (8-10),

and the viral double strand RNA (dsRNA) has been shown to serve as a conserved

MAMP (10). In addition, ETI also plays an important role in combating virus infection

in plants. For example, the N gene from *Nicotiana glutinosa* serves as an R protein to

specifically recognize the effector from *Tobacco mosaic virus* (TMV) and trigger the

ETI (11-14). SA plays important roles in triggering PTI and ETI (15). In addition, the

TMV-infected tobacco plant showing local necrotic lesions was found to become more

resistant against secondary virus infection in the distal leaves (16, 17). This systemic

immune response is known as systemic acquired resistance, which is a common plant

immune response and plays an important role in protecting plants from pathogen

infection (4).

In addition, RNA silencing or RNA interference (RNAi) has also been

demonstrated to play an important role in combating virus infection (18, 19). RNAi is

activated through the appearance of the viral dsRNA upon virus infection. The dsRNA

can be cleaved to short small-interfering dsRNA (siRNA) of 21–24 nucleotides (nt) in

size by the Dicer-like (DCL) nuclease (20). DCL2 and DCL4 in *Arabidopsis* play an

important role in generating the virus-derived siRNA (vsiRNA) (19, 21, 22). The

vsiRNA is further unwound and one strand (guide-strand) is incorporated into the RNA-

induced silencing complex (RISC) (19). This complex targets viral RNA with a

sequence that is complementary to the guide-strand and degrades the target RNA by

the catalytic component of the RISC, Argonaut (AGO) nucleases (19, 20, 23, 24).

Among the *Arabidopsis* AGOs, AGO1 plays an important role in the antiviral immunity

against RNA viruses. (24). More vsiRNA can be further generated *de novo* through the

cellular RNA-dependent RNA polymerases (RDRs) to trigger the secondary RNA

silencing against viruses (25-27). SA also plays important roles in the RNAi-pathway

as SA treatment can induce genes involved in the RNAi pathway including *RDRs* in

Arabidopsis, *Nicotiana tabacum and* Tomato , and *DCLs* and *AGOs* in Tomato (28-32).

SA induces the fluctuation of redox and serves as a signal to activate sets of

defense genes (33, 34). The SA-induced redox change can modify the NPR1

(nonexpressor of pathogenesis-related protein 1) from the multimeric protein complex

to a monomer via the oxidoreductases, Thioredoxin-3 and Thioredoxin-5 (35, 36). The

NPR1 monomer moves into the nucleus and activates multiple defense-related genes

including the *PR* (*pathogenesis-related*) gene in the SA-signaling pathway (37-39). In

addition to the NPR1-dependent immune pathway, some data also suggest that the

existence of a NPR1-independent pathway contributes to the virus resistance (40, 41).

Despite the importance of the SA governed immune pathway in antiviral immunity, the

regulation of this pathway remains largely elusive.

Proteins containing zinc-finger A20 and/or AN1 domains play an important role

in plant response against various abiotic stresses, and are known as stress associated

protein (SAPs). SAPs are conserved among different organisms (42, 43). Different

numbers of SAP homologs (range from 1 to 19) have been identified in organisms

including protists, fungi, animals, plants, and humans (42, 43). Compared to animals,

more of these proteins are found in plants. So far, 18, 14 and 14 SAPs have been

identified in rice, *Arabidopsis*, and *Phalaenopsis* orchid, respectively (42, 44).

Biochemical studies of proteins containing A20 and/or AN1 domains revealed that the

A20 domain confers E3 ligase and ubiquitin binding activity (43-49). The A20 domain

of human A20, Rabex-5 (guanine nucleotide exchange factor), *Arabidopsis* AtSAP5

and orchid Pha13 have been reported to exhibit E3 ligase activity (44, 45, 49, 50). In

addition, it has been shown that the A20 domain can also bind to various ubiquitin

chains (44, 46-48). In contrast to the A20 domain, the biochemical function of the AN1

domain is not fully understood.

Our recent study indicates that plant SAPs play a pivotal role in SA governed

antiviral immunity (44). Orchid *Pha13* and *Arabidopsis AtSAP5* are induced by SA

treatment at the early stage, and are involved in expression of orchid or *Arabidopsis*

NPR1 and *NPR1*-independent immune responsive genes including the induction of

plant *RdR1*. In addition to Pha13, our previous virus-induced gene silencing (VIGS)

assay also allowed us to identify the involvement of orchid *PhaTF21* (designated here

as *Pha21*) in SA–regulated immune response genes (51). Pha13 and Pha21 share 69.5%

amino sequence identity, and are most closely related among the orchid SAPs (44).

Whether Pha21 and Pha13 work in a cooperative manner in plant antiviral immunity

remains to be explored.

Here, we performed a detailed study of Pha21, and compared its biochemical and

physiological function to Pha13. Pha21 and Pha13 share similar biochemical properties

including E3-ligase and ubiquitin binding activities. Our results also indicate that

similar to *Pha13*, *Pha21* is early-induced by SA, involved in the expression of the

PhaNPR1 and *PhaNPR1*-independent genes, and plays an important role in antiviral

immunity. Importantly, our studies show that Pha21 is involved in expression of orchid

DCL4 and *AGO1*. Together with our previous data, these studies suggest that Pha13

and Pha21 coordinate the expression of genes important in the RNAi pathway including

orchid *RdR1*, *DCL4* and *AGO1*. In addition, our previous data and data presented in

this study also showed that *Arabidopsis* AtSAP5 is involved in expression of *RdR1*,

DCL4 and *AGO1*. Collectively, our data indicates that plant A20/AN1 proteins

involved antiviral immunity are conserved among plants, and the antiviral immunity is

partly through the RNAi pathway. Our findings suggest that A20/AN1 proteins may

serve as a link between SA and the RNAi pathway at the early stage of SA induced-

antiviral immunity.

RESULTS

Sequence and expression pattern analysis of Pha21

Previously, we reported that Pha21 (Orchidstra 2.0 database,

http://orchidstra2.abrc.sinica.edu.tw, accession number PATC144963) is involved in

the SA induced immune pathway in *P. aphrodite* (52). Pha21 contains dual zinc-finger

domains, A20 and AN1, in the N-terminal and C-terminal, respectively (Fig. 1A-C),

and belongs to the stress associated protein (SAP) family (43). Among the SAPs of *P.*

aphrodite, Pha21 is most related to our previously reported Pha13 and shared 69.5%

amino acid sequence identity (44). The A20 and AN1 zinc finger domains of Pha21

also shared high amino acid sequence identity among SAPs from different species

including AtSAP5 from *Arabidopsis,* and OsSAP3 and OsSAP5 from *Oryza sativa* (Fig.

1B). (44). Unlike Pha13, no nuclear localization signal was identified in Pha21 (Fig.

1A).

The expression of *Pha21* was analyzed in different tissues of *P. aphrodite* orchid

including root, leaf, septal, petal, lip, and column. The results revealed that *Pha21*

expression was higher in the column, root and leaf (Fig. 1D). In addition, we also

analyzed the absolute expression of *Pha21* and *Pha13* in the leaves of *P. aphrodite* by

use of droplet digital PCR. The results indicated that *Pha21* RNA expression is about

15 times lower than *Pha13* in leaves of *P. aphrodite* (Fig. 1E)

Pha21 is involved in the expression of SA responsive genes, *PhaNPR1* and *PhaPR1*

To further analyze the role of *Pha21* in the SA-induced immune pathway, we

transiently expressed two hairpin RNA (will generate 21 nt siRNA) of Pha21

(35S::hpPha21-1 and 35S::hpPha21-2) separately to knockdown Pha21 by

agroinfiltration in *P. aphrodite* carrying hairpin RNA expression construct, phpPha21-

1 and phpPha21-2 (Fig. 2A). Samples collected from the infiltrated site of orchid leaves

were used to detect the RNA level of Pha21, PhaNPR1, and PhaPR1 by quantitative

RT-PCR (qRT-PCR). As shown in Fig. 2A, the expression of *PhaNPR1* and *PhaPR1*

was decreased in both Pha21-silenced plants. In addition to transient knockdown of

Pha21, we also transiently overexpressed Pha21 (35S::FLAG-Pha21) through

agroinfiltration in the leaves of *P. aphrodite.* However, transient overexpression of

Pha21 in leaves did not significantly affect the RNA level of PhaNPR1 and PhaPR1

(Fig. 2B).

In addition, we also transiently silenced PhaNPR1 (phpPhaNPR1) in *P. aphrodite*

to analyze the expression of *Pha21*. The results showed that the silenced PhaNPR1

showed decreased expression of *PhaPR1*; however, no obvious effect was observed on

the expression of *Pha21* (Fig. 2C). These results suggest that Pha21 is involved in the

expression of *PhaNPR1*, but not vice versa.

Pha21 is induced by SA, jasmonic acid and ethylene

We also tested whether *Pha21* was induced by defense-related plant hormones

including SA, jasmonic acid (JA) and ethylene (ET). Orchid plants were treated with

SA, JA and ET and samples were collected after treatment at different time points (up

to 72 h) for analysis of the expression of *Pha21* and the marker genes of each plant

hormone by qRT-PCR. The induction of *Pha21* by SA was observed at 1 h post-

treatment (Fig. 2D). In addition, *Pha21* was also induced by JA and ET at 72 h and 12

h, respectively (Fig. 2E and F). Our results showed that *Pha21* can respond to multiple

defense-related plant hormones, and SA induced *Pha21* expression at the very early

stage.

Pha21 plays a positive role in virus resistance

To analyze whether Pha21 plays a role in virus resistance, we first analyzed the

expression of *Pha21* in response to virus infection. We detected the RNA level of Pha21

in mock- or CymMV-inoculated *P. aphrodite* using qRT-PCR, and the results showed

that the RNA of Pha21 is induced by CymMV infection (Fig. 3A). Furthermore, we

transiently silenced or overexpressed Pha21 in CymMV-infected *P. aphrodite* to assay

the effect on virus accumulation. Transient knockdown of Pha21 had no significant

effect on CymMV accumulation (Fig. 3B); whereas transient overexpression of Pha21

decreased CymMV accumulation (Fig. 3C). In addition, we also generated transgenic

P. equestris overexpressing Pha21 (35S::FLAG-Pha21). The expression of *Pha21* is

higher in the two individual asexually propagated progeny derived from two T0 lines

of transgenic *P. equestris* (Pha21#8 and Pha21#9) as compared to the non-transgenic

lines (WT) (Fig. 3D). We further inoculated CymMV into individual progenies derived

from two transgenic lines, and the result showed that the accumulation level of CymMV

is decreased to 21% and 34% in the two transgenic lines (Pha21#8 and Pha21#9) as

compared to the non-transgenic lines (WT) (Fig. 3D). Our data suggest that Pha21 plays

a positive role in virus resistance.

Transgenic *Arabidopsis* overexpressing Pha21 enhances antiviral resistance.

To understand whether Pha21-mediated antiviral immunity is conserved in plants,

we generated the transgenic *Arabidopsis* (Col-0) overexpressing Pha21 (35S::FLAG-

Pha21). Homozygous T3 plants derived from 3 T1 transgenic lines, At-Pha21#4, #5

and #6, were selected for further antiviral assay. The expression level of *Pha21* was

confirmed by qRT-PCR on the homozygote progenies (Fig. 4A).

We mechanically inoculated *Cucumber mosaic virus* (CMV) to wild-type (WT)

and Pha21 overexpressing *Arabidopsis* (At-Pha21#4, #5 and #6). Fourteen days post-

inoculation, the accumulation of CMV in WT and Pha21 overexpressing *Arabidopsis*

were detected by qRT-PCR. The results showed that Pha21 overexpressing *Arabidopsis*

decreased the accumulation of CMV (Fig. 4A) compared to the WT, and severe disease

symptoms were observed on the WT but not in the Pha21 overexpressing *Arabidopsis*

(Fig. 4B).

Subcellular localization of Pha21

To better understand the role of Pha21 within cells, we first analyzed the

subcellular localization of Pha21. We fused green fluorescent protein (GFP) in the N-

or C-terminal of Pha21 driven by 35S promoter (pG-Pha21 and pPha21-G). The GFP-

fused Pha21 was further transfected into protoplasts isolated from *P. aphrodite.*

Twenty-four hours post-transfection, the localization of Pha21 was observed by using

a confocal microscope. The results indicated that the C-terminal GFP fusion protein

(Pha21-G) was observed in the nucleus in about 50% of cells; whereas the N-terminal

GFP fusion protein (Pha21-G) showed a similar result to our GFP control vector, which

had no nucleus-specific GFP (Fig. 5A). Our data suggested that Pha21 can move into

the nucleus even without the predicated NLS signal.

Pha21 and Pha13 does not confer transcriptional activation ability in yeast two-

hybrid assay

Since both Pha21 and Pha13 have the ability to move into the nucleus (44), we tried

to understand whether Pha21 and Pha13 function as transcription factors. Therefore,

we analyzed its transcriptional activation ability through yeast two-hybrid (Y2H) assay.

We fused Pha21 and Pha13 to the Gal4 DNA binding domain to generate pGBKT7-

Pha21 and pGBKT7-Pha13. We transformed the pGBKT7-Pha21 or pGBKT7-Pha13

in the yeast strain AH109 without any Gal4 transcriptional activation domain (AD) and

further analyzed the transcriptional activation ability through the analysis of the

activation of reporter genes in yeast. As shown in Fig. 5B, yeast with pGBKT7-Pha21

or pGBKT7-Pha13 cannot grow on the -Trp/Aureobasidin and -Trp/-Ade/-His medium,

and the colonies did not turn blue on X-α-Gal containing medium.

Pha21 exhibits E3 ligase activity in which the A20 domain plays a major role

Several A20 and/or AN1 domain containing proteins confer E3 ligase activity (44-

46, 53). To determine whether Pha21 functions as an E3 ligase, *in vitro* self-

ubiquitination E3 ligase activity assay was performed using FLAG-Ubiquitin (FLAG-

Ub), human E1 (hE1), human E2 (UBCH2, hE2), and purified His-tagged Pha21

(Pha21-His). The poly-ubiquitinated Pha21 can be detected in the presence of FLAG-

Ub, hE1, and hE2 using anti-FLAG antibody to detect FLAG-Ubiquitin. The results

revealed that Pha21 confers E3 ligase activity (Fig. 6A).

Furthermore, we also analyzed the self-ubiquitination E3 ligase activity of the

A20 and/or AN1 domain of Pha21. We substituted the conserved cysteine and histidine

to glycine on the A20 and/or AN1 domains of Pha21-His (Fig 1B and C) to generate

A20 domain mutant (Pha21-A20m), AN1 domain mutant (Pha21-AN1m), and the

double mutant (Pha21-A20mAN1m) for E3 ligase activity analysis. As shown in Fig

6A, the major E3 ligase activity was conferred by the A20 domain. A20 domain mutant

and double mutant of Pha21 showed lower self-ubiquitination E3 ligase activity than

wild-type Pha21 and Pha21-AN1 mutant (Fig. 6A).

A20 domain of Pha21 confers ubiquitin binding ability

A20 and/or AN1 proteins have also been shown to confer ubiquitin binding ability

in animals and plants (44, 46, 47, 54, 55). To analyze whether Pha21 has ubiquitin

binding ability, Y2H assay was performed to verify the interaction using Pha21 as bait

and ubiquitin as prey. As shown in Fig 6B, Pha21 had a positive interaction with

ubiquitin, suggesting that Pha21 conferred ubiquitin binding ability. In addition, to

identify the ubiquitin binding domain of Pha21, a series of deletion mutants of Pha21

were generated, followed by Y2H assay. The result revealed that only truncated Pha21

fragments containing A20 domain (Pha21$^{1-136}$, Pha21$^{1-101}$, Pha21$^{1-40}$, Pha21$^{16-158}$,

Pha21$^{16-136}$, Pha21$^{16-101}$, Pha21$^{16-40}$) showed a positive interaction with ubiquitin,

suggesting that the A20 domain of Pha21 is responsible for the ubiquitin binding ability

(Fig. 6B).

Interaction analysis between orchid SAPs

The A20/AN1 proteins may confer self-interaction and also interact with each other

(56). Therefore, we also analyzed whether Pha21 and Pha13 confer self-interaction and

also interact with each other. The Y2H assay was performed to verify the interaction

between Pha21 and Pha13. Our Y2H results suggested that Pha21 and Pha13 did not

interact with each other and no self-interaction of Pha21 or Pha13 was observed (Fig.

7).

Pha21 is involved in the expression of SA-induced PhaNPR1-dependent and -

independent genes.

To further understand how Pha21 is involved in the antiviral immunity, we

performed transient silencing (35S::hpPha21-2) and overexpression assay

(35S::FLAG-Pha21) of Pha21 in *P. aphrodite*, and analyzed the expression of the

previously identified PhaNPR1-dependent and -independent antiviral genes,

Phalaenopsis homolog of *RNA dependent RNA polymerase 1* (*PhaRdR1)* and

Glutaredoxin C9 (PhaGRXC9), respectively (44). Our results indicated that transient

silencing of Pha21 RNA increased the expression of *PhaRdR1*, while *PhaGRXC9*

remained unchanged (Fig. 8A). Transient overexpression of Pha21 RNA decreased the

expression of *PhaGRXC9*, but *PhaRdR1* remain unchanged (Fig. 8B). Our data suggest

that Pha21 affects the expression of *PhaRdR1* and *PhaGRXC9*.

Pha21 is involved in the expression of *PhaDCL4* and *PhaAGO1*

Because the RNAi pathway plays an important role in antiviral immunity (18, 19),

we analyzed whether Pha21 is involved in the expression of core RNAi-related genes

including the *Phalaenopsis* homolog genes of *RdR2* (*PhaRdR2*, PATC124544), *RdR6*

(*PhaRdR6*, PATC131836), *DCL2* (*PhaDCL2*, PATC143544), *DCL4* (*PhaDCL4*,

PATC150652), *AGO1* (*PhaAGO1*, PATC157237), and *AGO10* (*PhaAGO10*,

PATC093469). Transient overexpression and silencing assay of Pha21 was performed

in *P. aphrodite*. Overexpression of Pha21 increased the RNA expression of *PhaDCL4*

and *PhaAGO1* (Fig. 9A); whereas transient silencing of Pha21 had no significant effect

on the RNA expression of RNAi-related genes (Fig. 9B).

For comparison, we also analyzed the effect of Pha13 on the RNAi-related genes.

The results showed that transient overexpression of Pha13 has no significant effect on

the expression of *PhaRdR2*, *PhaRdR6*, *PhaDCL2*, *PhaDCL4*, *PhaAGO1*, *and*

PhaAGO10 (Fig. 9C). Our results suggested that Pha21 is involved in the expression

of *PhaDCL4* and *PhaAGO1*.

To see whether orchid DCL4 (PhaDCL4) and AGO1 (PhaAGO1) also play a role

in the antiviral immunity, we transiently silenced PhaDCL4 and PhaAGO1 through

delivering the hairpin RNA into *P. aphrodite* by agroinfiltration carrying hairpin RNA

expression constructs, phpPhaDCL4 and phpPhaAGO1. Our results showed that

transient silencing of PhaDCL4 and PhaAGO1 increased the accumulation of CymMV

(Fig. 9D and E).

The *Arabidopsis* homolog gene of *Pha21*, *AtSAP5*, is involved in the expression

of *DCL4* and *AGO1*.

Our previous phylogenic analysis revealed that both Pha21 and Pha13 are closely

related to *Arabidopsis* AtSAP5 (accession number: AT3G12630) (44). Therefore, we

also analyzed whether AtSAP5 is involved in the expression of *DCL4* and *AGO1*. We

detected the expression of *DCL4* and *AGO1* in our previously generated transgenic

Arabidopsis overexpressing AtSAP5 (AtSAP5-oe-4 and 11) and RNAi lines of AtSAP5

(AtSAP5-RNAi-3 and 7) by qRT-PCR (44). The expression level of *AtSAP5* was

confirmed by qRT-PCR (Fig. 10A). Our results showed that slightly increased

expression of *DCL4* and *AGO1* were observed in overexpression transgenic lines,

AtSAP5-oe-11 and decreased expression of *DCL4* and *AGO1* were observed in the

RNAi lines, AtSAP5-RNAi-3 and AtSAP5-RNAi-7 (Fig. 10B and C). Our results

suggest that AtSAP5 is important in the expression of *DCL4* and *AGO1*.

A20 and AN1 domain of Pha21 play different roles in the expression of *PhaDCL4*,

***PhaAGO1*, *PhaGRXC9* and virus resistance.**

Our previous results indicated that Pha21 is involved in the expression of

PhaDCL4, *PhaAGO1*, and *PhaGRXC9* and virus accumulation (Figs 3C, 3D, 8B, and

9A). We further analyzed the roles of the A20 and/or AN1 domains of Pha21 in the

expression of *PhaDCL4*, *PhaAGO1*, *PhaGRXC9* and virus accumulation by

overexpression of Pha21 and Pha21 with mutation in the A20 and/or AN1 domain.

Therefore, we generated Pha21 with A20 domain mutant (Pha21A20m), AN1 domain

mutant (Pha21AN1m), and the double mutant (Pha21A20mAN1m) through the

substitution of the conserved cysteine and histidine to glycine on A20 and/or AN1

domain (Fig. 1B and C). We transiently overexpressed wild-type Pha21 or the mutants

(A20 and/or AN1 domain mutant) in healthy or CymMV pre-infected *P. aphrodite*, and

then detected *PhaDCL4, PhaAGRO1, PhaGRXC9,* and CymMV. The results indicated

that overexpression of Pha21 A20 mutant (Pha21A20m) but not AN1 mutant

(Pha21AN1m) or A20/AN1 mutant (Pha21A20mAN1m) increased the expression of

PhaDCL4, which is similar to the wild-type Pha21 (Fig. 11A). Overexpression of any

Pha21 A20 and/or AN1 mutant showed similar results to wild-type Pha21 with respect

to increased expression of *PhaAGO1* (Fig. 11A). Although overexpression of wild-type

and any Pha21 A20 and/or AN1 mutant resulted in decreased expression of *PhaGRXC9*

and CymMV accumulation compared to the vector control, overexpression of wild-type

Pha21 showed a greater effect on the expression of *PhaGRXC9* and CymMV

accumulation compared to any Pha21 A20 and/or AN1 mutant (Fig. 11A and B). Our

data suggest that both the A20 and AN1 domains of Pha21 are required for expression

of *PhaGRXC9* and the decrease of CymMV accumulation, and the AN1 domain plays

a major role in the expression of *PhaDCL4*. They also suggest that neither domain is

required for the expression of *PhaAGO1*.

DISCUSSION

Plant A20/AN1 proteins, Pha13, Pha21 and AtSAP5 share similar biochemical

properties and all involved in SA-mediated antiviral immunity

In this report, we performed detailed analysis of a previously identified orchid

gene, *Pha21*, involved in SA and RNAi-mediated antiviral immunity. Pha21 shares

high protein sequence identity to our previously identified Pha13 (Fig. S7 in Chang et

al, 2018), and exhibits similar biochemical properties including self-E3 ligase, ubiquitin

binding abilities (Fig. 6) and subcellular localization (Fig. 5A). Similar to Pha13, our

data suggests that Pha21 is involved in the SA-signaling pathway downstream of SA

and upstream of the expression of *PhaNPR1* (Fig. 2A, C and D). In addition, Pha13 and

Pha21 also positively mediate antiviral immunity (44). Transgenic *Arabidopsis*

overexpressing Pha21 also enhanced resistance to virus infection (Fig 4) (44). In

addition, our previous report and the data presented here indicate that *Arabidopsis*

AtSAP5 is also induced by SA at the early stage, and is involved in expression of similar

sets of SA mediated immune responsive genes, and positively regulates antiviral

immunity (44).

Pha13-mediated expression of *PhaGRXC9* and *PhaRdR1* are dominant during SA

treatment.

Our previous study indicated that Pha13 and AtSAP5 plays a positive role in the

expression of genes involved in the RNAi pathway and SA-governed oxidoreductases,

RdR1 and *GRXC9*, respectively (44). We demonstrated that both PhaRdR1 and

PhaGRXC9 play positive roles in antiviral immunity (44). In this study, we found that

Pha21 has an opposite role to Pha13 in the expression of *PhaRdR1* and *PhaGRXC9*

(Fig. 8). As our data indicates that the RNA level of Pha13 is approximately 15 times

higher than Pha21 in leaves of orchids as measured by digital PCR (Fig. 1E), we suggest

that the effect of Pha21 on the expression of *PhaRdR1* and *PhaGRXC9* is minor, and

Pha13-mediated positive expression of *PhaGRXC9* and *PhaRdR1* are dominant during

SA treatment.

Pha21 and AtSAP5 are all involved in expression of key genes in RNAi pathway,

but regulation of *PhaDCL4* and *PhaAGO1* by Pha21 in orchid is different from

regulation of *DCL4* and *AGO1* by AtSAP5 in *Arabidopsis*

Our data indicates that similar to *Arabidopsis* DCL4 and AGO1, orchid PhaDCL4

and PhaAGO1 also play a similar positive role in antiviral immunity in orchids as a

slight decrease in the RNA of PhaDCL4 or PhaAGO1 (approximately 40% decrease)

has a prominent effect on CymMV accumulation (approximately 100% increase) (Fig.

9D and E), (19). Interestingly, our data suggests that Pha21 regulates antiviral immunity

partly through the effect on the RNAi pathway, as transient overexpression of Pha21

(1.6-folds) but not Pha13 increased the expression of two *Phalaenopsis* orchid homolog

genes of *DCL4* (*PhaDCL4*, 2-fold) and *AGO1* (*PhaAGO1*, 1.6-fold) (Fig. 9A and C).

Although the effect of AtSAP5 on expression of *DCL4* and *AGO1* is not as

prominent as Pha21, our data shown here also indicates that AtSAP5 is involved in the

expression of *DCL4* and *AGO*1. In both of our silencing lines, AtSAP5-RNAi-3 and -7

showed noticeable effect on the expression of *DCL4* and *AGO1* (Fig. 10), but that only

one overexpression line, AtSAP5-oe-11 (expression increased about 7 times over

AtSAP5) slightly increased the expression of *DCL4* (40% increase) and *AGO1* (20%

increase). This suggests that AtSAP5 is still important for the maintenance of *DCL4*

and *AGO1* expression.

Collectively, our data suggests that although Pha13 and Pha21 both participate in

SA regulated immunity, they act differently in triggering downstream immune

responsive genes.

Pha13 and Pha21 showed no transcriptional activation activity by Y2H assay

Although Pha21 and Pha13 are involved in expression of *PhaGRXC9*, *PhaRdR1*,

PhaDCL4 and *PhaAGO1*, our Y2H analysis showed that Pha21 and Pha13 have no

transcriptional activation activity (Fig. 5B).

409   **A20 confers most E3 ligase and ubiquitin binding activity but the AN1 domain**

410   **also plays important roles in expression of immune responsive genes.**

411   Similar to other analyzed A20/AN1 proteins, our biochemical assay indicated that

412   both Pha21 and Pha13 confer E3 ligase activity and ubiquitin binding ability, and A20

413   domain of Pha21 and Pha13 is the major domain exhibiting self-E3 ligase and ubiquitin

414   binding activity (Fig. 6) (44). It has been well demonstrated that human A20 proteins

415   (without an AN1 domain) regulate the master immune transcription factor NF-kB

416   through ubiquitin editing activity including E3 ligase and ubiquitin binding ability on

417   different substrates (53). It is likely that Pha21 and Pha13 may function in a similar

418   manner to human A20 indirectly, rather than directly function as transcription factors.

419   Our previous domain functional analysis revealed that both the A20 and AN1

420   domains of Pha13 are required for expression of *PhaRdR1*, *PhaGRXC9*, and virus

421   accumulation, and the AN1 domain of Pha13 is involved in the expression of *PhaNPR1*

422   (44). Here we also showed that A20 and AN1 domains of Pha21 are both required for

423   expression of *PhaGRXC9* and CymMV accumulation, and the AN1 domain plays a

424   more important role in the expression of *PhaDCL4* (Fig. 11). This indicates that

425   although A20 confers most E3 ligase and ubiquitin binding activity, the AN1 domain

426   also plays important roles in expression of immune responsive genes. Interestingly, our

427   data suggests that neither domain of Pha21 is required for the expression of *PhaAGO1*,

428   which suggests that protein regions other than A20 and AN1 domain in Pha21 are also

429   involved in the expression of *PhaAGO1* (Fig. 11A).

430   **A20/AN1 proteins serve as important modulators in plant antiviral immunity**

Our previous data and the findings presented here indicate that A20/AN1 protein-

mediated antiviral immunity is conserved among plants, and A20/AN1 proteins may

work alone (AtSAP5) or in a cooperative manner (Pha13 and Pha21) in antiviral

immunity to induce the expression of the SA-mediated immune responsive genes,

including *NPR1*, NPR1-independent oxidoreductases gene (*GRXC9*), and genes

involved in the RNAi pathway *(RdR1*, *AGO1* and *DCL4*) (Fig. 12).

MATERIALS AND METHODS.

Plant materials and growth conditions

Orchid variety, *Phalaenopsis aphrodite var. Formosa*, was purchased from

Taiwan Sugar Research Institute (Tainan, Taiwan). All orchid plants we used including

P. aphrodite, P. equestris and transgenic *P. equestris* (35S::FLAG-Pha21) were first

analyzed for the infection with two prevalent orchid viruses, *Odontoglossum ringspot*

virus (ORSV) and CymMV, as detected by RT-PCR with primer pairs, ORSV-

F/ORSV-R and CymMV-F/CymMV-R (Table 1). Plants free from ORSV and CymMV

were maintained in greenhouse conditions with a controlled 12-h photoperiod (200

μmol m^{-2}s^{-2}) at 25°C/25°C (day/night). The wild-type (WT) *Arabidopsis* WT (Col-0)

and all transgenic *Arabidopsis* were maintained in a greenhouse with a controlled 12-h

photoperiod (200 μmol m-2s-2) at 22°C/22°C (day/night) for three to four weeks before

analysis. *Cucumber mosaic virus* (CMV) isolate 20 was maintained in the *Arabidopsis*

(Col-0) as inoculation source in our study.

Sequence analysis

The amino acid sequence of Pha21 was analyzed by the PROSITE database,

ExPASy Proteomics Server (http://ca.expasy.org/), and Conserved Domain Database

of NCBI database (http://www.ncbi.nlm.nih.gov/).

Phytohormone treatment

Sodium salicylate (50 mM) (Sigma, St. Louis, MO, USA), methyl jasmonate (45

μM) (Sigma), and aminocyclopropanecarboxylic acid (660 μM) (Sigma) were directly

rubbed on leaves of *P. aphrodite* by cotton swab. Leaf samples were collected at 0 h, 1

h, 3 h, 6 h, 12 h, 24 h, 48 h and 72 h after treatment.

RNA isolation and real-time quantitative RT-PCR (qRT-PCR) detection

Total RNA was extracted using the TOOLSmart RNA Extractor (BIOTOOLS,

Taiwan) as described previously (44). The cDNA template for qPCR was synthesized

from 500 ng of DNA-free RNA and oligo (dT) using PrimeScript RT Reagent Kit

(Perfect Real Time), following the manufacturer's instructions (Takara Bio, Shiga,

Japan). qPCR was performed using SYBR Green PCR Master Mix (Applied

Biosystems, Foster, CA, USA)) with ABI Prism 7500 sequence detection system

(Applied Biosystems). Target gene PCR products were sequenced to validate the

correct analysis of gene targets. *PhaUbiquitin 10* or *AtActin* was used as an internal

quantification control. Quantification of target gene expression was calculated

according to the manufacturer's instructions (Applied Biosystems). The primer pairs

used in this study are listed in Table 1.

Droplet digital PCR

Total RNA was extracted using the TOOLSmart RNA Extractor (BIOTOOLS,

Taiwan) as described previously (44). For droplet digital PCR (ddPCR), DNA-free

RNA (1 μg) and oligodT primer were used for cDNA synthesis with PrimeScript RT

Reagent Kit (Perfect Real Time) following the manufacturer's instructions (Takara

Bio). An amount of 5 μl of 25X dilute cDNA was used as a template for ddPCR reaction

(total 20 μl) following the manufacturer's instructions (Bio-Rad, Hercules, CA, USA).

The sequence of FAM-labeled Taqman probes and primer pairs of Pha13 and Pha21

for ddPCR are listed in Table 1.

Construction of transient silencing vector

For construction of transient silencing vector of Pha21, the oligonucleotide pairs

Pha21-hpRNA-1-F/Pha21-hpRNA-1-R   and   Pha21-hpRNA-2-F/Pha21-hpRNA-2-R

(Table 1) were used to amplify the hairpin dsDNA fragments. The hairpin dsDNA

fragments were cloned into the Gateway entry vector pENTR/D-TOPO (Thermo

Fisher-Scientific, Waltham, MA, USA) following the manufacturer's instructions to

generate pENTR-Pha21-hpRNA-1 and pENTR-Pha21-hpRNA-2. Then, LR Gateway

cloning reaction (Thermo Fisher-Scientific) was conducted to transfer the hairpin RNA

fragments from pENTR-Pha21-hpRNA into 35S promoter driven pB7GWIWG2(I) (57)

to obtain phpPha21-1 and phpPha21-2. The method for construction of the transient

silencing vector of *PhaNPR1, PhaDCL4, and PhaAGO1* (phpPhaNPR1, phpPhaDCL4,

and phpPhaAGO1) were similar to that described above, except the oligonucleotide

pairs PhaNPR1-hpRNA-F/PhaNPR1-hpRNA-R, PhaDCL4-hpRNA-F/PhaDCL4-

hpRNA-R, and PhaAGO1-hpRNA-F/ PhaAGO1-hpRNA-R (Table 1) were used to

generate the hairpin dsDNA fragments.

Construction of transient overexpression vector of Pha21, Pha21 A20 and/or AN1

domain mutant.

To construct Pha21 transient overexpression vector, total *P. aphrodite* RNA was

used as a template to amplify the N-terminus FLAG-tagged Pha21 by RT-PCR with the

primer pairs FLAG-Pha21ORF-F/Pha21ORF-R (Table 1). The amplified fragment was

cloned into Gateway entry vector pENTR/D-TOPO (Invitrogen) to generate pENTR-

FLAG-Pha21 following the manufacturer's protocol. Subsequently, LR Gateway

cloning reaction (Invitrogen) was performed to transfer the FLAG-Pha21 fragment

from pENTR-FLAG-Pha21 into the 35S promoter driven pK2GW7 (57), designated

pPha21-oe. For generation of A20 and/or AN1 mutant on pPha21-oe (Fig. 1), site-

directed mutagenesis was conducted by QuikChange Site-Directed Mutagenesis Kit

(Agilent Technologies, Santa Clara, CA, USA) and pPha21-oe was used as a template.

For A20 mutant, we substituted the conserved 3rd and 4th cysteine to glycine at A20

(C35G and C38G). For AN1 mutant, we substituted the conserved 3rd cysteine and 1st

histidine to glycine at AN1 (C113G and H123G). The A20 mutated clones, AN1

mutated clones, and the A20 and AN1 mutated clones, were designed as pPha21A20m,

pPha21AN1m, and pPha21A20mAN1m, respectively. Primer pairs used for site

directed mutagenesis are listed in Table 1.

Transgenic *Phalaenopsis* orchid

For construction of overexpression vector to generate transgenic *Phalaenopsis*

orchid, the FLAG-Pha21 fragment was transferred from pENTR-FLAG-Pha21

(described above) into 35S promoter driven binary vector, pH2GW7 (57), to obtain

pHPha21. pHPha21 was used to generate transgenic *P. equestris* orchid using the

method described by Hsing et al. (58).

Agroinfiltration

Agroinfiltration was conducted as previously described by (51) with some

modification. Briefly, *A. tumefaciens* C58C1 (pTiB6S3ΔT)^H competent cells were

transformed with pCambia-CymMV, pB7GWIWG2, pK2GW7 and their derivatives

using an electroporation system (Bio-Rad Laboratories, Hercules, CA, USA). Then, the

A. tumefaciens strains were incubated at 28°C until the optical density, OD_{600}, reached

0.8–1.0. After centrifugation, cells were resuspended in 20 ml AB-MES medium (17.2

mM K_2HPO_4, 8.3 mM NaH_2PO_4, 18.7 mM NH_4Cl, 2 mM KCl, 1.25 mM $MgSO_4$, 100

μM $CaCl_2$, 10 μM $FeSO_4$, 50 mM MES, 2% glucose (w/v), pH 5.5) with 200 μm

acetosyringone (59), and cultured overnight. The overnight culture was centrifuged

(3000 rpm, 10 min, in room temperature), supernatant was removed and the *A.*

tumefaciens culture was resuspended in 2 ml of infiltration medium containing 50%

MS medium (1/2 MS salt supplemented with 0.5% sucrose (w/v), pH 5.5), 50% AB-

MES and 200 μm acetosyringone (59). The infiltration medium containing the

transformed *A. tumefaciens* was applied for infiltration.

***Cymbidium mosaic virus* (CymMV) and *Cucumber mosaic virus* (CMV) inoculation**

and accumulation assay

To assay the effect of Pha21, PhaDCL4, or PhaAGO1 in CymMV accumulation,

we first inoculated CymMV in *P. aphrodite* through the infiltration of *A. tumefaciens*

carrying pCambia-CymMV (described above) in the leaf tip of *P. aphrodite*. The

CymMV-infected *P. aphrodite* was maintained at least 14 days before further analysis.

To assay the effect of transient silencing (Pha21, PhaDCL4, or PhaAGO1) or transient

overexpression (Pha21, or the derived mutants), *A. tumefaciens* carrying the control

vector pB7GWIWG2 (for silencing), pK2GW7 (for overexpression), silencing vectors

or overexpression vectors (described above) were infiltrated into the leaves. After

agroinfiltration, a pair of disks (6 mm diameter) were immediately (defined as 0 dpi)

collected from both the control and assay vector infiltrated regions. After 5 dpi, another

pair of disks were collected from the same infiltrated region. Total RNA extracted from

the samples was used as a template to analyze the accumulation of CymMV by use of

qRT-PCR. The fold change of CymMV accumulation at 0 dpi to 5 dpi was calculated

for relative quantification.

For inoculation with CMV, CMV-infected *Arabidopsis* leaves were ground with

0.01 M potassium phosphate buffer by pestle and mortar for use as the inoculation

source. Four-week-old *Arabidopsis* leaves were inoculated mechanically (pre-dusted

with 300-mesh Carborundum) with the CMV inoculation source. After 14 dpi, the

disease symptoms were observed, and three disks from three different distal leaves were

collected for CMV accumulation analysis by use of qRT-PCR. Primers used for CMV

detection are listed in Table 1.

Preparation and transfection of protoplasts

For construction of vectors used for subcellular localization analysis, the primer

pairs, Pha21-ORF-F/Pha21-ORF-R and Pha21-ORF-F/Pha21-ORF-non-stop-R (Table

1) were used to amplify 2 sets of Pha21 ORF (with or without a stop codon). The

amplified fragment was cloned into Gateway entry vector pENTR/D-TOPO (Invitrogen)

to generate pENTR-Pha21 and pENTR-Pha21-non-stop following the manufacturer's

protocol. Subsequently, LR Gateway cloning reaction (Invitrogen) was performed to

transfer ORF fragment of *Pha21* from pENTR-Pha21 into p2FGW7 driven by 35S

promoter (57) to obtain N-terminal GFP fused clones (pG-Pha21). To obtain C-terminal

GFP-fused clones (pPha21-G), we transferred and pENTR-Pha21-non-stop into

p2GWF7.

Protoplast isolation and transfection were as described by Lu et al. (51).

Transformed protoplasts were detected for florescence signals by confocal microscopy

(Zeiss LSM 780, plus ELYRA S.1) with excitation at 488 nm and emission at 500 to

587 nm for GFP, and excitation at 543 nm and emission at 600 to 630 nm for mCherry.

Transcriptional activation ability assay

The Pha13 and Pha21 ORF was amplified by PCR with gene specific primer pairs,

NdeI-Pha13-F/EcoR1-Pha13-R and NdeI-Pha21-F/BamH1-Pha21-R, respectively

(Table 1). The amplified ORF fragment was then cloned into the pGBKT7 vector

(Takara Bio) to generate the pGBKT7-Pha13 and pGBKT7-Pha21 constructs, which

fused to Gal4 BD sequence. These constructs were individually transformed into

AH109 yeast strain (Takara Bio) and the self-activation ability was analyzed. The

transformation and selection procedure was performed following the Yeast Protocols

HandBook (Takara Bio). Transformants were selected on SD/-Trp (tryptophan)

medium. The expression of reporter genes in yeast were tested on different selection

media; SD/-Trp/Aureobasidin was used to test for activation of the AUR1-C (inositol

phosphoryl ceramide synthase) reporter against Aureobasidin A, SD/-Trp/X-α-Gal

medium was used to test for activation of α-galactosidase, and SD/-Trp/-Ade/-His

medium was used to test for induction of the ADE2 and HIS3 reporters. The yeast

containing Gal4 DNA-BD fused with murine p53 (pGBKT7-53) and the Gal4 AD fused

with SV40 large T-antigen (pGADT7-T) from the Matchmaker Gold Yeast Two-

Hybrid System kit (Takara Bio) were used as positive control. The yeast containing

Gal4 DNA-BD fused with lamin (pGBKT7-Lam) and the pGADT7-T from the

Matchmaker Gold Yeast Two-Hybrid System kit (Takara Bio) were used as a negative

control.

603   Construction, expression, and purification of recombinant proteins

Full-length Pha21, or Pha21 A20 and/or AN1 domain mutant fragments were

amplified by PCR with primer pairs NdeI-Pha21-F/NotI-Pha21-R (Table 1) using

pPha21-oe, pPha21A20m, pPha21AN1m, and pPha21A20mAN1m as templates. The

amplified fragments were cloned into the pET24b expression vector (Merck, Darmstadt,

Germany) to produce fused C-terminal histidine tag (His-tag) expression plasmids,

pETPha21, pETPha21A20m, pETPha21AN1m and pETPha21A20mAN1m. The

constructed plasmids were transformed into *Escherichia coli* strain BL21 (DE3) for

protein expression. Bacteria were cultured at 37°C to an OD600 of 0.5, then transferred

to 25°C, for 1.5 h. Then, isopropylthio-β-galactoside (IPTG; Sigma) was added to a

final concentration of 1 mM for protein induction. His-tagged recombinant protein was

purified by TALON Superflow (GE Healthcare Life Sciences, Pittsburgh, PA, USA)

according to the manufacturer's description. His-tagged recombinant protein was

eluded with 250 mM imidazole (Sigma).

618   E3 ubiquitin ligase activity assay

In vitro ubiquitination assays were performed as described by (45) with

modification. An amount of 3 μg purified His-tagged recombinant proteins (*Pha21* or

derived mutants) was used for each ubiquitination reaction. Reactions were incubated

at 30°C for 3 hours and analyzed by SDS-PAGE followed by immunoblot analysis.

Blots were probed using anti-FLAG antibodies (Sigma) followed by HRP conjugated

goat anti-mouse antibodies (GE Healthcare Life Sciences).

626   Yeast two-hybrid assay

For the ubiquitin binding ability assay, the pGBKT7-Pha21 (described above)

was used as a bait vector. To map the region of Pha21 for ubiquitin binding, a deletion

mutant of Pha21 was generated by PCR amplification with the primer pairs described

in Table 1. The PCR-amplified fragment was individually cloned to pGBKT7 and used

as a bait vector (Fig. 6B). Full length of orchid ubiquitin was amplified by RT-PCR

with the primer pair, NdeI-PhaUBQ-F/BamHI-PhaUBQ-R, and cloned into pGADT7

vector (Takara Bio) to generate pGADT7-UBQ as a prey vector. pGBKT7-Pha21 and

pGADT7-UBQ was co-transformed into AH109 yeast strain for yeast two-hybrid assay

by using the Make Your Own "Mate & Plate" Library System (Takara Bio) following

the user manual. Yeast strains containing the appropriate bait and prey plasmids were

cultured in liquid 2-dropout medium (leucine$^-$ and tryptophan$^-$) overnight. The

overnight yeast culture was diluted to an OD600 of 0.06 and spotted on selection plates

(containing histidine$^-$, leucine$^-$, tryptophan$^-$ for growth assay.

For the interaction analysis between Pha21 and Pha13, full-length Pha13

fragment was amplified by PCR with primer pair NdeI-Pha13-F/EcoR1-Pha13-R

(Table 1). The Pha13 fragments were cloned into pGBKT7 and pGADT7 vector to

generate pGBKT7-Pha13 as the bait vector and pGADT7-Pha13 as the prey vector. The

full-length of Pha21 previously used to generate the pGBKT7-Pha21 (described above)

was also cloned into pGADT7 to generate pGADT7-Pha21 as prey vector. The

pGBKT7-Pha21, pGBKT7-Pha13, pGADT7-Pha21, and pGADT7-Pha13 were co-

transformed into AH109 yeast strain for yeast two-hybrid assay as described above.

The pGBKT7-53 and pGADT7-T from the Matchmaker Gold Yeast Two-Hybrid

System kit were used as a positive control.

Accession numbers

Pha21 (PATC144963)，Pha13 (PATC148746)，PhaPR1 (PATC126136)，PhaNPR1

(PATC135791)，PhadR1 (PATC143146)，PhaRdR2 (PATC124544)，PhaRdR6

(PATC131836)，PhaDCL2 (PATC143544)，PhaDCL4 (PATC150652)，PhaAGO1

(PATC157237)，PhaAGO10 (PATC093469)，PhaGRXC9 (PATC068819)，

PhaUBQ10 (PATC230548)，PhaJAZ1 (PATC141437)，PhaACO2 (PATC139319)，

AtActin (At3G18780)，AtSAP5 (AT3G12630)，DCL4 (AT5G20320)，AGO1

(AT1G48410)，OsSAP3 (LOC_Os01g56040.1)，OsSAP5 (LOC_Os02g32840.1)

ACKNOWLEDGMENTS.

We thank Shu-Chen Shen from the Confocal Microscope Core Facility at Academia

Sinica for assistance in confocal microscopy images and Chii-Gong Tong from the

Gene Transgenic Room at Academia Sinica for orchid transformation. This work was

supported by grants from Academia Sinica, Taipei, Taiwan and the Ministry of Science

and Technology (105-2313-B-001-004-MY3) of Taiwan. The funders had no role in

study design, data collection and interpretation, or the decision to submit the work for

publication

References

1. Pieterse CM, Leon-Reyes A, Van der Ent S, Van Wees SC. 2009. Networking by small-molecule hormones in plant immunity. Nat Chem Biol 5:308-16.

2. White R. 1979. Acetylsalicylic acid (aspirin) induces resistance to tobacco mosaic virus in tobacco. Virology 99:410-412.

3. Chivasa S, Murphy AM, Naylor M, Carr JP. 1997. Salicylic acid interferes with tobacco mosaic virus replication via a novel salicylhydroxamic acid-sensitive mechanism. The Plant Cell 9:547-557.

4. Fu ZQ, Dong X. 2013. Systemic acquired resistance: turning local infection into global defense. Annual review of plant biology 64:839-863.

5. Ausubel FM. 2005. Are innate immune signaling pathways in plants and animals conserved? Nature Immunology 6:973-979.

6. Boller T, Felix G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. Annual review of plant biology 60:379-406.

7. Jones JD, Dangl JL. 2006. The plant immune system. Nature 444:323-329.

8. Yang H, Gou X, He K, Xi D, Du J, Lin H, Li J. 2010. BAK1 and BKK1 in Arabidopsis thaliana confer reduced susceptibility to turnip crinkle virus. European journal of plant pathology 127:149-156.

9. Kørner CJ, Klauser D, Niehl A, Domínguez-Ferreras A, Chinchilla D, Boller T, Heinlein M, Hann DR. 2013. The immunity regulator BAK1 contributes to resistance against diverse RNA viruses. Molecular plant-microbe interactions 26:1271-1280.

10. Niehl A, Wyrsch I, Boller T, Heinlein M. 2016. Double-stranded RNAs induce a pattern-triggered immune signaling pathway in plants. New Phytologist 211:1008-1019.

11. Holmes FOJBG. 1929. Local lesions in tobacco mosaic.　87:39-55.

12. HoLMEs FOJP. 1937. Inheritance of resistance to tobacco-mosaic disease in the pepper (Capsicum sp.).　27:637-642.

13. Holmes FOJP. 1938. Inheritance of resistance to tobacco-mosaic disease in tobacco.　28.

14. Holmes FO. 1954. Inheritance of resistance to viral diseases in plants, p 1-30, Advances in virus research, vol 2. Elsevier.

15. Tsuda K, Katagiri FJCoipb. 2010. Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity.　13:459-465.

16. Ross AFJV. 1961. Localized acquired resistance to plant virus infection in hypersensitive hosts.　14:329-339.

17. Ross AFJV. 1961. Systemic acquired resistance induced by localized virus

infections in plants.   14:340-358.

18.   Ding S-W, Voinnet O. 2007. Antiviral immunity directed by small RNAs. Cell
130:413-426.

19.   Zhang C, Wu Z, Li Y, Wu J. 2015. Biogenesis, function, and applications of virus-
derived small RNAs in plants. Frontiers in microbiology 6:1237.

20.   Sharma N, Sahu PP, Puranik S, Prasad MJMb. 2013. Recent advances in plant–
virus interaction with emphasis on small interfering RNAs (siRNAs).   55:63-77.

21.   Blevins T, Rajeswaran R, Shivaprasad PV, Beknazariants D, Si-Ammour A, Park
H-S, Vazquez F, Robertson D, Meins Jr F, Hohn TJNar. 2006. Four plant Dicers
mediate viral small RNA biogenesis and DNA virus induced silencing.
34:6233-6246.

22.   Deleris A, Gallego-Bartolome J, Bao J, Kasschau KD, Carrington JC, Voinnet OJS.
2006. Hierarchical action and inhibition of plant Dicer-like proteins in antiviral
defense.   313:68-71.

23.   Vaucheret HJTips. 2008. Plant argonautes.   13:350-358.

24.   Carbonell A, Carrington JCJCoipb. 2015. Antiviral roles of plant ARGONAUTES.
27:111-117.

25.   Yu D, Fan B, MacFarlane SA, Chen Z. 2003. Analysis of the involvement of an
inducible Arabidopsis RNA-dependent RNA polymerase in antiviral defense.
Molecular Plant-Microbe Interactions 16:206-216.

26.   Yang S-J, Carter SA, Cole AB, Cheng N-H, Nelson RS. 2004. A natural variant of
a host RNA-dependent RNA polymerase is associated with increased
susceptibility to viruses by Nicotiana benthamiana. Proceedings of the National
Academy of Sciences of the United States of America 101:6297-6302.

27.   Liu Y, Gao Q, Wu B, Ai T, Guo X. 2009. NgRDR1, an RNA-dependent RNA
polymerase isolated from Nicotiana glutinosa, was involved in biotic and
abiotic stresses. Plant Physiology and Biochemistry 47:359-368.

28.   Li Y, Muhammad T, Wang Y, Zhang D, Crabbe MJC, Liang YJPJOB. 2018. Salicylic
acid collaborates with gene silencing to tomato defense against tomato yellow
leaf curl virus (TYLCV).

29.   Campos L, Granell P, Tárraga S, López-Gresa P, Conejero V, Bellés JM, Rodrigo I,
Lisón PJPp, biochemistry. 2014. Salicylic acid and gentisic acid induce RNA
silencing-related genes and plant resistance to RNA pathogens.   77:35-43.

30.   Alamillo JM, Saénz P, García JAJTPJ. 2006. Salicylic acid-mediated and RNA-
silencing defense mechanisms cooperate in the restriction of systemic spread
of plum pox virus in tobacco.   48:217-227.

31.   Xie Z, Fan B, Chen C, Chen Z. 2001. An important role of an inducible RNA-
dependent RNA polymerase in plant antiviral defense. Proceedings of the

749   National Academy of Sciences 98:6516-6521.

750   32.   Hunter LJR, Westwood JH, Heath G, Macaulay K, Smith AG, MacFarlane SA,
751   Palukaitis P, Carr JP. 2013. Regulation of RNA-dependent RNA polymerase 1 and
752   isochorismate synthase gene expression in Arabidopsis. PLoS ONE 8:e66530.

753   33.   Zhou M, Wang W, Karapetyan S, Mwimba M, Marqués J, Buchler NE, Dong XJN.
754   2015. Redox rhythm reinforces the circadian clock to gate immune response.
755   523:472.

756   34.   Mou Z, Fan W, Dong X. 2003. Inducers of plant systemic acquired resistance
757   regulate NPR1 function through redox changes. Cell 113:935-944.

758   35.   Tada Y, Spoel SH, Pajerowska-Mukhtar K, Mou Z, Song J, Wang C, Zuo J, Dong X.
759   2008. Plant immunity requires conformational charges of NPR1 via S-
760   nitrosylation and thioredoxins. Science 321:952-956.

761   36.   Herrera-Vásquez A, Salinas P, Holuigue LJFips. 2015. Salicylic acid and reactive
762   oxygen species interplay in the transcriptional control of defense genes
763   expression.   6:171.

764   37.   Cao H, Bowling SA, Gordon AS, Dong X. 1994. Characterization of an
765   Arabidopsis mutant that is nonresponsive to inducers of systemic acquired
766   resistance. The Plant Cell 6:1583-1592.

767   38.   Delaney T, Friedrich L, Ryals J. 1995. Arabidopsis signal transduction mutant
768   defective in chemically and biologically induced disease resistance.
769   Proceedings of the National Academy of Sciences 92:6602-6606.

770   39.   Shah J, Tsui F, Klessig DF. 1997. Characterization of a s alicylic a cid-i nsensitive
771   mutant (sai1) of Arabidopsis thaliana, identified in a selective screen utilizing
772   the SA-inducible expression of the tms2 gene. Molecular Plant-Microbe
773   Interactions 10:69-78.

774   40.   Wong CE, Carson RA, Carr JP. 2002. Chemically induced virus resistance in
775   Arabidopsis thaliana is independent of pathogenesis-related protein
776   expression and the NPR1 gene. Molecular plant-microbe interactions 15:75-81.

777   41.   Kachroo P, Yoshioka K, Shah J, Dooner HK, Klessig DF. 2000. Resistance to turnip
778   crinkle virus in Arabidopsis is regulated by two host genes and is salicylic acid
779   dependent but NPR1, ethylene, and jasmonate independent. The Plant Cell
780   12:677-690.

781   42.   Vij S, Tyagi AK. 2008. A20/AN1 zinc-finger domain-containing proteins in plants
782   and animals represent common elements in stress response. Functional &
783   integrative genomics 8:301-307.

784   43.   Giri J, Dansana PK, Kothari KS, Sharma G, Vij S, Tyagi AK. 2013. SAPs as novel
785   regulators of abiotic stress response in plants. Bioessays 35:639-648.

44.   Chang L, Chang H-H, Chang J-C, Lu H-C, Wang T-T, Hsu D-W, Tzean Y, Cheng A-P,

Chiu Y-S, Yeh H-HJPp. 2018. Plant A20/AN1 protein serves as the important hub
to mediate antiviral immunity.   14:e1007288.

45. Kang M, Fokar M, Abdelmageed H, Allen RD. 2011. Arabidopsis SAP5 functions
as a positive regulator of stress responses and exhibits E3 ubiquitin ligase
activity. Plant molecular biology 75:451-466.

46. Kang M, Lee S, Abdelmageed H, Reichert A, Lee HK, Fokar M, Mysore KS, Allen
RD. 2017. Arabidopsis stress associated protein 9 mediates biotic and abiotic
stress responsive ABA signaling via the proteasome pathway. Plant, cell &
environment 40:702-716.

47. Choi H, Han S, Shin D, Lee S. 2012. Polyubiquitin recognition by AtSAP5, an A20-
type zinc finger containing protein from Arabidopsis thaliana. Biochemical and
biophysical research communications 419:436-440.

48. Verhelst K, Carpentier I, Kreike M, Meloni L, Verstrepen L, Kensche T, Dikic I,
Beyaert R. 2012. A20 inhibits LUBAC-mediated NF-κB activation by binding
linear polyubiquitin chains via its zinc finger 7. The EMBO Journal 31:3845-3855.

49. Wertz IE, O'rourke KM, Zhou H, Eby M. 2004. De-ubiquitination and ubiquitin
ligase domains of A20 downregulate NF-kappaB signalling. Nature 430:694.

50. Mattera R, Tsai YC, Weissman AM, Bonifacino JS. 2006. The Rab5 guanine
nucleotide exchange factor Rabex-5 binds ubiquitin (Ub) and functions as a Ub
ligase through an atypical Ub interacting motif and a zinc finger domain. The
Journal of Biological Chemistry 281:6874-6883.

51. Lu HC, Hsieh MH, Chen CE, Chen HH, Wang HI, Yeh HH. 2012. A high-throughput
virus-induced gene-silencing vector for screening transcription factors in virus-
induced plant defense response in orchid. Molecular Plant-Microbe
Interactions 25:738-746.

52. Chang C-H, Wang H-I, Lu H-C, Chen C-E, Chen H-H, Yeh H-H, Tang CY. 2012. An
efficient RNA interference screening strategy for gene functional analysis. BMC
genomics 13:491.

53. Ma A, Malynn BA. 2012. A20: linking a complex regulator of ubiquitylation to
immunity and human disease. Nature reviews Immunology 12:774.

54. Hishiya A, Iemura Si, Natsume T, Takayama S, Ikeda K, Watanabe K. 2006. A
novel ubiquitin-binding protein ZNF216 functioning in muscle atrophy. The
EMBO Journal 25:554-564.

55. Miyata N, Okumoto K, Mukai S, Noguchi M, Fujiki Y. 2012. AWP1/ZFAND6
Functions in Pex5 Export by Interacting with Cys-Monoubiquitinated Pex5 and
Pex6 AAA ATPase. Traffic 13:168-183.

56. Giri J, Vij S, Dansana PK, Tyagi AK. 2011. Rice A20/AN1 zinc-finger containing
stress-associated proteins (SAP1/11) and a receptor-like cytoplasmic kinase

(OsRLCK253) interact via A20 zinc-finger and confer abiotic stress tolerance in

transgenic Arabidopsis plants. New Phytologist 191:721-732.

57.   Karimi M, Inzé D, Depicker A. 2002. GATEWAY™ vectors for Agrobacterium-

mediated plant transformation. Trends in plant science 7:193-195.

58.   Hsing H-X, Lin Y-J, Tong C-G, Li M-J, Chen Y-J, Ko S-S. 2016. Efficient and heritable

transformation of Phalaenopsis orchids. Botanical studies 57:30.

59.   Wu H-Y, Liu K-H, Wang Y-C, Wu J-F, Chiu W-L, Chen C-Y, Wu S-H, Sheen J, Lai E-

M. 2014. AGROBEST: an efficient Agrobacterium-mediated transient

expression method for versatile gene function analyses in Arabidopsis

seedlings. Plant methods 10:19.

FIGURE LEGENDS:

FIG 1 Sequence and expression analysis of Pha21 (A) A schematic representation of domain organization of Pha21. The open rectangle indicates the entire protein. A20 (dark grey rectangle) and AN1 (light grey rectangle) and 21-nucleotide position (short black line) used for designing hairpin RNA of Pha21 (hpPha21-1 and hpPha21-2) are indicated. (B) Amino sequence alignment of A20/AN1 zinc finger domains of Pha21 with stress-associated proteins from *Phalaenopsis aphrodite* (Pha13, accession number: PATC148746), *Arabidopsis thaliana* (AtSAP5, accession number: AT3G12630), and *Oryza sativa* (OsSAP3, accession number: LOC_Os01g56040.1; OsSAP5, accession number: LOC_Os02g32840.1) is shown. The black box indicates the conserved amino acid sequences. The black triangle indicates the conserved cysteine (C) and histidine (H) residues. (C) Primary sequence organization of A20 and AN1 domains. Xn: the number of amino acid residues between zinc ligands. (B-C) The mutation positions for domain functional analysis are indicated with a blue triangle. (D) Expression level of *Pha21* was analyzed by qRT-PCR from the root, leaf, petal, sepal, column, and lip of *P. aphrodite*. The RNA level of the root was set to 1. Data represent mean $\pm$ SD; n = 3 technical replicates; different letters indicate statistically significant differences analyzed by one-way analysis of variance (ANOVA) Tukey's test ($P < 0.05$). The experiment was repeated at least three times with similar results, and one representative

experiment is shown. *PhaUbiquitin 10* was used as an internal control for normalization.

(E) Absolute quantitative analysis of *Pha13* and *Pha21* expression. The mRNA copy

numbers of Pha13 and Pha21 in leaves of *P. aphrodite* were absolutely quantified by

droplet digital PCR. Total RNA (ng) was used for normalization. Data represent mean

± SD; n = 3 biological replicates; **, $P < 0.01$, Student's t test compared Pha13 with

Pha21

FIG 2 Pha21 is involved in the expression of *PhaPR1* and *PhaNPR1,* and responds to

phytohormone treatment. (A-B) qRT-PCR was performed to quantify the expression

level of *Pha21*, *PhaNPR1*, and *PhaPR1* from leaves of *P. aphrodite* infiltrated with

agrobacterium carrying control vector (Vector); hairpin RNA vector to knock down

Pha21 (hpPha21-1 and hpPha21-2; A); and overexpression vector of Pha21 (Pha21-oe;

B). The expression level of plants infiltrated with control vector was set to 1. (C)

Transient silencing of PhaNPR1. qRT-PCR was performed to quantify the expression

level of *PhaNPR1*, *PhaPR1*, and *Pha21* from leaves of *P. aphrodite* infiltrated with

agrobacterium carrying control vector (Vector) or hairpin RNA (hpRNA) vector to

knock down PhaNPR1 (hpPhaNPR1). The expression level of plants infiltrated with

control vector was set to 1. (A-C) Data represent mean ± SD; n = 3 biological replicates;

**, $P < 0.01$; *, $P < 0.05$, Student's t test compared to Vector. (D-F) Time-course

expression of *Pha21* under different plant hormone treatments in *P. aphrodite*. qRT-

PCR was performed to analyze the expression level of *Pha21* from leaves treated with

SA (D), JA (E), and ET (F) at different hours (h) post-treatment. Buffer treatment was

used as a mock control. Results of qRT-PCR were relative to that of mock at individual

time courses for relative quantification. The expression level at 0 h was set to 1 for

comparison between different time courses. *PhaPR1* and *PhaNPR1* were SA-

responsive marker genes. *PhaJAZ1* and *PhaACO2* were JA and ET–responsive marker

genes, respectively. Data represent mean ± SD; n = 3 biological replicates; **, $P < 0.01$;

*, $P < 0.05$, Student's t test compared to 0 h. (A-F) *PhaUbiquitin 10* was used as an

internal control for normalization.

FIG3 Pha21 is involved in virus accumulation. (A) qRT-PCR was performed to

quantify the CymMV accumulation level and expression level of *Pha21* in healthy

(Mock) and CymMV-infected (Cy-infected) *P. aphrodite.* The expression level in the

CymMV-infected (Cy-infected) *P. aphrodite* was set to 1. Data represent mean ± SD; n

= 3 biological replicates; **, $P < 0.01$; *, $P < 0.05$, Student's t test compared to mock.

(B-C) Transient silencing or overexpression of Pha21 in CymMV-infected plants. qRT-

PCR was performed to quantify the expression level of *Pha21* and CymMV

accumulation level from leaves of CymMV-infected *P. aphrodite* infiltrated with

agrobacterium carrying vector (Vector); hairpin RNA (hpRNA) vector to knockdown

Pha21 (hpPha21-1 and hpPha21-2; B); or overexpression vector of Pha21 (Pha21-oe;

C). The expression level of plants infiltrated with control vector was set to 1. Data

represent mean ± SD; n = 3 biological replicates; **, $P < 0.01$; *, $P < 0.05$, Student's t

test compared to Vector. (D) The wild-type (WT) *P. equestris* orchid and transgenic *P.*

equestris (Pha21#8 and #9) were used for analysis. expression level of *Pha21* and

CymMV were analyzed by qRT-PCR from leaves of WT or transgenic *P. equestris*

(Pha21#8 and #9) inoculated with CymMV. The expression level of WT was set to 1.

Data represent mean ± SD; n = 2 biological replicates; **, $P < 0.01$; *, $P < 0.05$,

Student's t-test compared to WT. (A-D) *PhaUbiquitin 10* was used as an internal

control for normalization.

FIG4 Overexpression of Pha21 in engineered transgenic *Arabidopsis* enhances

resistance against viruses. (A) Expression level of *Pha21* and accumulation level of

Cucumber mosaic virus (CMV) were analyzed by qRT-PCR from leaves of WT (Col-

0) or transgenic *Arabidopsis* (At-Pha21#4, #5, and #6). Data represent mean ± SD; n =

5 biological replicates; **, $P < 0.01$; *, $P < 0.05$, Student's t-test compared to WT.

AtActin was used as an internal control for normalization. (B) Disease symptoms of WT

(Col-0) or transgenic *Arabidopsis* (At-Pha21#4, #5, and #6) inoculated with CMV at

14 dpi. Images are one representative plant from five replicates; Scale bar, 1 cm.

FIG 5 Subcellular localization of Pha21 and transcriptional activation analysis of

Pha13 and Pha21. (A) Green fluorescent protein (GFP) or N- and C- terminal GFP-

fused Pha21 (G-Pha21 and Pha21-G) were co-transfected with nucleus localization

signal fused red fluorescence protein (NLS-RFP) into protoplasts of *P. aphrodite*.

Fluorescence was detected by confocal microscopy after transfection. Scale bars

represent 10 μm. (B) Yeast two-hybrid assay was used to analyze the transcriptional

activation. AH109 is the yeast strain used in this assay. The pGBKT7-Pha13 and

pGBKT7-Pha21 indicates that yeast encodes Pha13 or Pha21 fused to the Gal4 DNA

binding domain. The pGBKT7-53, pGADT7-T, and pGBKT7-Lam provided in

Matchmaker Gold Yeast Two-Hybrid System kit were used as a positive (pGBKT7-53

+ pGADT7-T) and negative (pGBKT7-Lam + pGADT7-T) control. The -Trp indicates

the medium without tryptophan. The -Trp/Aureobasidin was used to test for activation

of the inositol phosphoryl ceramide synthase (AUR1-C) for Aureobasidin A resistance.

The -Trp/X-α-Gal medium was used to test for activation of α-galactosidase, and -Trp/-

Ade/-His indicate the selective medium without tryptophan, adenine, and histidine

FIG 6 Pha21 confers self-ubiquitination E3 ligase activity and ubiquitin binding ability.

(A) The *in vitro* self-ubiquitination E3 ligase activity assay was conducted on wild-type

Pha21, A20 mutant (Pha21-A20m), AN1 mutant (Pha21-AN1m), A20/AN1 double

mutant (Pha21-A20mAN1m) with or without FLAG-Ub, human ubiquitin-activating

enzyme (hE1), or human ubiquitin-conjugating enzyme (hE2). Coomassie staining for

Pha21 was used as a loading control. Ubiquitinated proteins were analyzed with

immunoblotting using anti-FLAG antibodies. The ubiquitinated Pha21 are indicted as

Pha21-Ub. E1 conjugated with one Ub (E1-Ub), E2 conjugated with one or two Ub

(E2-Ub, E2-2Ub), and free Ub are indicated with a black arrow. (B) Yeast two-hybrid

assay was performed to analyze the interaction between full-length and deletion mutant

of Pha21 and ubiquitin. The schematic representations of full-length and deletion

mutant of Pha21 are indicated. The open rectangle indicates the entire Pha21 protein.

A20 domain and AN1 domain are indicated as a dark blue oval circle and a light blue

oval circle, respectively. The full-length and deletion mutant of Pha21 cloned in

pGBKT7 vector were co-transformed with pGADT7-PhaUBQ into yeast AH109 strain.

The 2DO indicates the Leucine and Trptophan dropout selective medium and 3DO

indicates the Leucine, Trptophan, and Histidine dropout selective medium. The

pGBKT7 was used as a negative control.

FIG 7 Interaction analysis of Pha21 and Pha13. Yeast two-hybrid assay was performed

to analyze the interaction between Pha21 and Pha13. The pGBKT7 and pGADT7 were

used as a negative control. The pGBKT7-53 and pGADT7-T (provided in the

Matchmaker Gold Yeast Two-Hybrid System kit), and pGBKT7-Pha21 and pGADT7-

PhaUBQ were used as a positive control. The 2DO indicates the dropout selective

medium without Leucine and Tryptophan. The 3DO indicates dropout selective

medium without the Leucine, Tryptophan, and Histidine.

FIG 8 Pha21 is involved in the expression of PhaNPR1-dependent gene, *PhaRdR1,* and

PhaNPR1-independent gene, *PhaGRXC9*. The expression levels of *Pha21, PhaRdR1*,

and *PhaGRXC9* were analyzed by qRT-PCR from leaves of *P. aphrodite* infiltrated with

agrobacterium carrying control vector (Vector) or hairpin RNA (hpRNA) vector to

knockdown Pha21 (hpPha21-2; A), or to overexpress Pha21 (Pha21-oe; B). The RNA

level of plants infiltrated with control vector was set to 1. Data represent mean ± SD; n

= 3 biological replicates; **, $P < 0.01$; *, $P < 0.05$, Student's t-test compared to Vector.

(A-B) *PhaUbiquitin 10* was used as an internal control for normalization.

FIG 9 Pha21 is involved in the expression of *PhaDCL4* and *PhaAGO1*. (A-C) The

expression levels of *Pha13*, *Pha21*, *PhaRdR2*, *PhaRdR6*, *PhaDCL2*, *PhaDCL4*,

PhaAGO1, and *PhaAGO10* were analyzed by qRT-PCR from leaves of *P. aphrodite*

infiltrated with agrobacterium carrying control vector (Vector); overexpression vector

of Pha21 (Pha21-oe; A); hairpin RNA vector to knock down Pha21 (hpPha21-2; B);

and overexpression vector of Pha13 (Pha13-oe; C). The expression level of plants

infiltrated with control vector was set to 1. Data represent mean ± SD; n = 3 biological

replicates; **, $P < 0.01$; *, $P < 0.05$, Student's t-test compared to Vector. (D-E) The

expression level of *PhaDCL4*, *PhaAGO1*, and CymMV accumulation level were

analyzed by qRT-PCR from leaves of CymMV-infected *P. aphrodite* infiltrated with

agrobacterium carrying vector (Vector); hairpin RNA (hpRNA) vector to knock down

PhaDCL4 (hpPhaDCL4, D), or PhaAGO1 (hpPhaAGO1, E). The expression level of

plants infiltrated with control vector was set to 1. Data represent mean ± SD; n = 3

biological replicates; **, $P < 0.01$; *, $P < 0.05$, Student's t test compared to Vector. (A-

E) *PhaUbiquitin 10* was used as an internal control for normalization.

FIG 10 AtSAP5 is involved in the expression of *DCL4* and *AGO1*. (A-C) The

expression levels of *AtSAP5, DCL4, AGO1* were analyzed by qRT-PCR in WT,

AtSAP5   overexpression (AtSAP5-oe-4 and AtSAP5-oe-11) and RNAi lines

(AtSAP5-RNAi-3 and AtSAP5-RNAi-7). Data represent mean ± SD; n = 5 biological

replicates; **, $P < 0.01$; *, $P < 0.05$, Student's t-test. Student's t test compared to WT.

AtActin was used as an internal control for normalization.

FIG 11 Pha21 A20 and AN1 domain play different roles in the expression of *PhaDCL4,*

PhaAGO1, PhaGRXC9 and virus resistance. The expression level of *Pha21, PhaDCL4,*

PhaAGO1, PhaGRXC9 and CymMV accumulation level were analyzed by qRT-PCR

in leaves of healthy *P. aphrodite* (A), or CymMV pre-infected *P. aphrodite* (B)

infiltrated with agrobacterium carrying control vector (Vector), overexpression clones

of Pha21 (Pha21-oe)*,* or the respective A20 and/or AN1 mutant clones (Pha21A20m,

Pha21AN1m, and Pha21A20mAN1m). The RNA level of plants infiltrated with control

vector was set to 1. Data represent mean ± SD; n = 3 biological replicates; **, $P < 0.01$;

*, $P < 0.05$; ns, no significant difference, Student's t test compared to Vector. (A-B)

PhaUbiquitin 10 was used as an internal control for normalization.

FIG 12 The role of A20/AN1 proteins in antiviral immunity.

A model illustrating the roles of SA-induced-A20/AN1 proteins Pha13 and Pha21 from

Phalaenopsis aphrodite, and AtSAP5 from *Arabidopsis thaliana* in antiviral immunity.

Virus infection caused accumulation of SA and led to the activation of *Pha13*, *Pha21*,

and *AtSAP5,* allowing it to trigger the SA-and RNAi-mediated immune responsive

genes including *NPR1*, *GRXC9*, *DCL4*, *AGO1*, and *RdR1* in antiviral immunity.

1005

1006 **Table 1.** Primers and probes used in this study

Name of primers and probes	Nucleotide sequence	Description
Semi-quantitative RT-PCR		
CymMV-F	5'-GAAATAATCATGGGAGAGCC-3'	Detection of *Cymbidium*
CymMV-R	5'-AGTTTGGCGTTATTCAGTAGG-3'	*mosaic virus* (CymMV)
ORSV-F	5'-ACGCACAATCTGATCCGTA-3'	Detection of *Odontoglossum*
ORSV-R	5'-ATCCGCAGTGAAAACCC-3'	*ringspot virus* (ORSV)
Real-time RT-PCR		
Pha13-qF	5'-TCATCTTAAAACAGAGCAGGATCTGA-3'	Detection of *Pha13*
Pha13-qR	5'-CGGCGTCTAAAGAAGAGGAAGAG-3'	(PATC148746[a])
Pha21-qF	5´-CCATCTTAAAACCCAGGATGCT-3´	Detection of *Pha21*
Pha21-qR	5´-TGGAGTCGACGAGGAACAGAA-3´	(PATC144963[a])
PhaNPR1-qF	5'-AGTTGATGGCACCTTGGAGTTT-3'	Detection of *PhaNonexpressor*
		of pathogenesis-related gene 1
PhaNPR1-qR	5'-AAGAAGCGTTTCCCGAGTTCA-3'	(*PhaNPR1*; PATC135791[a])
PhaPR1-qF	5'-GGATCATCGTCTTGCGATTT-3'	Detection of *PhaPathogenesis-*
		related gene 1 (*PhaPR1*;
PhaPR1-qR	5'-CCGCACAACTGTTACATGCAT-3'	PATC126136[a])
CymMV-qF	5'-TGATGCTGGCCACTAACGATC-3'	Detection of CymMV
CymMV-qR	5'-GGAATCAACGGCATCGAAGA-3'	
PhaRdR1-qF	5'-TGACAAGTACAAGGATGGCG-3'	Detection of *PhaRNA-*
		dependent RNA polymerase 1
PhaRdR1-qR	5'-ACTCTTCACCCCATCAACATC-3'	(*PhaRdR1*; PATC143146[a])
PhaRdR2-qF	5'-AAAATCCTTGTCTCCACCCTG-3'	Detection of *PhaRdR2*
PhaRdR2-qR	5'-GCAATCTACGAGTCCCATCTC-3'	(PATC124544[a])
PhaRdR6-qF	5'-AGAGCTGGGAATGGAAGTTG-3'	Detection of *PhaRdR6*
PhaRdR6-qR	5'-GTCGAATTTCTGCATGGTTGG-3'	(PATC131836[a])
PhaDCL2-qF	5'-CATTGCATCAGAAAGCCATCG-3'	Detection of *PhaDicer-like*
		nuclease 2 (*PhaDCL2*;
PhaDCL2-qR	5'-ACCTCAAACGTCTCCAACTTC-3'	PATC143544[a])
PhaDCL4-qF	5'-CATGACTTCGATAGGGTTGGG-3'	Detection of *PhaDCL4*
PhaDCL4-qR	5'-GAGTTTCGCCATTCTTTGCTG-3'	(PATC150652[a])
PhaAGO1-qF	5'-ACAGGAGACAAGCCAAAGAG-3'	Detection of *PhaArgonaut 1*
PhaAGO1-qR	5'-ACCACAAAAGTGACGAGAGG-3'	(*PhaAGO1*; PATC157237[a])

Table 1. Cont.

PhaAGO10-qF	5'-TGAACCAAACTACCAGCCAC-3'	Detection of *PhaAGO10*
PhaAGO10-qR	5'-TGTTTCCATTCCTGTCCGTAG-3'	(PATC093469[a])
PhaGRXC9-qF	5'-ACGTCGTAAAGCATCTCCTT-3'	Detection of *PhaGlutaredoxin*
		C9 (PhaGRXC9;
PhaGRXC9-qR	5'-CGGTAAGGTCACCGTAAAAC-3'	PATC068819[a])
PhaUBQ 10-qF	5'-CCGGATCAGCAAAGGTTGA-3'	Detection of (PhaUBQ10;
PhaUBQ 10-qR	5'-AAGATTTGCATCCCTCCCC-3'	PATC230548[a])
PhaJAZ1-qF	5'-CTCGGAGTGGTTGGCTTTATTAG-3'	Detection of *PhaJasmonate*
		ZIM-domain protein 1
PhaJAZ1-qR	5'-GTGATGCCTTCCTTGCGATT-3'	(*PhaJAZ1*; PATC141437[a])
PhaACO2-qF	5'-CGTCCAAAACGCTTGATTCC-3'	Detection of *PhaAmino-*
		cyclopropane-carboxylate
PhaACO2-qR	5'-GCGCAAATTCCTTCATGGTT-3'	*oxidase 2* (*PhaACO2*;
		PATC139319[a])
AtACT-qF	5'-GGCAAGTCATCACGATTGG -3'	Detection of *AtActin*
AtACT-qR	5'-CAGCTTCCATTCCCACAAAC -3'	(AT3G18780)
CMV-qF	5'-TTCCTGCCTCCTCGGACTTA-3'	Detection of CMV
CMV-qR	5'-GCTCCGTCCGCGAACATA-3'	
AtSAP5-qF	5'-ACCAGCTAAAGTCGTGATTCG-3'	Detection of *AtSAP5*
AtSAP5-qR	5'-AGCGGTTTTGTAGTCGTAGC-3'	
DCL4-qF	5'-GCCCCTCACTACTCGCAATA-3'	Detection of *DCL4*
DCL4-qR	5'-GAACAGCATCCCCAAGAAA-3'	
AGO1-qF	5'- GTTTTCGCATTCAGCACACTC-3'	Detection of *AGO1*
AGO1-qR	5'-CAATCTTTTGGAATACCGCTG-3'	

Droplet digital PCR

Pha13-ddPCR-P-FAM	5'-TCTCCACGGCGATCAGATCC-3'	FAM-labeled probe for
Pha21-ddPCR-P-FAM	5'-ATGCTCTGCTCGCTGTCGAA-3'	detection of *Pha13* or *Pha21*
Pha13-ddPCR-F	5'-AACCCTAAACCTCTGCTC-3'	Detection of *Pha13*
Pha13-ddPCR-R	5'-GAGGAGGAGGAGGAGAAG-3'	
Pha21-ddPCR-F	5'-CTGCTCTAAGTGTTACGG-3'	Detection of *Pha21*
Pha21-ddPCR-R	5'-TCGACGAGGAACAGAATA-3'	

Construction

Pha21-hpRNA-1-F	5'-CACCTTACGGAGACCATCTTAAAAC-3'	Construction of phpPha21-1
Pha21-hpRNA-1-R	5'-GTTTTAAGATGGTCTCCGTAA-3'	
Pha21-hpRNA-2-F	5'-CACCCCAGGATGCTCTGCTCGCTGT-3'	Construction of phpPha21-2
Pha21-hpRNA-2-R	5'-ACAGCGAGCAGAGCATCCTGG-3'	

Table 1. Cont.

PhaNPR1-hpRNA-F	5' -CACCCGAGATCATTGATTCCTTCCC-3'	Construction of phpPhaNPR1
PhaNPR1-hpRNA-R	5' -GGGAAGGAATCAATGATCTCG-3'	
PhaDCL4-hpRNA-F	5'-CACCGAGAATTTACTTGATGCAAAG-3'	Construction of phpPhaDCL4
PhaDCL4-hpRNA-R	5' -CTTTGCATCAAGTAAATTCTC-3'	
PhaAGO1-hpRNA-F	5'-CACCACATGGTGGTGTGCCACAAGG-3'	Construction of phpPhaAGO1
PhaAGO1-hpRNA-R	5'-CCTTGTGGCACACCACCATGT-3'	
Pha21-ORF-F	5'-CACCATGGCGGAGGATCAGCCATG-3'	Construction of pG-Pha21 or pPha21-G
Pha21-ORF-NONSTOP-R	5'-AATTTTGTCCAGCTTCTCCGCC-3'	
FLAG-Pha21ORF-F	5'-CACCATGGATTACAAGGATGACGACGATAAGATGGCGGAGGATCA-3'	Construction of pPha21-oe and pHPha21
Pha21ORF-R	5'-CTAAATTTTGTCCAGCTTCTCCG-3'	
Pha21A20m-F	5'-GCTACTCTCAATCTCGGCTCTAAGGGTTACGGAGACCATC-3'	Construction of pPha21A20m
Pha21A20m-R	5'-GATGGTCTCCGTAACCCTTAGAGCCGAGATTGAGAGTAGC-3'	
Pha21AN1m-F	5'- CTCCTTGGGTTCAAGGGTCGGTGCGGTGTGACGCACTGCGGAGACGGCCGGTACCCGGAGCA-3'	Construction of pPha21AN1m
Pha21AN1m-R	5'- TGCTCCGGGTACCGGCCGTCTCCGCAGTGGTCACACCGCACCGACCCTTGAACCCAAGGAG-3'	
NdeI-Pha21-F[b]	5'- ATTC<u>CATATG</u>GCGGAGGATCAGCCAT-3'	Construction of pETPha21, pETPha21A20m, pETPha21AN1m and pETPha21A20mAN1m
NotI-Pha21-R[b]	5'- ACAT<u>GCGGCCGC</u>AATTTTGTCCAGCTTCT -3'	
NdeI-Pha21-F[b]	5'- ATTC<u>CATATG</u>GCGGAGGATCAGCCAT-3'	
NdeI-Pha21-16-F[b]	5'-CTG<u>CATATG</u>GACCATCTTAAAACCCAGGATGCTC-3'	For construction of full-length and deletion mutant of Pha21 into pGBKT7 and construction of pGADT7-Pha21
NdeI-Pha21-41-F[b]	5'-CTG<u>CATATG</u>GACCATCTTAAAACCCAGGATGCTC-3'	
NdeI-Pha21-102-F[b]	5'-CTG<u>CATATG</u>TGCCGGAAGAAGGTGGG-3'	
NdeI-Pha21-137-F[b]	5'-CTG<u>CATATG</u>TCGGCCAGCCGCG-3'	
BamH1-Pha21-R[b]	5'-GAC<u>GGATCC</u>TAAATTTTGTCCAGCTTC-3'	
BamHI-Pha21-15-R[b]	5'-GAC<u>GGATCC</u>TATTCTTGGCATCTCCGTCCC-3'	
BamHI-Pha21-40-R[b]	5'-GAC<u>GGATCC</u>TATCCGTAACACTTAGAGCAGAGATTGAG-3'	

Table 1. Cont.

BamHI-Pha21-101-R[b]	5'-GAC<u>GGATCC</u>TACGCCGCGCACCG-3'	
BamHI-Pha21-136-R[b]	5'-GAC<u>GGATCC</u>TACTTGAAATCAAAGCAGCAGCC-3'	
NdeI-PhaUBQ-F[b]	5´-CTCT<u>CATATG</u>AGGGGAGGTATGCAGATTT TCG-3'	Construction of polyubiquitin into pGADT7
BamHI-PhaUBQ-R[b]	5'-CTAT<u>GGATCC</u>ACCACGAAGACGGAGCACA A-3'	
NdeI-Pha13-F[b]	5'-AGAG<u>CATATG</u>GCTGCCACCTTCTCCTCCT TTT-3'	Construction of pGBKT7-Pha13 and pGADT7-Pha13
EcoR1-Pha13-R[b]	5'-AGAG<u>GAATTC</u>AAATCTTCCGCAGCTTA-3'	

1007   [a]Accession number of Orchidstra database (http://orchidstra.abrc.sinica.edu.tw)

1008   [b]Underlined sequences indicate the sequence recognized by restriction enzyme. The random 4 nucleotides at the 5'

1009   end were added as recommended by the manufactur

1010

1011

1012

1013

FIG 1

1018   **FIG 2**

1019

1020

1021

1022   **FIG 3**

1023

1024

1025

1026 **FIG 4**

A

B

1027

1028

1029

1030

1031 **FIG 5**

A

B

1049

1050

1051

1052

1053 **FIG 6**

A

B

1054

1055

1056

1057   **FIG 7**

			2DO	3DO
pGBKT7	+	pGADT7-Pha21		
pGBKT7	+	pGADT7-Pha13		
pGBKT7-Pha21	+	pGADT7		
pGBKT7-Pha13	+	pGADT7		
pGBKT7-Pha21	+	pGADT7-Pha21		
pGBKT7-Pha21	+	pGADT7-Pha13		
pGBKT7-Pha13	+	pGADT7-Pha13		
pGBKT7-Pha13	+	pGADT7-Pha21		
pGBKT7-53	+	pGADT7-T		
pGBKT7-Pha21	+	pGADT7-PhaUBQ		

1072

1073

1074

1075

1076

1077

1078

A

B

1096

1097

1098

1099

1100

1101   **FIG 9**

1102

1103

1104 **FIG 10**

A

AtSAP5

B

DCL4

C

AGO1

1105

1106

1107

 FIG 11

A

B

1112 **FIG 12**

1113

1114

1115