Bridging evolutionary game theory and metabolic models for predicting microbial metabolic interactions

Jingyi Cai [1], Tianwei Tan[1*], Siu Hung Joshua Chan[2**],

1 National Energy R&D Center for Biorefinery, Beijing University of Chemical Technology, Beijing 100029, China

2 Department of Chemical and Biological Engineering, Colorado State University, Fort Collins, CO 80523, USA

*Tianwei Tan: twtan@mail.buct.edu.cn ** Siu Hung Joshua Chan (Lead contact): Joshua.Chan@colostate.edu,

ABSTRACT

Microbial metabolic interactions impact ecosystems, human health and biotechnological processes profoundly. However, their determination remains elusive, invoking an urgent need for predictive models that seamlessly integrate metabolic details with ecological and evolutionary principles which shape the interactions within microbial communities. Inspired by the evolutionary game theory, we formulated a bi-level optimization framework termed NECom for the prediction of Nash equilibria of microbial community metabolic models with significantly enhanced accuracy. NECom is free of a long hidden 'forced altruism' setup in previous static algorithm while allowing for 'sensing and responding' between microbial members that is missing in dynamic methods. We successfully predicted several classical games in the context of metabolic interactions that were falsely or incompletely predicted by existing methods, including prisoner's dilemma, snowdrift game and mutualism. The results provided insights into why mutualism is favorable despite seemingly costly cross-feeding metabolites, and demonstrated the potential to predict heterogeneous phenotypes among the same species. NECom was then applied to a reported algae-yeast co-culture system that shares typical cross-feeding features of lichen, a model system of mutualism. More than 1200 growth conditions were simulated, of which 488 conditions correspond to 3221 experimental data points. Without fitting any ad-hoc parameters, an overall 63.5% and 81.7% reduction in root-mean-square error in predicted growth rates for the two species respectively was achieved when compared with the standard flux balance analysis. The simulation results further show that growth-limiting crossfeeding metabolites can be pinpointed by shadow price analysis to explain the predicted frequency-dependent growth pattern, offering insights into how stabilizing microbial interactions control microbial populations.

INTRODUCTION

In nature, microorganisms seldom exist in isolation. Instead, they form communities governed by different types of interactions, which play essential roles in adaptation to environments (1-4) and evolution of species (5-9) . Among microbial interactions, exchange of metabolites is especially important and arguably primarily responsible for the fact that more than 99% of bacterial species are not cultivable (4, 10, 11). Understanding metabolic interactions is a fundamental task in microbiome sciences. Despite continual progresses in determining metabolite exchanges using experimental approach such as spatially separated apparatus designs (12, 13), isotope probing (14, 15) and tracing (16) techniques, 16s RNA (17) analysis and metabolome analysis (18), we still need governing principles to predict and understand these metabolic interactions.

Benefiting from the advancement of genome-scale metabolic models (19) and constraint based modeling techniques, modeling of microbial communities has been extended to the level of whole-cell metabolism. Since the first published work on constraint-based modeling of microbial communities more than a decade ago (20), new algorithms addressing various challenges have been proposed. Joint flux balance analysis (Joint-FBA) was first introduced to model a microbial community as a 'super' organism containing the compartments for each organism and an additional compartment for inter-cellular metabolite exchange (20, 21). The total community biomass was maximized to predict metabolism. Methods based on Joint-FBA for analyzing host-microbe and microbe-microbe interactions were widely applied to the gut microbiome (22-25). Later a dynamic simulation framework called dynamic multi-species metabolic modeling (DMMM) and other similar methods that iteratively perform FBA for each microbial species at each time step has been developed (26-28). Succeeding efforts include regularization to ensure a well-conditioned dynamical system (29) and incorporation of spatiotemporal elements (30-34). However, in most dynamic methods, the metabolic phenotype of an individual microbe is determined only by the metabolites present in the environment (by finding a particular FBA solution). Unlike Joint-FBA, there is no way for a microbe to directly 'sense and respond' to other microbes. These methods might therefore overlook some potential metabolic interactions favored by evolution. In order to reconcile the assumptions of fitness optimization for individuals vs. the entire community, OptCom (35) and its dynamic version d-OptCom (36), solve bilevel optimization problems to optimize the individual fitness in the inner problem, while optimizing the community fitness in the outer problem. To address the effect of viable abundance on the inter-cellular fluxes at population steady state, community FBA (cFBA) (37) generalized joint-FBA by adding parameterized abundances as weights to exchange fluxes. SteadyCom (38) as a reformulation of cFBA enabled efficient computation for flux variability analysis and other constraint based modeling techniques for communities with a large number of organisms. Recently, there have been efforts to simulate communities with a large number of species (39, 40) or handle a large quantity of cases (41-43). Joint-FBA and its linear derivatives are usually employed to avoid expensive computation. However, some constraints and/or the community

objective function in joint-FBA and other static FBA-based algorithms force individual organisms to produce a certain amount of metabolites for other members prior to optimizing its own fitness whenever doing so maximizes the community-level objective function value. Here we term this 'forced altruism'. We believe that forced altruism is potentially applicable to microbes that are physically connected, e.g., by nanotubes (44, 45) or to endosymbiosis, since inter-cellular exchange under these situations are likely governed by concentration gradients or regulation by the host, respectively. In general for microbes in a co-culture separate from each other by cell envelopes and not controlled by higher-level regulatory mechanisms (e.g., host-regulation, quorum sensing, etc.), however, the applicability of 'forced altruism' constraints should be questioned.

From evolutionary game theory, Nash equilibria (NE) and evolutionary stable strategies (ESSes) which are a subset of NE are more informative targets to identify for studying the community interactions (46). A microbial community is in a NE if the strategy of each microbe is maximizing its fitness function in the given environment and an ESS is a NE not invadable by other strategies (46). From the previous discussion, while joint-FBA and other static methods do not satisfy the condition of NE, dynamic methods simulating over a long time until equilibrium could reach a particular NE but may miss alternative strategies that are more evolutionarily favorable. In community metabolic networks, without introducing ad-hoc parameters, each microbe is represented as the sum of its constituent macromolecules synthesized from the network and the synthesis rate is usually used to represent fitness. The available strategies for each player are any possible flux distributions (the entire set of reaction fluxes) satisfying the mass balance principle, reaction directionality indicated by thermodynamics, and constrained by substrate availability, which depends not only on the nutrients in the environment but also the strategies of other players in terms of the cross-feeding metabolites they may produce (47). The metabolic details, continuous flux space and interdependence of available strategies between players characterize a unique class of microbial games in community metabolic networks (see Figure 1) that require novel game-theoretical methods. In this paper we introduce a bi-level mixed integer optimization framework free of forced altruism termed NECom to identify various NE. We demonstrate the ability of the proposed algorithms to identify all NE in toy community metabolic models that are analogous to classical games and predict reasonable interactions not captured by joint-FBA, OptCom and the dynamic method DMMM. We further validate NECom by analyzing a previously reported algae-yeast co-culture system, which shares a typical crossfeeding feature of lichen: fungi providing CO_2 for photobionts, which fix nitrogen for fungi. NECom is able to predict important trends observed from the experimental data with minimal input from the data and to provide insights into the metabolic states of the two organisms.

RESULTS AND DISCUSSION

A bi-level optimization approach for predicting global Nash Equilibrium of metabolic interactions

A bi-level optimization framework termed Nash equilibrium predictor for microbial community (NECom) was developed. The biggest advance of NECom (Figure *2*) in comparison with previous methods is the absence of 'forced altruism' constraint. While the inner problem is essentially the Flux Balance Analysis (FBA) problem (48), the primary innovation lies in modeling how community metabolites available to the member microbes depend on the exchange profile (strategy) of each microbe. We proved mathematically that NECom prediction can guarantee Nash equilibria for its community members (**Supplementary Note**). With community biomass production as the outer-level objective function, solutions consistent with ESS or strict Nash equilibria can be found (in some cases ESSes depend on the actual competition of substrates which in turn depends on uptake kinetics that is not modeled in this study). The community-level objective function was used in all NECom predictions in this study. The derivation of NECom is presented in **MATERIALS AND METHODS**

Predictions by NECom consistent with classical games analysis

Sub-optimal growth in a community of prototrophs (the prisoner's dilemma)

To test and compare the proposed algorithm with other algorithms in a simple and clear way, four toy models including classical cases in game theory were analyzed. Case 1 is a community model consisting of two individual species (or mutants) with equal abundance named sp1 and sp2 (Figure 3a). Each species can yield ATP through converting substrate S to intermediate P. ATP is subsequently required to convert P to biomass precursors A and B, which can be freely exchanged between the two species. The difference between them is that sp1 produces B more efficiently (consuming 1 ATP versus 3 ATP for sp2), while sp2 produces A more efficiently (consuming 1 ATP versus 3 ATP for sp1), reflecting the existence of pathways with various yields that are commonly observed in nature. Intuitively the optimal strategy for the community is that each species cross-feeds each other with the metabolites produced at higher efficiency, i.e., sp1 supplies B to sp2, while sp2 feeds A to sp1, coinciding with the prediction by joint-FBA and OptCom (the lower flux map in Figure 3a). However, it is not a NE since either species, say sp1, can have a mutant that does not excrete B but keeps consuming A and increases its own biomass production. Using NECom, we predicted that the two species will grow at a lower rate and exchange nothing (the upper flux map in Figure 3a). Since there is no way for each species to grow faster given no export by another species, this is a NE. The dynamic method DMMM predicted the same results. In analogy to classical matrix game, a payoff matrix

corresponding to different cross-feeding states (Figure 3a) was constructed. The payoff
matrix indicates that maximum growth without cross-feeding is a strict NE (the '[0, 0]' cell
on the top left of the payoff matrix in Figure 3a; see also the explanation on the top right
of Figure 3). A strict NE by the Maynard Smith's first condition is also an evolutionary
stable strategy (ESS), which is a NE not invadable by other mutants (or strategies) (49).
Any other strategies combinations including the complementary cross-feedings (the [4.17,
4.17] cell in Figure 3a) predicted by Joint-FBA/OptCom are not stable and will eventually
evolve to the NE. The interaction in the toy community model analyzed above is
characterized as a game called the 'prisoner's dilemma', because the NE is all members
choosing 'defect' instead of 'cooperate', which NECom and DMMM correctly predict.
The 'forced altruism' implied by the formulation is the cause for the mutualistic prediction
by joint-FBA and OptCom because the formulation artificially prioritizes the metabolite
export by each member for its partner over the optimization of its fitness in order to achieve
higher community-level fitness.

Differential biomass yields between members

Case 2 is a slight variant of case 1: the two organisms have different biomass yields on the
same precursors. This reflects a phenomenon common in nature that different species have
the same biomass precursors such as essential amino acidsbut the biomass yields on them
are different. NECom indicates that non-cross-feeding is the strict Nash equilibrium/ESS
(Figure 3b upper). However, Joint-FBA and OptCom predict that sp2 exports all of its
biomass precursors A and B to support the growth of sp1 (Figure 3b lower). Such results
are the consequence of formulations that invoke 'forced altruism', which determines that
doing so achieves the highest overall growth, despite that sp2 needs to completely sacrifice
its own fitness.

Evolutionarily stable mutualism between auxotrophs

Auxotrophy is common in microbial communities. Unlike the previous cases, auxotrophs
rely on complementary cross-feeding for growth while the cross-feeding metabolites such
as amino acids can be costly to produce. In such case how does cooperation become
favorable or would defect still be the outcome despite no growth? Can cheaters and donors
co-exist? To test how different algorithms predict this scenario, in case 3, we used the same
setup as in case 1 but removed the reaction for synthesizing metabolite A in sp1 and the
reaction for synthesizing B in sp2 (Figure 4). This mutualist-cheater case is a classical
question of interest in ecology (50-53). The payoff matrix in Figure 4 shows that all strategy
combinations, including mutualism, commensalisms, and non-crossfeeding are NE, and
Satisfying exactly the Maynard Smith's second condition (54), the mutualism (cell [5, 5])
is the ESS but there is no strict NE (see the explanation on the top right of Figure 4).
Therefore from the view of evolutionary game theory, the non-producing cheaters in this
case can still co-exist with producers although not being able to outgrow them. We checked
that all NE identified by the payoff matrix are feasible solutions of NECom. The ESS was
identical to the prediction of NECom, OptCom and Joint-FBA. However, DMMM predicts

no growth and no crossfeedings. This false prediction is caused by the algorithm design,
which is basically performing FBA for each organisms in each time-step. Each organism
is only indirectly connected via metabolites in the medium and has no way to 'know'
potentially more mutually favorable strategies from other organisms, which evolution
might select for. This example also explains at the level of metabolic resource why
mutualism or commensalism is possible even if it costs members some resource: because
when a particular metabolite is not growth limiting, exporting the metabolite does not
penalize the organism's fitness (e.g., sp1 is limited by A so exporting B does not penalizing
sp1's growth and increases sp2's growth at the same time). We applied NECom and DMMM
to all N pairs of *E. coli* mutants auxotrophic for different amino acids that were experimentally
tested in (55) and verified the same trend of predictions by each of the two methods (data not
shown). DMMM does not predict any E. coli mutant exporting the amino acid essential to its
partner organism, even when the amino acid it is auxotrophic for is available in the initial media.
The predicted growth rate for each member becomes zero after all amino acids initially present
in the media are depleted even if there is remaining glucose. In contrast, NECom predicts
mutualistic cross feeding between the auxotrophs.

Co-existence of mutualists and cheaters (snowdrift game)

Then we tested the algorithms applied to a community with more than two members in
case 4 (Figure 5a). The community was constructed by adding sp3, an exact copy of sp2,
to the toy model in case 3. sp2 and sp3 are dependent on the export of B by sp1 and sp1 is
dependent on the export of A by the other two members. An interesting question is whether
the symmetric pair sp2 and sp3 will both produce A at same time. Nash equilibrium
predicted by NECom indicates that sp2 and sp3 will always choose opposite strategies:
when sp2 produces A (cooperation), sp3 does not (cheating) and vice versa (note the
symmetry between sp2 and sp3 in Figure 5b). This defining feature categorizes the
interactions as a typical 'snowdrift game' between sp2 and sp3. The prediction explicitly
demonstrates the conclusion of our analysis of case 3 that mutualists and cheaters can co-
exist in a NE. This case also suggests that NECom has the potential to predict
species/strains level heterogeneity caused by evolution.

Analysis of co-culture of *C. reinhardtii* and *S. cerevisiae* using NECom

Through toy model analysis NECom was demonstrated to be able to predict results
consistent with traditional game-theoretical analysis. Next NECom was applied to a
realistic algae-yeast co-culture system. Hom and Murray reported that two
phylogenetically distant species *C. reinhardtii* and *S. cerevisiae* can establish obligatory
symbiosis in simple media (56). Their data showed that *C. reinhardtii* almost always had
a higher abundance than *S. cerevisiae* in all 80 experiments with different initial substrate
concentrations. Within a wide range of initial concentrations of two key substrates (nitrite
and glucose), these two species can co-exist for an extended time period ($\geq$10 days, Figure

6). To elucidate how nutrient availability and the underlying metabolic interactions
between the two organisms shaped the community growth pattern and the implication on
the system stability, we performed a series of analyses by applying NECom (Figure 6) to
a community model integrating the genome-scale metabolic models for *S. cerevisiae*
(Yeast 7.6) (57) and *C. reinhardtii* (*i*Cre1355) (58). First we show the predictive capability
of NECom for this system. Then from the simulation results we provide insights into
growth-limiting nutrients and cross feeding metabolites and how they contribute to the
predicted negative frequency dependent dynamics which stabilizes the community.

Improved growth prediction by NECom

In order to validate the applicability of the NECom method to the *C. reinhardtii-S.*
cerevisiae system, growth rates given uptake rates derived from experimental
measurements (see **Supplementary Note**) were predicted using both NECom and Joint-
FBA under a wide range of conditions (Figure 7a, b). The average root mean squared error
(RMSE) for NECom is 3.1e-3 (h^{-1}) for predicting the growth of *C. reinhardtii* and 8.5e-3
for the growth of *S. cerevisiae* respectively, achieving 63.5% and 81.7% reduction when
compared to those predicted by Joint-FBA (statistical significance in reduction verified by
Z-test, n =63 , p-value = 3.12e-7 and 1.56e-23 respectively). The growth rates of *S.*
cerevisiae predicted by Joint-FBA are generally significantly higher than the experimental
results, leading to *S. cerevisiae* predicted as a generally faster grower while the
experimental results indicate the opposite in majority of the cases (see Figure 7b). When
the glucose uptake of *S. cerevisiae* is low (<0.6 mmol/gDWh), Joint-FBA predicts positive
values instead of the experimental negative values. These errors are caused by 'forced
altruistic' constraints, which force *C. reinhardtii* model using photons to convert low
energy carbon containing metabolites (primarily CO_2) released by *S. cerevisiae* to various
energy richer substrates subsequently to be consumed by *S. cerevisiae* (see the example in
Figure 7c). We confirmed this by maximizing *S. cerevisiae*'s growth and *C. reinhardtii*'s
growth individually under the same condition using Joint-FBA and found *S. cerevisiae*'s
maximum growth rate 2.7 times greater than *C. reinhardtii*'s. Therefore, 'forced altruistic
constraints' unreasonably weaken the role of individual fitness and lead to inaccurate
predictions. For the prediction by NECom, although in most cases it agrees well with the
measurement-derived data, in some cases (e.g., nitrite uptake = 0.225 mmol/gDWh,
glucose uptake = 0.725 mmol/gDWh) *S. cerevisiae*'s growth is underestimated while *C.*
reinhardtii's growth is overestimated. One possible reason is over-estimated uptake rates
of CO_2 by *C. reinhardtii*. When nitrogen is in excess, a higher CO2 uptake causes more-
than-actual ammonia converted to biomass and less-than-actual ammonia exported for *S.*
cerevisiae's growth. The discrepancies are likely due to the absence of uptake kinetics in
the current study.

Impacts of substrate uptake rates on the relative growth between members

To the contrary of our impression that *S. cerevisiae* usually grows faster, experimental
data (see Figure *8*c) suggests that in 76.7% of the cases, *C. reinhardtii* is the faster grower

in this co-culture and usually has positive growth while *S. cerevisiae* could have negative
growth especially when its glucose uptake rate is low(see Figure *7*b). The lower the glucose
uptake is, the more the advantage of *C. reinhardtii* can have. Though at high glucose uptake
rates (v_{glc}^{uptake} >0.8 mmol/gDWh, see Figure *8*c), *S. cerevisiae* grows faster. We suspected
that the algae being able to catch up and surpass the yeast at the depletion of glucose is
important to the stabilization of the community and likely controlled by the change of
substrate uptake rates. NECom predictions capturing the aforementioned phenomena
(Figure *8*d) were then used to analyze the underlying mechanism.

Using NECom predictons, we examined the always-positive growth of *C. reinhardtii*.
Since the range of the experimental glucose uptake rate is wide enough for analysis only
at low nitrite uptake rates, two representative low nitrite uptake rates, 0.025 and 0.075
mmol/gDWh, were selected for further analysis (67% of the experimental cases has
estimated nitrite uptake rates < 0.1 mmol/gDWh). The maximum flux through the ATP
maintenance (ATPM) reaction for each species was computed. If the ATPM flux is lower
than a minimal value, which is the non-growth-associated maintenance (NGAM) value
(2.85 mmol/gDWh for *C. reinhardtii* and 7.5 mmol/gDWh for *S. cerevisiae*), the growth
rate will be negative. We found that the minimal ATPM flux is always satisfied for *C.*
reinhardtii. However, for *S. cerevisiae*, when the glucose uptake rate drops below 0.8
mmol/gDWh (Figure *8*b), the ATPM flux starts to drop below the minimal requirement,
the gap keeps increasing as the glucose uptake rate continues to decrease. One probable
reason for this difference lies in the different source of energy used for fueling ATPM by
the two species. For *C. reinhardtii* the energy source is photons whereas for *S. cerevisiae*
the energy originates from metabolizing glucose. In the experiment, illumination was
constant but the glucose concentration had different initial values and decreased over time,
supporting our analysis.

Then, we investigated the experimental trend that *C. reinhardtii* narrowed its growth
rate gap with *S. cerevisiae* and then surpassed it with the decrease of glucose uptake caused
by the depletion (see Figure 9). The carbon flux to the biomass of *S. cerevisiae* and nitrogen
flow to the biomass of *C. reinhardtii* have opposite trends. Based on NECom prediction,
in response to the decreasing glucose uptake rates *S. cerevisiae* has a decreasing ratio of
ATP for biomass synthesis to the less variant ATP for cellular maintenance, leading to a
decreasing growth rate (Figure *8*f). In the meantime, The ratio of carbon flow to *S.*
cerevisiae biomass over that to exported CO_2 decreases and is responsible for more carbon
available for *C. reinhardtii* to synthesize biomass (Figure *8*e). This is consistent with the
following limiting substrate analysis, where CO_2 is revealed to be a limiting substrate to *C.*
reinhardtii (see Figure 9d).

Predicted negative frequency-dependent dynamics caused by limiting cross-feeding
metabolites

The analysis thus far shows the ability of NECom to capture the growth trend of the
members and provides some metabolic insights into the potential reason for the observed

trend. A general question of interest for microbial communities is whether the community is stable and how the environmental conditions impact the stability. To complement the limitation that the measured data were concentrated in a small range of low substrate uptake rates, we extrapolated the predictions by NECom over a large range of relative abundance of *S. cerevisiae* and *C. reinhardtii* to address this question. NECom predicts that with higher glucose uptake rates (>0.75 mmol/gDWh) the growth rates of the two species will have opposite trends to the relative abundance of *C. reinhardtii* and often there exists a point of equal growth rate (black circles in Figure 9a), where the faster and slower growing species are swapped. Such a growth-abundance relationship is negative frequency-dependent, which stabilizes the co-existence of interacting species by dynamically evolving to the equal-growth-rate population steady state. An interesting observation is that the population steady state shifts toward the left on the abundance axis with the increase of glucose uptake rate for *C. reinhardtii* (Figure 9a), indicating that the growth of *S. cerevisiae* gains advantage over *C. reinhardtii* with an increasing glucose uptake rate. The prediction is consistent with the experimental data (Figure *8*c).

We speculated that the changing availability of some key substrates might be the reason for the negative frequency dependent dynamics and the shift of population steady state. To test our speculation, all cross-feeding metabolites and externally sourced substrates need to be considered. We asked which metabolites were the major substrates used by the two species. Among the simulation cases at various glucose uptake rates, nitrite uptake rates and abundances (represented by brown dots in Figure *8*a), cross-feeding metabolites observed include O_2, CO_2, ammonia, acetate and succinate. The former three metabolites occur in 93% of the total cases (Figure 9b). Therefore, O_2, CO_2, ammonia together with external sourced substrates including photon, glucose and nitrite were selected for further analysis. Since only limiting substrates can have impact on community members' growth, in order to specify which substrates are growth-limiting, shadow prices for the uptake of each metabolites were extracted from NECom solutions (see Supplementary Note). A positive shadow price indicates the corresponding metabolite has impact on cellular growth at a given state and the value indicates the intensity of such impact. We found that for *C. reinhardtii*, CO_2 is a limiting metabolite under almost all simulated conditions (Figure 9d), nitrite have significant limiting effect when both nitrite and glucose uptakes are low. Photon is generally not limiting. For *S. cerevisiae*, only ammonia exhibits significant limiting effect (Figure 9e).

We then focused on analyzing how CO_2 and ammonia affect the growth rates of this system. Figure 9f-i indicate that with increase in the abundance of *C. reinhardtii*, CO_2 supplied by *S. cerevisiae* decreases, causing decrease in the growth rate of *C. reinhardtii* since CO_2 is the limiting substrate confirmed by the positive shadow price curve. In the meantime, increase in ammonia supply from *C. reinhardtii* contributes to the increasing growth rates of *S. cerevisiae*, since ammonia uptake has increasingly positive shadow prices. Therefore, the opposite trends in the supply rate of each limiting substrates are responsible for the observed negative frequency-dependent dynamics. Shadow price

analysis can also explain why the population steady state changes with the glucose uptake
rate. Figure 9a shows that with the increase in glucose uptake rate, the two growth-
abundance curves becomes steeper at low *C. reinhardtii*'s abundance (<0.6mmol/gDWh),
causing the cross point shifting to the left. The steepness of a growth-abundance curve is
proportional to the shadow price for limiting substrate uptake. Comparison of Figure 9g
and Figure 9f reveals that, fixing the nitrite uptake rate at 0.09 mmol/gDWh and increasing
the glucose uptake rate from 0.93 mmol/gDWh to 1.93 mmol/gDWh result in an increase
in shadow price for CO_2 uptake by *C. reinhardtii* from 0 to 0.02-0.04 gDW/mmol when
the abundance of *C. reinhardtii* is below 0.6. Similarly Figure 9i and Figure 9h indicate
that if the nitrite uptake rate is fixed at 0.59 mmol/gDWh, shadow prices for ammonia
uptake by *S. cerevisiae* will increase from 0 to 0.25 gDW/mmol when the abundance of *C.*
reinhardtii is below 0.7.

Relations to previous methods

A game-theoretical method has been proposed recently to determine Nash equilibria
for interacting organisms using metabolic models(43). In that proposed method, the Black
Queen hypothesis (59) was tested by checking whether the existence of auxotrophic
mutants is favored by evolution if there is leakage of metabolites. Metabolite leakage was
predefined and uptake of leaky metabolites was determined using a distribution rule. FBA
was then performed for each genotype in the population to compute the payoff matrix
which was subsequently used for Nash equilibrium determination as well as dynamic
simulation using the replicator equation. While the results elegantly captured *S.*
cerevisiae's invertase system and suggested potential auxotrophic pairs that can stably co-
exist, similar to other dynamic methods, performing FBA for each individual genotype, or
more generally organism, does not allow individuals to 'sense and response' to each other
via changes in the metabolic environment. Therefore, the possible interactions observed
are confined to the cross-feeding of the predefined leaky metabolites. In the current study,
the Black Queen hypothesis is not invoked (though it can be). Also, by clearly defining the
concept of microbial metabolic games and the unique strategy space which is a continuous
flux space constrained by the strategies of other players, NECom is able to search the inter-
dependent continuous flux space to determine Nash equilibria among all available
interactions by using a bi-level framework, in which each organism optimizes its own
fitness, under the substrate uptake constraints shaped by the strategies of other community
members. In relation to previous algorithms for simulating community metabolic models,
OptCom (35) also employs a bi-level optimization framework but the main innovation of
NECom lies in the introduction of outer variables to constrain the uptake only for the inner
problems instead of both uptake and export (see Methods for the mathematical details).
SteadyCom (38) implements a population steady state but the single objective function also
unavoidably imposes forced altruism on the community the same way as joint-FBA does.
We have not implemented the population steady state in the formulation but it can still be
analyzed by fixing the relative abundances at different values as performed in the algae-

yeast case study. In Table 1 some key features of NECom and current algorithms are compared.

Algorithms	Static/Dynamic	Free of forced altruism?	Find multiple Nash equilibria?	Ref
NECom	Static	Yes	Yes	This sudy
OptCom	Static	No	No	(35)
cFBA	Static	No	No	(37)
Joint-FBA	Static	Yes	No	(20)
DMMM	Dynamic	Yes	No	(26)
COMETS	Dynamic	Yes	No	(30)

Table 1. Comparison of representative algorithms for predicting microbial metabolic interactions

In conclusion, we developed a bi-level non-convex mixed integer optimization algorithm called 'NECom' to predict the Nash equilibrium of multiple species interacting at the metabolic level. NECom was shown to predict Nash equilibria of different types of classical games correctly in toy models that previous methods' predictions cannot guarantee. The main reason for this difference is that NECom does not contains any 'forced altruism' that computationally forces individual species to fulfill other species' metabolic need before their own. As proof of concepts, NECom was used to analysis a well-documented co-culturing system of *C. reinhardtii* and *S. cerevisiae*. With minimal information from the experimental data, NECom is able to reveal that the decreasing glucose concentration vs. the constant photon availability provides significant growth advantage of *C. reinhardtii* over *S. cerevisiae*; CO_2 and ammonia are growth-limiting cross-feeding substrates for *C. reinhardtii* and *S. cerevisiae* respectively; and the negative frequency-dependent growth pattern is caused by opposite trends of the supply rates of each species' limiting substrates.

MATERIALS AND METHODS

Derivation of NECom

Let N be the set of organisms in a microbial community. For each organism $n \in N$, let I_n be the set of metabolites, J_n be the set of reactions, $\mathbf{c_n} = \left[c_{j,n}\right]_{j \in J_n}$ be the objective coefficient vector, and $v_{j,n}$ be the flux of reaction $j \in J_n$. The FBA problem for organism n can be formulated as:

$$Max \sum_{j \in J_n} c_{j,n} \, v_{j,n}$$

subject to:

$$\sum_{j \in J_n} S_{i,j,n} \, v_{j,n} = 0, \qquad \forall i \in I^n$$

$$-v_{j,n} \le -LB_{j,n} \qquad \forall j \in J_n$$

$$v_{j,n} \le UB_{j,n} \qquad \forall j \in J_n$$

$$v_{j,n} \in \mathbb{R} \qquad \forall j \in J_n$$

where $\left[S_{i,j,n}\right]_{i \in I_n, j \in J_n}$ is the stoichiometric matrix, LB_j and UB_j are the lower and upper bounds for the flux of reaction j respectively. The equality constraint represents metabolite balances at pseudo-steady state. The inequalities include bound constraints due to reaction directionality, substrate availability and the requirement of non-growth-associated-maintenance (NGAM) modeled as a minimum flux through an ATP hydrolysis reaction, which is usually called ATP maintenance (ATPM) in metabolic models.

The FBA problem will be the inner problem of NECom for optimizing the fitness of individual organisms. A set of outer-level variables to indicate the community metabolites available for uptake are then introduced and connected to the exchange flux variables of individual organisms. Let I_n^{ex} be the set of extracellular metabolites in the community, $J_n^{ex} \subseteq J_n$ be the set of exchange reactions between the community and organism n. Define an index mapping function $I_n^{ex} \to J_n^{ex}$ such that $com(i,n) \in J_n^{ex}$ map extracellular metabolite $i \in I_n^{ex}$ to its corresponding exchange reaction. For each individual organism, we split the exchange reaction flux for a possible cross-feeding metabolite $v_{com(i,n),n}$ into two non-negative continuous variables: the uptake rate $v_{com(i,n),n}^{ut}$ and the export rate $v_{com(i,n),n}^{ex}$:

$$v_{com(i,n),n} = -v_{com(i,n),n}^{ut} + v_{com(i,n),n}^{ex} \qquad \forall i \in I_n^{ex}, \forall n \in N$$

To explicitly model the mutual dependency in terms of inter-organism metabolite exchange, uptake of a metabolite by an organism is possible only if there is surplus

of the metabolite in the medium after the consumption/production by the rest of the
community:

$$X_n v_{com(i,n),n}^{ut} \leq \max\{v_i^{med} + \sum_{k \in N, k \neq n} X_k \, v_{com(i,n),n}, 0\} \quad \forall i \in I_n^{ex}, \forall n \in N \quad (eq.\,1)$$

where X_n is the relative abundance of organism n (a pre-set parameter), which is
multiplied by specific uptake rate $v_{com(i,n),n}^{ut}$ to correctly normalize the exchange with the
community, and v_i^{med} is the maximum community uptake of metabolite i from the
medium. Here to avoid forced altruism, $v_{com(i,n),n}^{ex}$ is designated as an inner variable
independent of the outer problem, while the uptake rate $v_{com(i,n),n}^{ut}$ is also an inner variable
but depends on the net availability of metabolite i in the community. In this way we can
ensure that individual organisms have 'autonomy' over their metabolite exports.
Maximization of the community fitness in the outer level cannot force an organism to
produce a certain metabolite unless FBA determines that the metabolite is necessary for,
or at least not undermining the maximum growth of the organism. This modeling constraint
is novel to our best knowledge. The max function in the *eq.1* is linearized by introducing
an outer continuous variable $\beta_{i,n}$ and an outer binary variable $\delta_{i,n}$:

462
$$\left[\begin{array}{l} \beta_{i,n} \geq v_i^{med} + \sum_{k \in N, k \neq n} X_k \, v_{com(i,n),n} \\[2ex] \beta_{i,n} \leq (1 - \delta_{i,n})M + v_i^{med} + \sum_{k \in N, k \neq n} X_k \, v_{com(i,n),n} \\[2ex] \beta_{i,n} \leq \delta_{i,n}M \\ \beta_{i,n} \geq 0 \end{array} \right] \quad \forall i \in I_n^{ex}, \forall n \in N$$

463 Where M is a large constant. Combining the inner problem and outer problem the complete
464 formulation of NECom can be written as:

466 $\quad$ max $\quad \displaystyle\sum_{n \in N} v_n^{biomass}$

467 $\quad$ subject to

$$
\left[
\begin{array}{ll}
\max \quad v_n^{biomass} & \\
\text{subject to} & \\
\quad \displaystyle\sum_{j \in J_n} S_{i,j,n}\, v_{j,n} = 0, & \forall i \in I_n \\[2mm]
\quad LB_{j,n} \leq v_{j,n} \leq UB_{j,n}\,, & \forall j \in J_n \\[2mm]
\quad v_{com(i,n),n} + v_{com(i,n),n}^{ut} - v_{com(i,n),n}^{ex} = 0, & \forall i \in I_n^{ex} \\[2mm]
\quad X_n v_{com(i,n),n}^{ut} \leq \beta_{i,n}\,, & \forall i \in I_n^{ex}
\end{array}
\right] \quad \forall n \in N
$$

$$\beta_{i,n} \geq v_i^{med} + \sum_{k \in N, k \neq n} X_k v_{com(i,k),k}\,, \qquad \forall i \in I_n^{ex}, n \in N$$

$$\beta_{i,n} \leq M\delta_{i,n} + v_i^{med} + \sum_{k \in N, k \neq n} X_k v_{com(i,k),k}\,, \qquad \forall i \in I_n^{ex}, n \in N$$

$$\beta_{i,n} \leq M\left(1 - \delta_{i,n}\right), \qquad \forall i \in I_n^{ex}, n \in N$$

$$v_{j,n} \in \mathbb{R}, \qquad \forall j \in J_n, n \in N$$

$$v_{com(i,n),n}^{ut}, v_{com(i,n),n}^{ex} \geq 0, \qquad \forall i \in I_n^{ex}, n \in N$$

$$\delta_{i,n} \in [0,1], \qquad \forall i \in I_n^{ex}, n \in N$$

The only situation where the above algebraic system is not feasible is that one or more
members cannot satisfy the minimal NGAM, which is represented as a direct ATP
hydrolysis reaction with a pre-defined lower bound in model. Biologically this suggests
the occurrence of cell death. In order to enable NECom to predict this phenomenon, we
separate community members into two sets based on NECom solution: let N^G be the set
of community members which have adequate substrates for uptake to satisfy their NGAM,
N^D be the set of the rest of the community members that are dying due to insufficient
nutrients. For the latter case, the default FBA problem is infeasible. Instead of maximizing
biomass production, we relax the NGAM constraint and maximize the ATP hydrolysis
activity that an organism can generate to minimize cell death, resulting in NECom enabling
negative growth prediction (NEComNG) as:

$\max \ \sum_{n \in N} v_n^{biomass}$

$$
\begin{bmatrix}
\max \quad v_n^{biomass} \\
\text{subject to} \\
\quad \sum_{j \in J_n} S_{i,j,n}\, v_{j,n} = 0, \qquad\qquad\qquad \forall i \in I_n \\
\quad LB_{j,n} \le v_{j,n} \le UB_{j,n}\,, \qquad\qquad\quad \forall j \in J_n \\
\quad v_{com(i,n),n} + v_{com(i,n),n}^{ut} - v_{com(i,n),n}^{ex} = 0, \qquad \forall i \in I_n^{ex} \\
\quad X_n v_{com(i,n),n}^{ut} \le \beta_{i,n}\,, \qquad\qquad\qquad \forall i \in I_n^{ex}
\end{bmatrix} \forall n \in N^G
$$

$$
\begin{bmatrix}
\max \quad v_n^{ATPM} \\
\text{subject to} \\
\quad \sum_{j \in J_n} S_{i,j,n}\, v_{j,n} = 0, \qquad\qquad\qquad \forall i \in I_n \\
\quad LB_{j,n} \le v_{j,n} \le UB_{j,n}\,, \qquad\qquad\quad \forall j \in J_n \\
\quad v_{com(i,n),n} + v_{com(i,n),n}^{ut} - v_{com(i,n),n}^{ex} = 0, \qquad \forall i \in I_n^{ex} \\
\quad X_n v_{com(i,n),n}^{ut} \le \beta_{i,n}\,, \qquad\qquad\qquad \forall i \in I_n^{ex}
\end{bmatrix} \forall n \in N^D
$$

$$
\beta_{i,n} \ge v_i^{med} + \sum_{k \in N, k \ne n} X_k v_{com(i,k),k}\,, \qquad \forall i \in I_n^{ex}, n \in N
$$

$$
\beta_{i,n} \le M\delta_{i,n} + v_i^{med} + \sum_{k \in N, k \ne n} X_k v_{com(i,k),k}\,, \qquad \forall i \in I_n^{ex}, n \in N
$$

$$
\beta_{i,n} \le M(1 - \delta_{i,n}), \qquad\qquad\qquad\qquad \forall i \in I_n^{ex}, n \in N
$$

$$
v_{j,n} \in \mathbb{R}, \qquad\qquad\qquad\qquad\qquad \forall j \in J_n, n \in N
$$

$$
v_{com(i,n),n}^{ut}, v_{com(i,n),n}^{ex} \ge 0, \qquad\qquad\qquad \forall i \in I_n^{ex}, n \in N
$$

$$
\delta_{i,n} \in [0,1], \qquad\qquad\qquad\qquad\qquad \forall i \in I_n^{ex}, n \in N
$$

The transformation of the bilevel optimization problem into a mixed integer non-linear problem is detailed in **Supplementary Note.**

Software used

Baron (60) was employed to solve all non-linear and mixed integer non-linear optimization problems. CPLEX and GLPK were used to solve linear problems. Pathway maps were originally generated in Escher Map (61). All scripts were written in Matlab.

Code and data availability

Data files and example codes will be available on https://github.com/Jingyi-Cai/NECom.git

Acknowledgments

We acknowledge Dr. Wenyan Yuan, Dr. Erik F. Y. Hom, Dr. Guang Cheng, Dr. Xueying Ni and Ying Zhu for their useful comments.

Author contributions

Conceptualization, J.C and S.H.J.C; Methodology, J.C and S.H.J.C; Software, J.C; Analysis and Investigation, J.C and S.H.J.C; Writing – Original Draft, J.C and S.H.J.C; Writing – Review & Editing, S.H.J.C, T.T, and J.C; Visualization, J.C and S.H.J.C; Supervision, T.T and S.H.J.C.

Competing interests

The authors declare no competing interests

REFERENCES

1. Zhou J, Deng Y, Luo F, He Z, Tu Q, Zhi X. Functional Molecular Ecological Networks. mBio. 2010;1(4).

2. Fiegna F, Moreno-Letelier A, Bell T, Barraclough TG. Evolution of species interactions determines microbial community productivity in new environments. The ISME Journal. 2014;9(5):1235-45.

3. Lawrence D, Fiegna F, Behrends V, Bundy JG, Phillimore AB, Bell T, et al. Species Interactions Alter Evolutionary Responses to a Novel Environment. PLoS Biology. 2012;10(5):e1001330-e.

4. Pande S, Kost C. Bacterial Unculturability and the Formation of Intercellular Metabolic Networks. Trends in Microbiology. 2017;25(5):349-61.

5. D'Souza G, Kost C. Experimental Evolution of Metabolic Dependency in Bacteria. PLOS Genetics. 2016;12(11):e1006364-e.

6. Hillesland KL, Lim S, Flowers JJ, Turkarslan S, Pinel N, Zane GM, et al. Erosion of functional independence early in the evolution of a microbial mutualism. Proceedings of the National Academy of Sciences. 2014;111(41):14822-7.

7. Hillesland KL, Stahl DA. Rapid evolution of stability and productivity at the origin of a microbial mutualism. Proceedings of the National Academy of Sciences. 2010;107(5):2124-9.

8. McCutcheon JP, Moran NA. Parallel genomic evolution and metabolic interdependence in an ancient symbiosis. Proceedings of the National Academy of Sciences. 2007;104(49):19392-7.

9. Paterson S, Vogwill T, Buckling A, Benmayor R, Spiers AJ, Thomson NR, et al. Antagonistic coevolution accelerates molecular evolution. Nature. 2010;464(7286):275-8.

10. Kaeberlein T. Isolating "Uncultivable" Microorganisms in Pure Culture in a Simulated Natural Environment. Science. 2002;296(5570):1127-9.

11. Pham VHT, Kim J. Cultivation of unculturable soil bacteria. Trends in Biotechnology. 2012;30(9):475-84.

12. Moutinho TJ, Panagides JC, Biggs MB, Medlock GL, Kolling GL, Papin JA. Novel co-culture plate enables growth dynamic-based assessment of contact-independent microbial interactions. PLOS ONE. 2017;12(8):e0182163-e.

13. Zhang Z, Ji H, Gong G, Zhang X, Tan T. Synergistic effects of oleaginous yeast Rhodotorula glutinis and microalga Chlorella vulgaris for enhancement of biomass and lipid yields. Bioresour Technol. 2014;164:93-9.

14. Abraham W-R. Applications and impacts of stable isotope probing for analysis of microbial interactions. Applied Microbiology and Biotechnology. 2014;98(11):4817-28.

15. Wang Y, Song Y, Tao Y, Muhamadali H, Goodacre R, Zhou N-Y, et al. Reverse and Multiple Stable Isotope Probing to Study Bacterial Metabolism and Interactions at the Single Cell Level. Analytical Chemistry. 2016;88(19):9443-50.

16. Gebreselassie NA, Antoniewicz MR. 13 C-metabolic flux analysis of co-cultures: A novel approach. Metabolic Engineering. 2015;31:132-9.

17.   Zhang C, Derrien M, Levenez F, Brazeilles R, Ballal SA, Kim J, et al.
Ecological robustness of the gut microbiota in response to ingestion of transient
food-borne microbes. The ISME Journal. 2016;10(9):2235-45.
18.   Medlock GL, Carey MA, McDuffie DG, Mundy MB, Giallourou N, Swann
JR, et al. Inferring Metabolic Mechanisms of Interaction within a Defined Gut
Microbiota. Cell Systems. 2018;7(3):245-57.e7.
19.   Kim TY, Sohn SB, Kim YB, Kim WJ, Lee SY. Recent advances in
reconstruction and applications of genome-scale metabolic models. Current Opinion
in Biotechnology. 2012;23(4):617-23.
20.   Stolyar S, Van Dien S, Hillesland KL, Pinel N, Lie TJ, Leigh JA, et al.
Metabolic modeling of a mutualistic microbial community. Mol Syst Biol. 2007;3:92.
21.   Freilich S, Zarecki R, Eilam O, Segal ES, Henry CS, Kupiec M, et al.
Competitive and cooperative metabolic interactions in bacterial communities.
Nature Communications. 2011;2:589.
22.   Heinken A, Sahoo S, Fleming RMT, Thiele I. Systems-level
characterization of a host-microbe metabolic symbiosis in the mammalian gut. Gut
Microbes. 2013;4(1):28-40.
23.   Heinken A, Thiele I. Systematic prediction of health-relevant human-
microbial co-metabolism through a computational framework. Gut Microbes.
2015;6(2):120-30.
24.   Heinken A, Thiele I. Anoxic Conditions Promote Species-Specific
Mutualism between Gut MicrobesIn Silico. Applied and Environmental Microbiology.
2015;81(12):4049-61.
25.   Magnúsdóttir Sa, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A,
et al. Generation of genome-scale metabolic reconstructions for 773 members of the
human gut microbiota. Nature Biotechnology. 2016;35(1):81-9.
26.   Zhuang K, Izallalen M, Mouser P, Richter H, Risso C, Mahadevan R, et
al. Genome-scale dynamic modeling of the competition between Rhodoferax and
Geobacter in anoxic subsurface environments. The ISME Journal. 2010;5(2):305-16.
27.   Xu X, Zarecki R, Medina S, Ofaim S, Liu X, Chen C, et al. Modeling
microbial communities from atrazine contaminated soils promotes the development
of biostimulation solutions. The ISME Journal. 2018;13(2):494-508.
28.   Hanly TJ, Henson MA. Dynamic flux balance modeling of microbial co-
cultures for efficient batch fermentation of glucose and xylose mixtures. Biotechnol
Bioeng. 2011;108(2):376-85.
29.   Gomez JA, Höffner K, Barton PI. DFBAlab: a fast and reliable MATLAB
code for dynamic flux balance analysis. BMC Bioinformatics. 2014;15(1).
30.   Harcombe WR, Riehl WJ, Dukovski I, Granger BR, Betts A, Lang AH, et
al. Metabolic resource allocation in individual microbes determines ecosystem
interactions and spatial dynamics. Cell Rep. 2014;7(4):1104-15.
31.   Henson M, Phalak P. Byproduct Cross Feeding and Community
Stability in an In Silico Biofilm Model of the Gut Microbiome. Processes.
2017;5(4):13-.

32.   Henson MA, Phalak P. Microbiota dysbiosis in inflammatory bowel
diseases: in silico investigation of the oxygen hypothesis. BMC Systems Biology.
2017;11(1).
33.   Bauer E, Zimmermann J, Baldini F, Thiele I, Kaleta C. BacArena:
Individual-based metabolic modeling of heterogeneous microbes in complex
communities. PLOS Computational Biology. 2017;13(5):e1005544-e.
34.   Chan SHJ, Friedman ES, Wu GD, Maranas CD. Predicting the
Longitudinally and Radially Varying Gut Microbiota Composition Using Multi-Scale
Microbial Metabolic Modeling. Processes. 2019;7(7):394-.
35.   Zomorrodi AR, Maranas CD. OptCom: A Multi-Level Optimization
Framework for the Metabolic Modeling and Analysis of Microbial Communities.
PLoS Computational Biology. 2012;8(2):e1002363-e.
36.   Zomorrodi AR, Islam MM, Maranas CD. d-OptCom: Dynamic Multi-
level and Multi-objective Metabolic Modeling of Microbial Communities. {ACS}
Synthetic Biology. 2014;3(4):247-57.
37.   Khandelwal RA, Olivier BG, Röling WFM, Teusink B, Bruggeman FJ.
Community Flux Balance Analysis for Microbial Consortia at Balanced Growth. PLoS
ONE. 2013;8(5):e64567-e.
38.   Chan SHJ, Simons MN, Maranas CD. SteadyCom: Predicting microbial
abundances while ensuring community stability. PLOS Computational Biology.
2017;13(5):e1005539-e.
39.   Shoaie S, Ghaffari P, Kovatcheva-Datchary P, Mardinoglu A, Sen P,
Pujos-Guillot E, et al. Quantifying Diet-Induced Metabolic Changes of the Human Gut
Microbiome. Cell Metab. 2015;22(2):320-31.
40.   Henson MA, Phalak P. Suboptimal community growth mediated
through metabolite crossfeeding promotes species diversity in the gut microbiota.
PLOS Computational Biology. 2018;14(10):e1006558-e.
41.   Zelezniak A, Andrejev S, Ponomarova O, Mende DR, Bork P, Patil KR.
Metabolic dependencies drive species co-occurrence in diverse microbial
communities. Proc Natl Acad Sci U S A. 2015;112(20):6449-54.
42.   Pacheco AR, Moel M, Segrè D. Costless metabolic secretions as drivers
of interspecies interactions in microbial ecosystems. Nature Communications.
2019;10(1).
43.   Zomorrodi AR, Segrè D. Genome-driven evolutionary game theory
helps understand the rise of metabolic interdependencies in microbial communities.
Nature Communications. 2017;8(1).
44.   Dubey GP, Ben-Yehuda S. Intercellular Nanotubes Mediate Bacterial
Communication. Cell. 2011;144(4):590-600.
45.   Pande S, Shitut S, Freund L, Westermann M, Bertels F, Colesie C, et al.
Metabolic cross-feeding via intercellular nanotubes among bacteria. Nature
Communications. 2015;6.
46.   Nowak MA. Evolutionary dynamics: exploring the equations of life:
Harvard University Press; 2006.

47.   Schuster S, Kreft J-U, Schroeter A, Pfeiffer T. Use of Game-Theoretical
Methods in Biochemistry and Biophysics. Journal of Biological Physics. 2008;34(1-
2):1-17.
48.   Orth JD, Thiele I, Palsson BO. What is flux balance analysis? Nature
Biotechnology. 2010;28(3):245-8.
49.   Smith JM. The theory of games and the evolution of animal conflicts. J
Theor Biol. 1974;47(1):209-21.
50.   Ferriere R, Bronstein JL, Rinaldi S, Law R, Gauduchon M. Cheating and
the evolutionary stability of mutualisms. Proceedings of the Royal Society of London
Series B: Biological Sciences. 2002;269(1493):773-80.
51.   Ferrière R, Gauduchon M, Bronstein JL. Evolution and persistence of
obligate mutualists and exploiters: competition for partners and evolutionary
immunization. Ecology Letters. 2007;10(2):115-26.
52.   Medaney F, Dimitriu T, Ellis RJ, Raymond B. Live to cheat another day:
bacterial dormancy facilitates the social exploitation of β-lactamases. The ISME
Journal. 2015;10(3):778-87.
53.   Waite AJ, Shou W. Adaptation to a new environment allows
cooperators to purge cheaters stochastically. Proceedings of the National Academy
of Sciences. 2012;109(47):19079-86.
54.   Smith JM, Price GRJN. The logic of animal conflict.
1973;246(5427):15-8.
55.   Mee MT, Collins JJ, Church GM, Wang HH. Syntrophic exchange in
synthetic microbial communities. Proceedings of the National Academy of Sciences.
2014;111(20):E2149-E56.
56.   Hom EFY, Murray AW. Niche engineering demonstrates a latent
capacity for fungal-algal mutualism. Science. 2014.
57.   Aung HW, Henry SA, Walker LP. Revising the Representation of Fatty
Acid, Glycerolipid, and Glycerophospholipid Metabolism in the Consensus Model of
Yeast Metabolism. Industrial Biotechnology. 2013;9(4):215-28.
58.   Imam S, Schäuble S, Valenzuela J, de Lomana ALGa, Carter W, Price
ND, et al. A refined genome-scale reconstruction of Chlamydomonas metabolism
provides a platform for systems-level analyses. The Plant Journal. 2015;84(6):1239-
56.
59.   Morris JJ, Zinser REL, Erik R. The Black Queen Hypothesis: Evolution
of Dependencies through Adaptive Gene Loss. mBio. 2012;3(2).
60.   Liu J, Ploskas N, Sahinidis NV. Tuning BARON using derivative-free
optimization algorithms. Journal of Global Optimization. 2018.
61.   King ZA, Dräger A, Ebrahim A, Sonnenschein N, Lewis NE, Palsson BO.
Escher: A Web Application for Building, Sharing, and Embedding Data-Rich
Visualizations of Biological Pathways. PLOS Computational Biology.
2015;11(8):e1004321-e.

Figure 1. **Schematic diagram of interactive strategy space of two community members**. An available strategy for an organism is any feasible flux distribution satisfying the pseudo-steady-state condition and reaction directionality constrained by the availability of extracellular metabolites, which can be affected by other community members. Interactive strategy space causes the complexity of microbial metabolic games, distinguishing microbial metabolic game from most traditional games.

682
683 Figure 2. **Schematic diagram of the NECom formulation.**

Figure 3. **Flux distribution for toy models 1 and 2 predicted by different algorithms.** The corresponding flux distributions predicted by Joint-FBA, OptCom, DMMM and NECom (NC) respectively. Growth rates of species 1 (sp1) and species 2

(sp2) calculated with FBA are shown in each cell in the payoff matrices for each case.
(a) Toy community model 1 consisting of two members, both capable of producing
biomass precursors A and B, but at different ATP costs. **(b)** Toy community model 2
consisting of two members, both capable of producing biomass precursors A and B,
but has different biomass yields on these two precursors. Only NECom and DMMM
can predict the ESS derived from the traditional payoff matrix approach in these two
cases where the ESS is not mutualistic.

Case 3

Figure 4. **Flux distribution for toy model 3 predicted by different algorithms.** Toy community model 3 consists of two members, one capable of producing biomass precursors A but auxotrophic for B, the other capable of producing biomass precursors B but auxotrophic for A. Only NECom, OptCom and Joint-FBA are able to predict the ESS derived from the traditional payoff matrix approach in this case where the ESS is mutualistic.

Figure 5. **Toy community model 4**. **(a)** Toy community consisting of three organisms.
sp1 and sp2 are the same as in Toy model 3 and sp3 is a copy of sp2. sp2 and sp3 both
depend on B produced by sp1, and sp1 depends on A produced by either or both of
sp2 and sp3. **(b)** Growth rates and the cross-feeding reaction rates for each species
under different relative abundance distributions. Note the symmetry in the column
of ternary plots for sp1 and the antisymmetry between the columns for sp2 and sp3.

Figure 6. **Workflow of analysis of co-culture of** *C. reinhardtii* **and** *S. cerevisiae* **using NECom.**

a

b

NECom	Joint-FBA	NECom	Joint-FBA

Specific growth rates h^{-1}

Glucose:0.225 Nitrite:0.025 — RMSE:0.0025,0.0136 / RMSE:0.0017,0.0470

Glucose:0.525 Nitrite:0.075 — RMSE:0.0012,0.0069 / RMSE:0.0095,0.0569

Glucose:0.375 Nitrite:0.025 — RMSE:0.0013,0.0102 / RMSE:0.0023,0.0425

Glucose:0.625 Nitrite:0.075 — RMSE:0.0016,0.0034 / RMSE:0.0091,0.0433

Glucose:0.525 Nitrite:0.025 — RMSE:0.0011,0.0023 / RMSE:0.0028,0.0392

Glucose:0.725 Nitrite:0.075 — RMSE:0.0048,0.0151 / RMSE:0.0076,0.0551

Glucose:0.675 Nitrite:0.025 — RMSE:0.0058,0.0190 / RMSE:0.0036,0.0578

Glucose:0.825 Nitrite:0.075 — RMSE:0.0110,0.0266 / RMSE:0.0049,0.0538

Glucose:0.675 Nitrite:0.175 — RMSE:0.0012,0.0069 / RMSE:0.0095,0.0569

Glucose:0.725 Nitrite:0.225 — RMSE:0.0048,0.0151 / RMSE:0.0076,0.0551

Glucose:0.775 Nitrite:0.175 — RMSE:0.0016,0.0034 / RMSE:0.0091,0.0433

Glucose:0.725 Nitrite:0.325 — RMSE:0.0110,0.0266 / RMSE:0.0049,0.0538

Relative abundance of *C. reinhardtii*

c

S. cerevisiae Growth Rate(h^{-1})

Experimental derived	-0.021
NECom	-0.030
Joint-FBA	0.013

C. reinhardtii Growth Rate(h^{-1})

Experimental derived	0.002
NECom	0.004
Joint-FBA	0

O$_2$ 0.304
CO$_2$ 0.304

L-glutamine 0.003
L-arginine 0.130
Ethanol 0.990
Formate 0.141

O2 1.958
Succinate 0.197
L-leucine 0.0008
Hypoxanthine 0.0003
L-histidine 0.0003

Acetate 0.990
CO2 1.268
NH4 0.250
L-ornithine 0.128

metabolites exchange in mmol/(community)gDWh

Uptake rate of glucose: 0.225 mmol/(*S. cerevisiae*)gDWh
Uptake rate of nitrite: 0.025mmol/(*C. reinhardtii*)gDWh
Abundance Ratio(*S. cerevisiae*: *C. reinhardtii*)=9:31

Joint-FBA · **NECOM** · **Reduction**

| S. C | 0.0480 | 0.0088 | ▼ 81.7% |
| C. R | 0.0085 | 0.0031 | ▼ 63.5% |

Root mean squared error(RMSE) of predictions

Experimental mean growth rates (1/hr)

○ Measurement derived data (*S. cerevisiae*)
○ Measurement derived date (*C. reinhardtii*)
● Simulation data(*S. cerevisiae*)
✴ Simulation data(*C. reinhardtii*)

Figure 7. **Verification of NECom and comparison with Joint-FBA. (a)** Distribution of experimental and predicted cases in terms of the substrate uptake rates. Hallow circles represent the 3221 experimental cases (see Supplementary Note, noting that an axis of abundance is not presented in the figure, one circle can represent multiple cases with different abundances). Glucose and nitrite uptake rates were estimated based on the elemental composition of biomass and time series of cell counts (see Supplementary Note). (Dark and light) blue filled circles represent 488 simulation cases, which represent the experimental cases and with representativeness scores assigned (see Supplementary Note). **(b)** Simulation results of example cases (dark blue filled circles in (a)) using NECom and Joint-FBA compared with experimental cases. Rooted squared mean error (RSME) are calculated in each case. Weighted average RSMEs for both simulation methods are shown at the bottom. **(c)** Example demonstrating the difference of prediction between NECom and Joint-FBA in terms of predicted cross-feeding metabolites and its associated fluxes. RSMEs are labeled.

Figure 8. **Mass and energy flow analysis. (a)** Experimental and simulation data distribution. Hollow circles and light blue filled circles are set up in the same way as Figure 7a. Brown filled circles represent NECom predictions at various *C. reinhardtii*'s relative abundances, 631 cases in total (see **Supplementary Note**). Dark blue asterisks represent simulation cases for ATP analysis shown in (b). For each point, simulations were performed at two different *C. reinhardtii* abundances (0.725 and 0.925). **(b)** ATP consumed for the cellular maintenance of *C. reinhardtii* and *S. cerevisiae* as a function of the glucose uptake rate for *S. cerevisiae* with the nitrite uptake rate and abundance of *C. reinhardtii* fixed at different values. **(c)** Average specific growth rate of *C. reinhardtii* and *S. cerevisiae* from experimental data (represented by the hallow circles, which are grouped into five intervals of glucose uptake rate, each interval includes 20% of the experimental data. **(d)** Weighted

average specific growth rate of *C. reinhardtii* and *S. cerevisiae* over simulations
represented by light blue filled circles (488 cases in total). The weight is the
representativeness score explained in **Supplementary Note**. **(e - h)** NECom-
predicted distribution of elemental flows and energy flows, using simulation cases
(represented by brown filled circles in **a**) covering a wide range of glucose and nitrite
uptake rates where experimental cases are sparsely distributed.

Figure 9 **Shadow price based limiting substrate analysis**. Simulation cases referred in this figure are those represented by brown filled circles in Figure 8a. **(a)** Growth rates of *C. reinhardtii* and *S. cerevisiae* as functions of the relative abundance of *C. reinhardtii*, at medium to high glucose uptake rates. **(b - c)** Prevalence and average flux of each cross-feeding metabolites across all simulation cases. **(d - e)** Average shadow price of major substrate uptake (including cross-feeding metabolites) by *C. reinhardtii* and *S. cerevisiae* under various nitrite and glucose uptake rates. **(f - g)** Shadow prices and supply rates of cross-feeding CO_2 for *C. reinhardtii*, and ammonia for *S. cerevisiae* as functions of the relative abundance of *C. reinhardtii*.