Within-population genome size variation is mediated by multiple genomic

elements that segregate independently during meiosis

Authors: Claus-Peter Stelzer[1,2], Maria Pichler[1], Peter Stadler[1], Anita Hatheuer[1] & Simone Riss[1]

[1] University of Innsbruck, 5310-Mondsee, Austria

[2] Corresponding Author

University of Innsbruck, Research Department for Limnology, Mondseestr. 9, 5310 Mondsee, Austria

Tel.: +43 512507 50201

Fax: +43 512507 50299

Email: claus-peter.stelzer@uibk.ac.at

ORCiD: https://orcid.org/0000-0002-6682-0904

Abstract

Within-species variation in genome size has been documented in many animals and plants. Despite its importance for understanding eukaryotic genome diversity, there is only sparse knowledge about how individual-level processes mediate genome size variation in populations. Here we study a natural population of the rotifer *Brachionus asplanchnoidis* whose members differ up to 1.9-fold in genome size, but were still able to interbreed and produce viable offspring. We show that genome size is highly heritable and can be artificially selected up or down, but not beyond a minimum diploid genome size. Analyses of segregation patterns in haploid males reveal that large genomic elements (several megabases in size) provide the substrate of genome size variation. These elements, and their segregation patterns, explain the generation of new genome size variants, the short-term evolutionary potential of genome size change in populations, and some seemingly paradoxical patterns, like an increase in genome size variation among highly inbred lines. Our study suggests that a conceptual model involving only two variables, (1) a minimum genome size of the population, and (2) a vector containing information on additional elements that may increase genome size in this population (size, number, and meiotic segregation behaviour), can effectively address most scenarios of short-term evolutionary change of genome size in a population.

Introduction

Despite an exponential increase of genomic information during the last two decades, there is still no consensus about the ultimate causes of genome size variation in eukaryotes [1-3]. At the core of this controversy is the puzzling genome size variation across eukaryotic taxa, spanning approximately five orders of magnitude. Genome sequencing has revealed that this variation is primarily caused by gene-, chromosome-, or genome duplications, by variation in the length of introns, number of transposons, and the amount of simple repetitive DNA [4-7]. Since most of these sequences make no substantial contribution to the phenotype, at least not through their information content, genome size is only a poor predictor of organismal complexity [1]. On the other hand, ubiquitous correlations of genome size and cell size, or other phenotypic traits such as metabolic- or developmental rates, and body size [8-12], suggest the sheer amount of DNA in a genome can affect the phenotype[13]. Collectively, this might explain why current theories on genome size variation in eukaryotes differ strongly in their emphasis on selection, mutation and drift [2, 3, 13-16].

Theories invoking selection for increased genome size have been criticized for assuming causal links behind the correlations between genome size and phenotypic traits [2]. Indeed, such correlations often involve species that have been separated for long evolutionary timespans and thus differ in many other aspects than genome size. According to the mutational hazard hypothesis, non-coding DNA is never beneficial, but it may accumulate as a consequence of genetic drift [17]. Thus, this hypothesis offers an alternative, neutral explanation to the observed genome size - phenotype correlations by stating that the accumulation of non-coding DNA in organisms with large body size might be due to their smaller effective population sizes [17, 18]. Testing whether large genome size can sometimes be beneficial, or whether it is at least conditionally deleterious, ideally requires a model system that exhibits substantial genome size differences across a relatively homogeneous genomic background. This requirement appears to be best fulfilled in species with intraspecific genome size variation, where individuals share their genomic background and evolutionary history.

Cases of intraspecific genome size variation are well-documented in plants [19], with cultivated maize and its close relatives being probably one of the best-studied examples [20, 21]. In animals, intraspecific genome size variation has been found in snapping shrimp [22] and in grasshoppers [23]. Interestingly, two intensively studied model species with comparably small genomes, *Arabidopsis thaliana* and *Drosophila melanogaster,* have also turned out to exhibit substantial levels of intraspecific genome size variation [24, 25], suggesting that this phenomenon might be more widespread than previously assumed. Intraspecific genome size variation is sometimes associated with variation in chromosome numbers, for instance due to supernumerary

(B-)chromosomes [19], but there are also documented cases where genome size variation is not

reflected the karyotype [22, 26].

Despite its importance, surprisingly little is known about the basic mechanisms and

inheritance of intraspecific genome size variation, and the links to population-level phenomena, i.e.:

What characterizes those parts of a genome that account for the differences in genome size between

individuals? How is genome size inherited by offspring? How (fast) can the trait 'genome size' change

through generations, e.g., if it is directionally selected? Even in the best-studied systems, researchers

typically rely on assumptions and draw analogies to models of quantitative genetic variation. For

example, the trait 'genome size' is often considered a quantitative trait influenced by a large number

'loci', with 'alleles' that increase or decrease genome size [27, 28]. We are not aware of any empirical

study that has yet assessed the appropriateness of such a model.

Here we study the basic mechanisms of genome size variation in a population of the rotifer

Brachionus asplanchnoidis. This species is characterized by an almost two times larger genome size

relative to its sister species, by a high 44% genomic content of repetitive elements [29], and by

intraspecific genome size variation [30, 31]. Using crossbreeding experiments, selfed lines, and

artificial selection, we disentangle the basic mechanism by which variation for genome size is

mediated in this population, and how it is inherited by offspring. We capitalize on several advantages

of our model system, such as short generation times, sexual and asexual reproduction, and a haploid-

diploid lifecycle, which allows us to probe into meiotic patterns associated with intraspecific genome

size variation. We could identify the size and number of individual elements that contribute to

increases in genome size, and thus account for the gradual differences among individuals in the

population. Our findings suggest that intraspecific genome size variation can be conceptualized in

terms of a 'minimum genome size' (representing the smallest genomes in a population), and

additional 'elements' found in individuals with larger genomes (each characterized by size, number,

and meiotic segregation behaviour). This difference in perspective, compared to a conventional

quantitative trait-model, has significant implications on the genome size distribution of populations,

and the short-term evolutionary potential of the trait 'genome size'.

Results

Within-population genome size variation in _B. asplanchnoidis_

To quantify genome size variation within populations, we examined 118 _B. asplanchnoidis_ clones sampled from four geographic populations (Fig. 1, Supplementary table 1). Of these, two Austrian populations from 'Obere Halbjochlacke' (OHJ, 74 clones) and 'Runde Lacke' (RL, 29 clones) show highly significant intrapopulation variation (OHJ: ANOVA $F_{52,194}$ = 112, P < 0.001; RL: $F_{28,84}$ = 23.14, P < 0.001). The OHJ-population spans a genome size range of 1.33-fold, from 414 to 552 Mbp (Fig. 1a), with one outlier at 792 Mbp (i.e., 1.91-fold). The RL-population is also variable (Fig. 1b), spanning a range of 1.24-fold across all sampled clones. In contrast, the Lake Nakuru population (11 clones sampled) was not significantly variable ($F_{10,21}$ = 8.64, P = 0.078), and most genomes were close to 420 Mbp (Fig 1c). Previously, we reported two conspecific clones isolated from a Mongolian lake, which have relatively large genome sizes of 652 and 732 Mbp (Fig. 1c, data from Riss et al. 2016). Genome size is mitotically stable, since the genome sizes of clones, as well as the differences among clones, were highly reproducible over a period of more than five years (i.e., approximately 600 asexual generations, Supplementary Fig. 1).

Inheritance of within-population genome size variation

Clones with divergent genome size can mate with each other and produce viable and fertile offspring, which are intermediate in genome size between their parents but show some variation (Supplementary Fig. 4, see also [32]). Genome size responds to artificial selection with extremely high heritability. We applied truncation selection to the OHJ population by crossing clones with the 10% largest genome sizes among each other (excluding the outlier clone at 792Mbp), and by crossing clones with the 10% smallest genomes, respectively. In the up-selection treatment, we obtained genome sizes exceeding the range of the parental OHJ population (Fig. 2). We could select genome sizes of up to 640 Mbp, with a narrow-sense heritability h^2 of 0.905 in the first generation, and 0.912 in the second generation. Likewise, heritability was high in the first generation of the down-selection treatment (h^2 = 0.924). However, in contrast to selection for large genome size, it did not extend the range of the original population. In fact, we could not select genome sizes below 414 Mbp. Additionally, h^2 of the second generation of the down-selection treatment collapsed to zero. A parent-offspring regression including all our crosses yields an overall estimate for h^2 of 0.96 (Supplementary Fig. 5).

Combining all available genome size estimates of *B. asplanchnoidis* reveals a positively

skewed distribution, with a high number of observations at 410-430 Mbp and an elongated tail of

large genome sizes (Supplementary Fig. 6). The striking absence of genome sizes smaller than 410

Mbp, despite high sampling effort and intentional selection for small genome size, suggests that this

genome size might be a true biological limit. By contrast, there was no constraint in terms of

increases in genome size, as suggested by the more or less continuous rise of genome sizes to 792

Mbp. Interestingly, this clone with the largest genome size was not artificially selected, but hatched

from a resting egg of the natural population (c.f., Fig. 1a).

To get additional insights into the inheritance of genome size, we analysed two selfed lines

that were derived from a clone with large and small genome size, respectively. Theoretically, one

would expect trait variation to decrease upon selfing as this causes 50% reduction of heterozygosity

each generation. In contrast, we found that genome size variation remained high in a selfed line that

descended from a large-sized clone (524 Mbp). Genome sizes of selfed offspring ranged between

values of 522 Mbp and 644 Mbp, representing a 23% increase (Fig. 3). Among-clone variation in the

large selfed line was statistically significant, even after three generations of selfing (ANOVA $F_{9,24}$ =

53.61, $P < 0.001$). Variation in the large line was also significantly higher than in the line descending

from the small genome ($p < 0.001$; R package *cvequality*, version 0.2.0, [33]). After three generations

of selfing, we performed a sexual cross between both lines. As was the case in the cross between two

natural clones, the cross of the selfed lines yielded offspring that were variable and intermediate

between their parents (Fig. 3).

Evidence for independently segregating genomic elements

To gain mechanistic insights into segregation of genome size variation during meiosis, we

analysed haploid rotifer males. In a diploid organism with equally sized chromosome pairs and no

extra-chromosomal elements, all meiotic products should contain exactly half the DNA of a diploid

cell. Consistently, haploid males of the rotifer *B. calyciflorus* have half the genome size of females

([34]; see Appendix S2 in this publication). We also find this pattern in some of our *B. asplanchnoidis*

clones (e.g. Fig. 4a). However, in many others we obtained striking variation in male genome size,

which manifested in multiple discrete "male peaks" (Fig. 4b-h, Supplementary table 2). In the

simplest case, we observed two male peaks spaced symmetrically around the expected 1C-value (Fig.

4b). In general, the male peak pattern of a clone could be characterized by (i) the number of peaks,

(ii) an odd/even number of peaks, indicating the presence/absence of a central male peak at exactly

1C, and (iii) the relative abundance of certain male genome size classes, as inferred from the area

under each male peak. In most cases, the central male genome sizes (the ones close to 1C) were

more abundant than the ones in the periphery (Supplementary Fig. 7, Supplementary table 2).

Overall, male peak patterns were clone-specific and highly repeatable (Supplementary Fig. 8).

In some clones, we could not resolve individual male peaks, but instead obtained an

extremely broad distribution of genome sizes around the 1C value (e.g., Fig 4f). This pattern likely

reflects real genome size variation among males of a clone, rather than a lack of measurement

precision, since the coefficient of variation of male genome sizes in this example was 12%, while

variation of the female peak ~2%. Interestingly, this clone with the extremely broad male peak was

the same as the "outlier" with the largest genome size of 792 Mbp (Fig. 1a). Although this clone was

the extreme case, we obtained broad distributions of male genome sizes in other clones as well (data

not shown). Finally, in some clones we observed characteristically unequal distances between male

peaks (Fig. 4 g,h), such that some male peaks were more narrow together than others.

All the subsequent results and analyses build on the following *ad hoc* hypothesis, which is

based on our observations so far: We hypothesized that large genomic elements, several Mbp in size,

are segregating independently from each other, and thus give rise to discrete classes of male genome

size. By independent, we mean that each element has an approximately equal chance of segregating

into any of the four gametes during gametogenesis. Note that this is different from normal

chromosome segregation, where homologous chromosomes will always end up in opposite gametes.

The presence of one independently segregating element can explain the simplest pattern of two

male peaks (Fig 4b), being present in the large male genome size but absent in the small one.

Similarly, two equally sized elements can explain a pattern of three male peaks (Fig. 4c), which

correspond to genome size classes with zero, one, or two elements. Likewise, patterns with higher

numbers can be explained by *n*-1 elements. Assuming that independently segregating elements have

identical size, we can predict the relative ratios of the male genome size classes to 1:1 (one element),

1:2:1 (two elements), or 1:3:3:1 (three elements). Indeed, some of our clones closely follow these

predicted frequencies, whereas others showed some deviations (Supplementary fig. 7). For example,

in clones with two male peaks, the smaller male genome size was often at higher frequency than the

expected value of 0.5 (Supplementary fig. 7), suggesting that males without an element were more

frequent among the hatched males. Likewise, in clones with three and four male peaks, the central

male genome sizes were sometimes at a higher frequency than expected. One clone showing four

male peaks with almost identical heights (Supplementary fig. 7c) presents a particularly interesting

deviation. In this same clone, the two central peaks were closer together (cf. Fig. 4g). The most

parsimonious explanation seems to be that this clone carries two differentially sized elements, a

small and a large one, and that the four male peaks correspond to: (1) zero elements, (2) the small

element, (3) the large element, and (4) both elements. Likewise, two large and one small element

can explain the "three double-peaks" pattern in the clone depicted in Fig. 4h.

With this in mind, we estimated the size and number of independently segregating elements,

based on the distance between male peaks in a clone and its 2C genome size (Fig. 5a). We find that

the natural OHJ-population harbours a large diversity of elements (Fig. 5b). Many clones contain

elements of ~34Mbp size, but smaller elements down to 15 Mbp were also present in this

population. In contrast, the clones of our "large" selfed line apparently contained only elements in

the 34 Mbp size range (Fig. 5b), and they exhibited significantly less variation in element size than the

natural population (p<0.001; R package *cvequality*, version 0.2.0, [32]). Interestingly, the founding

clone of this line shows four male peaks (Fig. 4d), and indicated three elements of ~34Mbp size. Thus,

it appears that this same 34Mbp element causes all the observed genome size variation in the "large"

selfed line, being present in different numbers in different offspring.

Link between independently segregating elements and genome size variation

The hypothesis of independently segregating elements offers an explanation to many

patterns in our data. However, does it provide a *sufficient* explanation for within-population genome

size variation? To explore this idea, we formalized our findings in a quantitative model of multiple

independently segregating elements (for details and code, see Supplementary Methods). We

assumed a minimum diploid genome size at 414 Mbp and that each additional element in a clone

would proportionally increase its genome size. Thus, the main input variable of the model is a vector

specifying the size and number of elements in a clone. We assumed that all elements segregate

completely independently from each other. Thus, if a clone contains, for example, five elements,

there is a small chance that it produces males with zero or five elements, while the majority of males

will contain two or three elements. In the model, we also defined a parameter for the precision of

the flow cytometry measurement, in terms of the coefficient of variance (CV). We set its default

value to 2.7%, reflecting the overall mean in our male peak data (Supplementary table 3), but we also

explored a CV of 2%, which we obtained in some of our best samples. On the whole, our model can

reproduce virtually all male peak patterns found in *B. asplanchnoidis*, and it illustrates some

technical issues, e.g., how measurement precision limits the detection of the fine patterns produced

by small elements or when a mixture of different element sizes are present in a clone

(Supplementary Figs. 9, 10). We could also reproduce the peculiar case of clone OHJ72, our outlier of

the natural OHJ-population, with its extremely broad male peak. According to our model, this clone

might harbour eleven 34Mbp-elements, or even more, if these elements are smaller.

One method to test whether independently segregating elements provide a sufficient
explanation to genome size variation is to "predict" genome size of a clone based on the number and
size of its elements. Although this prediction is recursive, since it requires knowledge of female
genome size for sizing of the elements, it is not circular, because the distance between male peaks is
free to vary. Thus, if we should find that predicted and observed genome sizes differ greatly, this
would suggest that independently segregating elements are *not* a sufficient explanation for genome
size variation. To explore this idea, we analysed the crossed offspring of the two selfed lines (cf., Fig.
3). This cross was especially suitable because of their low diversity of segregating elements (Fig. 5b,
Supplementary table 2). We obtained striking agreement between predicted and observed genome
sizes, and could account for 96% of the diploid genome size, on average, just based on the number of
34Mbp elements (Fig. 6a). Nevertheless, all observed genome sizes were approximately 18 Mbp
higher than the predicted ones, indicating that there might still be some variation that the model
does not account for.

Identifying the exact size and number of independently segregating elements is not possible
for the natural OHJ population as a whole, due to the higher diversity of segregating elements in this
population (Fig. 5b), and due to limited resolution of male peak patterns in clones containing
multiple differentially sized elements (Supplementary Fig. 9). To circumvent these limitations, we
tested whether the amount of meiotically segregating genome size variation is correlated with
genome size. To this end, we defined the variable RCV (Relative Coefficient of Variation), which is the
CV of all combined male genome sizes of a clone divided by the CV of the female genome size. Thus,
RCV accounts for differences in measurement precision across samples (indicated by the CV of the
female peak), while it requires only that a clone can produce males in order to be measured, thus
extending our analyses to many more clones (n=72). We found that the RCVs of OHJ-clones were
considerably higher than those of outgroup rotifer species without genome size variation (Fig. 6b),
and that genome size and RCV were positively correlated (Spearman rank correlation, $\rho = 0.89$, *P* <
0.001). Consistently, RCVs in the "large" selfed line were significantly higher than in the "small"
selfed line (Fig. 6c; Mann Whitney U-Test, W = 88, *P* < 0.001). Our model predicted the same
curvilinear relationship between genome size and RCV as we observed in *B. asplanchnoidis*, assuming
either 20Mbp or 34 Mbp elements (Supplementary Fig. 11). In particular, the prediction based on 20
Mbp elements seemed to match the majority of clones of the OHJ population. Altogether, the
positive correlation of RCV with genome size as well as the congruence of the model predictions and
the natural OHJ-population suggest that independently segregating elements are largely responsible
for genome size variation in *B. asplanchnoidis*.

Discussion

Extensive within-population genome size variation in B. asplanchnoidis

Here we report a case of within-species genome size variation in the rotifer *B. asplanchnoidis* and show that even individuals *within* a natural population can differ by 1.33-fold, exhibiting genome sizes ranging from 414 to 552 Mbp. Moreover, artificial selection for large genome size can extend this range to 1.55-fold (640 Mbp) within just two generations. To our knowledge, these ranges represent some of the most pronounced cases of intraspecific variation in animals and plants (Supplementary Table 4).

Claims of intraspecific genome size variation have sometimes met scepticism, due to possible methodological artefacts [35], or the species concept underlying the study [36]. Thus, we used two different internal standards in our flow cytometry measurements, either *Drosophila melanogaster* flies or other rotifer clones that were co-prepared with a focal sample, to demonstrate genome size differences among rotifer clones. Both internal standards yield virtually identical results, confirming that our genome size estimates are accurate and reflect real differences among clones (Supplementary Figs. 2, 3). Previous studies have documented the species status of *B. asplanchnoidis* [32, 37], and have shown that the populations from Austria, Mongolia and Lake Nakuru are genetically distinct, but also experience natural gene flow on a larger geographic scale [32]. Thus, in the case of *B. asplanchnoidis* we can exclude any unrecognized cryptic species.

Independently segregating elements are responsible for genome size variation

Within-population variation of genome size in *B. asplanchnoidis* is mediated by multiple, large genomic elements that segregate independently during meiosis. These elements range between 15 and 34 Mbp in size and are present in variable numbers in individuals of the natural population. Thus, they account for the full range of genome size variation in this population, including the apparent "outlier" at 792 Mbp. Individuals with the smallest observed genome size are free of such elements, while individuals with larger genomes contain progressively more elements, in different numbers and combinations. Altogether, this allows for a seemingly gradual distribution of genome sizes across the whole population.

The meiotic behaviour of these elements is unusual in that they segregate independently from each other. This has important implications for maintenance of genome size variation in this population, because it facilitates the (re-)generation of a large number of genome size variants in sexual offspring, even if the two parents have identical genome size. For example, consider two

parents with three 20 MBp and two 30 Mbp elements, respectively. Both have the same genome size

(414 + 60 = 474MBp), but due to independent meiotic segregation of these elements, and

recombination, their crossed offspring can have one of ten different genome sizes and range

between 414 and 534 Mbp. There is some resemblance to allelic recombination in a genetically

controlled trait determined by multiple loci. However, in the case of genome size, variation can even

be generated in genetically highly homogeneous lines, such as selfed lines.

On a chromosome level, independently segregating elements in *B. asplanchnoidis* may

resemble supernumerary chromosomes (B-chromosomes) that do not pair during meiosis. This is

suggested by our results from the selfed line, which seems to have accumulated up to six identical

elements derived from an ancestor with three elements. If so, the B-chromosomes are quite large in

their relative size, since one 34Mbp element corresponds to 8.2% of the minimum diploid base

genome observed in this population. B-chromosomes reported in previous studies were often

smaller than normal chromosomes [38], but there are also reports where they reach 3-5% of the

diploid genome size [23, 38]. On the other hand, regarding absolute size, B-chomosomes of other

species can be twice as large the whole genome size of *B. asplanchnoidis* [23]. A second possible

mechanism is that the extra DNA is integrated into the normal set of chromosomes, thus resulting in

homeologous chromosomes of different length [e.g., 26]. According to this hypothesis, simple

patterns of two or three male peaks could be caused by one or two elements located on different

chromosomes, being present only once per chromosome pair. Unfortunately and despite much

effort, we could not obtain reliable information on the karyotypes of *B. asplanchnoidis*, which might

have allowed us to distinguish between these two possibilities, likely because cell divisions in this

organism are limited to a short period when the embryo is encapsulated in a robust eggshell.

However, even though we cannot completely rule out the second mechanism, supernumerary

chromosomes appear to be the most plausible explanation based on our selfed lines. After all, the

two mechanisms are not mutually exclusive. For example in maize, B-chromosomes and

"heterochromatic knobs" can both contribute to intraspecific genome size variation [19]. Regardless

of whether these independently segregating elements are located on supernumerary chromosomes

or integrated in the regular set of chromosomes, we can draw important general conclusions about

intrapopulation genome size variation.

Towards a general model of within-species genome size variation

The model developed here requires as parameters only a minimum genome size (for individuals free

of elements), a vector describing size and number of all elements within a clone, and information

about their meiotic segregation patterns. This model is consistent with high heritability of genome

size and the strong response to selection, while it accounts for the fact that we could not select

below a minimum genome size. It is worth mentioning that the minimum genome size that we

defined for our population was not obvious from the distribution of the natural population (Fig. 1a),

but could only be established by experimental manipulation. Interestingly, previous reports of

intraspecific genome size variation in other organisms also report positively skewed distributions (22,

34), which could be an indication of a minimum genome size. Our model could reproduce virtually all

observed meiotic segregation patterns ("male peaks"), and it is consistent with the increase of male

segregational variation (RCV) with genome size that we observe in the OHJ population (Fig. 6b).

Finally, this model can explain the unexpectedly high genome size variation in a selfed line derived

from a clone with large genome size.

Our model differs in several notable aspects from previous conceptual models on genome

size that were adopted from a perspective of quantitative trait (QT) models. Importantly, QT-models

are defined in terms of the mean genome size of a population governed by a large number of

insertion and deletion alleles (24). According to our model, QT-models thus refer to the special case

where all members of the population have an elevated genome size. However, observations like the

minimum genome size at ~414MBp in the OHJ population, or the fact that genome size variation

remained high in the selfed line, suggest that the QT-model might not capture all aspects of the

genome size dynamics. In addition, a focus on minimum genome size and DNA-additions, rather than

on deletions, is more consistent with the notion that the largest fraction in eukaryotic genomes is

excess junk DNA [39]. Nevertheless, it is important to realize that our concept of a 'minimum genome

size' is not equivalent to a minimal possible genome, but that it is operationally defined, and includes

all DNA-additions that are invariably "fixed" in a specific population, or species.

Our present model may be generalized further. Currently, it is strongly oriented towards a

mechanism that involves completely independent segregation of extra DNA, which might be best

represented in the case of B-chromosomes. However, it can be extended to include other

mechanisms, e.g., where extra genomic material is located on normal chromosomes. Accounting for

the latter would require some constraints regarding segregation patterns of extra DNA, particularly if

these elements are located on the same chromosome. Other variables and parameters, such as

minimum genome size and size of elements, would be largely unaffected. Such a generalization has

interesting consequences. For example, extra genomic material may become 'fixed' in a population

once all members carry the same extra DNA element on both chromosomes. Ultimately, a general

model would incorporate both mechanisms, integration to normal chromosomes and supernumerary

chromosomes, and perhaps even an exchange between both pools [40-43]. Thus, genome size could

be highly dynamic on short time scales, due to mechanisms that involve independent segregation,

while fixation/loss of extra genomic material on the normal set of chromosomes could explain the

long-term changes in genome size across populations or species. Overall, this model does not even

contradict long-term gradual (Brownian motion) changes in genome size, while it adds a population-

level perspective to taxa that seem to have undergone "saltations" in genome size over

macroevolutionary time scales (e.g., [44])

Conclusions

We have shown that within-population genome size variation in the rotifer *B. asplanchnoidis*

is mediated by relatively large genomic elements that segregate independently of each other during

meiosis. Our data on short-term artificial selection and inbred line variation suggest that a model that

involves only two variables, a minimum genome size and a vector specifying the number, size, and

segregation behaviour of the elements that increase genome size is sufficient for capturing most

aspects of genome size dynamics in this population. Collectively, our study closes an important gap in

our knowledge of how intraspecific genome size variation in populations is mediated by processes at

the individual level. Since the genome size variations in this model system are realized across a

relatively homogeneous genomic background, this has general implications for identifying the

evolutionary forces that are responsible for the immense variation of genome sizes seen across

eukaryotes. Most notably, future studies in this system should allow disentangling whether larger

genome size can be beneficial, or whether it is always slightly deleterious. In the future, it will also be

interesting to elucidate in more detail the mechanisms behind these independently segregating

elements, for example, their underlying genomic architecture, or how they interact with the "non-

dispensable" parts of the genome.

Methods

Resting eggs of rotifers were collected in the field from Obere Halbjochlacke (OHJ, N 47°47'11",E 16°50'31") and from Runde Lacke (N 47°47'08", E 16°47'34"), two small alkaline playa lakes in Burgenland (Austria) in 2011. Resting eggs from Lake Nakuru (Kenya) and the two Mongolian clones were obtained from colleagues and have been previously described in detail [30, 32]. All rotifers were cultured as clones, consisting of the asexual descendants of the female that initially hatched from a resting egg (for details, see Supplementary methods).

Genome size measurements were performed with flow cytometry using a detergent-trypsin method and propidium iodide (PI) staining according to [30] with minor modifications (for details, see Supplementary methods). As an internal standard of known genome size, we used the fruit fly, *Drosophila melanogaster* (strain ISO-1, diploid nuclear DNA content: 0.35 pg; [45]).

For sexual crosses between two rotifer clones, we used freshly hatched virgin females and males, which were harvested as eggs from dense rotifer cultures that had initiated sexual reproduction (for details, see Supplementary methods). To analyse inheritance of genome size within a population, we crossed two clones with divergent diploid genome size (414 versus 524 Mbp; called ohj22 and ohj7, respectively) and analysed 27 of their sexual offspring. Two selfed lines were established from the same two clones by growing mass cultures until they produced resting eggs (which were the product of self-fertilization, i.e., males fertilizing females of the same clone). Selfed lines were propagated for three sexual generations by randomly selecting one offspring clone each generation. Finally, we randomly selected one offspring clone from the "large" and "small" line and cross-mated them to produce an inter-line cross.

For the artificial selection experiments, we applied truncation selection to the natural OHJ population by crossing eight clones representing the 10% largest genome sizes among each other (excluding the outlier clone at 792Mbp), and by crossing eight clones representing the 10% smallest genomes, respectively. In the first generation, we used 14 combinations of parental clones for each selection treatment and analysed 1-9 of their offspring. In total, the F1-generation encompassed 31 and 26 offspring clones for the large and small selection-line, respectively. We repeated this selection procedure in the F1-generation to produce a F2-generation. In the small selection line, we used five clone combinations to produce 15 F2-offspring clones. In the large selection line, sexual propensity of the F1-clones was extremely low, limiting us to just one clone combination and 14 of their offspring in F2. The exact genealogy of clones in the F1- and F2-generation can be inferred from their names listed in supplementary table 1. Narrow sense heritabilities h^2 were estimated using to the "Breeder's equation" $\Delta Z = h^2 \cdot S$, where S is the selection differential, and ΔZ is the response to selection. Additionally, we calculated h^2 from the slope of the best-fit line for a plot of mid-offspring versus

mid-parent genome size of all our available data, which included all crosses and the self-fertilized

clones.

To determine the genome size of males relative to (diploid) females, we grew clones from

low to high population densities until they started to produce males. We co-prepared males and

females from the same clone and subjected them to the same protocol as above, except that we did

not use a *Drosophila* internal standard. To better visualize male peaks, we used an excess of males,

from 200 males + 100 females to 300 males + 60 females, depending on genome size of the clone. In

the flow-cytometry analysis, we quantified the following variables: number of male peaks (up to 6),

position of each individual male peak (as the median of the YL-2A value), position of the female peak,

standard deviation (YL-2A value) of all combined male peaks, and standard deviation of the female

peak. In contrast to the previous genome size measurements, we applied a slightly stricter precision

cut-off at 3.5% CV of the diploid female peak, and discarded all samples with higher CVs.

We performed two types of analyses depending on the quality our male peak data. First, in

clones that showed multiple discrete male peaks, we counted and sized these peaks (relative to the

female peak). We also determined the area under each peak, as a measure for the frequency of each

male genome size class. Second, for these and for all remaining clones, we calculated relative

coefficient of variation (RCV) according to

$$RCV = CV_{allMP}/CV_{FP}$$

, where CV_{allMP} is the CV across all male peaks and CV_{FP} is the CV of the female peak. Thus, RCV is a

measure for male genome size variation within a sample, corrected for its measurement error

(indicated by the CV of the female peak). To obtain a point of reference for RCV values in species

without intraspecific genome size variation, we conducted the same analyses in four sister species (*B.*

rotundiformis, *B. plicatilis*, *B. manjavacas*, B. 'Nevada').

Fig. 1

Fig. 1 Genome size variation in four natural *B. asplanchnoidis* populations. **a** Obere Halbjochlacke

(Austria). **b** Runde Lacke (Austria). **c** Lake Nakuru (Kenya) and Chuluutyn Tsagaan Nuur (Mongolia).

Genome size was highly variable in the two Austrian populations (genome sizes ranging from 414 to

792 Mbp), whereas clones isolated from Lake Nakuru sediments differed little from each other (426 ±

5.1 Mbp, mean and standard deviation). The two clones from the Mongolian site had distinct and

relatively large genome sizes of 652 and 732 Mbp (Data from [32]).

Fig. 2

Fig. 2 Artificial selection for increased or reduced genome size. The parental population consisted of clones isolated from the natural population from Obere Halbjochlacke (cf., Fig. 1a) and was the same for both treatments. Dashed black lines indicate population mean genome sizes for each generation. Solid lines indicate the mean genome size of the selected top 10% (in red) or bottom 10% (in blue) for the parental and F1-generation.

Fig. 3

Fig. 3 Genome size variation in two selfed lines and their inter-line cross. Two selfed lines were

established from two natural rotifer clones, one with large genome size (524 Mbp, yellow) and one

with small genome size (414 Mbp, purple), and were propagated for three sexual generations. Clones

used for breeding are indicated by triangles, their siblings are displayed as circles. After three

generations of selfing, the two lines were crossed with each other to produce hybrid offspring

(orange).

Fig. 4

Fig. 4 Representative examples of genome size variation in haploid males. Each histogram corresponds to a different clone. X-axis corresponds to the YL2-A fluorescence on a linear scale. Only the section surrounding the 1C and 2C values is shown. The 1C value was set at 0.5x the median fluorescence of the 2C (female) peak. **a** One single "male peak" (MP), indicates that male genome size is not variable. Note that the MP is located at exactly one half (=1C) of the diploid female genome size (2C). **b** Two MPs located at 0.46 and 0.54 of female genome size, indicating that there are two discrete classes of male genome size. **c** Three MPs, located at 0.44, 0.5, and 0.56. **d** Four MPs, located at 0.41, 0.47, 0.53, and 0.59. **e** Five MPs, located at 0.42, 0.46, 0.5, 0.54, and 0.58 (the MP on the far right is very faint). **f** No discrete MPs, but extremely broad distribution of male genome sizes around the 1C value. However, the 2C peak of this sample is very narrow (~2% CV), indicating that the broad MP is due to real variation in genome size rather than a lack of precision. **g** Four MPs. Note that in contrast to **d**, the MPs are approximately in the same height (i.e., relative frequency), and that there is unequal spacing between the MPs (the middle MPs are close together). **h** Six MPs with unequal spacing, showing some resemblance to the case of three MPs, but each MP appears to be split up into two sub-peaks.

Fig. 5

Fig. 5 Sizing of genomic elements responsible for male genome size variation. **a** Three examples of size inference, based on the distance between two male peaks (MPs) and knowledge of diploid genome size. Top: Example of a clone with two MPs, which can be explained by segregation of one ~34Mbp element. Males corresponding to peak 2 contain the 34 Mbp element, while males corresponding to peak 1 are lacking it. Middle: Example of a clone with 3 MPs, which can be explained by two elements of 34Mbp size: Males corresponding to peak 1 are free of elements, males of peak 2 contain one element, and males of peak 3 contain both elements. Bottom: A more complicated case with two, apparently differently sized elements (20Mbp and 35Mbp): Peak 1 corresponds to males without any element, peak 2 to males with the 20Mbp element, peak 3 to males with the 35Mbp element, and peak 4 to males with both elements. **b** Beeswarm plot showing the estimated sizes of elements in the natural population (including their outbred offspring) and in the selfed line (yellow).

Fig. 6

Fig. 6 Mechanistic explanations of intraspecific genome size variation. *a* Prediction of genome size based on the size and number of independently segregating elements, assuming a minimum genome size of 414 Mbp (i.e., free of elements). This figure shows seven clones of the inbred line cross (orange), from which we could determine the exact number of 34Mbp-elements based on their MP patterns. The two parental clones, one with four elements (yellow) and the other with zero elements (purple), are also shown. **b** Genome size *vs*. "relative coefficient of variation" (RCV) in clones of the natural population (Fig. 1a) and their outbred descendants. We used RCV as a proxy for the amount of genomic material that independently segregates during meiosis. RCV was calculated by dividing the CV of all combined male genome sizes (in a clone) by the CV of the female genome size. Horizontal dashed lines are the 95% confidence intervals of the RCV in four outgroup species, which have no intraspecific genome size variation (*B. rotundiformis*, *B. plicatilis*, *B. manjavacas*, B. 'Nevada'). **c** Genome size *vs*. RCV in selfed lines. Selfed line with small genome size in purple, line with large genome size in yellow, and the cross between the two inbred lines in orange.

References

1.	Gregory TR. Synergy between sequence and size in large-scale genomics. Nature Reviews Genetics. 2005;6(9):699-708. doi: 10.1038/nrg1674. PubMed PMID: WOS:000231539300013.

2.	Lynch M. The origins of genome architecture. Sunderland, MA: Sinauer; 2007.

3.	Cavalier-Smith T. Economy, speed and size matter: Evolutionary forces driving nuclear genome miniaturization and expansion. Annals of Botany. 2005;95(1):147-75. doi: 10.1093/aob/mci010. PubMed PMID: WOS:000226370900010.

4.	Elliott TA, Gregory TR. What's in a genome? The C-value enigma and the evolution of eukaryotic genome content. Philosophical Transactions of the Royal Society B-Biological Sciences. 2015;370(1678). doi: 10.1098/rstb.2014.0331. PubMed PMID: WOS:000360552800011.

5.	Bennetzen JL, Ma JX, Devos K. Mechanisms of recent genome size variation in flowering plants. Annals of Botany. 2005;95(1):127-32. doi: Doi 10.1093/Aob/Mci008. PubMed PMID: WOS:000226370900008.

6.	Bennetzen JL, Wang H. The Contributions of Transposable Elements to the Structure, Function, and Evolution of Plant Genomes. Annual Review of Plant Biology. 2014;65(1):505-30. doi: 10.1146/annurev-arplant-050213-035811.

7.	Kidwell MG. Transposable elements and the evolution of genome size in eukaryotes. Genetica. 2002;115(1):49-63. doi: 10.1023/A:1016072014259. PubMed PMID: 12188048.

8.	Gregory TR. The bigger the C-value, the larger the cell: Genome size and red blood cell size in vertebrates. Blood Cell Mol Dis. 2001;27(5):830-43. doi: DOI 10.1006/bcmd.2001.0457. PubMed PMID: WOS:000172785800005.

9.	Gregory TR. Genome size and developmental complexity. Genetica. 2002;115(1):131-46. PubMed PMID: WOS:000176413900010.

10.	Beaulieu JM, Leitch IJ, Patel S, Pendharkar A, Knight CA. Genome size is a strong predictor of cell size and stomatal density in angiosperms. New Phytologist. 2008;179(4):975-86. doi: 10.1111/j.1469-8137.2008.02528.x.

11.	Realini MF, Poggio L, Cámara-Hernández J, González GE. Intra-specific variation in genome size in maize: cytological and phenotypic correlates. AoB Plants. 2016;8:plv138-plv. doi: 10.1093/aobpla/plv138.

12.	Dufresne F, Jeffery N. A guided tour of large genome size in animals: what we know and where we are heading. Chromosome Res. 2011;19(7):925-38. doi: 10.1007/s10577-011-9248-x. PubMed PMID: WOS:000300089700007.

13.	Bennett MD. Nuclear DNA Content and Minimum Generation Time in Herbaceous Plants. Proceedings of the Royal Society of London Series B Biological Sciences. 1972;181(1063):109-35. doi: 10.1098/rspb.1972.0042.

14.	Petrov DA. Mutational Equilibrium Model of Genome Size Evolution. Theor Popul Biol. 2002;61(4):531--44. doi: 10.1006/tpbi.2002.1605. PubMed PMID: Petrov2002.

15.	Hessen DO. Noncoding DNA as a Phenotypic Driver. Evolutionary Biology. 2015;42(4):427-31.

16.	Hessen DO, Jeyasingh PD, Neiman M, Weider LJ. Genome streamlining and the elemental costs of growth. Trends Ecol Evol. 2010. doi: 10.1016/j.tree.2009.08.004. PubMed PMID: Hessen2010.

17.	Lynch M, Conery JS. The origins of genome complexity. Science. 2003;302(5649):1401-4. PubMed PMID: ISI:000186683500054.

18.	Lynch M, Bobay L-M, Catania F, Gout J-F, Rho M. The Repatterning of Eukaryotic Genomes by Random Genetic Drift. Annual review of genomics and human genetics. 2011;12:347-66. doi: 10.1146/annurev-genom-082410-101412. PubMed PMID: PMC4519033.

19.	Smarda P, Bures P. Understanding intraspecific variation in genome size in plants. Preslia. 2010;82(1):41-61. PubMed PMID: ISI:000277220000004.

20.	Díez CM, Gaut BS, Meca E, Scheinvar E, Montes-Hernandez S, Eguiarte LE, et al. Genome size variation in wild and cultivated maize along altitudinal gradients. New Phytologist. 2013;199(1):264-76. doi: 10.1111/nph.12247.

21.   Chia J-M, Song C, Bradbury PJ, Costich D, de Leon N, Doebley J, et al. Maize HapMap2
identifies extant variation from a genome in flux. Nature Genetics. 2012;44:803. doi:
10.1038/ng.2313

https://www.nature.com/articles/ng.2313#supplementary-information.
22.   Jeffery NW, Hultgren K, Chak STC, Gregory R, Rubenstein DR. Patterns of genome size
variation in snapping shrimp. Genome. 2016. doi: 10.1139/gen-2015-0206.
23.   Ruiz-Ruano FJ, Ruiz-Estevez M, Rodriguez-Perez J, Lopez-Pino JL, Cabrero J, Camacho JPM.
DNA Amount of X and B Chromosomes in the Grasshoppers Eyprepocnemis plorans and Locusta
migratoria. Cytogenet Genome Res. 2011;134(2):120-6. doi: 10.1159/000324690. PubMed PMID:
WOS:000290305100007.
24.   Huang W, Massouras A, Inoue Y, Peiffer J, Ràmia M, Tarone AM, et al. Natural variation in
genome architecture among 205 Drosophila melanogaster Genetic Reference Panel lines. Genome
Research. 2014;24(7):1193-208. doi: 10.1101/gr.171546.113.
25.   Long Q, Rabanal FA, Meng D, Huber CD, Farlow A, Platzer A, et al. Massive genomic variation
and strong selection in Arabidopsis thaliana lines from Sweden. Nature Genetics. 2013;45:884. doi:
10.1038/ng.2678

https://www.nature.com/articles/ng.2678#supplementary-information.
26.   Šmarda P, Bureš P, Horová L, Rotreklová O. Intrapopulation Genome Size Dynamics in Festuca
pallens. Annals of Botany. 2008;102(4):599-607. doi: 10.1093/aob/mcn133.
27.   Bilinski P, Albert PS, Berg JJ, Birchler JA, Grote MN, Lorant A, et al. Parallel altitudinal clines
reveal trends in adaptive evolution of genome size in Zea mays. PLOS Genetics.
2018;14(5):e1007162. doi: 10.1371/journal.pgen.1007162.
28.   Šmarda P, Horová L, Bureš P, Hralová I, Marková M. Stabilizing selection on genome size in a
population of Festuca pallens under conditions of intensive intraspecific competition. New
Phytologist. 2010;187(4):1195-204. doi: 10.1111/j.1469-8137.2010.03335.x.
29.   Blommaert J, Riss S, Hecox-Lea B, Mark Welch DB, Stelzer CP. Small, but surprisingly
repetitive genomes: transposon expansion and not polyploidy has driven a doubling in genome size
in a metazoan species complex. BMC Genomics. 2019;20(1):466. doi: 10.1186/s12864-019-5859-y.
30.   Stelzer CP, Riss S, Stadler P. Genome size evolution at the speciation level: The cryptic species
complex *Brachionus plicatilis* (Rotifera). BMC evolutionary biology. 2011;11. PubMed PMID:
ISI:000290202100001.
31.   Mills S, Alcántara-Rodríguez AJ, Ciros-Pérez J, Gómez A, Hagiwara A, Galindo KH, et al. Fifteen
species in one: deciphering the Brachionus plicatilis species complex (Rotifera, Monogononta)
through DNA taxonomy. Hydrobiologia. 2016:1-20. doi: 10.1007/s10750-016-2725-7.
32.   Riss S, Arthofer W, Steiner FM, Schlick-Steiner BC, Pichler M, Stadler P, et al. Do genome size
differences within Brachionus asplanchnoidis (Rotifera, Monogononta) cause reproductive barriers
among geographic populations? Hydrobiologia. 2017;796(1):59-75. doi: 10.1007/s10750-016-2872-x.
33.   Marwick B, Krishnamoorthy K. cvequality: Tests for the Equality of Coefficients of Variation
from Multiple Groups. R package version 020 [Internet]. 2019.
34.   Stelzer CP, Schmidt J, Wiedlroither A, Riss S. Loss of sexual reproduction and dwarfing in a
small metazoan. PLoS One. 2010;5(9):e12854. doi:10.1371/journal.pone.0012854. doi:
10.1371/journal.pone.0012854. PubMed PMID: 20862222; PubMed Central PMCID: PMC2942836.
35.   Greilhuber J. Intraspecific variation in genome size in angiosperms: Identifiying its existence.
Annals of Botany. 2005;95(1):91-8. doi: 10.1093/aob/mci004. PubMed PMID:
WOS:000226370900004.
36.   Murray BG. When does intraspecific C-value variation become taxonomically significant?
Annals of Botany. 2005;95(1):119-25. doi: 10.1093/aob/mci007. PubMed PMID:
WOS:000226370900007.
37.   Michaloudi E, Mills S, Papakostas S, Stelzer C-P, Triantafyllidis A, Kappas I, et al.
Morphological and taxonomic demarcation of Brachionus asplanchnoidis Charin within the

Brachionus plicatilis cryptic species complex (Rotifera, Monogononta). Hydrobiologia.
2017;796(1):19-37. doi: 10.1007/s10750-016-2924-2.
38. Jones RN, Viegas W, Houben A. A Century of B Chromosomes in Plants: So What? Annals of
Botany. 2008;101(6):767-75. doi: 10.1093/aob/mcm167.
39. Doolittle WF, Brunet TDP. On causal roles and selected effects: our genome is mostly junk.
BMC Biology. 2017;15(1):116. doi: 10.1186/s12915-017-0460-9.
40. Jones N, Houben A. B chromosomes in plants: escapees from the A chromosome genome?
Trends Plant Sci. 2003;8(9):417-23. doi: 10.1016/S1360-1385(03)00187-0. PubMed PMID: 13678908.
41. Houben A, Banaei-Moghaddam AM, Klemme S, Timmis JN. Evolution and biology of
supernumerary B chromosomes. Cellular and Molecular Life Sciences. 2014;71(3):467-78. doi:
10.1007/s00018-013-1437-7.
42. Navarro-Domínguez B, Ruiz-Ruano FJ, Cabrero J, Corral JM, López-León MD, Sharbel TF, et al.
Protein-coding genes in B chromosomes of the grasshopper Eyprepocnemis plorans. Scientific
Reports. 2017;7:45200. doi: 10.1038/srep45200

https://www.nature.com/articles/srep45200#supplementary-information.
43. Araujo S, Pompolo SG, Perfectti F, Camacho JPM. Integration of a B chromosome into the A
genome of a wasp. Proceedings of the Royal Society B-Biological Sciences. 2001;268(1472):1127-31.
PubMed PMID: WOS:000169356900004.
44. Liedtke HC, Gower DJ, Wilkinson M, Gomez-Mestre I. Macroevolutionary shift in the size of
amphibian genomes and the role of life history and climate. Nature Ecology & Evolution.
2018;2(11):1792-9. doi: 10.1038/s41559-018-0674-4.
45. Gregory TR. Animal Genome Size Database. 2019:http://www.genomesize.com.
46. Guillard RRL, Ryther JH. Studies of marine planktonic diatoms. I. *Cyclotella nana* Husted, and
Detonula confervacea Cleve. Canadian Journal of Microbiology. 1962;8:229-39.

Summary of all supplementary information

Supplementary information

Supplementary Figures 1–11, Supplementary Table 4, and Supplementary Methods.

Supplementary file

Supplementary table 1: Genome size estimates of all clones used in this study; 2: Clones used for sizing of independently segregating elements; 3: Clones used for RCV (relative coefficient of variance) calculations.

Acknowledgements

We thank Alois Herzig for providing access to the sampling sites and help during sampling of Obere Halbjochlacke and Runde Lacke. Sediments from Lake Nakuru were kindly supplied by Steve Omondi and Michael Schagerl. Julie Blommaert helped in preparation of media in some of the experiments. We thank David Mark Welch and Julie Blommaert for helpful comments on the manuscript.

Authors' contributions

CPS conceived the study, designed and participated in all experiments. MP and PS contributed the flow-cytometry analysis. SR contributed the line crosses and the selection experiment. AH and MP did most of the culturing work and prepared rotifer biomass for flow cytometry. CPS analysed the data and drafted the manuscript. All authors read and approved the final manuscript.

Supplementary Information

Supplementary Figures

Supplementary figure 1. Long-term constancy of genome size during clonal propagation. Time

courses of genome size for six *B. asplanchnoidis* clones over a period of > five years (i.e.,

approximately 600 asexual generations). Colours denote different clones: *ohj22* (red), *ohj7* (blue),

ohj13 (green), *mnchu24* (cyan), *mnchu008* (pink), *ohj72* (black). Each symbol represents an

independent genome size measurement.

Supplementary figure 2. Comparison of two variants of the flow cytometry method for measuring

intraspecific genome size variation. a Three clones of different genome size were co-prepared with

an internal standard of known genome size (*D. melanogaster*, strain ISO 1, 352 Mbp). **b** Pairs of the

same three clones were combined into a sample. **C** A comparison of the genome size ratios (= large

GS / small GS) determined by the second method shows that these closely matched those predicted

by the measurements of the individual rotifer clones, as determined by the first method.

Supplementary figure 3. Genome size differences among selected pairs of clones in the OHJ-population. Error bars are standard deviations based on 3-4 replicates (measured on different dates). In each replicate the clonal pairs were co-prepared and stained in the same tube. Kruskal-Wallis test: H= 8.95, $d.f.$=2, P=0.011.

697 **Supplementary figure 4. Sexual cross between a clone with large genome size and a clone with**

698 **small genome size.** Parental genome sizes (414 and 524 Mbp) are indicated by the vertical red lines.

699 The dashed red line indicates the parental mean, while the dashed black line indicates the offspring

700 mean genome size. Overall, the 27 offspring genome sizes were intermediate between their parents,

701 even though their distribution was not unimodal.

Supplementary figure 5. Narrow-sense heritability (h^2) of genome size in *B. asplanchnoidis*. Each symbol represents the mean of parents and offspring genome sizes of a family. This figure combines our results from the artificial selection experiment and crossings between clones with large and small genome size.

Supplementary figure 6. Distribution of genome sizes found in *B. asplanchnoidis*. Gray: All *B.*

asplanchnoidis clones measured so far (including natural populations, selection lines, and selfed

lines). **Green**: Only the natural OHJ population. **Red arrow** highlights genome size classes absent in

the natural population, which could be artificially selected in the laboratory.

Supplementary figure 7. Relative frequency of the different male genome sizes. a two malepeaks;

prediction (thick line) based on one independently segregating element **b** three malepeaks;

prediction based on two equally sized elements, **c** four male peaks; prediction based on three equally

sized elements. Coloured lines denote different clones. Dashed vertical line indicates 0.5x diploid

genome size.

Supplementary figure 8. Constancy of "male peak" (MP) patterns in asexually propagated clones.

The number of male peaks and their heights (relative to each other) are characteristic for a clone.

This figure shows the 2C-female peaks and the corresponding male peaks (peaks scattering around

1C) for three different clones on different dates. **Blue**: Clone *k13x110n2* (two MPs); **Green**: Clone

ohj76 (three MPs); **Orange**: Clone *ohj85* (three MPs). In some of these measurements we used

Drosophila melanogaster (**D.m.**) as internal standard. Please note that the height of MPs relative to

the 2C-female peak might differ among dates, since the number of males per female in a culture is

not always the same.

Supplementary figure 9. Model simulations of "male peak" (MP) patterns with only one type of independently segregating element. a No element; **b** one element of 34 Mbp size; **c** one element of 20 Mbp; **d** two elements of 34 Mbp; **e** three elements of 34 Mbp; **f** four elements of 34 Mbp; **g** five elements of 34 Mbp; **h** five elements of 20 Mbp. The model-parameter 'measurement precision' was set to 2.7% (i.e., the coefficient of variance) in all cases.

Supplementary figure 10. Model simulations of "male peak" (MP) patterns with different types of independently segregating elements. a two elements with 20 and 34Mbp; **b** three elements with 15, 34, and 34 Mbp; **c** three elements with 15, 20, and 34 Mbp. The model-parameter measurement precision was set to 2.7% (i.e., the coefficient of variance). **d-f** are identical to **a-c**, except that measurement precision was set to 2.0%.

Supplementary figure 11. Model simulations of the relationship of RCV and genome size. RCV
("relative coefficient of variation") was calculated by dividing the CV of all combined male genome
sizes (in a clone) by the CV of the female genome size. Red line shows model predictions for genomes
with multiple 34 Mbp elements. Blue line shows model predictions for genomes with multiple 20
Mbp elements. . The model-parameter 'measurement precision' was set to 2.7%. Dots are all
measurements of RCV in this study (cf. Fig. 6 b & c)

Supplementary Tables

Supplementary table 4. Examples of natural intraspecific genome size variation in animals and plants

Species name	Common name	1C-value (pg)	Fold-variation in GS	Reference
Zea mays	Maize	2.30	1.37	Diez et al. (2013)
Festuca pallens	Grass	2.53	1.19	Smarda et al. (2008)
Microseris douglasii	Coastal silverpuff	1.20	1.20	Smarda & Bures (2010)
Hordeum spontaneum	Wild barley	5.50	1.13	Smarda & Bures (2010)
Arabidopsis thaliana	Thale cress	0.16	1.14	Long et al. (2013)
Eyprepocnemis plorans	Grasshopper	10.39	1.19*	Ruiz-Ruano et al. (2011)
Synalpheus duffyi	Snapping shrimp	17.38	1.35	Jeffery et al. (2016)
Drosophila melanogaster	Fruit fly	0.18	1.15†	Huang et al. (2014)
Brachionus asplanchnoidis	Rotifer	0.21	1.33 (1.91)	**this study**

* assuming individuals with zero *vs*. six B-chromosomes of 0.64pg size

† variation between inbred strains

 Supplementary Methods

Rotifer cultures

Rotifers were cultured in F/2 medium [46] at 16 ppt salinity and with *Tetraselmis suecica* algae as

food source (500–1000 cells μl^{-1}). Continuous illumination was provided with daylight LED lamps

(SunStrip, Econlux) at 30–40 μmol quanta m^{-2} s^{-1} for rotifers, and 200 μmol quanta m^{-2} s^{-1} for algae.

During the experiments, rotifers were cultured at 23°C. Stock cultures were kept either at 18 °C, re-

inoculated once per week by transferring 20 asexual females to 20ml fresh culture medium, or they

were kept for long-term storage at 9°C, replacing approximately 80% of the medium with fresh food

suspension every 4 weeks.

Genome size measurements

To determine the genome size of rotifer clones, we used flow cytometry and propidium iodide

staining of nuclei. Rotifer clones were grown in 250-1000 ml flasks until their population size reached

several hundreds to a few thousand individuals. On the day before preparation, rotifers were

harvested with 60 μm sieves, and starved overnight in 11 ppt 0.2μm-filtered medium, followed by

two washes with filtered medium on the next day. For each sample, we isolated 350 females and

subjected them to the flow cytometry protocol. Briefly, rotifers were homogenized in a citrate buffer

(3.4 mM Trisodium citrate dihydrate, Nonidet P40 at 0.1% v/v, 1.5 mM Sperminetetrahydrochloride,

0.5 mM Trishydroxymethylaminomethane, pH 7.6), and were then transferred to 750 μl stock

solution in a 1 ml Dounce tissue homogenizer. Rotifers were homogenized on ice with 20 strokes

using the "tight" pestle of the homogenizer. As internal standard of known genome size we used the

fruit fly, *Drosophila melanogaster* (strain ISO-1, nuclear DNA content: 0.35 pg, [45]). After adding two

female *Drosophila* heads, the sample was further homogenized with ten strokes. Large debris was

removed by filtration through a 40 μm mesh nylon sieve. After addition of 100 μl of 0.021% Trypsin

(dissolved in stock solution) the sample was incubated for exactly 10 min at 37°C. To prevent further

degradation, 75μl of 0.25% trypsin inhibitor was added (this solution also included 0.05% RNAse A)

and the samples were incubated for another 10 min at 37°C. Finally, samples were stained with

propidium iodide at a concentration of 50 μg/ml. Stained samples were kept overnight on ice in the

dark. Flow cytometric analysis was performed on the next day on an Attune NxT® acoustic focusing

cytometer (Thermo Fisher) with an excitation wavelength of 561 nm and a custom-made 590–

650 nm bandpass filter (yellow, YL-2) for detection of propidium iodide fluorescence. Flow cytometric

data were analyzed using FlowJo software version 10.0.7r2 (FlowJo LLC). To exclude doublets (i.e.,

nuclei that pass the detector too close together, thus being recorded as a single "event") we

employed YL2-A *vs*. YL2-H gating. Coefficients of variance (CVs) of individual peaks typically ranged

between 1.5% and 4% for both *Drosophila* and rotifers. Very few measurements had CVs higher than

5%, and those replicates were discarded. Conversion from picograms DNA to base pairs were made

with the factor: 1 pg =978 Mbp [45]. In most cases, we obtained at least three replicate

measurements for a clone, usually on different days and sometimes with several weeks or months

apart.

Crossings between rotifer clones

For sexual crosses between two rotifer clones, we used freshly hatched virgin females and males,

which were harvested as eggs from dense rotifer cultures that had initiated sexual reproduction. Eggs

were detached from the females by vigorously vortexing the rotifer culture in 50-ml Falcon tubes for

ten minutes. Crossings between clones were accomplished by placing 100 female eggs and 50 male

eggs together into the same well of a 24-well plate filled with 750 µl of F/2 medium. After 24 h, when

all viable eggs had hatched and animals had time to mate, females were transferred to new wells

with fresh food suspension. After a few days, when females started producing eggs, we classified

them into asexual females, sexual male-producing females (which were unfertilized), and sexual

resting egg producing females. All resting egg producing females were isolated and stored at 7 °C in

the dark for at least 2 weeks. To induce hatching, resting eggs were incubated with food suspension

at 23 °C and high light intensities (200 µmol quanta m^{-2} s^{-1}). Usually after 48 h, the first hatchlings

started to emerge, and clonal cultures were initiated.

Theoretical model of independently segregating elements

We conceptualized a theoretical model to summarize our finding of independently segregating

genomic elements. The main purpose of this model was to test the hypothesis that these genomic

elements are *sufficient* as the mechanism of genome size variation. In particular, we examine

whether such a model is consistent with our observations of genome size variations in males from

clones of different diploid genome size. The basic input parameters and variables of the model are:

(i) an assumed minimum diploid genome size of 414 Mbp (for a clone that contains no additional

genomic elements), and (ii) a vector describing the size and number of individual elements in a clone

(e.g., [34 34 20] for two 34 Mbp and one 20 Mbp element, respectively). The output variable of the

model was the predicted diploid genome size of females, calculated as minimum genome size + the

summed contribution of all elements. The second output variable was the genome size distribution

of haploid males of this same clone (i.e., its "male-peak pattern"). Male genome sizes were

calculated as ½ minimum genome size+ the summed contribution of all elements in a particular male.

To this end, we assumed that all elements segregate completely independently from each other

during meiosis, i.e., each element having an equal chance of ending up in one out of four gametes

after the two meiotic divisions. For instance, a clone with two 34 MBp elements should produce 25%

gametes/males containing no element, 50% gametes containing one element (50%), and 25%

containing two elements. We calculated the male genome size distributions for a total of 10,000

gametes/males per clone. We also simulated measurement errors by assuming a coefficient of

variance of 2.7%. This value is based on the average precision of our flow-cytometry measurements

of male genome size variation (Supplementary table 3). We also explored CV values of 2%, which we

obtained in some of our best samples. The model was written in the MATLAB programming

environment (MathWorks®, version R2017A) with the code as follows.

```
845  function [relCV, gsFem, allGametes, allGametesERR] = ...
846      estimateRelativeCV(Bvec, measErr)
847
848  % MATLAB FUNCTION "estimateRelativeCV" for calculation of (diploid) female
849  % genome size and male genome size distribution, based on a vector
850  % specifying the size and number of independently segregating elements in a
851  % clone.
852  %
853  % Claus-Peter Stelzer (2019)
854
855  % Input variables
856  % Bvec       = [20]; % Vector of elements (contains size in MBp)
857  % measErr    = 2.7; % measurement error in flow cytometry (coefficient of
858  %                     variance in %)
859
860  % Model parameters
861  baseGS      = 207;                    % Genome size of male without elements
862  gsFem       = baseGS*2 + sum(Bvec); % Genome size of diploid female
863  nGametes    = 10000;                  % Number of gametes/males simutated
864  allGametes = nan(nGametes,1);
865
866  for i = 1 : 2 : nGametes-1
867      GameteA = baseGS;
868      GameteB = baseGS;
869      for j = 1:length(Bvec)
870          if rand>0.5
871              GameteA = GameteA + Bvec(j);
872          else
873              GameteB = GameteB + Bvec(j);
874          end
875      end
876      allGametes(i)   = GameteA;
877      allGametes(i+1) = GameteB;
878  end
879
880  % Convert coefficient of variance into standard deviation
881  maleMeanGS = gsFem/2;
882  sdMales = measErr/100*maleMeanGS;
```

```matlab
% Simulate with measurement error
allGametesERR = allGametes + sdMales*randn(size(allGametes));
mpCV = std(allGametesERR)/mean(allGametesERR)*100;
relCV = mpCV/measErr;
```