Survey of intertidal ecosystem reveals a legacy of potentially toxic elements from industrial activity in the Skeena Estuary, British Columbia, Canada

Tom Sizmur[1*], Lily Campbell[2], Karina Dracott[3], Megan Jones[1], Nelson J. O'Driscoll[4] and Travis Gerwing[2,5]

[1] Department of Geography and Environmental Science, University of Reading, Reading, UK

[2] Department of Biology, University of Victoria. Victoria, British Columbia, Canada.

[3] North Coast Cetacean Research Initiative, Ocean Wise Conservation Association, Prince Rupert, British Columbia, Canada.

[4] Department of Earth & Environmental Sciences, Acadia University, Wolfville, Nova Scotia, Canada.

[5] Ecosystem Science and Management Program, University of Northern British Columbia, Prince George, British Columbia, Canada.

*Corresponding author

Email: t.sizmur@reading.ac.uk

Abstract

Relationships between concentrations of Potentially Toxic Elements (PTEs) in estuarine sediments and their impact benthic invertebrate communities are poorly understood. We sampled and analysed PTEs in sediments and benthic invertebrates from five sites surrounding the Skeena Estuary, including sites adjacent to an abandoned cannery and a decommissioned papermill. There was no indication that sediments of the salmon cannery are polluted, but acidic sediments adjacent to the papermill contained elevated concentrations of Cd, Cr, Hg and Pb. Benthic invertebrate community assemblages confirm that sediments have recovered from prior disturbances associated with discharge of papermill sludge. Oregon pill bugs (*Gnorimosphaeroma oregonensis*), observed at all five sites, feed on the fibers associated with the papermill discharge. Thus, *G. oregonensis* are useful biomonitors for quantifying the impact of the decommissioned papermill, and similar industrial development projects, on intertidal ecosystems along the north coast of British Columbia, Canada.

Introduction

Conflict between the economic benefits of industrial development and the potential impact of such developments on the natural environment is most clearly expressed in remote communities, entirely dependent on natural capital for survival [1-3]. Anthropogenic activities have impacted the biogeochemical cycling of Potentially Toxic Elements (PTEs) in even the most remote ecosystems on our planet [4-6]. Quantifying the disruption of natural biogeochemical cycles of these elements by anthropogenic activities is difficult if the concentrations prior to industrial development are unknown and may be redundant if investigations are only conducted after environmental damage has occurred. Therefore, it is important to conduct baseline surveys to assess the bioavailability and bioaccumulation of PTEs by organisms in remote locations to predict the likely impact of further pollution on the ecosystem.

Intertidal estuarine sediments support ecologically and economically important ecosystems that provide nursery habitats to several species (some that support important commercial fisheries), and often host a diverse and abundant fauna [7-11]. Benthic invertebrates are often used as indicators of environmental pollution since they live in sediments and are prey for many commercially important fish species [12-14]. Benthic invertebrates inhabiting estuaries are inherently resistant to physical and chemical change as they have adapted to living in a dynamic environment with wide spatial and temporal ranges of chemical and physical properties, such as pH, redox potential, salinity, and particle size [15-17]. Because the chemical and physical properties of estuarine sediment are temporally and spatially dynamic, it is difficult to predict the impact of industrial development on the fate and impact of PTEs [18]. We do not currently understand the relationship between sediment properties, food web structure and PTE bioaccumulation by sediment-dwelling invertebrates well enough to predict the impact of future emissions to the estuarine environment. The objective of this study was to explore the relationship between sediment properties and the fate of PTEs from historic industrial development in the Skeena Estuary and quantify the impact of PTEs on the diversity of benthic invertebrates inhabiting intertidal sediments.

The Skeena is British Columbia's second largest river and provides important habitat for Pacific salmon populations, upon which Canadian First Nations communities rely [2, 8, 19]. The estuary and the intertidal areas towards the north of the Estuary along the coast of British Columbia are important nursery habitats for juvenile salmon [8, 19]. Coastal areas to the north of the estuary

surrounding the small port cities of Prince Rupert and Port Edward have been extensively developed

[3-5, 20]. Industrial developments include an international port, a papermill, and several canneries.

Findings of previous surveys of the benthic invertebrates inhabiting the intertidal sediment in the

Skeena estuary reveal an infaunal community that is relatively undisturbed at the estuary scale, but

which still shows the scars of historic disturbance at finer-grained scales [10, 15, 21, 22]. Amphipods

are powerful indicator species [23-27] whose high densities throughout the Skeena estuary [10, 15]

suggests that current disturbances to intertidal areas are relatively limited. Similarly, we observe 40

intertidal species in this area, including multiple species at all trophic levels within the foodweb [10,

15], and such a complex community is often associated with non-disturbed habitats [4, 28-30].

Conversely, disturbed sites are often more easily invaded by invasive species [31, 32], and

Capitellidae (Capitella capitate species complex) polychaetas are often observed in disturbed

habitats, particularly areas that have been organically enriched [4, 27, 28, 33]. Capitellidae

polychaetas have a clumped distribution within the Skeena estuary and can be locally abundant at

the scale of a $1m^2$ plot [10, 15]. Similarly, abundant but localized populations of the invasive

Cumacea *Nippoleucon hinumensis*, have been observed in the Skeena estuary [10, 15], and on other

historically impacted mudflats along BC's north coast [27]. While the universal presence and high

abundances of amphipods, coupled with the complexity of the intertidal community at the estuary

scale strongly suggest that these intertidal communities are relatively undisturbed, the locally

abundant populations of invasive species and Capitellidae polychaetas offer contradictory evidence.

These indicators of disturbance suggest that either current small-scale disturbances are occurring, or

this biological signal is a remnant of past disturbances.

Persistent localised biological signals of historic disturbances in mudflats that are currently relatively

undisturbed have been observed before on the north coast of BC [27]. To the best of our knowledge,

no current disturbances are occurring in the Skeena estuary that can explain the small-scale

indicators of disturbance observed in the biological community [10, 15, 22, 27]. Therefore, we

hypothesise that fine-grained biological indicators of disturbance in the Skeena estuary are a legacy

of past disturbances [10, 15]. It is likely that these past disturbances are at least partially related to

discharge from the papermill, which was released into the immediate near shore area (Porpoise

Bay), strongly depressing the invertebrate communities in this area during the 1970s [20, 34, 35]. As

such, we postulate that these intertidal areas have been passively recovering from this disturbance
for ~50 years.

Considerable developments have been proposed in the Skeena estuary, including oil and gas
pipelines, super-tanker routes, potash loading facilities, and a liquid natural gas (LNG) terminal. The
Skeena salmon run contributes an estimated $110 million dollars annually to the local economy [36],
so pollution of the intertidal nursery habitat of the Skeena Estuary could have devastating
consequences on both the economy and ecosystem [37, 38]. It is therefore critical to predict the
impact that future developments will have on the bioaccumulation of PTEs by benthic invertebrates
inhabiting intertidal sediments of the Skeena Estuary. This work presents a survey of intertidal
sediments and invertebrates at five locations north of the Skeena estuary to examine baseline levels
of PTE concentrations at sites that may be recovering from previous disturbances and identify
organisms that could be used to biomonitor the impact of future industrial developments.

108   Methods

109   Field sampling

Sediment cores and benthic invertebrates were collected from five locations north of the Skeena
Estuary (Fig 1) during July 2017: Cassiar Cannery (CC), Inverness Passage (IP), Tyee Banks (TB),
Papermill Bay (PB), and Wolfe Cove (WC). Cassiar Cannery (N54° 10' 40.4, W130° 10' 40.4) is a
former salmon cannery that closed in 1983 and is now an ecotourism lodge. Inverness passage (N54°
10' 05.9, W130° 09' 40.4), a mudflat ~3 km closer to the Skeena mouth than Cassiar Cannery, has
intertidal habitat similar to Cassiar Cannery, but was never directly impacted by a cannery or other
anthropogenic development. Tyee Banks (N54° 11' 59.1, W129° 57' 36.7) is located 20km upstream
from the mouth of the Skeena River on a large intertidal mudflat. At some point in the past, this area
had a small-scale sawmill operating and accumulations of sawdust and woodchips are still present in
the upper intertidal sediment. Papermill Bay (N54° 13' 59.3, W130° 17' 07.5) is a small bay located
directly adjacent to a large decommissioned papermill. Finally, Wolfe Cove (N54° 14' 33.0, W130° 17'
34.5) is a mudflat located approximately 1 km from the papermill. The papermill was closed in 2001,
ceasing all operations and discharge [39-41].

Fig 1. Map showing sites adjacent to the Skeena Estuary, British Columbia, Canada where
intertidal mudflats at Papermill Bay, Wolfe Cove, Cassiar Cannery, Inverness Passage, and Tyee
Banks were sampled during this project.

Sediments were sampled by pushing polycarbonate cores (5 cm diameter, 20 cm length) into the
mud with a rubber mallet, digging out with a spade, and wrapping with cling film, before extruding
and dividing into strata; 0-5 cm, 5-10 cm, 10-15 cm and 15-20cm. Sediment samples from each strata
were dried at 40 ℃ for 48 hours and shipped to the UK for analysis. A total of 15 cores were
collected from each site along five transects comprising three samples taken from the upper, mid
and lower shore in a random stratified sampling design [42]. Sediments from the upper, mid and
lower shore (but the same strata) were pooled and homogenized prior to analysis. At each location
that a sediment core was collected, two more cores (7 cm diameter) were taken to a depth of at
least 5 cm (well into the anoxic layer), transferred to a plastic bag in the field and then passed
through a 250μm stainless steel sieve to retain sediment-dwelling benthic infauna. This infauna
sampling strategy was supplemented by opportunistic digging in areas of the site in which infauna
were clearly present (e.g. burrow openings or surface casting) to collect macrofauna. All
invertebrates found in sufficient quantities for chemical analysis were identified to the species level
[10, 15], pooled into a single sample for each site, rinsed in deionised water, frozen, shipped to the
UK, and freeze-dried prior to analysis.

Laboratory analysis

The particle size distribution of sediments was determined using laser granulometry (Malvern
Mastersizer 3000). Sub-samples were then ground to a fine powder using an agate ball mill and
analysed for total organic carbon and nitrogen content using a Thermo Scientific Flash 2000 Organic
Elemental Analyser. Details of quality control can be found in the supporting information file.
Sediment pH was determined in a soil-water suspension after shaking with water for 15 min at a
1:10 w/v ratio based on BS7755-3.2 [43]. The total concentration of PTEs in sediments was
determined by ICP-MS (Inductively Coupled Plasma Mass Spectrometry) analysis of 0.5 g sediment
samples digested in reverse aqua regia (9 ml of nitric acid and 3 ml of hydrochloric acid) using a
MARS 6 microwave digestion system, based on EPA [44]. After preliminary analysis of a large range
of PTE concentrations, the elements selected for further investigation were Cadmium (Cd), Cobalt
(Co), Chromium (Cr), Nickel (Ni), Lead (Pb) and Zinc (Zn). Details of quality control can be found in

the supporting information file. The total concentration of mercury in sediments was determined
using thermal degradation – gold amalgamation atomic absorbance spectroscopy as outlined in EPA
[45] using a Nippon MA-3000 analyser. Details of quality control can be found in the supporting
information file. The bioavailability of PTEs to biota was estimated by extracting 2.5 g of sediments
with 25 ml of 0.05M EDTA (Ethylenediaminetetraacetic acid) at 20°C for one hour, centrifuging,
filtering and analysing PTEs (As, Cd, Co, Cr, Cu, Ni, Pb and Zn) in the extract using ICP-OES
(Inductively Coupled Plasma Optical Emission Spectrometry). The total concentrations of PTEs in
invertebrates were determined by ICP-MS analysis of 0.5 g of sample digested in slightly diluted
nitric acid (2 ml of ultra-pure water and 8 ml of nitric acid) using a MARS 6 microwave digestion
system.

Statistical analysis

The influence of site and sediment depth on sediment PTE concentrations and properties was
quantified using analysis of variance (ANOVA) and permutational multivariate analysis of covariance
(PERMANCOVA) [46, 47]. The relationship between PTE concentrations and sediment properties was
first tested using PRIMER's RELATE function [48]. This function compares two resemblance matrices
looking for any relationships. Relationships were further explored using principal component
analyses (PCA) on the variance-covariance matrix of all sediment PTE and sediment property data.
Relationships between the PTE concentrations observed in the sediments and in the collected
invertebrates were examined in several configurations using RELATE and plotted using non-metric
multidimensional (nMDS) scaling plots. A full description of the statistical analysis undertaken is
provided in the supporting information file and outputs provided in S1, S2 and S3 Tables.

Results and Discussion

Inverness Passage, Wolfe Cove and Tyee Banks are suitable reference sites

The PERMANCOVA analysis indicates that properties of sediments (pH, median particle diameter, C
and N) are significantly influenced by both site and sediment depth, with site explaining more than
50% of the observed variance in the data (S1 Table). There is no significant ($p > 0.05$) difference in

the median sediment particle diameter between Inverness Passage and Cassiar Cannery, or between
Wolfe Cove and Papermill Bay (Fig 2 and S3 Table). This observation supports our assumption that
the sediment deposited in the sites potentially contaminated by industrial emissions (Cassiar
Cannery and Papermill Bay) and their respective proximal reference sites (Inverness Passage and
Wolfe Cove) have the same geogenic origin and are subject to a similar depositional environment.
Thus, any differences in sediment chemical properties between potentially contaminated sites and
their proximal reference sites we infer is due to anthropogenic influences.

Fig 2. Sediment properties; Sediment pH (A); Median sediment particle diameter (B); Percentage
Organic Carbon content (C); and Percentage Total Nitrogen content (D) presented at four depths in
sediments sampled from Cassiar Cannery, Inverness Passage, Papermill Bay, Wolfe Cove and Tyee
Banks intertidal mudflats.

Tyee Banks has a significantly ($p < 0.05$) larger median particle size than Inverness Passage and
Cassiar Cannery and a significantly ($p < 0.05$) smaller median particle size than Wolfe Cove and
Papermill Bay (Fig 2 and S3 Table). The location of Tyee Banks is closer to the mouth of the Skeena
River and, therefore, only receives sediment from the Skeena, whereas the other four sites may also
receive sediment from the Nass River, to the north of the Skeena Estuary [49]. However, because
Tyee Banks is much further from potential emission sources, the site can be used to determine
whether proximal reference (Inverness Passage and Wolfe Cove) sites are contaminated with PTEs
from the adjacent industrial activity.

Papermill Bay sediments are contaminated with Cd, Cr, Hg and Pb

The PERMANCOVA analysis (S2 Table) indicates significant differences in sediment total and EDTA
extractable PTE concentrations between different sites, but not between different depths. We
observed significantly ($p < 0.05$) greater concentrations of As, Cr, Hg and Pb in the Papermill Bay
sediments than the proximal reference sediments at Wolfe Cove (Fig 3 and S3 Table). However, Co,
Cu and Ni concentrations were significantly ($p < 0.05$) lower in the Papermill Bay sediments (Fig 3
and S3 Table). A Geoaccumulation index, calculated following [50], using Wolfe Cove as an
uncontaminated reference site, indicates that Papermill Bay is 'unpolluted' with As, Cd, Co, Cr, Cu, Ni
and Zn, but 'unpolluted to moderately polluted' with Pb and 'moderately polluted' with Hg (Table 1).
When Tyee Banks is used as the reference site, Papermill Bay is classified as 'unpolluted to

moderately polluted' with Cd, Cr and Hg. Thus, there is evidence to suggest that the sediments in

Papermill Bay are contaminated with Cd, Cr, Hg and Pb, most likely emanating from the sludge [51]

discharged by the decommissioned papermill on Watson Island (Fig 1).

Fig 3. Total concentrations of Potentially Toxic Elements (As, Cd, Co, Cr, Cu, Hg, Ni, Pb and Zn) with
depth in sediments sampled from Cassiar Cannery, Inverness Passage, Papermill Bay, Wolfe Cove
and Tyee Banks intertidal mudflats

Table 1. Geoaccumulation index (Igeo)[a] for potentially toxic elements in sediments from Cassiar
Cannery (CC) and Papermill Bay (PB) using reference sites Tyee Banks (TB), Wolfe Cove (WC) and
Inverness Passage (IP).

Site	Reference site	As	Cd	Co	Cr	Cu	Hg	Ni	Pb	Zn
CC	IP	-0.99	-0.71	-0.78	-0.57	-0.69	-0.69	-0.67	-0.58	-0.63
	TB	-0.48	-0.33	-0.44	-0.27	-0.01	-0.13	-0.26	0.08	-0.24
PB	WC	-0.10	-0.25	-1.08	-0.32	-1.14	1.61	-0.82	0.74	-0.60
	TB	-1.51	0.39	-0.90	0.35	-0.19	0.20	-1.01	-0.02	-0.45

[a] Geoaccumulation index is calculated as $Igeo = Log_2 \frac{C_n}{1.5 \times B_n}$

when C_n = Concentration measured in sediment at site and B_n = Concentration measured in sediment at
reference (background) site.

Sites are classified as unpolluted if Igeo ☐ 0, unpolluted to moderately polluted if 0 ☐ Igeo ☐ 1, moderately
polluted if 1 ☐ Igeo ☐ 2, moderate to strongly polluted if 2 ☐ Igeo ☐ 3, strongly polluted if 3 ☐ Igeo ☐ 4,
strongly to extremely polluted if 4 ☐ Igeo ☐ 5, and extremely polluted if Igeo ☐ 5.

Discharges by papermills have previously been associated with pollution of the environment with

PTEs, including Cr, Cd, Cu, Pb and Zn [52, 53]. This pollution associated with papermill operations can

originate from a number of sources, such as metals in the wood entering the papermill, or

atmospheric deposition of metals from the smoke stacks [52]. PTE-containing materials may also be

used during the pulp making processes. For example, chromate bricks are used in the recovery

furnace of kraft paper mills to reduce chemical attack by the spent liquors [54]. However, Cr

concentrations observed in the sediments of Papermill Bay (Fig 3) are considerably lower than the

52.4 mg kg^{-1} observed in a deposit of pulp covering the sediments of a Swiss lake [55], or

concentrations of up to 197 mg kg^{-1} observed in a fibrous deposit in the Ångermanälven river
estuary, on the east coast of Sweden (Apler et al 2019).

The carbon and nitrogen content of the sediments at Papermill Bay were significantly ($p < 0.05$) and
considerably greater than all of the other sites (Fig 2 and S3 Table). It was evident from visual
inspection of the sediments themselves that they contained a high proportion of organic fibers,
presumably discharged into the bay from the papermill during the operational phase. The PTEs
observed to be elevated in the Papermill Bay sediments (Cd, Cr, Hg and Pb) all bind strongly to
organic matter in sediments [56-58] and so their presence could be due to (i) the discharge and
deposition of PTE contaminated organic material, (ii) the discharge of organic material alongside the
discharge of PTEs in contaminated effluent that later bind with the organic material, or (iii) the
discharge of organic material which binds to naturally present PTEs in the sediment and prevents
their export to the overlying water.

In contrast to the observations made in the Papermill Bay sediments, none of the PTEs analysed in
the Cassiar Cannery sediments were found to be present in significantly ($p > 0.05$) greater
concentrations than the Inverness Passage sediments (Fig 3 and S3 Table). In fact, Co and Ni
concentrations were significantly lower in Cassiar Cannery sediments than at Inverness Passage.
When Inverness Passage is used as a reference site, all the PTEs analysed can be classified by the
Geoaccumulation index as 'unpolluted' (Table 1). This observation is a clear indicator that any impact
resulting from either the former use of the Cassiar Cannery site (up until 1983) as a salmon cannery,
or its current use as an ecotourism lodge, can no longer be observed in the concentrations of PTEs in
the top 20 cm of sediment at this site. Any sediment contaminated by cannery waste at this site may
have been buried by fresh sediment, rendering the PTEs inaccessible to benthic invertebrates.

Relationship between sediment properties and total and labile PTE
concentrations

Alongside greater total concentrations of Cd, Cr, Hg and Pb in the Papermill Bay sediments,
compared to Wolfe Cove, we observed significantly ($p < 0.05$) greater EDTA extractable
concentrations of Cd, Co, Cr, Ni, Pb and Zn (Fig 4 and S3 Table), indicating a greater availability of

these elements to benthic invertebrates [59]. The significantly (p < 0.05) greater carbon content and
the slightly finer texture of the Papermill Bay sediments, compared to Wolfe Cove (Fig 2 and S3
Table) should provide more sites for PTEs to bind to [58, 60, 61]. Thus, the greater concentrations of
EDTA extractable PTEs is largely due to the Papermill Bay sediments being significantly (p < 0.05)
more acidic than the Wolfe Cove sediments. Lower pH results in greater competition between PTEs
and hydrogen ions for the binding sites on the sediment surface and leads to greater PTE lability [62,
63].

Fig 4. EDTA extractable (potentially bioavailable) Potentially Toxic Elements (Cd, Co, Cr, Cu, Ni, Pb
and Zn) with depth in sediments sampled from Cassiar Cannery, Inverness Passage, Papermill Bay,
Wolf Cove, and Tyee Banks intertidal mudflats

The RELATE revealed a statistically significant correlation between the concentration of potentially
toxic elements and the sediment physicochemical properties resemblance matrices (Rho: 0.46; p =
0.001, Permutations = 9999). The relationship between total and labile PTE concentrations and
sediment properties was further explored using PCA (Fig 5). The principal component scores plot (Fig
5B) reveals a clear separation between the Papermill Bay sediments and the proximal and remote
references sites of Wolfe Cove and Tyee Banks, but no separation between Cassiar Cannery and
Inverness Passage, re-enforcing the conclusion that the sediments of Cassiar Cannery show little
evidence of anthropogenic contamination. Principal component 2 (Fig 5), separates the Inverness
Passage and Cassiar Cannery sediments from the other three sites (Papermill Bay, Wolfe Cove, and
Tyee Banks), which is attributed to a different geochemical matrix composition at Cassiar Cannery
and Inverness Passage, as also observed by DelValls et al. [64]. Principal component 1 (Fig 5) reveals
a separation with depth in the Papermill Bay sediments. Papermill Bay sediments have greater total
and EDTA extractable concentrations of several PTEs, higher C and N, and lower pH, all of which also
increase with depth in the Papermill Bay sediments. This observation of deeper layers of sediment
with a lower pH and higher availability of PTEs indicates that contaminated sediment is overlain by
less contaminated sediment, deposited since discharge of sludge ceased. Although most benthic
invertebrates live in surficial sediments, it is important to consider the likelihood that deeper
sediment will be exposed and mobilized either by development projects [65], or by bioturbation
from deep burrowing invertebrates [66, 67].

Fig 5. Principal Components Analysis of sediment properties (pH median sediment particle
diameter, Percentage Organic Carbon content and Percentage Total Nitrogen content) and total
and EDTA extractable Potentially Toxic Elements (As, Cd, Co, Cr, Cu, Hg, Ni, Pb and Zn) in
sediments sampled 0-5 cm, 5-10 cm, 10-15 cm and 15-20cm from Cassiar Cannery, Inverness
Passage, Papermill Bay, Wolf Cove and Tyee Banks intertidal mudflats. (A) is the latent vectors for
each variable plotted in the plane of principal component one and principal component two, (B) is
the principal component scores of all the samples plotted in the plane of principal component one
and principal component two. Error bars represent the standard deviations of 5 replicate samples
taken from 5 different transects.

Sediments host a healthy invertebrate population with no

relationship between PTE concentrations in sediments and

invertebrates

Overall, 22 different taxa (listed in S4 Table, along with PTE concentrations) were observed in this

study, with Wolf Cove and Cassiar Cannery the most biodiverse sites, with 16 and 14 species

collected, respectively. These taxa represent a subset of the 40 intertidal species that are commonly

observed in the Skeena Estuary [10, 15]. Findings of previous studies in the Skeena Estuary reveal an

infaunal community predominantly dominated by cumaceans, polychaetes, oligochaetes,

nematodes, copepods, amphipods, and bivalves [15, 68]. The 40 intertidal species observed in this

area include multiple species at all trophic levels within the food web [10, 15], and such a complex

community is often associated with non-disturbed habitats [4, 28]. This complexity was also

observed in our study, as *Alitta brandti, Neries vexillosa, Glycinde picta,* ribbon worms and crabs can

act as predators, while *Macoma balthica*, isopods, amphipods, and sessile polychaete worms are

primary consumers [47, 69, 70]. We therefore provide evidence to support our previous research

that indicates that the intertidal ecosystem has been passively recovering for ~50 years from past

disturbances related to discharge from the papermill, which was released into the immediate near

shore area (Porpoise Bay), strongly depressing the invertebrate communities in this area during the

1970s [20, 34, 35] and currently exhibit a community that is relatively healthy.

When all invertebrates collected at all five sites are considered, we found no relationship between

PTE concentrations in invertebrates and either total (Rho: 0.27; p = 0.32, Permutations = 9999),

EDTA extractable (Rho: 0.24; p = 0.79, Permutations = 9999), or both total and EDTA extractable

(Rho: 0.78; p = 0.08, Permutations = 9999) PTEs in the sediment. This finding is confirmed by

inspecting a nMDS plot of invertebrate PTE loadings, which includes all organisms collected at all five

sites (Fig 6A). Bivalves of the same species (e.g. *M. balthica*, *Mytilus edulis*, and *Mya arenaria*),

collected at different sites, seem to cluster in multidimensional space. There is slightly more

separation in the PTE concentrations measured in polychaete worms (e.g. *Abarenicola pacifica*, *N.*

vexillosa, *Paranemertes peregrina*, *G. picta*, *A. brandti*, *Nephtys caeca*, *Neotrypaea californiensis*, and

Streblospio benedicti) which are the cause of much of the dissimilarity in the dataset.

Fig 6. Non-metric multidimensional scaling plots (nMDS) of invertebrate PTE concentrations and
the vector overlay (left hand side) of five intertidal mudflats along the north coast of British
Columbia, Canada. CC: Cassiar Cannery. WC: Wolfe Cove. IP: Inverness Passage. PB: Papermill Bay.
TB: Tyee Banks with (A) the entire dataset considered or (B) only including the two benthic
invertebrates found at all five sites; Baltic clams (*Macoma balthica*) and Oregon pill bugs
(*Gnorimosphaeroma oregonensis*).

The suitability of *G. oregonensis* as biomonitors of sediment PTE bioavailability

The selection of one or more of benthic invertebrate as a biomonitor of PTE pollution needs to

consider how cosmopolitan their distribution is and the pathway by which they are exposed to the

PTE [71]. Mussels and clams acquire food from the water column by suspension feeding, and clams

are also able to acquire food from surface sediments by way of their siphon. Isopods are more

motile and acquire food by ingesting organic debris, usually on the surface of the sediment.

Polychaete worms, as a class, have a more diverse range of feeding strategies, including suspension

feeding with tentacles or a mucus net, deposit feeding selectively at the surface or at depth, and

predation of other infauna [72, 73]. This large diversity of feeding strategies may have led to the

greater range of PTE concentrations observed in Fig 6A, compared to the bivalves and crustacea.

Furthermore, the soft tissues of polychaete worms may only represent exposure to PTEs in the

recent past due to large temporal fluctuations [74-76]. The whole body (including soft tissues and

shells) of bivalves may better represent a record of time-averaged bioavailability of PTEs in the water

column over the lifetime of the organism, albeit not without difficulties in interpretation [77, 78].

There were only two benthic invertebrate species observed at all five sites; Baltic clams (*M. balthica*)

and Oregon pill bugs (*G. oregonensis*). When PTE concentrations in *M. balthica*, and *G. oregonensis*

was compared to sediment concentrations, no relationship was observed with either total (Rho:

0.31; p = 0.19, Permutations = 9999), EDTA extractable (Rho: -0.0.; p = 0.48, Permutations = 9999),

or both total and EDTA extractable (Rho: 0.21; p = 0.28, Permutations = 9999) sediment PTE

concentrations. These findings contrast to numerous articles in the literature quantifying

relationships between sediment and benthic invertebrate PTE concentrations [58, 71, 79, 80]. When

we plot the PTE concentrations in *M. balthica*, and *G. oregonensis* in multidimensional space (Fig

6B), we reveal that the elemental profiles of *M. balthica*, and *G. oregonensis* contrast greatly. *M.*

Bathica collected at Papermill Bay contained lower concentrations of Cr, Co, Ni, Zn, Cd, As and Pb

than those collected at the other four sites, but higher concentrations of Cu. In contrast, *G.*

oregonensis collected at Papermill Bay have higher concentrations of Cr, Ni, Zn, and Pb than those

collected at the other four sites.

Biota-Sediment Accumulation Factors (BSAFs) were calculated by dividing the concentrations of PTEs

in the invertebrate tissues by the average concentration in the sediments from the site from which

the invertebrates were collected (average across all transects and depths). Relationships between

sediment properties and BSAFs reveal the importance of pH in explaining the difference in the

bioaccumulation of PTEs by *M. balthica*, and *G. oregonensis*. For most PTEs there is a positive

relationship between pH and the BSAF for *M Balthica* (Table 2), including a significant relationship

with the Cr BSAF (Fig 7A). This is an unexpected finding that does not have an immediately obvious

explanation. However, we observe a negative relationship between sediment pH and the BAF of

most PTEs for *G. oregonensis* (Table 2), including Cr (Fig 7B). This relationship is intuitive since metal

cations dissociate from mineral surfaces at lower pH levels [62, 63]. Because *G. oregonensis* feed on

organic material on the surface of the sediments, the concentrations of PTEs associated with acidic

sediments contaminated by papermill sludge at Papermill Bay, are more likely to be assimilated by

G. oregonensis and become bioaccumulated in their tissues. A survey of intertidal areas adjacent to

papermills on the British Columbia coastline (including the papermill on Watson island) identified *G.*

oregonensis as tolerant of papermill impacted shorelines [81]. The greater abundance of isopods

(including *G. oregonensis*) in close proximity to British Columbian papermills, also observed by Robin

et al. [82], is attributed to the provision of pulp fibers from the mill as a food source creating a

stressed environment which enables these isopods to thrive.

Fig 7. The relationship between Chromium (Cr) Biota-Sediment Accumulation Factor (BSAF) and
sediment pH for (A) Baltic clams (*Macoma balthica*) and (B) Oregon pill bugs (*Gnorimosphaeroma*
***oregonensis*) sampled at five intertidal mudflats along the north coast of British Columbia, Canada.**

Table 2. Correlation coefficients for the relationship between sediment pH and Biota-Sediment Accumulation Factors of 8 potentially toxic elements in Baltic clams (*Macoma balthica*) and Oregon pill bugs (*Gnorimosphaeroma oregonensis*) at all five sampling locations in the Skeena Estuary.

Species	As	Cd	Co	Cr	Cu	Ni	Pb	Zn
M Balthica	0.4397	0.2564	0.9162	0.8255*	-0.8177	0.9461	0.5788	0.896
G. oregonensis	-0.1744	0.3758	-0.7362*	-0.8688*	-0.1895	-0.8117	-0.2083	-0.7732*

* = p < 0.05

Conclusions

We found evidence to indicate that the discharge of papermill sludge from the decommissioned papermill on Watson Island has changed the sediment geochemistry at Papermill Bay by reducing the pH, and increasing the total and EDTA extractable concentrations of Cd, Cr, and Pb. However, the benthic invertebrate community composition confirms that the population has recovered from previous disturbance. Oregon pill bugs (*G. oregonensis*) were one of only two benthic invertebrate species observed at all five sites we visited. *G. oregonensis* are a cosmopolitan species along the north coast of British Columbia and are tolerant of sites contaminated with papermill effluent because they use the fibers discharged as a food source. Thus, we conclude here that *G. oregonensis* make an excellent candidate biomonitor species to assess recovery from the environmental impact of the papermill on Watson Island and monitor the future impacts of similar industrial developments in the region.

Acknowledgements

Sean Allen is acknowledged for fieldwork assistance. Ilse Kamerling is acknowledged for assisting C and N analysis. Anne Dudley is acknowledged for assisting ICP-OES analysis. Andy Dodson is acknowledged for assisting ICP-MS analysis. Accommodation and access to the shoreline were graciously provided by Cassiar Cannery Ecotourism lodge.

References

References

1. Parlee BL. Avoiding the Resource Curse: Indigenous Communities and Canada's Oil
Sands. World Development. 2015;74:425-36. doi:
http://dx.doi.org/10.1016/j.worlddev.2015.03.004.

2. Gerwing TG, Cox K. Erosion of trust in government consultation will impede the creation
of environmental policy. Marine Policy. 2017;83:126-7.

3. Waldichuk M, Bousfield EL. Amphipods in low-oxygen marine waters adjacent to a
sulphite pulp mill. Journal of the Fisheries Board of Canada. 1962;19(6):1163-5.

4. Pearson TH, Rosenberg R. A comparative study of the effects on the marine environment
of wastes from cellulose industries in Scotland and Sweden. Ambio. 1976:77-9.

5. Tuominen TM, Sekela MA. Dioxins and furans in sediment and fish from the vicinity of
four inland pulp and/or paper mills and one petroleum refinery in British Columbia:
Environment Canada. Conservation and Protection; 1992.

6. Yunker MB, Lachmuth CL, Cretney WJ, Fowler BR, Dangerfield N, White L, et al. Biota–
sediment partitioning of aluminium smelter related PAHs and pulp mill related diterpenes by
intertidal clams at Kitimat, British Columbia. Marine Environmental Research.
2011;72(3):105-26.

7. Kritzer JP, DeLucia M-B, Greene E, Shumway C, Topolski MF, Thomas-Blate J, et al. The
Importance of Benthic Habitats for Coastal Fisheries. BioScience. 2016;66(4):274-84. doi:
10.1093/biosci/biw014.

8. Carr-Harris C, Gottesfeld AS, Moore JW. Juvenile Salmon usage of the Skeena River
Estuary. PloS One. 2015;10(3):e0118988.

9. Schindler DE, Scheuerell MD, Moore JW, Gende SM, Francis TB, Palen WJ. Pacific
salmon and the ecology of coastal ecosystems. Frontiers in Ecology and the Environment.
2003;1(1):31-7.

10. Gerwing TG, Gerwing AMA, Macdonald T, Cox K, Juanes F, Dudas SE. Intertidal soft-
sediment community does not respond to disturbance as postulated by the intermediate
disturbance hypothesis. Journal of Sea Research. 2017;129:22-8.

11. Gerwing TG, Allen Gerwing AM, Drolet D, Barbeau MA, Hamilton DJ. Spatiotemporal
variation in biotic and abiotic features of eight intertidal mudflats in the Upper Bay of Fundy,
Canada. Northeastern Naturalist. 2015;22(12):1–44.

12. Amoozadeh E, Malek M, Rashidinejad R, Nabavi S, Karbassi M, Ghayoumi R, et al.
Marine organisms as heavy metal bioindicators in the Persian Gulf and the Gulf of Oman.
Environmental Science and Pollution Research. 2014;21(3):2386-95.

13. Gómez Gesteira JL, Dauvin J-C. Amphipods are good bioindicators of the impact of oil
spills on soft-bottom macrobenthic communities. Marine Pollution Bulletin.
2000;40(11):1017-27.

14. Gerwing TG, Hamilton DJ, Barbeau MA, Haralampides KA, Yamazaki G. Short-term
response of a downstream marine system to the partial opening of a tidal-river causeway.
Estuaries and Coasts 2017;40(3):717-25. doi: 10.1007/s12237-016-0173-2.

15. Gerwing TG, Allen Gerwing AM, Macdonald T, Cox K, Juanes F, Dudas SE. Assessing
the relationship between community dispersion and disturbance in a soft-sediment ecosystem.
Marine Ecology. 2018;39(4):e12505.

16. Scholz B, Liebezeit G. Microphytobenthic dynamics in a Wadden Sea intertidal flat–Part
II: Seasonal and spatial variability of non-diatom community components in relation to
abiotic parameters. European Journal of Phycology. 2012;47(2):120-37.

17. van Proosdij D, Milligan T, Bugden G, Butler K. A tale of two macro tidal estuaries:
differential morphodynamic response of the intertidal zone to causeway construction. Journal
of Coastal Research. 2009;56:772–6.

18. de Souza Machado AA, Spencer K, Kloas W, Toffolon M, Zarfl C. Metal fate and effects
in estuaries: A review and conceptual model for better understanding of toxicity. Science of
The Total Environment. 2016;541:268-81. doi:
http://dx.doi.org/10.1016/j.scitotenv.2015.09.045.

19. Moore JW, Gordon J, Carr-Harris C, Gottesfeld AS, Wilson SM, Russell JH. Assessing
estuaries as stopover habitats for juvenile Pacific salmon. Marine Ecology Progress Series.
2016;559:201-15.

20. Wilkes B, Dwernychuk LW. Environment studies in the marine receiving environment at
the Skeena Cellulose pulp mill, Watson Island, BC. Pulp & Paper Canada. 1991:92:10.

21. Gerwing TG. Preliminary report of intertidal research along the north coast of British
Columbia: Summer 2016. Report to the Kitsumkalum First Nations. 34 p. 2016.

22. Gerwing TG, Cox K, Gerwing AMA, Carr-Harris CN, Dudas SE, Juanes F. Depth to the
apparent redox potential discontinuity (aRPD) as a parameter of interest in marine benthic
habitat quality models. International Journal of Sediment Research. 2018;33(2):149-56.

23. Sánchez-Moyano JE, García-Gómez JC. The arthropod community, especially Crustacea,
as a bioindicator in Algeciras Bay (Southern Spain) based on a spatial distribution. Journal of
Coastal research. 1998;14(3):1119-33.

24. Thomas JD. Biological monitoring and tropical biodiversity in marine environments: A
critique with recommendations, and comments on the use of amphipods as bioindicators.
Journal of Natural History. 1993;27(4):795-806.

25. Conlan K. Amphipod crustaceans and environmental disturbance: a review. Journal of
Natural History. 1994;28(3):519-54.

26. Picone M, Bergamin M, Novelli Alessandra A, Noventa S, Delaney E, Barbanti A, et al.
Evaluation of Corophium orientale as bioindicator for Venice Lagoon: Sensitivity assessment
and toxicity-score proposal. Ecotoxicology and Environmental Safety. 2008;70(1):174-84.

27. Gerwing TG, Campbell L, Allen Gerwing AM, Allen S, Cox K, Rogers M, et al. Potential
impacts of logging on intertidal infaunal communities within the Kitimat River estuary.
Journal of Natural History. 2018;52(43-44):2833-55.

28. Pearson TH, Rosenberg R. Macrobenthic succession in relation to organic enrichment and
pollution of the marine environment. Oceanography and Marine Biology: An Annual
Review. 1978;16:229-331.

29. Allen L. The application of biodiversity indicators to infer ecosystem health in
regenerating tropical forest: University of Glasgow; 2019.

30. Zhang W, Dulloo E, Kennedy G, Bailey A, Sandhu H, Nkonya E. Biodiversity and
Ecosystem Services. Sustainable Food and Agriculture: Elsevier; 2019. p. 137-52.

31. Smith MD, Knapp AK. Exotic plant species in a C 4-dominated grassland: invasibility,
disturbance, and community structure. Oecologia. 1999;120(4):605-12.

32. Burke MJW, Grime JP. An experimental study of plant community invasibility. Ecology.
1996;77(3):776-90.

33. Guerra-García JM, García-Gómez JC. Polychaete assemblages and sediment pollution in
a harbour with two opposing entrances. Helgoland Marine Research. 2004;58(3):183-91.

34. HC. Hatfield Consultants Ltd. Skeena environmental effects monitoring (EEM) pre-
design reference document. Prepared for Skeena Cellulose Inc. 1994.

35. Hoos LM. The Skeena River Estuary: status of environmental knowledge to 1975: Report
of the Estuary Working Group, Department of the Environment, Regional Board, Pacific
Region. Environment Canada, 1975 Contract No.: 3.

36. NIBR. Northwest Institute for Bioregional Research: Valuation of the wild Salmon
economy of the Skeena River watershed. 1-30 2006.

37. Hilborn R, Walters CJ. Differing goals of salmon management on the Skeena River.
Journal of the Fisheries Board of Canada. 1977;34(1):64-72.

38. Higgins RJ, Schouwenburg WJ. A biological assessment of fish utilization of the Skeena
River Estuary, with special reference to port development in Prince Rupert. Technical Report
1973-1. Department of Environment, Fisheries and Marine Services. Vancouver, BC, 1973.

39. Faggetter BA. Review of the Environmental and Socioeconomic

Impacts of Marine Pollution in the

North and Central Coast Regions of British

Columbia. 2008.

40. Yunker MB, Cretney WJ, Ikonomou MG. Assessment of chlorinated dibenzo-p-dioxin
and dibenzofuran trends in sediment and crab hepatopancreas from pulp mill and harbor sites
using multivariate- and index-based approaches. Environmental Science & Technology.
2002;36(9):1869-78. doi: 10.1021/es0112893. PubMed PMID: WOS:000175311900006.

41. Akenhead S. A review of the oceanography and marine ecology of Prince Rupert
Harbour a propos sewage outfalls. 1992.

42. Cox K, Black M, Filip N, Miller M, Mohns K, Mortimor J, et al. Comparison of
community assessment techniques and implications for diversity indices and species
accumulation curves. Ecology and Evolution. 2017;DOI: 10.1002/ece3.3580.

43. BS7755-3.2. Soil Quality. Part 3: Chemical Methods. Section 3.2: Determination of pH
British Standards Institution, London, UK. 1995.

44. EPA US. Method 3051A (SW-846): Microwave Assisted Acid Digestion of Sediments,
Sludges, and Oils, Revision 1. Washington, DC. 2007.

45. EPA US. Method 7473 (SW-846): Mercury in Solids and Solutions by Thermal
Decomposition, Amalgamation, and Atomic Absorption Spectrophotometry, Revision 0.
Washington, DC. 1998.

46. Anderson M, Gorley RN, Clarke RK. Permanova+ for Primer: Guide to software and
statistical methods. Plymouth, United Kingdom: PRIMER-E Ltd; 2008.

47. Gerwing TG, Drolet D, Hamilton DJ, Barbeau MA. Relative importance of biotic and
abiotic forces on the composition and dynamics of a soft-sediment intertidal community
PLoS One. 2016;11(1):11:e0147098.

48. Clarke KR, Gorley RN. PRIMER v7: user manual/tutorial 3rd ed. Plymouth, United
Kingdom: Primer-E Ltd; 2015.

49. McLaren P. The environmental implications of sediment transport in the waters of Prince
Rupert, British Columbia, Canada: A comparison between kinematic and dynamic
approaches. Journal of Coastal Research. 2016;32(3):465-82.

50. Muller G. Index of geoaccumulation in sediments of the Rhine River. Geojournal.
1969;2:108-18.

51. Monte MC, Fuente E, Blanco A, Negro C. Waste management from pulp and paper
production in the European Union. Waste management. 2009;29(1):293-308.

52. Hoffman E, Lyons J, Boxall J, Robertson C, Lake CB, Walker TR. Spatiotemporal
assessment (quarter century) of pulp mill metal (loid) contaminated sediment to inform
remediation decisions. Environmental monitoring and assessment. 2017;189(6):257.

53. Borah P, Singh P, Rangan L, Karak T, Mitra S. Mobility, bioavailability and ecological
risk assessment of cadmium and chromium in soils contaminated by paper mill wastes.
Groundwater for Sustainable Development. 2018;6:189-99.

54. Nriagu JO, Nieboer E. Chromium in the natural and human environments: John Wiley &
Sons; 1988.

55. Kienle C, Langer-Jaesrich M, Baumberger D, Hohmann D, Santiago S, Köhler H-R, et al.
Integrated toxicity evaluation of a pulp deposit using organisms of different trophic levels.
Journal of soils and sediments. 2013;13(9):1611-25.

56. Alvarez MB, Malla ME, Batistoni DA. Comparative assessment of two sequential
chemical extraction schemes for the fractionation of cadmium, chromium, lead and zinc in
surface coastal sediments. Fresenius' journal of analytical chemistry. 2001;369(1):81-90.

57. Chakraborty P, Babu PVR, Sarma VV. A study of lead and cadmium speciation in some
estuarine and coastal sediments. Chemical Geology. 2012;294:217-25.

58. Sizmur T, Canário J, Gerwing TG, Mallory ML, O'Driscoll NJ. Mercury and
methylmercury bioaccumulation by polychaete worms is governed by both feeding ecology
and mercury bioavailability in coastal mudflats. Environmental Pollution. 2013;176:18-25.

59. Weimin Y, Batley G, Ahsanullah M. The ability of sediment extractants to measure the
bioavailability of metals to three marine invertebrates. Science of the Total Environment.
1992;125:67-84.

60. Burton ED, Phillips IR, Hawker DW. Geochemical partitioning of copper, lead, and zinc
in benthic, estuarine sediment profiles. Journal of environmental quality. 2005;34(1):263-73.

61. Luoma SN. Processes affecting metal concentrations in estuarine and coastal marine
sediments. Heavy metals in the marine environment: CRC Press; 2017. p. 51-66.

62. Riba I, Delvalls TÁ, Forja JM, Gómez-Parra A. The influence of pH and salinity on the
toxicity of heavy metals in sediment to the estuarine clam Ruditapes philippinarum.
Environmental toxicology and chemistry. 2004;23(5):1100-7.

63. Zhang C, Yu Z-g, Zeng G-m, Jiang M, Yang Z-z, Cui F, et al. Effects of sediment
geochemical properties on heavy metal bioavailability. Environment International.
2014;73:270-81. doi: https://doi.org/10.1016/j.envint.2014.08.010.

64. DelValls TÁ, Forja J, González-Mazo E, Gómez-Parra A, Blasco J. Determining
contamination sources in marine sediments using multivariate analysis. TrAC Trends in
Analytical Chemistry. 1998;17(4):181-92.

65. Chapman PM, Anderson J. A decision-making framework for sediment contamination.
Integrated environmental assessment and management. 2005;1(3):163-73.

66. Ciutat A, Boudou A. Bioturbation effects on cadmium and zinc transfers from a
contaminated sediment and on metal bioavailability to benthic bivalves. Environmental
toxicology and chemistry. 2003;22(7):1574-81.

67. Sizmur T, Canário J, Edmonds S, Godfrey A, O'Driscoll NJ. The polychaete worm Nereis
diversicolor increases mercury lability and methylation in intertidal mudflats. Environmental
Toxicology and Chemistry. 2013;32(8):1888-95.

68. Gerwing TG, Allen Gerwing AM, Macdonald T, Cox K, Juanes F, Dudas SE. Intertidal
soft-sediment community does not respond to disturbance as postulated by the intermediate
disturbance hypothesis. Journal of Sea Research. 2017;129:22-8.

69. Light SF. The Light and Smith manual: intertidal invertebrates from central California to
Oregon. 4 ed. Cameron JT, editor: University of California Press, Berkely, CA, USA; 2007.

70. Gerwing TG, Kim JH, Hamilton DJ, Barbeau MA, Addison JA. Diet reconstruction using
next-generation sequencing increases the known ecosystem usage by a shorebird. The Auk.
2016;133(2):168–77.

71. Rainbow PS. Biomonitoring of heavy metal availability in the marine environment.
Marine pollution bulletin. 1995;31(4-12):183-92.

72. Fauchald K, Jumars PA. The diet of worms: a study of polychaete feeding guilds.
Oceanography and Marine Biology Annual Review. 1979;17:193-284.

73. Pagliosa PR. Another diet of worms: the applicability of polychaete feeding guilds as a
useful conceptual framework and biological variable. Marine Ecology. 2005;26:246-54.

74. Howard L, Brown B. Natural variations in tissue concentration of copper, zinc and iron in
the polychaete Nereis diversicolor. Marine biology. 1983;78(1):87-97.

75. Saha M, Sarkar S, Bhattacharya B. Interspecific variation in heavy metal body
concentrations in biota of Sunderban mangrove wetland, northeast India. Environment
International. 2006;32(2):203-7.

76. Saiz-Salinas J, Francés-Zubillaga G. Nereis diversicolor: an unreliable biomonitor of
metal contamination in the 'Ría de Bilbao'(Spain). Marine Ecology. 1997;18(2):113-25.

77. Schöne BR, Krause Jr RA. Retrospective environmental biomonitoring–Mussel Watch
expanded. Global and Planetary Change. 2016;144:228-51.

78. Zuykov M, Pelletier E, Harper DAT. Bivalve mollusks in metal pollution studies: from
bioaccumulation to biomonitoring. Chemosphere. 2013;93(2):201-8.

79. Caçador I, Costa J, Duarte B, Silva G, Medeiros J, Azeda C, et al. Macroinvertebrates and
fishes as biomonitors of heavy metal concentration in the Seixal Bay (Tagus Estuary): which
species perform better? Ecological Indicators. 2012;19:184-90.

80. Sarkar SK, Cabral H, Chatterjee M, Cardoso I, Bhattacharya AK, Satpathy KK, et al.
Biomonitoring of heavy metals using the bivalve molluscs in Sunderban mangrove wetland,
northeast coast of Bay of Bengal (India): possible risks to human health. CLEAN–Soil, Air,
Water. 2008;36(2):187-94.

81. Bard S, M,. A biological index to predict pulp mill pollution levels. Water environment
research. 1998;70(1):108-22.

82. Robin J, Harger E, Nassichuk MD. Marine intertidal community responses to Kraft pulp
mill effluent. Water, Air, and Soil Pollution. 1974;3(1):107-22.

645 Supporting information captions

646 The supporting information file contains additional text that elaborates on the quality control and
647 statistical analysis undertaken in addition to the following tables:

649 **S1 Table. PERMANCOVA showing sediment properties (pH, median particle diameter, C and N)**
650 **varied by site, depth, and transect.**
651

652 **S2 Table. PERMANCOVA showing that sediment total and available (EDTA extractable) PTEs varied**
653 **by site and transect.**
654

655 **S3 Table. Analysis of Variance for sediment properties. F-statistics of a two-way ANOVA with 'site'**
656 **and 'depth' as the two factors. The last two columns indicate whether reference sites; Tyee Banks**
657 **(TB), Wolfe Cove (WC) and Inverness Passage (IP) are significantly ($p < 0.05$) different from**
658 **potentially contaminated sites (Cassiar Cannery and Papermill Bay).**
659

660 **S4 Table. Concentrations (mg kg^{-1}) of Cr, Co, Ni, Cu, Zn, As, Cd and Pb in benthic invertebrates**
661 **sampled from five intertidal mudflats along the north coast of British Columbia, Canada. CC:**
662 **Cassiar Cannery. WC: Wolfe Cove. IP: Inverness Passage. PB: Papermill Bay. TB: Tyee Banks**
663

Cassiar Cannery
Inverness Passage
Papermill bay
Wolfe Cove
Tyee Banks

Cassiar Cannery ■
Inverness Passage □
Papermill bay ▲
Wolf Cove △
Tyee Banks ●

A

2D Stress: 0.03

Taxa
- ▲ Abarenicola pacifica
- ▲ Americorophium salmonis
- ◆ Eogammarus confervicolus
- ● Gnorimosphaeroma oregonensis
- ◆ Hemigrapsus oregonensis
- ✕ Macoma balthica
- ✕ Modiolus rectus
- ✶ Mya arenaria
- ✦ Mytilus edulis
- ▼ Nephtys caeca
- ◇ Nereis vexillosa
- ◇ Paguras hirsutiusculus
- ◇ Paranemertes peregrina
- □ Streblospio benedicti
- ● Idotea wosnesenskii
- ✶ Alita brandti
- ✶ Aricidea hartleyi
- ✚ Cerebratulus californiensis
- ✚ Glycinde picta
- ✕ Macoma nasuta
- ★ Neotrypaea californiensis

B

2D Stress: 0

Site
- ▲ CC
- ◆ IP
- ■ PB
- ▼ TB
- ● WC

$R^2 = 0.8631$

$R^2 = 0.7548$

Sediment pH