Uncovering and characterizing splice variants associated with survival in lung cancer patients

Sean West[1], Sushil Kumar[2], Surinder K. Batra[2], Hesham Ali[1*], Dario Ghersi[1*]

1 College of Information and Science Technology, University of Nebraska at Omaha, Omaha, NE, USA **2** Department of Biochemistry and Molecular Biology, University of Nebraska Medical Center, Omaha, NE, USA

* hali@unomaha.edu, dghersi@unomaha.edu

Abstract

Splice variants have been shown to play an important role in tumor initiation and progression and can serve as novel cancer biomarkers. However, the clinical importance of individual splice variants and the mechanisms by which they can perturb cellular functions are still poorly understood. To address these issues, we developed an efficient and robust computational method to: (1) identify splice variants that are associated with patient survival in a statistically significant manner; and (2) predict rewired protein-protein interactions that may result from altered patterns of expression of such variants. We applied our method to the lung adenocarcinoma dataset from TCGA and identified splice variants that are significantly associated with patient survival and can alter protein-protein interactions. Among these variants, several are implicated in DNA repair through homologous recombination. To computationally validate our findings, we characterized the mutational signatures in patients, grouped by low and high expression of a splice variant associated with patient survival and involved in DNA repair. The results of the mutational signature analysis are in agreement with the molecular mechanism suggested by our method. To the best of our knowledge, this is the first attempt to build a computational approach to systematically identify splice variants associated with patient survival that can also generate experimentally testable, mechanistic hypotheses. Code for identifying survival-significant splice variants using the Null Empirically Estimated P-value method can be found at https://github.com/thecodingdoc/neep. Code for construction of Multi-Granularity Graphs to discover potential rewired protein interactions can be found at https://github.com/scwest/SINBAD.

Author summary

In spite of many recent breakthroughs, there is still a pressing need for better ways to diagnose and treat cancer in ways that are specific to the unique biology of the disease. Novel computational methods applied to large-scale datasets can help us reach this goal more effectively. In this work we shed light on a still poorly understood biological process that is often aberrant in cancer and that can lead to tumor formation, progression, and invasion. This mechanism is alternative splicing and is the ability of one gene to code for many different variants with distinct functions. We developed a fast and statistically robust approach to identify splice variants that are significantly associated with patient survival. Then, we computationally characterized the protein

products of these splice variants by identifying potential losses and gains of protein interactions that could explain their biological role in cancer. We applied our method to a lung adenocarcinoma dataset and identified several splice variants associated with patient survival that lose biologically important interactions. We conducted case studies and computationally validated some of our results by finding mutation signatures that support the molecular mechanism suggested by our method.

Introduction

Large-scale cancer sequencing initiatives have opened up a window into the genome of individual cancers, offering unprecedented opportunities for studying the functional consequences of molecular alterations in human cancers [1]. However, despite new evidence highlighting the importance of cancer-specific splice variants in tumor initiation and progression, the role played by splice variants in the biology of human cancers and patient survival is still poorly understood.

To address this need, we developed a novel, statistically robust, multi-granular method to identify splice variants that are significantly associated with cancer patient survival. After clinically important splice variants are detected, our approach identifies variant-specific losses and gains of protein domains that can potentially rewire the cellular interactome, enabling a mechanistic interpretation of the role played by these splice variants. We applied this approach to Lung Adenocarcinoma (LUAD) using data from The Cancer Genome Atlas (TCGA), based on the clinical importance of the disease, the large number of observed point mutations that can lead to aberrant splicing, and the spread in patient survival time.

With lung cancer as the leading cause of cancer-related deaths [2], novel treatments are constantly developed to improve patient survival while reducing the toxicity and side-effects of conventional chemotherapy. In this regard, molecular targeted therapy is arguably one of the most investigated new paradigms in oncology. The general strategy in targeted therapy is to use small molecules to inhibit specific pathways important for cancer cell survival and invasiveness. [3]. However, for molecular targeted therapy to be successful, it is important to identify highly specific molecular mechanisms that lead to tumor survival and metastasis. For this reason, cancer-specific alternative splicing has received increasing attention for its role in tumor formation [4,5], and methods to target splicing in cancer have recently been proposed [6] using RNAseq data.

While some argue in favor of the robustness of gene level analysis of RNAseq data [7], splice variant analysis offers unique information [8]. Sebestyén et al. developed a computational method to identify splicing signatures in cancers using TCGA data, showing that splicing signatures can be used to distinguish tumor from normal samples [9]. Yang and co-workers developed a web server called ISOexpresso to identify predominantly expressed isoforms in TCGA data [10]. Recent work by Climente-González has investigated the potential functional consequences of splice variants in TCGA datasets, highlighting Gene Ontology terms that are enriched in isoform switches and protein interactions that might be lost or gained [11]. We suggest that more detailed understanding of the influence of transcriptional expression on disease mechanisms in lung cancer may be gleaned from granularity-aware integration of RNASeq data with other sources of domain knowledge.

The strategy we propose here is to combine established RNAseq processing methods with patient survival analysis at the gene and splice variant level. Exon-skipping and splice variant level survival analysis have been shown to outperform gene level analysis at predicting patient survival [12]. Many methods for integration of RNAseq data exist and usually occur during downstream processing [13]. However, to the best of our knowledge this is the first attempt at building a high-throughput approach for survival

analysis at the gene or splice variant level for early integration with domain sources. 45

Another important requirement for confident knowledge extraction is that of 46
robustness, defined as stability of the results with respect to small changes in the 47
parameters of the method. Non-parametric survival analysis requires a threshold that 48
splits patients into subpopulations (e.g. an expression threshold). Previously, 49
high-throughput survival analysis using arbitrary thresholds were shown to be sensitive 50
to changes in patient groups [14]. One known approach for increasing this robustness is 51
to find the minimum p-value by sampling multiple thresholds. However, this produces 52
p-values that are non-uniform and skewed. Lausen and Schumacher examined this 53
problem for the logrank test, developing an equation for estimating the corrected 54
p-values [15]. Yet, this type of correction method was not developed for high-throughput 55
analysis, where multiple hypothesis testing is required and is dependent on very small 56
changes of individual p-values. To overcome these issues, we developed a reliable and 57
high-throughput survival analysis method called NEEP (null empirically estimated 58
p-values). This method requires no distribution assumptions, guarantees statistically 59
valid p-values, and may be implemented in a high-throughput way. 60

We applied this method to the LUAD RNAseq dataset from TCGA to conduct 61
survival analysis at the splice variant level, identifying variants that are significantly 62
associated with patient survival. Then, we integrated these results with protein-domain 63
annotation and protein-protein interaction data to provide a mechanistic interpretation 64
for the potential role played by these clinically relevant splice variants in tumor 65
formation and progression. More specifically, we identified variants that lose or gain 66
protein domains involved in mediating protein-protein interactions. We refer to this 67
integration approach as a multi-granularity graph (MGG). Fig. 1 shows the overall 68
workflow of the method. 69

In the absence of a gold standard, we conducted case study analysis with literature 70
review to assess the biological plausibility of the inferred graphs. In summary, our main 71
contributions are the following: (1) we use the splice variant granularity level as a way 72
to identify changes in protein mechanisms that can affect patient survival; (2) we 73
present a novel, computationally efficient, and robust method to perform 74
high-throughput survival analysis; and (3) we integrate multiple data types for inferring 75
biologically plausible and potentially actionable molecular targets in lung cancer. 76

Results 77

NEEP identifies splice variants significantly associated with 78
patient survival 79

One of the contributions of this work is the development of a statistically robust and 80
computationally efficient method to identify optimal expression thresholds that yield 81
minimum p-values when performing survival analysis over a large number of transcripts. 82
The efficiency of the NEEP method is due to the fact that we can estimate empirical 83
p-values for all transcripts by computing only one null distribution, as opposed to 84
having to perform sample permutations for every transcript (as discussed in detail in 85
Materials and Methods). To visually show this, we plot the p-value distributions of 100 86
randomized gene expression patterns of size 1000, against 20 simulated null 87
distributions of size 1000 obtained by shuffling samples (Fig 2A). The overlap between 88
these distributions visually confirms that the same null distribution can be utilized 89
across all expression patterns without the need for expensive sample permutations for 90
every transcript. 91

We ran NEEP using 10 randomly generated null distributions of 1,000,000 isoform 92
simulations. In all 10 repetitions, there were 181 NEEP significant (FDR < 0.1) splice 93

Fig 1. Workflow for generating multi-granularity graphs (MGGs). Using the NEEP method significant splice variants were calculated, utilizing clinical data, the complementary-DNA reference library, and RNA-Seq data from Ensembl and TCGA. Multi-granular graphs were constructed for known protein-protein interactions from BioGRID, mapped from Ensembl to at least one significant splice variant. Domains were predicted for each splice variant using HMMER3 and Pfam. Gained and ghost domains were identified in splice variants associated with survival. Final MGGs represent potential lost or gained protein interactions associated with patient survival.

variants from the LUAD dataset. Since the order of splice variants in NEEP does not depend on the null but only their statistical significance does, these 181 splice variants were identical each time. The mean hazard ratio of the significant splice variants was 3.122 and the minimum was 2.18.

The distribution of all the p-values is nearly uniform, as can be seen in Fig 2B. The small deviation of low p-values from the uniform distribution is due (as expected) to the significant splice variants.

Case study of multi-granular graphs linked to DNA repair

After identifying splice variants significantly associated with survival, we followed the procedure described in Materials and Methods and summarized in Fig. 1 to generate multi-granular graphs (MGGs). Briefly, MGGs show splice variant specific protein-protein interactions that might be gained or lost as a result of gains or losses of protein domains in splice variants significantly associated with patient survival.

After constructing the MGGs using the NEEP significant splice variants, we obtained 50 MGGs (the complete list is available in S3 File). We first checked whether

Fig 2. NEEP yields uniformly distributed *p*-values. (A) The *p*-value distribution is shown for the minimum *p*-value method and NEEP. An ideal statistical test produces *p*-values that are uniform under the null. (B) Density line plots were constructed for each of 100 simulations of 1000 random abundance patterns (blue) and 20 sets of 1000 values chosen at random from the initial 1,000,000 null distribution (red). The overlap between the two groups of density plots empirically confirms that the null distribution is the same for all expression patterns, given identical clinical data.

the MGGs we obtained were representative of the full set of significant splice variants   109
identified by NEEP. To do that, we performed Gene Ontology enrichment analysis on   110
NEEP significant genes, NEEP significant isoforms, and genes represented in MGGs, as   111
discussed in S1 File. All GO enrichment term groups obtained at the gene or isoform   112
levels were also present in the MGGs, as shown in S1 Fig.   113

Using these MGGs and information available from the literature, we investigated   114
possible mechanisms that can explain why these splice variants are correlated with   115
survival of lung cancer patients. Here we discuss three case studies linked to DNA   116
repair. Additional MGGs are discussed in the supplemental materials S2 File. All   117
discussed MGGs are visualized in Fig 3.   118

RAD51C-202 loses ability to bind to the Holliday junction, XRCC3, and HELQ   119 120

NEEP suggests that the increased expression of RAD51C-202 is associated with lower   121
patient survival. Further, the interaction between RAD51C-202 and XRCC3 is   122
prevented because of several missing domains. Upon deeper inspection using RCBP   123
PDB and Ensembl, we see that RAD51C-202 is missing two exons that are present in   124
the remaining RAD51C isoforms. The first exon allows RAD51C to bind to the Holliday   125
junction, a relationship missed by the MGG. The second exon contains the domains   126
which allows RAD51C to bind to XRCC3. RAD51C and XRCC3 have long been known   127
to form a complex (CX3 complex) which regulates genetic recombination and DNA   128
repair [31]. Dysruption of this complex (perhaps via increased expression of   129
RAD51C-202) may promote cancer in two ways. First, loss of RAD51C has been   130
implicated in tumorigenesis [32]. Second, it was suggested that the CX3 complex   131
increases mitochondrial stability, reducing the risk of cancer [33]. In addition, HELQ   132
interacts with RAD51C within the BCDX2 complex [34]. Disruption of the BCDX2   133
complex increases cellular susceptibility to agents that induce DNA-interstrand   134
crosslinks [35]. Thus, the additional transcription of the RAD51C-202 isoform which is   135
missing the domain necessary to bind to HELQ may result in DNA-interstrand   136
crosslinks.   137

Fig 3. Multi-granularity graphs. Splice variants are in blue solid boxes and domains are in dotted boxes. The significant splice variant (left box) is linked to its gained (green) and ghost (white) domains. Domains in gray belong to the non-significant splice variant (right box). Splice variants of the same gene that have identical domain connections in the MGG are stacked and their identifiers are listed below the boxes. RAD51C-202 has two MGGs which are separated to simplify visualization.

RAD23A-201 loses interaction with CHEK1 proteins 138

The RAD23A-201 isoform does not include the UBA_2 domain, which is found in all of 139
the other RAD23A isoforms. The survival analysis indicates that a high expression of 140
RAD23A-201 is associated with lower patient survival. RAD23A operates by bringing 141
proteins marked for ubiquitination to the proteasome, avoiding its own degradation. 142
Without UBE_2 RAD23A is ubiquitinized, preventing its role in nucleotide excision 143
DNA repair and protein degredation [36]. Further, without the UBA_2 domain, the PPI 144
between RAD23A and CHEK1 via the DDI between UBE_2 and Pkinase will not occur. 145
It may be possible that RAD23A phosphorylates CHEK1 since its cousin, RAD17, is a 146
known phosphorylator of CHEK1 [37]. However, this is unlikely since RAD23A is 147
known to function primarily through its association with the proteasome [38]. The 148
inhibition of the PPI with CHEK1 may prevent degradation of CHEK1. Under several 149
circumstances CHEK1 promotes tumor growth and increases therapy resistance [37]. 150
Thus the reduction of RAD23A with longevity could inhibit CHEK1 degradation and 151
allow promotion of tumor growth and therapy resistance. 152

INTS7-202 has altered structure and number of TPR domains ₁₅₃

INTS proteins serve as members of the Integrator complex (INT). The roles of this ₁₅₄
complex are diverse but suggestive of potential roles in cancer, such as mediation of ₁₅₅
RNA polymerase II or DNA damage response [39]. Yet the INT is 'modular', containing ₁₅₆
different subunits of INTS genes and causing different effects on the role of INT [40]. ₁₅₇
The MGG model suggests that the interaction between INTS7 and INTS8 may be ₁₅₈
effected by the loss of a few TPR domains in INTS7-202. The TPR domains are ₁₅₉
common PPI domains which allow the construction of many protein complexes, ₁₆₀
including INT. Losses or disruptions of these domains in INTS7-202 may cause ₁₆₁
instability and prevent the formation of INT. Alternatively, given that many other TPR ₁₆₂
domains are uneffected by the alternative splicing of INTS7-202, INT may still form but ₁₆₃
function differently. In addition, INTS proteins also function outside of the INT ₁₆₄
complex [39]. ₁₆₅

RAD51C-202 expression is linked to a characteristic mutational signature and lower patient survival _{166 167}

The biological models discussed above suggest a role played by these splice variants in ₁₆₈
DNA repair. Alterations in DNA repair mechanisms result in characteristic mutations ₁₆₉
signatures. We hypothesized that patients in the RAD51C-202 low expression and high ₁₇₀
expression groups might have different mutation signatures reflecting different abilities ₁₇₁
to perform DNA repair via homologous recombination. ₁₇₂

As shown in Table 1, RAD51C-202 is associated with COSMIC mutation signatures ₁₇₃
2, 3, 11, 12, 16, 17, and 23. Of particular interest, signature 3 (Fig 4) is reported by ₁₇₄
COSMIC as associated with failure in DNA double-strand breaks in homologous ₁₇₅
recombination. Indeed, the results of the Mood's test indicate that signature 3 is more ₁₇₆
prevalent in patients with high expression of RAD51C-202 than in patients with low ₁₇₇
expression ($p < 0.0232$, Fig 4). RAD51C-202 had the highest association with COSMIC ₁₇₈
signature 2, which is known to be caused by AID/APOBEC activity. However, we could ₁₇₉
not find more information from the literature regarding this particular signature. ₁₈₀

Table 1. COSMIC mutation signatures associated with RAD51C-202 expression.

	High - Low (Mean Contribution)	Description from COSMIC
$Signature2$	11.032	Activity of AID/APOBEC deaminases
$Signature3$	3.864	DNA double-strand break-repair by homologous recombination
$Signature11$	0.809	Similar mutation pattern to alkylating agents
$Signature12$	2.323	Unknown
$Signature16$	3.194	Unknown
$Signature17$	-0.016	Unknown
$Signature23$	0.533	Unknown

The difference between the mean contributions of *low* from *high* RAD51C-202 expression groups to each significant mutation
signature is reported. Signature 4, associated with smoking, is not significant.

Since smoking is commonly associated with lung cancer and may cause particular ₁₈₁
mutation patterns, the relationship between RAD51C-202 with signature 3 could be ₁₈₂
confounded by smoking. However, smoking does not appear to be a confounding factor ₁₈₃
in RAD51C-202 association with COSMIC gene signatures. As can be seen in Table 2, ₁₈₄
no smoking variable was significantly different between patients with low and high ₁₈₅
expression of RAD51C-202. Further, signature 4 contribution (found in smokers) was ₁₈₆
not significantly associated with RAD51C-202 abundance. In addition, survival does not ₁₈₇
appear to be a confounding factor. The survival Cox-PH p-value for RAD51C-202 was ₁₈₈
significant with and without using signature 3 contribution as a confounding factor ₁₈₉

Fig 4. Mutation signatures associated with RAD51C and smoking effects.
(A) The mean contribution of each patient in the high RAD51C-202 expression group
was subtracted from the low group. A positive number indicates more contribution of
the COSMIC mutation signature to the patients in the high expression group. Listed
signatures have a significant Mood's test p-value for RAD51C-202 expression. (B)
Potential smoking confounding variables were separated by RAD51C-202 expression and
tested according to their data type. (C) Box-plots were generated for the signature-3
contribution to patients in low (red) and high (blue) expression of RAD51C-202.

(p-values: 0.010 and 0.036). The proportion of NEEP significant isoforms that were   190
associated with signature 3 was insignificant (p-value = 0.6403).   191

Table 2. Statistical test results comparing smoking variables to RAD51C-202 Expression

Smoking Variable	Data Type	Statistical Test	p-value
Cigarettes Per Day	Metric	Welch Two Sample T-test	0.086
Years Smoked	Integer	Wilcoxon Rank Sum Test	0.098
Is Smoker	Binary	2-Sample Test for Equal Proportions	0.383

Robustness of NEEP to changes in patient population   192

We conducted 100 simulations by removing 5% and 10% of the patients at random prior   193
to running NEEP. We compared the rank of the significant isoforms observed in the full   194
dataset to their rank in the simulations. Fig 5A shows the range of the middle 50%   195
ranks from the simulations, and it also plots the original, observed values. Of the 181   196
observed significant splice variants, few had ranks greater than 400 out of 69,561.   197
Further, splice variants with lower observed rank had lower variance in the simulations.   198
In addition and as expected, ranks in the 5% simulations were much lower than the   199
ranks in the 10% simulations. Fig 5B is the density plot of the observed ranks paired   200
with the 10% simulation ranks. The density nearly follows the $y = x$ line, where the   201
observed rank is the same as the simulated rank. Once again we see that splice variants   202
with low observed rank have lower simulation variance than variants of higher rank.   203
Mean Spearman rank correlations between the entire set of observed splice variants and   204
the 5% and 10% simulation rounds were 0.861 and 0.931, respectively. Overall, these   205
results suggest that the NEEP method is fairly robust to removal of patient samples.   206

Fig 5. Robustness of single threshold methods and NEEP (A) The central 50 ranks for the 100 simulations of each of the 181 significant splice variants are plotted as a shaded range according to their mean simulated rank, separately for the 5% simulations (red) and the 10% simulations. Each of these ranges correspond to an observed (non-simulated) rank, which is plotted as a dot along the same x-axis. The 5% simulation ranges have lower means of simulation rank than the 10% simulation ranges; thus the red dots are closer to the identity line. The maximum mean of the simulation rank is lower for the 5% simulations; thus the red shade ends much sooner than the blue shade. (B) The density plot of the observed vs. simulation pairs for the 181 significant splice variants.

Discussion   207

An exploratory survival analysis must be high-throughput to span all genes/splice   208
variants (in this study, 21,100 genes and 69,561 splice variants). Cox regression and   209
Kaplan-Meier (KM) curves are two options. We chose to avoid Cox regression for two   210
reasons. First, we violate the Cox regression assumption that the mechanism of   211
censoring is not related to patient survival, as patients who survive to the end of the   212
study will have a higher survival rate than the overall population at the outset. As   213
discussed in the Methods, we can choose a threshold range which prevents loss of   214
statistical power due to this phenomena on the KM test. Second, Cox regression firmly   215
assumes the proportional hazards assumption, where the assumption is looser in KM   216
analysis. The effect of many splice variants on survival is not constant over time,   217
resulting in varying hazard rates. Driver genes are just one example that violate this   218
assumption.   219

The hypothesis of the KM logrank test is also better suited to this problem than Cox   220
regression: Cox regression tests against the null that the variability of the survival   221
cannot be explained by the variability of the splice variant, whereas the KM logrank   222
test compares *low* against *high* expression. Therefore, each KM hypothesis is associated   223
with the maximum difference between two survival curves, and exact delineations of *low*   224
and *high* expression are extracted.   225

A current limitation of our approach is that the multi-granularity graphs only   226
highlight splice variants that can gain or lose protein-protein interactions as a result of   227
gains/losses of protein domains. Protein-DNA, protein-RNA, and protein-ligand   228
interactions are not considered. We plan to extend our work in this direction in the near   229
future.   230

In conclusion, we presented NEEP, a novel computational method to systematically   231
identify splice variants significantly associated with patient survival. An important   232
advantage of NEEP over the existing methods is that the optimal expression threshold   233
to separate patients into better vs. worse survival groups is automatically detected by   234
the method, and the resulting *p*-values are uniformly distributed under the null   235
hypothesis, allowing for proper multiple hypothesis correction.   236

By applying our method to the TCGA lung adenocarcinoma dataset, we identified several splice variants that significantly correlate with patient survival. We then built multiparted, multi-granular graphs (splice variant-domain-domain-splice variant) that allowed us to identify putative losses and gains of protein-protein interactions associated with these splice variants. Several of these variants turned out to be implicated in DNA repair through homologous recombination. To computationally verify this, we performed a mutational signature analysis that supported the molecular mechanism suggested by the multi-granular graphs.

Materials and methods

Data retrieval and transcriptome profiling

We downloaded the TCGA Lung Adenocarcinoma (LUAD) dataset from the Cancer Genomics Hub (CGHub). Of the 515 patients with RNA-Seq data, 75 were removed due to clinical annotations suggesting possible confounding factors like "Prior malignancy" or "Synchronous malignancy". Eight patients were removed due to missing survival data, resulting in 432 total patients. To reconstruct splice variant abundance estimates, the TCGA BAM files were reverted back to fastq format using SAMtools [16].

We used Salmon [17] with default parameters to estimate transcript-level abundances based on the Ensembl 91 cDNA library. Non-alignment methods such as Salmon offer accurate abundance estimates at the splice variant level [18]. In addition, Salmon performed best during a comparison of alignment-free methods [19]. Using Ensembl gene/splice variant mapping, we summed splice variant abundances to estimate gene abundance. TPM values were used to represent the gene/splice variant abundances. We note that since non-parametric survival analysis utilizes two Kaplan-Meier curves for each gene/splice variant (one for *high* abundance and one for *low* abundance), the choice of using counts, FPKM or TPM will result in identical logrank p-values.

High-throughput survival analysis

We performed non-parametric survival analysis at the gene and splice variant level using a previously published approach ("minimum p-value") and our new NEEP ("null empirically estimated p-values") method, as described below. The clinical files containing survival data were obtained from TCGA. The *days_to_death* and *days_to_last_followup* were used as uncensored and censored survival values, respectively.

Minimum p-value approach

Choosing the threshold which maximizes a test statistic avoids the choice of an arbitrary threshold, while increasing reliability. Minimum p-values were calculated for each gene and splice variant. The logrank test was performed on each possible split between high and low abundance within a threshold range of 0.15 to 0.85. This method results in a distribution of p-values that is non-uniform and skewed to the left.

Empirical estimation of p-values

To overcome the limitations of choosing a single threshold (arbitrary threshold) and minimum p-value methods (non-uniform p-values under the null hypothesis), we developed a new approach resulting in null empirically estimated p-values (NEEP). The method works by estimating the null distribution of the minimum p-value approach, so that p-values under the null hypothesis are uniformly distributed, allowing for p-value adjustment.

When using the logrank test with a single expression threshold t, the set of all possible p-values is finite, and it corresponds to all possible patient partitions into the *low* and *high* expression groups. By extension, the null distribution of the minimum p-value method is defined as the entire discrete distribution of possible p-values across all values of t. The exact number of possible p-values can be calculated as a function of s (the number of samples), l (the minimum percentile threshold), and h (the maximum percentile threshold), according to the following equation:

$$\text{size of null} = \sum_{i=\lfloor l*s \rfloor}^{\lfloor h*s \rfloor} \binom{\text{s}}{i} \tag{1}$$

where $\lfloor x \rfloor$ is the greatest integer less than or equal to x. To determine a viable range of thresholds, we first chose minimum power and effect size. Given a sample size of 432 patients with 151 recorded death events and an alpha of 0.1, to obtain a hazard ratio greater than 2 (or less than 0.5) and a minimum power of 0.75, the survival curve with fewest patients must have at least 62 members. This corresponds to a minimum percentile threshold of 0.143, which we round in this study to 0.15 for simplicity. So, in this study we set l to 0.15 and h to 0.85.

Considering the size of s, enumerating all possible p-values is computationally unfeasible. Therefore, we built a null distribution with 1,000,000 Monte Carlo simulations performed by randomly partitioning samples into the two groups using all possible thresholds and extracting the smallest logrank test p-value. We estimated p-values as the fraction of simulated values less than or equal to the minimum p-values obtained from the actual data. As expected, the NEEP procedure yielded uniformly distributed p-values. Benjamini-Hochberg adjustment was conducted on the NEEP p-values, and adjusted values less than 0.1 were considered significant.

It is noteworthy to mention that individual expression values do not impact the KM survival curve as long as they result in the same split between high and low expression. Therefore, the NEEP procedure needs to be carried out only once for the whole set of genes/splice variants, provided the fraction of patients expressing the gene/splice variant is as large as or greater than $1 - l$, the minimum percentile threshold. In our dataset, this criterion was met by 21,100 genes and 69,561 splice variants. Thus, we empirically estimated the p-values of these genes/splice variants using a single sampled null distribution. While effect sizes of the logrank test are not as naturally interpretable as many other statistical tests, we calculated the hazard rate ratio of *high* over *low* expression curves as well as the mortality rates at 1, 2, and 5 years. Code for constructing NEEP values from clinical and expression data and calculating effect size statistics is available at https://github.com/thecodingdoc/neep.

Robustness

We measured robustness of the NEEP method by examining sensitivity of significant splice variants to changes in the set of patients. We conducted two rounds of 100 simulations, where 5% and 10% of patients were randomly removed, respectively. Since the number of patients in each simulation is then compared to the actual values, we expect the adjusted NEEP p-value change is due to a smaller sample size. So we compared ranks instead of p-values to isolate the sensitivity of NEEP p-values to resampling. We compared ranks of the significant isoforms in the full dataset against their rank in the simulations.

Multi-granular graphs

We utilized data across multiple granularities to construct plausible graphs for the association between splice variant expression and lung cancer survival. Specifically, we used protein-protein interactions (PPI) from BioGRID [20], domain-domain interactions (DDI) from 3did [21], splice variant domain prediction using HMMER3.1 [22], and the survival analysis significance values using NEEP, with the goal of identifying cancer-specific changes in PPI interactions.

The general structure of these multi-granular graphs (MGG) can be seen in Fig 1. The outcome of the workflow consists of quadripartite networks, connecting splice variants with their potential interaction partners via their domain-domain interaction, as *variant-domain-domain-variant* paths.

To build these graphs, we extracted protein-protein interactions from BioGRID v3.4.156 [20], retaining only physical interactions in human, resulting in 377,228 relationships. Of these interactions, 348,609 involved proteins with an active Ensembl ID. Because PPIs are reported at the gene product level, we expanded the interactions to include all possible variant-variant interactions (73,588,825). To identify graphs associated with survival, the left-most splice variant is required to be both survival-significant (NEEP FDR < 0.1) and protein-coding, leaving 655,407 unique possible interactions.

We identify two domain patterns of interest, "ghost" and "gained". Ghost domains are those that are present in survival-insignificant splice variants within the same gene, but are not found in the survival-significant splice variant. Gained domains are found in the survival-insignificant splice variants but present in the survival-significant splice variants. By incorporating these gained and ghost domains direct into their PPI context, we can consider MGGs as potential interaction changes caused by domain exclusion or inclusion. Genes without multiple splice variants were removed from further analysis. We downloaded the 11,200 domain-domain interactions (DDI) from 3did [21], where inclusion requires a known protein-protein structure to support the interaction. Variant-domain membership was determined using HMMER3.1 [23] with default parameters and the Pfam database [24]. After removal of interactions without DDI support, 5912 interactions remained. Of these, only 1208 had gained or ghost domains. There were 50 genes with an isoform significant for survival for these 1208 interactions. Code for constructing MGG interaction paths from variable data sources is available at https://github.com/scwest/SINBAD.

Identification of significant COSMIC mutation signatures

To examine if the splice variant RAD51C-202 was associated with homologous recombination mutations, we utilized COSMIC gene signatures [25]. We used the optimal NEEP threshold of RAD51C-202 to separate the patients into a low expression (344 patients) and high expression (88 patients) group. Mutation data for each patient was downloaded from TCGA for all 4 variant callers: MuSE [26], MuTect [27], SomaticSniper [28], and VarScan [29]. Mutation profiles for each patient were constructed as described as the 6 substitution types along with the neighboring bases for a total of 96 mutation types using only mutations found by all four variant callers. The Catalogue of Somatic Mutations In Cancer (COSMIC) has identified a set of 30 commonly identified mutation signatures deconvolved from cancer patient mutation profiles [25]. We used MutationalPatterns [30] in R to find the linear contributions of each of the 30 signatures to a patient profile. We used Mood's test for median to determine if the contributions were significantly different between low and high RAD51C-202 expression groups for each signature.

To check for possible confounding factors, we considered three smoking variables

reported by TCGA and checked whether they were different between the low and high ₃₇₄
RAD51C-202 expression groups: *cigarettes per day* was tested using the Welch two ₃₇₅
sample t-test; *years smoked* was tested using the Wilcoxon rank sum test; and the ₃₇₆
binary *smoker* variable was tested using the two-sample test for equal proportions. In ₃₇₇
addition, we checked if survival confounded the relationship between RAD51C-202 and ₃₇₈
signature 3. Because the causality between mutations and survival must be directional, ₃₇₉
we conducted Cox-PH survival analysis of RAD51C-202 with and without the ₃₈₀
contribution of Signature 3 as a confounding variable. Further, Mood's test was also ₃₈₁
conducted for the relationships between Signature 3 and the remaining splice variants ₃₈₂
with MGGs. The exact binomial test was used to determine if the proportion of splice ₃₈₃
variants significant for Signature 3 was greater than expected by chance. ₃₈₄

Supporting information ₃₈₅

S1 File. Enrichment analysis methods. ₃₈₆

S1 Fig. Enrichment term cluster membership across granularities. Counts ₃₈₇
of genes belonging to each enrichment cluster for the gene, isoform (splice variant), and ₃₈₈
MGG granularities are displayed as bar lengths. Missing bars does not indicate no ₃₈₉
membership, just insignificant enrichment of any term in the cluster. ₃₉₀

S2 File. Additional case studies of MGGs. ₃₉₁

S3 File. List of multi-granularity graphs. This table reports the entire list of ₃₉₂
MGGs for the TCGA LUAD dataset. The file does not contain the exact paths, only ₃₉₃
the members of each component of the MGG. ₃₉₄

Acknowledgments ₃₉₅

This research was partly funded by a University of Nebraska Collaboration ₃₉₆
Initiative/System Science Seed Grant to SK, HA, and DG and by the NIH AA026428 ₃₉₇
R21 grant to SK. ₃₉₈

References

1. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW. Cancer genome landscapes. Science. 2013;doi:10.1126/science.1235122.

2. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2017. CA: a cancer journal for clinicians. 2017;67(1):7–30.

3. Lee YT, Tan YJ, Oon CE. Molecular targeted therapy: Treating cancer with specificity. European journal of pharmacology. 2018;.

4. Germann S, Gratadou L, Dutertre M, Auboeuf D. Splicing programs and cancer. Journal of Nucleic Acids. 2012;2012. doi:10.1155/2012/269570.

5. Oltean S, Bates DO. Hallmarks of alternative splicing in cancer. Oncogene. 2014;doi:10.1038/onc.2013.533.

6. Lee SCW, Abdel-Wahab O. Therapeutic targeting of splicing in cancer. Nature Medicine. 2016;doi:10.1038/nm.4165.

7. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. F1000Research. 2015;4.

8. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, et al. Transcript assembly and abundance estimation from RNA-Seq reveals thousands of new transcripts and switching among isoforms. Nature biotechnology. 2010;28(5):511.

9. Sebestyén E, Zawisza M, Eyras E. Detection of recurrent alternative splicing switches in tumor samples reveals novel signatures of cancer. Nucleic Acids Research. 2015;doi:10.1093/nar/gku1392.

10. Yang IS, Son H, Kim S, Kim S. ISOexpresso: A web-based platform for isoform-level expression analysis in human cancer. BMC Genomics. 2016;doi:10.1186/s12864-016-2852-6.

11. Climente-González H, Porta-Pardo E, Godzik A, Eyras E. The Functional Impact of Alternative Splicing in Cancer. Cell Reports. 2017;20(9):2215–2226. doi:10.1016/j.celrep.2017.08.012.

12. Shen S, Wang Y, Wang C, Wu YN, Xing Y. SURVIV for survival analysis of mRNA isoform variation. Nature communications. 2016;7:11548.

13. Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, et al. A survey of best practices for RNA-seq data analysis. Genome biology. 2016;17(1):13.

14. Sehgal V, Seviour EG, Moss TJ, Mills GB, Azencott R, Ram PT. Robust selection algorithm (RSA) for multi-omic biomarker discovery; integration with functional network analysis to identify miRNA regulated pathways in multiple cancers. PloS one. 2015;10(10):e0140072.

15. Lausen B, Schumacher M. Maximally selected rank statistics. Biometrics. 1992; p. 73–85.

16. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence alignment/map format and SAMtools. Bioinformatics. 2009;25(16):2078–2079.

17. Patro R, Duggal G, Kingsford C. Salmon: accurate, versatile and ultrafast quantification from RNA-seq data using lightweight-alignment. Biorxiv. 2015; p. 021592.

18. Jin H, Wan YW, Liu Z. Comprehensive evaluation of RNA-seq quantification methods for linearity. BMC bioinformatics. 2017;18(4):117.

19. Zhang C, Zhang B, Lin LL, Zhao S. Evaluation and comparison of computational tools for RNA-seq isoform quantification. BMC genomics. 2017;18(1):583.

20. Chatr-Aryamontri A, Oughtred R, Boucher L, Rust J, Chang C, Kolas NK, et al. The BioGRID interaction database: 2017 update. Nucleic acids research. 2017;45(D1):D379.

21. Mosca R, Céol A, Stein A, Olivella R, Aloy P. 3did: a catalog of domain-based interactions of known three-dimensional structure. Nucleic acids research. 2013;42(D1):D379.

22. Johnson LS, Eddy SR, Portugaly E. Hidden Markov model speed heuristic and iterative HMM search procedure. BMC bioinformatics. 2010;11(1):431.

23. Finn RD, Clements J, Arndt W, Miller BL, Wheeler TJ, Schreiber F, et al. HMMER web server: 2015 update. Nucleic acids research. 2015;43(W1):W38.

24. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: the protein families database. Nucleic acids research. 2013;42(D1):D230.

25. Forbes SA, Bindal N, Bamford S, Cole C, Kok CY, Beare D, et al. COSMIC: mining complete cancer genomes in the Catalogue of Somatic Mutations in Cancer. Nucleic acids research. 2010;39(suppl_1):D950.

26. Fan Y, Xi L, Hughes DS, Zhang J, Zhang J, Futreal PA, et al. MuSE: accounting for tumor heterogeneity using a sample-specific error model improves sensitivity and specificity in mutation calling from sequencing data. Genome biology. 2016;17(1):178.

27. Cibulskis K, Lawrence MS, Carter SL, Sivachenko A, Jaffe D, Sougnez C, et al. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. Nature biotechnology. 2013;31(3):213.

28. Larson DE, Harris CC, Chen K, Koboldt DC, Abbott TE, Dooling DJ, et al. SomaticSniper: identification of somatic point mutations in whole genome sequencing data. Bioinformatics. 2011;28(3):311–317.

29. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. Genome research. 2012;.

30. Blokzijl F, Janssen R, van Boxtel R, Cuppen E. MutationalPatterns: comprehensive genome-wide analysis of mutational processes. Genome medicine. 2018;10(1):33.

31. Liu Y, Tarsounas M, O'Regan P, West SC. Role of RAD51C and XRCC3 in genetic recombination and DNA repair. Journal of Biological Chemistry. 2007;282(3):1973–1979.

32. Kuznetsov SG, Haines DC, Martin BK, Sharan SK. Loss of Rad51c leads to embryonic lethality and modulation of Trp53-dependent tumorigenesis in mice. Cancer research. 2009;69(3):863–872.

33. Mishra A, Saxena S, Kaushal A, Nagaraju G. RAD51C/XRCC3 facilitates mitochondrial DNA replication and maintains integrity of the mitochondrial genome. Molecular and cellular biology. 2018;38(3):489.

34. Pelttari LM, Kinnunen L, Kiiski JI, Khan S, Blomqvist C, Aittomäki K, et al. Screening of HELQ in breast and ovarian cancer families. Familial cancer. 2016;15(1):19–23.

35. ichi Takata K, Reh S, Tomida J, Person MD, Wood RD. Human DNA helicase HELQ participates in DNA interstrand crosslink tolerance with ATR and RAD51 paralogs. Nature communications. 2013;4:2338.

36. Heessen S, Masucci MG, Dantuma NP. The UBA2 domain functions as an intrinsic stabilization signal that protects Rad23 from proteasomal degradation. Molecular cell. 2005;18(2):225–235.

37. Zhang Y, Hunter T. Roles of Chk1 in cell biology and cancer therapy. International journal of cancer. 2014;134(5):1013–1023.

38. Wade SL, Auble DT. The Rad23 ubiquitin receptor, the proteasome and functional specificity in transcriptional control. Transcription. 2010;1(1):22–26.

39. Federico A, Rienzo M, Abbondanza C, Costa V, Ciccodicola A, Casamassimi A. Pan-cancer mutational and transcriptional analysis of the integrator complex. International journal of molecular sciences. 2017;18(5):936.

40. Baillat D, Wagner EJ. Integrator: surprisingly diverse functions in gene expression. Trends in biochemical sciences. 2015;40(5):257–264.