1 **Title: Developmental exposure to perfluorooctane sulfonate (PFOS) and perfluorooctanoic**

2 **acid (PFOA) selectively decreases brain dopamine levels in northern leopard frogs**

4 **Authors:** Rachel M. Foguth[1,2], R. Wesley Flynn[3], Chloe de Perre[4], Michael Iacchetta[3], Linda S.

5 Lee[4], Maria S. Sepúlveda[3], Jason R. Cannon[1,2,*]

7 **Affiliations:**   1. School of Health Sciences, Purdue University, West Lafayette, IN 47907

8   2. Purdue Institute for Integrative Neuroscience, Purdue University, West

9   Lafayette, IN 47907

10   3. Department of Forestry and Natural Resources, Purdue University, West

11   Lafayette, IN 47907

12   4. Department of Agronomy, Purdue University, West Lafayette, IN 47907

14 ***Corresponding author**:

15   Jason Cannon, Ph.D.

16   School of Health Sciences

17   Purdue University

18   207 S. Martin Jischke Dr.

19   West Lafayette, IN 47907

20   Tel: (765) 494-0794

21   Fax: (765) 496-1377

Abstract

Per- and polyfluoroalkyl substances (PFAS) are synthetic compounds that are a major public health concern due to widespread use, long environmental and biological half-lives, detection in most human plasma samples, and links to multiple adverse health outcomes. The literature suggests that some PFAS may be neurotoxic. However, there are major gaps in the literature with respect to how environmentally-relevant doses during development may influence the nervous system. To address this gap, we utilized a sentinel species, Northern leopard frogs (*Lithobates pipiens)* to determine the effects of developmental exposure to environmentally relevant perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA) on major neurotransmitter systems. Frog larvae at Gosner stage 25 were exposed to 10, 100, or 1000 ppb PFOS or PFOA for 30 days before neurochemical analysis. High performance liquid chromatography (HPLC) with electrochemical detection or fluorescent detection assays was used to measure neurotransmitter levels, which were normalized to protein levels in each sample. Dopamine (DA) decreased significantly in the brains of frogs treated with PFOA (1000 ppb) and PFOS (100 and 1000 ppb). Significant increases in DA turnover also resulted from PFOA and PFOS treatment. Neither PFOS, nor PFOA produced detectable alterations in serotonin (nor its metabolite), norepinephrine, gamma-amino butyric acid (GABA), glutamate, or acetylcholine. PFAS body burdens showed that PFOS accumulated relative to dose, while PFOA did not. These data suggest that DArgic neurotransmission is selectively affected in developmentally exposed amphibians and that PFAS should be evaluated for a potential role in diseases that target the DA system.

Keywords: Per- and polyfluoroalkyl substances; perfluorooctanoic acid; perfluorooctane sulfonate; neurotoxicity; dopamine

1. Introduction

Per- and polyfluoroalkyl substances (PFAS) are compounds that have been extensively used in manufacturing of surfactants, paper and packaging treatments, anti-static agents, corrosion inhibitors, insecticides, shampoos, and firefighting foams (Posner, 2012; Stahl *et al.*, 2011). PFAS are of significant concern due in part to the stable carbon-fluorine bond and, therefore, long half-life and potential for accumulation (Moody *et al.*, 2003). Perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA) have been extensively investigated for adverse health effects. While PFOS and PFOA have largely been phased out in favor of shorter chain length PFAS, they are ubiquitously detected in the global population (Jian *et al.*, 2018; Olsen *et al.*, 2009).

PFOS and PFOA both cross the blood-brain barrier, with PFOS generally accumulating 10-100x more in mammals than many other major PFAS (Maestri *et al.*, 2006). Studies in large mammals suggest that PFAS accumulate in the brain and are potentially neurotoxic. In polar bears, brain PFAS levels (including PFOS) were found to correlate with neurotransmitter alterations (Dassuncao *et al.*, 2019). Further, PFAS in North Atlantic pilot whales (*Globicephala melas*) were found to accumulate in brain, where only the liver had higher levels. Here, PFAS were strongly correlated with the high phospholipid content in the brain (Dassuncao et al., 2019). Laboratory studies in rodents on PFAS neurotoxicity have suggested neurobehavioral alterations. For example, mice exposed to various PFAS on post-natal day 10 showed that exposure increased spontaneous activity and increased proteins important for making synapses (Johansson *et al.*, 2009; Lee and Viberg, 2013). After pregnant dams were treated with PFOA, male offspring had significantly shorter latency to fall on the wire-hang test when they were 5-8 weeks old, suggesting a potential motor deficit (Onishchenko *et al.*, 2011). This implies that PFAS

exposure may affect brain development. Exposures to invertebrates have shown that *C. elegans*

have decreased motor function after 48 hour exposure to PFOS (Chen *et al.*, 2014). Finally, *in*

vitro exposures suggest that PFAS may influence neuronal differentiation, where, in PC12 cells,

PFOS promoted differentiation to cholinergic neurons at the expense of differentiation to

dopaminergic (DA)rgic neurons, whereas other tested PFAS suppressed or enhanced overall

differentiation (Slotkin *et al.*, 2008). This research suggests that developmental exposure could

lead to a long-term deficit in the number of DA neurons.

In general, there is a lack of data on PFAS neurotoxicity and potential relevance to

neurological diseases. Neurobehavioral alterations have been examined in a number of

developmental and adult studies, with some behavioral endpoints suggesting the cholinergic

system as a target (Johansson *et al.*, 2008). In vitro studies have suggested glutamatergic

signaling as a target (Liao *et al.*, 2009). However, environmental relevance of the tested doses is

generally lacking and to our knowledge no studies measured neurotransmitter levels from

multiple systems in response to PFAS. The primary goal of this report was to determine the

effects of developmental PFAS exposure at doses which include those detected in environmental

samples in a sentinel species. The long-term goal is to prompt specific research into how

exposure to PFAS during development may affect specific neurotransmitter systems and

influence neurological disease.

2. Methods

2.1 Animals, nominal exposure, and measured water and sediment PFAS levels.

All animal studies were approved by Purdue Animal Care and Use Committee standards.

The Northern leopard frog (*Lithobates pipiens*) was chosen as a relevant sentinel species

(Hoover *et al.*, 2017). Eggs were collected and reared to Gosner stage 25 in outdoor tanks prior

to the start of the experiment. Treatment tanks were prepared with 75 L of aged well water

(containing algae and zooplankton) and 10 kg of sediment spiked with PFOS or PFOA to

achieve nominal sediment concentrations of 0, 10, 100, or 1000 ppb. We chose to spike

sediments, instead of simple addition of chemicals to the water to represent a more ecologically

relevant exposure scenario. Tadpoles unintentionally consume sediment while grazing for algae

and detritus. Thus, spiking sediments provided more realism in incorporating all pertinent routes

of exposure amphibian larvae would likely experience in the environment. The doses were

selected to include concentrations detected at contaminated sites (Moody *et al.*, 2003). Measured

water and sediment concentrations, along with body burdens were determined by tandem mass

spectrometry from representative samples as previously reported (Hoover *et al.*, 2017). Exposure

lasted 30 days and tadpoles were sacrificed immediately using tricaine (MS-222). For this short

initial report on neurotoxicity, 6-14 frogs/treatment (PFOA or PFOS at 0 to1000 ppb) were

randomly selected from 32 treatment tanks (~20 tadpoles housed/tank) for detailed

neurotransmitter analysis (81 total samples analyzed for each neurotransmitter), where the

majority of the animals were devoted to a long-term, non-neuronal study on systemic

ecotoxicity. Brains were removed, frozen in liquid nitrogen, and stored at -80°C until processed.

2.2 Neurotransmitter Quantification

Six neurotransmitter systems were assessed. The majority of neurotransmitters were

analyzed by high performance liquid chromatography (HPLC) with electrochemical detection as

we have previously reported (Agim & Cannon, 2018; Cannon et al., 2009; Horowitz et al., 2011;

Wirbisky, Weber, Lee, Cannon, & Freeman, 2014; Wirbisky et al., 2015). Neurotransmitter

levels in the samples were quantified by measuring the area under the curve and comparing it to

a standard curve. Neurotransmitter levels were then normalized to protein levels in the sample

(ng neurotransmitter/mg protein). DA turnover was calculated as [(DOPAC + HVA)/DA where

DOPAC and HVA refer to DA metabolites 4-dihydroxyphenylacetic acid and homovanillic acid,

respectively] (Agim and Cannon, 2018).

Acetylcholine (ACh) was quantified using the Invitrogen Molecular Probes Amplex

ACh/Acetylcholinesterase Assay Kit (A12217). Briefly, samples were diluted 1:1 in the reaction

buffer. Samples and standards were incubated in 200 μM amplex red, 1 U/mL horseradish

peroxidase, 0.1 U/mL choline oxidase and 0.5 U/mL acetylcholinesterase 1:1 reaction buffer.

Concentrations were determined with fluorescence spectroscopy (530 nm excitation and 590 nm

emission). Acetylcholine levels were normalized to protein levels in the sample (ng

neurotransmitter/mg protein).

2.4 Statistical Analysis

Differences in bioaccumulation of PFAS between treatments were tested using linear

regression on log_{10}-transformed PFAS body burdens and comparisons of estimated marginal

means post-hoc analysis. Neurotransmitters were analyzed by one-way ANOVA with Sidak's

post-hoc test to compare all treatment groups within an exposure (PFOA or PFOS) and to

controls. Whole brain analyses of lower order species (i.e., zebrafish) produce considerably more

variable neurochemistry data than micro-dissected brain analysis in rodents (Dukes *et al.*, 2016;

Lee *et al.*, 2015). Thus, outlier analysis was conducted. Statistical analyses were conducted both

before and after outlier analysis. Importantly, outlier analyses for raw neurotransmitter data had

no effect on statistical significance, where no new significant comparisons were revealed, nor

were any significant comparisons eliminated after outlier analysis. Outliers were identified using

stringent ROUT analysis (Q = 0.1%) (Motulsky and Brown, 2006). Outlier analyses resulted in

no more than 3 samples/group being excluded, with the following total exclusions/endpoint: DA

(0/81 data points omitted), DOPAC (7/81 data points omitted), HVA (0/81 data points omitted),

5-HT (0/81 data points omitted), 5-HIAA (0/81 data points omitted), NE (1/81 data points

omitted), DA turnover (9/81), 5-HT turnover (1/81 data points omitted), glutamate (0/81 data

points omitted), GABA (11/81 data points omitted), and acetylcholine (0/81 data points omitted).

All data were expressed as mean ± standard error of the mean (SEM) and $p < 0.05$ used to

identify significant differences.

3. Results

3.1 Water, sediment, and body burden levels

Quantified PFAS exposure and body burden levels are shown in ***Table 1*** and ***Figure 1***.

150
151 **Table 1. PFAS applied and measured doses and total body burden**. Measured concentrations include sediment and water
152 analysis, along with whole body burden determination at the end of the 30-day exposure ($n = 4$-5). Control values = (PFOA + PFOS).

Exposure	Nominal sediment concentration (ppb)	Measured water concentration at initiation (ppb ± SEM)	Measured water concentration at completion (ppb ± SEM)	Measured sediment concentration at initiation (ppb ± SEM)	Measured sediment concentration at completion (ppb ± SEM)	Body burden (ppb ± SEM)
Control	0	0.000 ± 0.000	0.000 ± 0.000	0.102 ± 0.060	0.0778 ± 0.0768	27.7 ± 9.2
PFOA	10	3.22 ± 0.12	2.58 ± 0.17	18.9 ± 1.0	7.61 ± 0.40	205 ± 26
PFOA	100	8.34 ± 0.73	6.33 ± 0.13	42.0 ± 1.6	14.3 ± 1.1	161 ± 17
PFOA	1000	76.911 ± 2.603	55.379 ± 0.575	423 ± 22	140 ± 15	169 ± 24
PFOS	10	0.0812 ± 0.0390	0.0326 ± 0.0295	7.89 ± 0.34	5.94 ± 0.67	196 ± 41
PFOS	100	0.771 ± 0.078	1.15 ± 0.13	89.1 ± 4.7	45.6 ± 2.8	853 ± 46
PFOS	1000	16.4 ± 0.4	14.8 ± 0.3	739 ± 29	456 ± 45	4500 ± 660

153
3.2. Neurotransmitter levels

Quantification of catecholamine neurotransmitters showed a significant decrease in DA

in frogs treated with 1000 ppb PFOA and 100 or 1000 ppb PFOS, (3.61 ± 0.61 ng/mg protein,

1.62 ± 0.18 ng/mg protein, 2.41 ± 0.15 ng/mg protein, 2.09 ± 0.24 ng/mg protein for the control,

1000 ppb PFOA, 100 ppb PFOS, and 1000 ppb PFOS, respectively) (***Figure 2A***), but no

significant changes in the DA metabolites DOPAC or HVA (***Figure 2B, C***). DA decreases,

relative to metabolite levels led to a significant increase in DA turnover [(DOPAC + HVA)/DA]

for both 1000 ppb PFOS and PFOA (0.332 ± 0.051, 1.22 ± 0.20, 0.693 ± 0.085, 1.02 ± 0.10 for

the control, 1000 ppb PFOA, 100 ppb PFOS, and 1000 ppb PFOS, respectively) (*Figure 2D*).

Neither PFOS nor PFOA caused significant changes in norepinephrine (NE) levels (*Figure 2E*).

Neither serotonin (5-HT) nor its metabolite 5-hydroxyindoleacetic acid (5-HIAA) were

significantly affected by treatment with either PFAS, leading to no changes in 5-HT metabolism

(*Figure 2F - H*).

The amino acid neurotransmitters glutamate and gamma-aminobutyric acid (GABA)

were measured and no significant differences were found in treatment groups versus the control

for either glutamate or GABA (*Figure 3A, B*). In addition, there was no significant change in

ACh levels in any treatment group compared to the control (*Figure 3C*).

4. Discussion

There are major gaps in the literature related to PFAS neurotoxicity. Here, we have

shown that developmental exposure within the concentration range documented at contaminated

sites produce selective total brain DA depletion in an amphibian sentinel species. Further, in

general, we observed that PFAS exposure increased DA turnover, which is typically a

compensatory mechanism in response to DArgic neurotoxicity and DA depletion; and an

additional metric of damage to DArgic function (Sossi *et al.*, 2002). This short report is expected

to prompt research on the neurochemical and neuropathological effects of PFAS on the brain.

Further, our findings suggest that the potential role of PFAS in neurological diseases affecting

DArgic neurotransmission should be examined.

DA depletion in the frog brain could have several implications, especially given that there

are some significant advantages of specific frog species as a model. Leopard frogs are unique in

that they have neuromelanin-containing DArgic neurons in the ventral motor neurons which

could be related to the DArgic neurons in the nigrostriatal DA system that are affected in

Parkinson's disease (PD). Importantly, these pigmented neurons have been shown to bind PD-

relevant toxicants, where high-neuromelanin expressing species exhibit heightened sensitivity

with respect to low-neuromelanin expressing species (Karlsson *et al.*, 2009; Lindquist *et al.*,

1988; Sokolowski *et al.*, 1990; Spencer *et al.*, 2004). Taken together, the literature suggests

that, with respect to neuromelanin expression and toxicant binding to neuromelanin, certain

amphibian species are more representative of human DArgic neurons than low-expressing rodent

models. Thus, the potential relevance of DA depletion in PFAS-treated amphibians to PD should

be further examined. An additional common DArgic pathway in mammals is the reward system

with DArgic neurons extending from the ventral tegmental area to the ventral striatum (Lammel

et al., 2014). Frogs have a similar DArgic pathway for reward that contains DArgic neurons in

the anteroventral tegmental area, thus the DA depletion that we observed could potentially result

in suppression of the reward pathway (O'Connell *et al.*, 2010). DArgic neurons are also

important for the hypothalamic function and multiple neuroendocrine axes that control well-

conserved behaviors such as metabolic rate stress, digestion, and mood (Hu *et al.*, 2008). Thus,

neuroendocrine function in response to PFAS should also be tested.

Another important aspect of these findings potentially relates to developmental origins of

adult health and disease (DOHaD). It will be important to examine whether developmental PFAS

exposures result in increased susceptibility to neurodegenerative diseases. PD, for example is a

disease of aging, where, once the nigrostriatal DA reserve is depleted (i.e., loss of DA neurons

reaches a symptomatic threshold), the onset of clinical symptoms appear (Tabbal *et al.*, 2012).

Testing whether a developmental PFAS exposure may lower such reserve and increase

susceptibility later in life is of importance.

Relationships between PFAS treatments and body burden show log-linear

bioaccumulation of PFOS, whereas PFOA did not, which is consistent with the bioaccumulative

properties of PFOS, especially in the brain (Maestri *et al.*, 2006). Differences between nominal

and measured sediment PFAS levels are due to redistribution of the PFAS between sediment and

water, and to a lesser part from a mass balance perspective, to uptake by algal communities and

the frogs. Mass balance checks (water, sediment, and frogs) are generally within 15% of the

nominal mass added to the system and calculated sediment-water partition coefficients are in the

range reported in the literature (Higgins and Luthy, 2006). PFAS levels observed in control

animals were low and to be expected, given the ubiquitous global distribution of PFAS. We

hypothesize that PFAS detected in control animals could be derived from the natural water

source used to fill tanks, algae, zooplankton and/or leaching from the plastic.

There are some notable limitations in the present study. Neurochemical quantification is a

frequent endpoint studied for effects of compounds on the brain in rodents, but has also been

measured in aquatic species, both in whole zebrafish larva and brains (Cannon *et al.*, 2009;

Dukes *et al.*, 2016; Milanese *et al.*, 2012; Wirbisky *et al.*, 2014; Wirbisky *et al.*, 2015).

However, quantification in aquatic species is more difficult and more variable than specific

region neurotransmitter quantification in rodents. Clearly, our samples exhibited increased

variability relative to micro-dissected rodent studies. Indeed, GABA and ACh analyses exhibited

potential nonsignificant trends resulting from PFAS treatment, suggesting further examination.

Future studies will require an increased number of animals/group, along with sub-regional

analysis.

In summary, our findings suggest developmental PFAS exposure in a sentinel species at

environmentally-relevant doses produce selective DA depletion. These findings imply that the

role of PFAS in neurological diseases involving DArgic neurotransmission should be evaluated.

5. Funding

Ralph W. and Grace M. Showalter Research Trust (to J.R.C.). Strategic Environmental Research

and Development Program (ER-2626) (to M.S.S and L.S.L.).

6. References

Agim, Z. S., and Cannon, J. R. (2018). Alterations in the nigrostriatal dopamine system after

acute systemic PhIP exposure. *Toxicology letters* **287**, 31-41, 10.1016/j.toxlet.2018.01.017.

Cannon, J. R., Tapias, V., Na, H. M., Honick, A. S., Drolet, R. E., and Greenamyre, J. T. (2009).

A highly reproducible rotenone model of Parkinson's disease. *Neurobiol Dis* **34**(2), 279-90.

Chen, N., Li, J., Li, D., Yang, Y., and He, D. (2014). Chronic exposure to perfluorooctane

sulfonate induces behavior defects and neurotoxicity through oxidative damages, in vivo and in

vitro. *PloS one* **9**(11), e113453, 10.1371/journal.pone.0113453.

Dassuncao, C., Pickard, H., Pfohl, M., Tokranov, A. K., Li, M., Mikkelsen, B., Slitt, A., and

Sunderland, E. M. (2019). Phospholipid Levels Predict the Tissue Distribution of Poly- and

Perfluoroalkyl Substances in a Marine Mammal. *Environ Sci Tech Let* **6**(3), 119-125,

10.1021/acs.estlett.9b00031.

Dukes, A. A., et al. (2016). Live imaging of mitochondrial dynamics in CNS dopaminergic

neurons in vivo demonstrates early reversal of mitochondrial transport following MPP+

exposure. *Neurobiol Dis* **95**, 238-249, 10.1016/j.nbd.2016.07.020.

Higgins, C. P., and Luthy, R. G. (2006). Sorption of perfluorinated surfactants on sediments.

Environmental science & technology **40**(23), 7251-6.

Hoover, G. M., Chislock, M. F., Tornabene, B. J., Guffey, S. C., Choi, Y. J., De Perre, C.,

Hoverman, J. T., Lee, L. S., and Sepulveda, M. S. (2017). Uptake and Depuration of Four

Per/Polyfluoroalkyl Substances (PFASS) in Northern Leopard Frog Rana pipiens Tadpoles.

Environ Sci Tech Let **4**(10), 399-403, 10.1021/acs.estlett.7b00339.

Hu, F., Crespi, E. J., and Denver, R. J. (2008). Programming neuroendocrine stress axis activity

by exposure to glucocorticoids during postembryonic development of the frog, Xenopus laevis.

Endocrinology **149**(11), 5470-81, 10.1210/en.2008-0767.

Jian, J. M., Chen, D., Han, F. J., Guo, Y., Zeng, L., Lu, X., and Wang, F. (2018). A short review

on human exposure to and tissue distribution of per- and polyfluoroalkyl substances (PFASs). *Sci*

Total Environ **636**, 1058-1069, 10.1016/j.scitotenv.2018.04.380.

Johansson, N., Eriksson, P., and Viberg, H. (2009). Neonatal exposure to PFOS and PFOA in

mice results in changes in proteins which are important for neuronal growth and synaptogenesis

in the developing brain. *Toxicological sciences : an official journal of the Society of Toxicology*

108(2), 412-8, 10.1093/toxsci/kfp029.

Johansson, N., Fredriksson, A., and Eriksson, P. (2008). Neonatal exposure to perfluorooctane

sulfonate (PFOS) and perfluorooctanoic acid (PFOA) causes neurobehavioural defects in adult

mice. *Neurotoxicology* **29**(1), 160-9, 10.1016/j.neuro.2007.10.008.

Karlsson, O., Berg, C., Brittebo, E. B., and Lindquist, N. G. (2009). Retention of the

cyanobacterial neurotoxin beta-N-methylamino-l-alanine in melanin and neuromelanin-

containing cells--a possible link between Parkinson-dementia complex and pigmentary

retinopathy. *Pigment Cell Melanoma Res* **22**(1), 120-30, 10.1111/j.1755-148X.2008.00508.x.

Lammel, S., Lim, B. K., and Malenka, R. C. (2014). Reward and aversion in a heterogeneous

midbrain   dopamine   system.   *Neuropharmacology*   **76**   **Pt**   **B**,   351-9,

10.1016/j.neuropharm.2013.03.019.

Lee, I., and Viberg, H. (2013). A single neonatal exposure to perfluorohexane sulfonate (PFHxS)

affects the levels of important neuroproteins in the developing mouse brain. *Neurotoxicology* **37**,

190-6, 10.1016/j.neuro.2013.05.007.

Lee, J. W., Tapias, V., Di Maio, R., Greenamyre, J. T., and Cannon, J. R. (2015). Behavioral,

neurochemical, and pathologic alterations in bacterial artificial chromosome transgenic G2019S

leucine-rich   repeated   kinase   2   rats.   *Neurobiology   of   aging*   **36**(1),   505-18,

10.1016/j.neurobiolaging.2014.07.011.

Liao, C., Wang, T., Cui, L., Zhou, Q., Duan, S., and Jiang, G. (2009). Changes in synaptic

transmission, calcium current, and neurite growth by perfluorinated compounds are dependent on

the chain length and functional group. *Environmental science & technology* **43**(6), 2099-104.

Lindquist, N. G., Larsson, B. S., and Lyden-Sokolowski, A. (1988). Autoradiography of

[14C]paraquat or [14C]diquat in frogs and mice: accumulation in neuromelanin. *Neurosci Lett*

93(1), 1-6.

Maestri, L., Negri, S., Ferrari, M., Ghittori, S., Fabris, F., Danesino, P., and Imbriani, M. (2006).

Determination of perfluorooctanoic acid and perfluorooctanesulfonate in human tissues by liquid

chromatography/single quadrupole mass spectrometry. *Rapid Commun Mass Spectrom* **20**(18),

2728-34, 10.1002/rcm.2661.

Milanese, C., Sager, J. J., Bai, Q., Farrell, T. C., Cannon, J. R., Greenamyre, J. T., and Burton, E.

A. (2012). Hypokinesia and reduced dopamine levels in zebrafish lacking beta- and gamma1-

synucleins. *The Journal of biological chemistry* **287**(5), 2971-83, 10.1074/jbc.M111.308312.

Moody, C. A., Hebert, G. N., Strauss, S. H., and Field, J. A. (2003). Occurrence and persistence

of perfluorooctanesulfonate and other perfluorinated surfactants in groundwater at a fire-training

area at Wurtsmith Air Force Base, Michigan, USA. *J Environ Monit* **5**(2), 341-5.

Motulsky, H. J., and Brown, R. E. (2006). Detecting outliers when fitting data with nonlinear

regression - a new method based on robust nonlinear regression and the false discovery rate.

BMC Bioinformatics **7**, 123, 10.1186/1471-2105-7-123.

O'Connell, L. A., Matthews, B. J., Ryan, M. J., and Hofmann, H. A. (2010). Characterization of

the dopamine system in the brain of the túngara frog, Physalaemus pustulosus. *Brain Behav Evol*

76(3-4), 211-25, 10.1159/000321715.

Olsen, G. W., Butenhoff, J. L., and Zobel, L. R. (2009). Perfluoroalkyl chemicals and human

fetal development: an epidemiologic review with clinical and toxicological perspectives. *Reprod*

Toxicol **27**(3-4), 212-30, 10.1016/j.reprotox.2009.02.001.

Onishchenko, N., Fischer, C., Wan Ibrahim, W. N., Negri, S., Spulber, S., Cottica, D., and

Ceccatelli, S. (2011). Prenatal exposure to PFOS or PFOA alters motor function in mice in a sex-

related manner. *Neurotoxicity research* **19**(3), 452-61, 10.1007/s12640-010-9200-4.

Posner, S. (2012). Perfluorinated Compounds: Occurrence and Uses in Products. In

Polyfluorinated Chemicals and Transformation Products (T. P. Knepper, and F. T. Lange, Eds.)

doi: 10.1007/978-3-642-21872-91st ed., pp. 25-39. Springer-Verlag Berlin Heidelberg, Berlin,

Heidelberg.

Slotkin, T. A., MacKillop, E. A., Melnick, R. L., Thayer, K. A., and Seidler, F. J. (2008).

Developmental neurotoxicity of perfluorinated chemicals modeled in vitro. *Environmental health*

perspectives **116**(6), 716-22, 10.1289/ehp.11253.

Sokolowski, A. L., Larsson, B. S., and Lindquist, N. G. (1990). Distribution of 1-(3H)-methyl-4-

phenyl-1,2,3,6-tetrahydropyridine (3H-MPTP) in the frog: uptake in neuromelanin. *Pharmacol*

Toxicol **66**(4), 252-8.

Sossi, V., de La Fuente-Fernandez, R., Holden, J. E., Doudet, D. J., McKenzie, J., Stoessl, A. J.,

and Ruth, T. J. (2002). Increase in dopamine turnover occurs early in Parkinson's disease:

evidence from a new modeling approach to PET 18 F-fluorodopa data. *J Cereb Blood Flow*

Metab **22**(2), 232-9, 10.1097/00004647-200202000-00011.

Spencer, S. J., Ebner, K., and Day, T. A. (2004). Differential involvement of rat medial

prefrontal cortex dopamine receptors in modulation of hypothalamic-pituitary-adrenal axis

responses to different stressors. *The European journal of neuroscience* **20**(4), 1008-16,

10.1111/j.1460-9568.2004.03569.x.

Stahl, T., Mattern, D., and Brunn, H. (2011). Toxicology of Perfluorinated compounds. *Environ*

mental Sciences Europe **23**(38), 1-52.

Tabbal, S. D., Tian, L., Karimi, M., Brown, C. A., Loftin, S. K., and Perlmutter, J. S. (2012).

Low nigrostriatal reserve for motor parkinsonism in nonhuman primates. *Exp Neurol* **237**(2),

355-62, 10.1016/j.expneurol.2012.07.008.

Wirbisky, S. E., Weber, G. J., Lee, J. W., Cannon, J. R., and Freeman, J. L. (2014). Novel dose-

dependent alterations in excitatory GABA during embryonic development associated with lead

(Pb) neurotoxicity. *Toxicology letters* **229**(1), 1-8, 10.1016/j.toxlet.2014.05.016.

Wirbisky, S. E., Weber, G. J., Sepulveda, M. S., Xiao, C., Cannon, J. R., and Freeman, J. L.

(2015). Developmental origins of neurotransmitter and transcriptome alterations in adult female

zebrafish exposed to atrazine during embryogenesis. *Toxicology* **333**, 156-167,

10.1016/j.tox.2015.04.016.

Figure Legends:

Figure 1. Total body burden in PFAS-treated frogs 30 days after treatment (mean +/-

SEM). A regression shows that all PFAS treatments have greater body burdens than controls.

Post-hoc analysis showed that: 1) the two highest PFOS treatment were significantly greater than

the low PFOS treatment and all PFOA treatments, 2) the 1000ppb PFOS treatment was greater

than the 100ppb PFOS, and 3) that none of the PFOA treatments differed from each other. Note,

y-axis is $\log_{10}$-transformed.

Figure 2. Developmental exposure to PFOA or PFOS selectively decreases dopamine levels.

Northern leopard frog (*Lithobates pipiens*) tadpoles at Gosner stage 25 were exposed to PFOS or

PFOA for 30 days. Whole brain monoamines and major metabolites were quantified using HPLC

with electrochemical detection and normalized to amount of protein per sample: Dopamine (DA)

(*A*); 3,4-dihydroxyphenylacetic acid (DOPAC) (*B*); homovanillic acid (HVA) (*C*); DA turnover

[(DOPAC + HVA)/DA] (*D*); norepinephrine (NE) (*E*); serotonin (5-HT) (*F*); 5-

hydroxyindoleacetic acid (5-HIAA) (*G*); 5-HT turnover (5-HIAA/5-HT) (*H*). Data was analyzed

using one-way ANOVA followed by Sidak's post hoc test. *p<0.05, **p<0.005, ***p<0.0001;

n=6-14/group.

Figure 3. Developmental exposure to PFOA or PFOS does not affect amino acid or

cholinergic neurotransmitter levels. Northern leopard frog (*Lithobates pipiens*) tadpoles at

Gosner stage 25 were exposed to PFOS or PFOA for 30 days. Whole brain amino acid

neurotransmitters were quantified using HPLC with electrochemical detection: glutamate (*A*);

and gamma-aminobutyric acid (GABA) (*B*). Whole brain acetylcholine was quantified using the

Invitrogen Molecular Probes Amplex Acetylcholine/Acetylcholinesterase Assay Kit with

fluorometric detection. Neurotransmitters were normalized to amount of protein per sample.

Data was analyzed using one-way ANOVA followed by Sidak's post hoc test; n=6-14/group.

A

DA

B

DOPAC

C

HVA

D

DA Turnover

E

NE

F

5-HT

G

5-HIAA

H

5-HT Turnover