**Tmem119-EGFP and Tmem119-CreERT2 transgenic mice for labeling and manipulating**

**microglia**

Tobias Kaiser[1,2] and Guoping Feng[1,2,3,4]

[1] McGovern Institute for Brain Research
[2] Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology,
Cambridge, MA 02139, USA
[3] Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA
02142, USA
[4] Correspondence: fengg@mit.edu
10

**Abstract**

Microglia are specialized brain-resident macrophages with important functions in health

and disease. To improve our understanding of these cells, the research community needs genetic

tools to identify and control them in a manner that distinguishes them from closely related cell-

types. We have targeted the recently discovered microglia-specific *Tmem119* gene to generate

knock-in mice expressing EGFP (JAX#031823) or CreERT2 (JAX#031820) for the

identification and manipulation of microglia, respectively. Genetic characterization of the locus

and qPCR-based analysis demonstrate correct positioning of the transgenes and intact expression

of endogenous *Tmem119* in the knock-in mouse models. Immunofluorescence analysis further

shows that parenchymal microglia, but not other brain macrophages, are completely and

faithfully labeled in the EGFP-line at different time points of development. Flow cytometry

indicates highly selective expression of EGFP in CD11b$^+$CD45lo microglia. Similarly,

immunofluorescence and flow cytometry analyses using a Cre-dependent reporter mouse line

demonstrate activity of CreERT2 in microglia upon tamoxifen administration. Finally, flow

cytometric analyses reveal absence of EGFP expression and minimal activity of CreERT2 in

blood monocytes of the *Tmem119-EGFP* and *Tmem119-CreERT2* lines, respectively. These new

transgenic lines extend the microglia toolbox by providing the currently most specific genetic

labeling and control over these cells in the myeloid compartment of mice.

**Visual abstract**

**Significance statement**

Tools that specifically label and manipulate only microglia are currently unavailable, but

are critically needed to further our understanding of this cell type. Complementing and

significantly extending recently introduced microglia-specific immunostaining methods that have

quickly become a new standard in the field, we generated two mouse lines that label and control

gene expression in microglia with high specificity and made them publicly available. Using these

readily accessible mice, the research community will be able to study microglia biology with

improved specificity.

**Introduction**

Microglia are specialized brain-resident macrophages that comprise 5 to 12% of the glial cells in the adult mouse brain (Lawson et al., 1990). Under physiological conditions, phagocytic and secretory activity of these glia support neurogenesis, development of neuronal connectivity, and survival of neurons (Stevens et al., 2007; Sierra et al., 2010; Paolicelli et al., 2011; Schafer et al., 2012; Ueno et al., 2013; Weinhard et al., 2018). Complementing these homeostatic functions, microglia react to perturbations, which has been shown in the context of vascular injury, multiple sclerosis lesions, and neurodegeneration (Itagaki et al., 1989; Davalos et al., 2005; Ransohoff, 2016; Aguzzi and Zhu, 2017; Keren-Shaul et al., 2017; Mathys et al., 2017; O'Loughlin et al., 2018). In the future, many of these processes and their potential for therapeutic targeting will be further examined, and understanding of both homeostatic and disease-related contributions of microglia will critically depend on means to specifically identify and control them. Several mouse lines are currently available for either fluorescent labeling or Cre-expression that harness the loci of putative microglia signature genes *Tie2, Runx1, Csf1r, Aif1, Lyz2, Itgam, Sall1,* or *CX3cr1* (Clausen et al., 1999; Sasmono et al., 2003; Ferron and Vacher, 2005; Hirasawa et al., 2005; Samokhvalov et al., 2007; Ginhoux et al., 2010; Parkhurst et al., 2013; Yona et al., 2013; Buttgereit et al., 2016). Collectively, these mouse lines have been instrumental in gaining insights on microglia. However, furthering our understanding of microglia using the currently available lines is complicated by the fact that it is difficult to distinguish microglia from other closely related peripheral and central cell types such as blood monocytes as well as perivascular, choroid plexus, and meningeal macrophages (Wieghofer and Prinz, 2016; Haimon et al., 2018).

Recent advances in RNA sequencing and other cell profiling technologies have enabled the discovery of cell type-specific signature genes (Butovsky et al., 2013). Amongst these, transmembrane protein 119 (*Tmem119*) is highly and exclusively expressed in microglia in the brains of mice and humans (Bennett et al., 2016; Satoh et al., 2016). While TMEM119-specific antibodies are already widely used as a gold standard for immunohistochemical methods, faithful fluorescent reporter lines and inducible Cre-lines that allow *in vivo* observation and manipulation are currently not available. Here, we report the generation and characterization of knock-in mouse lines *Tmem119-EGFP* (JAX#031823) and *Tmem119-CreERT2* (JAX#031820), where microglia express EGFP and CreERT2, respectively, while preserving endogenous *Tmem119*

expression. We demonstrate that EGFP is expressed throughout the brain and that the tag is

confined to microglia only, without significantly labeling other brain macrophages. We further

provide evidence that the inducible Cre is specifically active in microglia by crossing to the

conditional fluorescent reporter mouse line Ai14. Finally, we demonstrate minimal to absent

transgene expression in monocytes of the mice. These publicly available mouse lines provide

valuable tools for the functional study of bona fide microglia.

**Materials and methods**

*Animal work*

All animal procedures were performed in accordance with the [Author University] animal

care committee's regulations. Overall, more than 30 animals of more than 10 litters were

generated per line and mice were crossed to C57BL/6J up to generation N3. For all experiments,

at least three independent mice were analyzed, which included both sexes and no apparent sex

differences were observed.

*Generation of transgenic animals using CRISPR/Cas9*

To generate *Tmem119* knock-in mice, donor DNA templates encoding ribosome-skipping

peptide *porcine teschovirus-1 polyprotein* and *EGFP* (*P2A-EGFP*) or *CreERT2* (*P2A-CreERT2*)

were synthesized. These sequences were flanked by sequences of 55-300 bp for EGFP and 1.5

kb for CreERT2 homologous to 5' and 3' regions around the *Tmem119* stop codon. These

templates were injected into fertilized mouse oocytes together with a single crRNA

(AGUCUCCCCAGUGUCUAAC, Synthego) that cuts at the stop codon.

Donor DNA templates were generated by digesting pAAV-P2A-EGFP (sequence below)

with XbaI and EcoRI (New England Biolabs) and inserting three gblocks (LHA, P2A-EGFP or

P2A-CreERT2, RHA, purchased from IDT, sequences below) using Gibson cloning (HIFI

assembly mix, New England Biolabs) according to the manufacturers protocol. Resulting

plasmids were purified and sequenced. For CreERT2, the highly purified dsDNA plasmid was

directly used as donor DNA in injections. For EGFP, single strand DNA (ssDNA) was produced

using PCR with forward primers for left homology arms between 55 and 300 bp (5'-

A*G*C*AACTGGTCCTCCTGAAA-3' and 5'-CAAAGCCTGTGAAGGGTGGG-3',

respectively, * denoting phosphorothioate) and a reverse primer for a 55 bp right homology arm

(5'-CAAAGAGGTGACCCTCAAGG-3', with 5' phosphorylation for lambda digest of antisense

strand). Using these primers, large scale PCR with Takara PrimeStar was performed to obtain

40-60 µg of product. The PCR product was highly purified with the Qiagen PCR purification kit

and subject to digestion of the antisense strand. Lambda exonuclease (New England Biolabs)

was used to digest 20 µg dsDNA at 37 °C for 60 min. Complete digestion of dsDNA was

confirmed by agarose gel electrophoresis and Sanger sequencing with sense- and antisense-

binding primers.

Mixtures for injection of zygotes were prepared freshly on the morning of the day of

injection. Briefly, water to a final volume of 100 µl was mixed with final concentrations of 10

mM TrisHcl buffer, 0.61 µM crRNA, 0.61 µM tracrRNA (sequence proprietary to Synthego) and

heated to 95 °C for 5 min. Heated mixtures were cooled to room temperature for 10 min and 30

ng/µl EnGen Cas9 protein (New England Biolabs) was added. Mixtures were incubated at 37 °C

for 15 min to form Cas9 to crRNA-tracrRNA complexes. Final concentrations of 5 ng/µl donor

DNA (P2A-EGFP ssDNA or P2A-CreERT2 plasmid dsDNA) and 10 ng/µl recombinant RAD51

protein were added. Mixtures were kept on ice until use, when they were incubated at 37 °C for

15 min followed by centrifugation at 10.000 rpm for 1 min to prevent clogging of the

micropipette.

Injection of mixtures was carried out using standardized protocols of the transgenics

facility in zygotes obtained from pure C57BL/B6N mice (Taconic).

*Genetic analysis and genotyping of Tmem119 mice and Ai14*

Founder mice were genetically examined by amplifying sequences spanning the 5' and 3'

junction and including the entire inserted transgene. Specifically, high quality DNA was obtained

from ear punches using a tissue DNA extraction kit according to the manufacturer's instructions

(Macherey-Nagel). One amplicon spanning approximately 1.5 kb (open arrows, Figure 1) of

these sequences was generated using Q5 polymerase with primers 5'ofLHA-F

(GCCTCTGTCACTTAAGTTGG) and P2A-R (GCTTCAGCAGGCTGAAGTTA). A second

amplicon of about 2.5-3.5 kb (closed arrows, Figure 1) was generated using primers Tmem-

LHA-P2A-F (CAGTGTCGGAAGCGGAGCTA) and 3'ofRHA-R

(GAAAGAGGAAGCTAGAAGGG). Both amplicons were purified and then sequenced using

primers spanning the entire length using Sanger sequencing (Genewiz). Trace files were aligned

to *in silico* assemblies and analyzed using Snapgene.

To genotype mice, ear tissue was prepared using an alkaline buffer (25 mM NaOH, 0.2

mM $Na_2EDTA$, pH=12) at 95 °C for 30 min. The lysed tissue solution was neutralized using an

acidic buffer (40 mM TrisHcl, pH=5). PCR using primers WT-F

GTCAGGAGGAGGCCCAGGAA, EGFP-F CTGCTGCCCGACAACCACTA, CreERT2-F

ACCGCCTACATGCGCCCACT, and Common-R GTTTCCTGGGGTGCACCAGA yielded

products of 400 bp for the wildtype allele and 320 bp for the EGFP or CreERT2 allele.

Ai14(RCL-tdT) mice were obtained from JAX (Stock 007914) and maintained in house.

Homozygous Ai14 mice were bred to heterozygous *Tmem119-CreERT2* mice to generate

offspring for the analysis. Ai14 mice were genotyped according to the protocol provided by JAX.

*RNA isolation and quantitative PCR (qPCR)*

Brain hemispheres from adult *Tmem119-EGFP$^{+/-}$* mice were dissected and snap-frozen in

liquid nitrogen. Highly pure RNA was isolated using an RNeasy purification kit (Qiagen)

following the manufacturer's instructions. Equal amounts of RNA (1.5 µg per sample) were

reverse transcribed using an iScript Advanced cDNA synthesis kit (Bio-Rad). Resulting cDNA

was diluted 1:50 in ultrapure water. qPCR was carried out using SsOAdvanced Universal SYBR

Green Supermix (Bio-Rad) on a CFX96 realtime system. Primers used were exon-spanning

whenever possible and of the following sequences: Tmem119-F

CCTTCACCCAGAGCTGGTTC, Tmem119-R GGCTACATCCTCCAGGAAGG, GAPDH-F

GCCTTCCGTGTTCCTACC, GAPDH-R CCTCAGTGTAGCCCAAGATG, b-actin-F

CTAAGGCCAACCGTGAAAAG, b-actin-R ACCAGAGGCATACAGGGACA. Differential

gene expression analysis was performed using built-in software for the Bio-Rad CFX96 realtime

system.

*Tamoxifen administration*

Tamoxifen (Sigma, T5648) was dissolved in corn oil at 20 mg/ml under agitation for

several hours in the dark and kept at room temperature for 2–3 days. Using needles from Harvard

Apparatus (52-4025), separately housed adult animals were fed via oral gavage for 3 days.

Needles were rinsed between days and syringes were replaced. At time of administration, the

mice weighted approximately 19-26 g and received 250-400 µl of the 20 mg/ml tamoxifen

solution, corresponding to 0.2 mg per g body weight. Postnatal day 2 mice weighing

approximately 2 g received 5 µl of the 20 mg/ml solution, corresponding to 50 µg per g body

weight per day for three consecutive days. The health of the mice was monitored daily. At least 9

days week after the last dose, mice were sacrificed by transcardiac perfusion.

*Immunofluorescence staining and imaging*

Adult mice were deeply anesthetized and perfused with 25 ml phosphate buffered saline

(PBS) followed by 25 ml 4% paraformaldehyde (PFA) in PBS. Early postnatal mice (P3) were

perfused with 8 ml PBS and 8 ml PFA. Brains were surgically removed and postfixed in the

fixative at 4 °C for 24 h. Fixed brain were washed in PBS once and sliced into 100 µm-thick

sagittal slices using a Leica VT1000S. Slices were washed twice in PBS, permeabilized in 1.2%

TX100 in PBS for 15 min, washed twice in PBS, and subject to incubation in blocking solution

(5% normal goat serum, 2% bovine serum albumin, 0.2% TX100 in PBS). Blocked sections were

incubated with primary antibodies for IBA1 (1:500, Synaptic Systems, 234006), GFP (1:1000,

Invitrogen, A11122, 1:500, Aves Labs, GFP-1020), TMEM119 (1:1000, Abcam, ab209064),

Olig2 (1:1000, Millipore, AB9610), NeuN (1:1000, Millipore, MAB377), GFAP (1:1000, Sigma,

G9269), or CD163 (1:200, Abcam, ab182422, antigen retrieval required) for 24 h at 4 °C.

Primary antibody incubation was followed by three washes in PBS and incubation with species-

matched and Alexa fluorophore-conjugated secondary antibodies raised in goat (Invitrogen,

1:1000) for 2 h. DAPI (1:10.000) was included in a washing step or secondary antibody

incubation. Slices were washed three times in PBS and mounted and coverslipped using

vectashield H-1000 mounting medium.

For immunostaining of CD163, an antigen retrieval step was included. Briefly, vibratome

slices were washed twice in PBS, incubated in a retrieval buffer (10 mM Sodium citrate, pH 8.5)

for 5 min, followed by incubation in the retrieval buffer at 80 °C for 30 min. Sections were

cooled to room temperature and washed twice in PBS. Blocking and immunostaining were then

carried out as described above.

For imaging, slides were scanned on an Olympus Fluoview FV1000 fixed stage confocal

microscope (high power magnifications) or Olympus BX61 epifluorescence microscope (sagittal

section montage) using built-in software. For co-expression analysis in the *Tmem119-EGFP* line,

maximum projections of 4 z-stacks from three 20X high-power fields (approximately 100 cells

per region) were counted manually.

For microglia morphology analysis, z-stacks were acquired at 20X magnification with 2X

digital zoom and 1.2 µm step size. Images were imported as stacks into Neurolucida software.

Cell bodies in the middle of the stacks with complete process arbors were selected and manually

traced. Raw trace files were imported into Neurolucida explorer software and convex hull area,

number of processes and total process length were determined using the convex hull analysis and

branched structure analysis functions. Rendered traces were exported as monochrome vector

graphics. Cell body areas were separately determined in FIJI using the polygon tool.

*Cell suspension preparation*

Microglia and blood monocytes for single cell suspensions for flow cytometry were prepared as

follows. Mice were deeply anesthetized with isofluorane and transcardially perfused with ice-

cold HBSS. During perfusion, 2 ml of whole blood was collected from the right atrium into tubes

containing 40 µl of 10% (w/v) EDTA. The 2 ml of whole blood were added to 40 ml of red

blood cell lysis buffer (Abcam, ab204733) and incubated for 10 min at room temperature to lyse

red blood cells. The suspension containing lysed RBCs and white blood cells was spun down at

300 g for 5 min at 4 °C, resuspended in 10 ml ice-cold HBSS, and pelleted again at 300 g for 5

min at 4 °C. WBCs were resuspended in 500 µl ice-cold FACS buffer (HBSS, 0.5% BSA, 1 mM

EDTA) and subject to staining. In parallel with the WBC enrichment, brains were rapidly

dissected into 2 ml ice-cold HBSS and cerebella and brainstem were removed. Brains were

minced into small pieces and transferred to a dounce-homogenizer containing 5 ml ice-cold

HBSS with 20 µg/ml DNase I (Worthington, DPRF, LS006343). Tissue chunks were

homogenized with 15 loose and 15 more tight strokes and the homogenate was transferred to a

50 ml falcon tube through a pre-wet 70 µm strainer. The strainer was rinsed with HBSS to top

off the volume of each sample to 10 ml. Filtered homogenates were transferred to 15 ml falcon

tubes and spun at 300 g for 5 min at 4 °C. Supernatants were carefully removed, and pellets

resuspended in 10 ml of ice-cold 40% Percoll in HBSS. Samples were spun at 500 g for 30 min

at 4 °C with full acceleration and deceleration. Myelin and debris from the supernatant were

carefully removed and pellets resuspended in 10 ml ice-cold HBSS. Following another spin at

300 g for 5 min at 4 °C, the supernatant was removed and the microglial pellet resuspended in 1

ml FACS buffer (0.5% BSA HBSS, 1 mM EDTA).

*Staining for flow cytometry*

Suspensions of white blood cells and microglia were transferred to 2 ml Eppendorf

microcentrifuge tubes, and a small fraction of sample was removed for single-color controls. To

stain dead cells, live/dead violet (1:500 to 1:1000, Thermofisher, L34955) was added and

incubated for 5 min on ice. Tubes with live/dead-stained cells were topped off to 2 ml with

FACS buffer (0.5% BSA HBSS, 1 mM EDTA) and spun down at 300 g for 5 min at 4 °C. Pellets

were resuspended and incubated with 1:200 Mouse Fc Block (BD, 2.4G2, 553142) on ice for 15

min. Samples were incubated with 1:200 rat anti CD45-APC/Cy7 (Biolegend, 103115) and rat

anti Cd11b-PE/Cy5 (Biolegend, 101209) at 4 °C. Tubes were topped off to 2 ml with ice-cold

FACS buffer and microglia pelleted at 300 g for 5 min at 4 °C. Supernatants were removed and

microglia resuspended in 500 µl. Resuspended microglia were filtered through corning strainer

polystyrene tubes (Corning, 352235). Flow cytometry data was acquired on Aria II, Fortessa

HTS, and LSRII HTS, and analyzed using Flowjo.

*Statistical analysis*

Quantitative data from the qPCR, immunofluorescence, and microglia morphology

experiments were analyzed using GraphPad prism. Testing for normality was not possible due to

N=4-6 (four to six independent mice were used per group). Not assuming Gaussian distribution

of the data, a nonparametric test (Mann-Whitney) was used to compare median expression in

both groups.

*DNA sequcnes*

*pAAV-P2A-EGFP*

CCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCGGG

CGTCGGGCGACCTTTGGTCGCCCGGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGG

AGTGGCCAACTCCATCACTAGGGGTTCCTGCGGCCGCACGCGTTTAATTAAGTGTCT

AGACTGCAGAGGGCCCTGCGTATGAGTGCAAGTGGGTTTTAGGACCAGGATGAGGC

GGGGTGGGGGTGCCTACCTGACGACCGACCCCGACCCACTGGACAAGCACCCAACC

CCCATTCCCCAAATTGCGCATCCCCTATCAGAGAGGGGGAGGGGAAACAGGATGCG

GCGAGGCGCGTGCGCACTGCCAGCTTCAGCACCGCGGACAGTGCCTTCGCCCCCGC

CTGGCGGCGCGCGCCACCGCCGCCTCAGCACTGAAGGCGCGCTGACGTCACTCGCC

GGTCCCCGCAAACTCCCCTTCCCGGCCACCTTGGTCGCGTCCGCGCCGCCGCCGGC

CCAGCCGGACCGCACCACGCGAGGCGCGAGATAGGGGGGCACGGGCGCGACCATC

TGCGCTGCGGCGCCGGCGACTCAGCGCTGCCTCAGTCTGCGGTGGGCAGCGGAGGA

GTCGTGTCGTGCCTGAGAGCGCAGTCGAGAAACCGGCTAGAGGATCCTTCGAAACC

GGTGCTAGCAGCGCTGTTAACGGAAGCGGAGCCACTAACTTCTCCCTGTTGAAACAA

GCAGGGGATGTCGAAGAGAATCCCGGGCCAATGGTGAGCAAGGGCGAGGAGCTGTT

CACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGT

TCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAG

TTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGCCCACCCTCGTGACCACCTTC

GGCTACGGCCTGCAGTGCTTCGCCCGCTACCCCGACCACATGAAGCAGCACGACTTC

TTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGAC

GACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAA

CCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACA

AGCTGGAGTACAACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAG

AACGGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCA

GCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGC

CCGACAACCACTACCTGAGCTACCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAG

CGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGGGATCACTCTCGGCATG

GACGAGCTGTACAAGTAAGGCGCGCCCCTGCAGGGAATTCGATATCAAGCTTATCG

ATAATCAACCTCTGGATTACAAAATTTGTGAAAGATTGACTGGTATTCTTAACTATG

TTGCTCCTTTTACGCTATGTGGATACGCTGCTTTAATGCCTTTGTATCATGCTATTGCT

TCCCGTATGGCTTTCATTTTCTCCTCCTTGTATAAATCCTGGTTGCTGTCTCTTTATGA

GGAGTTGTGGCCCGTTGTCAGGCAACGTGGCGTGGTGTGCACTGTGTTTGCTGACGC

AACCCCCACTGGTTGGGGCATTGCCACCACCTGTCAGCTCCTTTCCGGGACTTTCGC

TTTCCCCCTCCCTATTGCCACGGCGGAACTCATCGCCGCCTGCCTTGCCCGCTGCTGG

ACAGGGGCTCGGCTGTTGGGCACTGACAATTCCGTGGTGTTGTCGGGGAAATCATCG

TCCTTTCCTTGGCTGCTCGCCTATGTTGCCACCTGGATTCTGCGCGGGACGTCCTTCT

GCTACGTCCCTTCGGCCCTCAATCCAGCGGACCTTCCTTCCCGCGGCCTGCTGCCGG

```
295   CTCTGCGGCCTCTTCCGCGTCTTCGCCTTCGCCCTCAGACGAGTCGGATCTCCCTTTG
296   GGCCGCCTCCCCGCATCGATACCGAGCGCTGCTCGAGAGATCTACGGGTGGCATCCC
297   TGTGACCCCTCCCCAGTGCCTCTCCTGGCCCTGGAAGTTGCCACTCCAGTGCCCACC
298   AGCCTTGTCCTAATAAAATTAAGTTGCATCATTTTGTCTGACTAGGTGTCCTTCTATA
299   ATATTATGGGGTGGAGGGGGGTGGTATGGAGCAAGGGGCAAGTTGGGAAGACAAC
300   CTGTAGGGCCTGCGGGGTCTATTGGGAACCAAGCTGGAGTGCAGTGGCACAATCTT
301   GGCTCACTGCAATCTCCGCCTCCTGGGTTCAAGCGATTCTCCTGCCTCAGCCTCCCG
302   AGTTGTTGGGATTCCAGGCATGCATGACCAGGCTCAGCTAATTTTTGTTTTTTGGTA
303   GAGACGGGGTTTCACCATATTGGCCAGGCTGGTCTCCAACTCCTAATCTCAGGTGAT
304   CTACCCACCTTGGCCTCCCAAATTGCTGGGATTACAGGCGTGAACCACTGCTCCCTT
305   CCCTGTCCTTCTGATTTTGTAGGTAACCACGTGCGGACCGAGCGGCCGCAGGAACCC
306   CTAGTGATGGAGTTGGCCACTCCCTCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGG
307   CGACCAAAGGTCGCCCGACGCCCGGGCTTTGCCCGGGCGGCCTCAGTGAGCGAGCG
308   AGCGCGCAGCTGCCTGCAGGGGCGCCTGATGCGGTATTTTCTCCTTACGCATCTGTG
309   CGGTATTTCACACCGCATACGTCAAAGCAACCATAGTACGCGCCCTGTAGCGGCGC
310   ATTAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCG
311   CCCTAGCGCCCGCTCCTTTCGCTTTCTTCCCTTCCTTTCTCGCCACGTTCGCCGGCTTT
312   CCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGG
313   CACCTCGACCCCAAAAAACTTGATTTGGGTGATGGTTCACGTAGTGGGCCATCGCCC
314   TGATAGACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCT
315   TGTTCCAAACTGGAACAACACTCAACCCTATCTCGGGCTATTCTTTTGATTTATAAGG
316   GATTTTGCCGATTTCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAA
317   CGCGAATTTTAACAAAATATTAACGTTTACAATTTTATGGTGCACTCTCAGTACAAT
318   CTGCTCTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAACACCCGCTGACGC
319   GCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTC
320   CGGGAGCTGCATGTGTCAGAGGTTTTCACCGTCATCACCGAAACGCGCGAGACGAA
321   AGGGCCTCGTGATACGCCTATTTTTATAGGTTAATGTCATGATAATAATGGTTTCTTA
322   GACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTC
323   TAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAA
324   TAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCC
325   TTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAA
```

AAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTACATCGAACTGGATCTCAAC

AGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACT

TTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGACGCCGGGCAAGAGCAA

CTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCACA

GAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAAC

CATGAGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGG

AGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAACTCGCCTTGATCGTTGGG

AACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTA

GCAATGGCAACAACGTTGCGCAAACTATTAACTGGCGAACTACTTACTCTAGCTTCC

CGGCAACAATTAATAGACTGGATGGAGGCGGATAAAGTTGCAGGACCACTTCTGCG

CTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGG

GTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGT

TATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTG

AGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCATATA

TACTTTAGATTGATTTAAAACTTCATTTTTAATTTAAAAGGATCTAGGTGAAGATCCT

TTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCA

GACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATC

TGCTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTTTGCCGGATCAA

GAGCTACCAACTCTTTTTCCGAAGGTAACTGGCTTCAGCAGAGCGCAGATACCAAAT

ACTGTCCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCG

CCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAG

TCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTCG

GGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTTGGAGCGAACGACCTACACCGA

ACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAA

AGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGA

GCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTCGGGTTTCGCCACCTCTG

ACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGC

CAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTGCTCACATGT

*LHA*

GCGGCCGCACGCGTTTAATTAAGTGTCTAATTGGACAGGGAGATGGCTCAGGTGTG

GAAAGCCAATATAGCGGTGCATGCTTGCAAGGCCAGGAGTAATGATGGGGACATAG

GAGGATTCTGGGAGCCTGGCCGGATAACCTATCCCCTAACACATAAGCACCAGGTC

CCACCCAGTAAGAGGCCCTGCCTCGAAGAAGAAGTGGAGGGTCCCTGAGGAGGAA

GGGCATTTGGGGTTATCCTCTGGTCTCCAAATGCATCCCTCCTCCCACACAAACACA

CAAAGACACTTTCACTTGTATCTGATGCCTGGTTAGCCATGAGCGTAAGCTAGGATT

GTGAATTCAGCCACCCTGGTACAGCAACTATTGAGTGCTTAGTGTATACTCTGGGTT

CGAAACACCGAGGACAGAAATGAATAAGATGCAGTTTCTGTCCTCAGGACCACCAG

AAGGAAACCAGGACGGGTGGCATCCTGTGTAGACAGACCCCGGAGTGTGTGCGTGT

TGCTTAGCTAAGTGATTTTCTGGAGTGTGTTCTGGCTCCCCGGCCGTCCTCCCGAACT

AGCAGACCACTCGTGCTGCCGGAGTCCTGGAGCGACAGGAGACTCAGCCTGACTTC

TCTCTTAACTCTCCCTCCGCGCAGGTTCCATAGCTCAACATGGTCCCCTGGTTCCTCC

TGTCTCTGCTGCTACTTGCGAGGCCTGTGCCTGGGGTGGCCTACTCTGTGTCACTCCC

GGCCTCCTTCCTGGAGGATGTAGCCGGCAGCGGGGAAGCTGAGGGTTCTTCAGCCTC

TTCCCCGAGCCTGCCGCCGCCTGGGACTCCAGCCTTCAGTCCCACACCGGAGAGACC

CCAGCCCACAGCTCTGGACGGCCCCGTGCCACCCACCAACCTCCTGGAAGGGATCA

TGGATTTCTTCCGGCAGTACGTGATGCTCATCGCGGTGGTGGGCTCGCTGACCTTCCT

CATCATGTTCATAGTCTGCGCCGCCCTCATCACGCGCCAGAAGCACAAGGCCACAGC

CTACTACCCATCCTCGTTCCCTGAAAAGAAGTATGTGGACCAGAGAGACCGGGCTG

GGGGACCCCGTACCTTCAGCGAGGTCCCTGACAGGGCACCTGACAGCCGGCATGAA

GAAGGCCTGGACACCTCCCATCAGCTCCAGGCTGACATTCTGGCTGCTACCCAGAAC

CTCCGGTCTCCAGCTAGAGCCCTGCCAGGCAATGGGGAGGGAGCAAAGCCTGTGAA

GGGTGGGTCGGAGGAGGAGGAGGAAGAGGTGCTCAGCGGTCAGGAGGAGGCCCAG

GAAGCCCCAGTATGTGGGGTCACTGAAGAGAAGCTGGGGGTCCCAGAGGAGTCGGT

CTCAGCAGAGGCTGAAGGGGTTCCTGCCACCAGTGAGGGCCAAGGGGAAGCAGAA

GGGTCTTTCTCCTTAGCCCAGGAATCCCAGGGAGCAACTGGTCCTCCTGAAAGTCCC

TGTGCCTGCAACAGAGTCTCTCCCAGTGTC

*P2A-CreERT2*

TGTGCCTGCAACAGAGTCTCTCCCAGTGTCGGAAGCGGAGCTACTAACTTCAGCCTG

CTGAAGCAGGCTGGAGACGTGGAGGAGAACCCTGGACCTATGGCCAATTTACTGAC

CGTACACCAAAATTTGCCTGCATTACCGGTCGATGCAACGAGTGATGAGGTTCGCAA

GAACCTGATGGACATGTTCAGGGATCGCCAGGCGTTTTCTGAGCATACCTGGAAAAT

GCTTCTGTCCGTTTGCCGGTCGTGGGCGGCATGGTGCAAGTTGAATAACCGGAAATG

GTTTCCCGCAGAACCTGAAGATGTTCGCGATTATCTTCTATATCTTCAGGCGCGCGG

TCTGGCAGTAAAAACTATCCAGCAACATTTGGGCCAGCTAAACATGCTTCATCGTCG

GTCCGGGCTGCCACGACCAAGTGACAGCAATGCTGTTTCACTGGTTATGCGGCGGAT

CCGAAAAGAAACGTTGATGCCGGTGAACGTGCAAAACAGGCTCTAGCGTTCGAAC

GCACTGATTTCGACCAGGTTCGTTCACTCATGGAAAATAGCGATCGCTGCCAGGATA

TACGTAATCTGGCATTTCTGGGGATTGCTTATAACACCCTGTTACGTATAGCCGAAA

TTGCCAGGATCAGGGTTAAAGATATCTCACGTACTGACGGTGGGAGAATGTTAATCC

ATATTGGCAGAACGAAACGCTGGTTAGCACCGCAGGTGTAGAGAAGGCACTTAGC

CTGGGGGTAACTAAACTGGTCGAGCGATGGATTTCCGTCTCTGGTGTAGCTGATGAT

CCGAATAACTACCTGTTTTGCCGGGTCAGAAAAAATGGTGTTGCCGCGCCATCTGCC

ACCAGCCAGCTATCAACTCGCGCCCTGGAAGGGATTTTTGAAGCAACTCATCGATTG

ATTTACGGCGCTAAGGATGACTCTGGTCAGAGATACCTGGCCTGGTCTGGACACAGT

GCCCGTGTCGGAGCCGCGCGAGATATGGCCCGCGCTGGAGTTTCAATACCGGAGAT

CATGCAAGCTGGTGGCTGGACCAATGTAAATATTGTCATGAACTATATCCGTAACCT

GGATAGTGAAACAGGGGCAATGGTGCGCCTGCTGGAAGATGGCGATCTCGAGCCAT

CTGCTGGAGACATGAGAGCTGCCAACCTTTGGCCAAGCCCGCTCATGATCAAACGCT

CTAAGAAGAACAGCCTGGCCTTGTCCCTGACGGCCGACCAGATGGTCAGTGCCTTGT

TGGATGCTGAGCCCCCCATACTCTATTCCGAGTATGATCCTACCAGACCCTTCAGTG

AAGCTTCGATGATGGGCTTACTGACCAACCTGGCAGACAGGGAGCTGGTTCACATG

ATCAACTGGGCGAAGAGGGTGCCAGGCTTTGTGGATTTGACCCTCCATGATCAGGTC

CACCTTCTAGAATGTGCCTGGCTAGAGATCCTGATGATTGGTCTCGTCTGGCGCTCC

ATGGAGCACCCAGTGAAGCTACTGTTTGCTCCTAACTTGCTCTTGGACAGGAACCAG

GGAAAATGTGTAGAGGGCATGGTGGAGATCTTCGACATGCTGCTGGCTACATCATCT

CGGTTCCGCATGATGAATCTGCAGGGAGAGGAGTTTGTGTGCCTCAAATCTATTATT

TTGCTTAATTCTGGAGTGTACACATTTCTGTCCAGCACCCTGAAGTCTCTGGAAGAG

AAGGACCATATCCACCGAGTCCTGGACAAGATCACAGACACTTTGATCCACCTGAT

GGCCAAGGCAGGCCTGACCCTGCAGCAGCAGCACCAGCGGCTGGCCCAGCTCCTCC

TCATCCTCTCCCACATCAGGCACATGAGTAACAAAGGCATGGAGCATCTGTACAGC

ATGAAGTGCAAGAACGTGGTGCCCCTCTATGACCTGCTGCTGGAGGCGGCGGACGC

CCACCGCCTACATGCGCCCACTAGCCGTGGAGGGGCATCCGTGGAGGAGACGGACC

AAAGCCACTTGGCCACTGCGGGCTCTACTTCATCGCATTCCTTGCAAAAGTATTACA

TCACGGGGGAGGCAGAGGGTTTCCCTGCCACAGCTTGACATGCCCCAGAACTGCTG

GGACCCGAATGT

*P2A-EGFP*

TGTGCCTGCAACAGAGTCTCTCCCAGTGTCGGAAGCGGAGCTACTAACTTCAGCCTG

CTGAAGCAGGCTGGAGACGTGGAGGAGAACCCTGGACCTATGGTGAGCAAGGGCG

AGGAGCTGTTCACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAAC

GGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCT

GACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGCCCACCCTCGT

GACCACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGC

AGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCT

TCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGAC

ACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACAT

CCTGGGGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTATATCATGGCCG

ACAAGCAGAAGAACGGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGAC

GGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCC

CGTGCTGCTGCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACC

CCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGGGATC

ACTCTCGGCATGGACGAGCTGTACAAGTAACATGCCCCAGAACTGCTGGGACCCGA

ATGT

*RHA*

CATGCCCCAGAACTGCTGGGACCCGAATGTTGGGTCCTTGAGGGTCACCTCTTTGGT

CAAGAAAGGCATTCAGCTCTAACTGCTCCTTGATACCACGTGGCTTGGCCATTGCTG

GTGCCAAGGCTGACCCCGAACTGGCAGAGCCGATGCCCTCTGGTGCACCCCAGGAA

ACATCTCCCCAAGTTCCAGCGCCCTTAATGACTCTTGCCACCCTGGGGGCTTCACCC

TAACGCACCACTTCTCTGGAAGGGGAAGGCCAGACACATGCCAGTTGGGGCTGCAT

GAGGCAGTCCTCAGAGCAGAAGGGGACCAGGCCAGAGGCCACCTGTGACGGGGCA

AACTGCATCTCGGCTGTGGAGACCAGAGGGGCTGTTAGATTTGGAAGACATCAATG

ACTGGGCCTGCGGCGCAGCCCGTGTCTGGTAATACCAGGGACGGCAGAGGCGTTTG

CATCTTCCCATCACCTGCAATGTCGCTGTCACTCTGCCCCTGTTCAGTGGACTTGATT

AGTTAGGAAACTTCTGGAAGGGGCCCCCTACTTTATATCACAGAGTTTGCCCTAGAC

ACCCCGTGGAAACACAAACTCAAAATCAAGTGGCTTTAGGAGCCGCTGTGCCCCTC

CACAAGCTGACATGGCTACTCTAAGGGTTCCTGCTGGGCTGGCTTTGCTACGCTTTC

CTCAAGCTGCTTTCTTATTACCAGGATGCCTCACAGCTACAAAGTCCAATCTCACAG

CACCCGCACTGGAAAATACTGTTCTCCATTTTTTTCCAAGGCACCCTGCTTTATGATT

GGCTCAGAGTTGGAATATGGGATGCAGGGCGTCTGGCTCTCTCAAGTGTCAAGCAA

CCCTGGCTTCCGCATGTGGGGCGAGGGGAGTGAGCAGAACTCTTCTCTGTCAGTCCA

CGAAGTAGACTAGAAGCCAACGGGTGCCTGGGGGCAACTGACGGTCTGGATTCTGA

CGCCCAGCCTGGAGCAGGGGCCTGGCCCATCCTATGCTCACACAGTGGTCTGGCAG

CCTGAGCCCACACAACTCCTCAGTCCTTGACAATGCTTAGGCTCTGTTCTGAGGGAC

TCCCCACACCTCCATTCAGGGCTCCCCGAGTCCTCAGCTCTTCGGATGTGGATATAT

GACACACTAGCATAATAAATCTTGATCTGGCTTGAGTCTCTCTGGATTATCTGCACA

GGTCTGCAGGAATGGGGTGTTCATGGGCTGGGGGCAGCATTGGGGGGAGGGCATCA

GCCAGGTCACAGAGCTGAGACTGTGTGTGGGTGCTCACAGGTGTTGGACCCACGTA

CCTGAGGAGTCCTGGAGGTGGTGTGTCAAACGTACAGACACTTTGCTGCCTGGCTCT

GCAGCTGGTGCAGCCATGCCACAGTCACTTCTTTGTGCTGTGGGACAAAGCTGGGGC

AAAGTCATGGCTGCCAATTCCCTACAGATGGGTGAGAACTGACTGAGCCTCAGGGC

GATATCAAGCTTATCGATAATCAACCTCT

**Results**

*Generation and genetic confirmation of Tmem119-EGFP and Tmem119-CreERT2 knock-in lines*

Access to highly specific fluorescent reporter lines for imaging and Cre-driver lines for

functional manipulation is critical in our quest for understanding the roles of microglia in health

and disease. Previously generated lines efficiently target microglia, but also other cells of the

monophagocytic system (Wieghofer and Prinz, 2016). Amongst a group of more recently

discovered microglia-specific genes, we chose targeting the Tmem119 gene to generate such

mouse lines (Bennett et al., 2016). Specifically, we opted for a transgenic approach that

harnesses the Tmem119 locus to drive expression, while preserving endogenous Tmem119

expression. We designed a CRISPR/Cas9 strategy to insert EGFP or CreERT2, preceded by

ribosome-skipping peptide porcine teschovirus-1 polyprotein (P2A), into the stop codon of

murine *Tmem119* (Figure 1A). Two founder lines were established for Tmem119-EGFP and 5

for Tmem119-CreERT2.

To determine whether the donor DNA was inserted correctly, a PCR-based approach was

used with primers spanning each junction. For both of the lines generated, specific 5' and 3'

junction PCR products were obtained, while no bands were observed with DNA from wildtype

animals (Figure 1B). This indicates specific insertion of the transgenes into the *Tmem119* locus.

To further examine scarless integration at the junctions, PCR products from the CreERT2 line

were subject to Sanger sequencing covering the junctions. Both 5' and 3' sequencing traces show

endogenous bases corresponding to the WT sequence in intron 1 and the 3' UTR (Figure 1C, D).

Additional sequencing confirmed absence of mutations across the entire targeted allele in both

lines. Together, these results confirm the generation of *Tmem119-EGFP* and *Tmem119-CreERT2*

lines.

While the function of endogenous *Tmem119* is currently unknown, both its temporally

distinct pattern and high expression level suggest functional importance of this gene (Bennett et

al., 2016). We thus designed a knock-in approach with the objective of leaving endogenous

expression intact. To test whether *Tmem119* expression is actually unaffected, we quantified

*Tmem119* transcript levels in the *Tmem119-EGFP* line by qPCR. Examination of *Tmem119*

expression in WT and *Tmem119-EGFP*[+/-] mice revealed that transcript levels are comparable

between genotypes (Figure 1E, WT median 0.9717, n=4, EGFP[+/-] median 0.9306, n=6, p=

0.5667[a]). We further examined TMEM119 protein expression using immunofluorescence and

found no difference between wildtype and *Tmem119-EGFP*[+/-] knock-in mice (Figure 1F-G, WT

median 0.9835, n=4, EGFP[+/-] median 0.117, n=4, p=0.20[b]) Together, these data support the

conclusion that *Tmem119* knock-in mice were generated with precision.

To further test whether the genetic engineering could affect the properties of microglia,

we examined basic morphological features of reconstructed immunostained microglia in

wildtype and *Tmem119-EGFP*[+/-] knock-in mice (Figure 1H-I). Using this approach, we

determined that the cell body area (Figure 1J, WT median 50.67 $\mu m^2$, EGFP[+/-] median 46.82

$\mu m^2$, n=12 microglia from 4 mice per genotype, p=0.3429[c]), number of processes (Figure 1K,

WT median 6.167, EGFP[+/-] median 5.667, n=12 microglia from 4 mice per genotype,

p=0.4571[d]), convex hull area (Figure 1L, WT median 2413 $\mu m^2$, EGFP[+/-] median 1871 $\mu m^2$,

n=12 microglia from 4 mice per genotype, p=0.4857[e]), and total process length (Figure 1M, WT

median 351.2 µm, EGFP[+/-] median 419.6 µm, n=12 microglia from 4 mice per genotype,

p=0.3428[f]) are comparable between microglia from control and knock-in mice. These data

suggest that the genetic modification does not affect basic microglial properties.

*EGFP is faithfully expressed in microglia*

Microglia populate the parenchyma in virtually all regions across the brain. To determine

the extent of EGFP expression in the newly generated Tmem119-EGFP line, we used confocal

microscopy on brain slices. Native EGFP fluorescence and antibody-enhanced signal were

visible across the brain in sagittal brain sections prepared from postnatal day 25 (P25) EGFP+/-

mice (Figure 2A). To further assess whether EGFP is expressed in microglia,

immunofluorescence labeling of the endogenous microglia proteins IBA1 and TMEM119 was

employed. High power confocal microscopy shows that EGFP-positive cells in the parenchyma

are IBA1 and TMEM119-positive (Figure 2B-E). Completion and fidelity of expression were

further assessed quantitatively in several brain regions. We found that 96.1 ± 2.0 % of the EGFP-

expressing cells in the somatosensory cortex are IBA-1 expressing microglia and 98.2 ± 3.3 % of

IBA1-positive cells in this region are labeled by EGFP (Table 1). In the striatum, 99.2 ± 2.1 % of

the cells labeled with EGFP were IBA1-expressing and 97.2 ± 1.8 % of IBA1-positive cells

displayed EGFP fluorescence. Labeling of microglia in the thalamus and hippocampus was 98.5

± 0.8 % and 98.9 ± 1.9 % complete and 96.9 ± 2.3 % and 98.5 ± 3.5 % specific, respectively. To

further confirm EGFP expression in microglia across the brain, we used the *Tmem119-EGFP*

line in flow cytometry (Figure 2F-N). We gated for single, live cells (Figure 2F-I) and detected

EGFP[+] cells among CD45[+] cells (Figure J-M). We determined that all CD11b[+]CD45lo microglia

(Figure 2K) expressed EGFP (Figure 2N). In addition, to more comprehensively assess

specificity of EGFP expression, we immunostained oligodendrocytes, neurons, and astrocytes.

We found no evidence for expression of EGFP in these major CNS celltypes (Figure 2O-Q).

Together, these data indicate highly complete and specific labeling of parenchymal microglia in

the *Tmem119-EGFP* reporter line.

All microglia express IBA1, but not all IBA1-expressing cells are parenchymal

macrophages. In fact, non-parenchymal macrophages such as meningeal, perivascular, and

choroid plexus macrophages express the myeloid- and macrophage marker IBA1 (Prinz and

Priller, 2014). To determine whether EGFP is expressed in these non-parenchymal macrophages,

we examined EGFP immunofluorescence in IBA1-expressing cells in the meninges and choroid

plexus, which can be readily discerned by their morphology and IBA1 immunoreactivity. High

power confocal images from both regions, including adjacent parenchyma, show that, while

parenchymal microglia express the EGFP label, neither IBA1-positive meningeal (Figure 3A, B)

nor choroid plexus macrophages express EGFP (Figure 3C, D).

In contrast to the clear anatomical location of choroid plexus and meningeal

macrophages, sites of perivascular macrophages are not easily discernible anatomically. To

examine whether perivascular macrophages are labeled in the *Tmem119-EGFP* line, we stained

for CD163, a well-characterized marker for perivascular macrophages (Kim et al., 2006). CD163

readily identified perivascular macrophages, which did not express EGFP (Figure 3E-F).

Adjacent microglia were clearly labeled with EGFP.

To further assess the specificity of EGFP expression in microglia beyond anatomical

location-based immunofluorescence analysis, we used flow cytometry (Figure 3G-I). Examining

$CD45^+EGFP^+$ cells, we determined that 99.6% are $CD11b^+CD45lo$ cells corresponding to

microglia (Figure 3H). We further detected that 153 of 2212 $CD11b^+CD45int/hi$ cells (6.9%)

corresponding to non-microglial cells express EGFP (Figure 3G-I). Together, these data indicate

the capacity of the *Tmem119-EGFP* line to distinguish parenchymal microglia from other brain

macrophages with good selectivity.

*EGFP is expressed in microglia of Tmem119-EGFP mice at early postnatal stages*

Microglia populate the brain during early embryonic development (Ginhoux et al., 2010).

*Tmem119* mRNA is expressed throughout microglial development, but antibody-based methods

currently consistently label TMEM119 only as of later postnatal stages, typically postnatal day

14 (P14) and later (Bennett et al., 2016). This makes it difficult to identify and discern microglia

from other monocytes at earlier time points. To determine whether the *Tmem119-EGFP* line

would specifically label microglia at early postnatal stages, we carried out immunofluorescence

microscopy at postnatal days 1, 3, and 6. EGFP expression in IBA1-positive cells was readily

observed in several regions examined, suggesting translation of *Tmem119-P2A-EGFP* transcript

at this stage (Figure 4A-L). Surprisingly, we also found EGFP expression associated with blood

vessels of P1 mice (Figure 4A-C), but not P3 or later, indicating that the transcript is also

processed in cells of the intracerebral microvascular compartment at this stage. Together, these

data indicate utility of the *Tmem119-EGFP* line for early developmental studies of microglia.

*Cre activity is specifically inducible in microglia of Tmem119-CreERT2 mice*

580           Cre activity in CreERT2 fusion protein is induced by the estrogen receptor antagonist

tamoxifen and can be assessed in the presence of a reporter construct. To assess whether

*Tmem119-CreERT2* mice display Cre activity in parenchymal microglia upon induction, we

crossed *Tmem119-CreERT2* mice to the Ai14 reporter line (Madisen et al., 2010), which

expresses tdTomato in the presence of Cre (Figure 5A). We administered tamoxifen daily for 3,

5, or 10 days to adult mice and 3 days to postnatal day 2 (P2) pups, and compared recombination

efficiency and fidelity for these different paradigms (Figure 5A). The mice were sacrificed 9-10

days after the last dose and subjected to immunofluorescence staining. High power images from

confocal microscopy showed tdTomato-labeling of IBA1-expressing microglia in mice that

received tamoxifen (Figure 5B-D) and absence of tdTomato expression without tamoxifen

(Figure 5E-G). Confocal micrographs also indicated absence of EGFP expression in other major

cell types of the CNS, such as oligodendrocytes, neurons, and astrocytes (Figure 5H-J).

Assessment of the completeness of labeling as the ratio of tdTomato$^+$IBA1$^+$- double positive

microglia (tdT$^+$IBA1$^+$) to all parenchymal microglia (IBA1$^+$) revealed very high efficiency of

recombination (Figure 5K). Similarly, we examined the fidelity of labeling which represents the

fraction of actual parenchymal microglia (tdT$^+$IBA1$^+$) amongst all tdTomato-labeled cells (tdT$^+$)

in different parenchymal regions, and found high specificity close to 100 percent (Figure 5L). Of

note, increasing the days of tamoxifen administered to mice appeared to increase the fraction of

microglia undergoing recombination. At the same time, prolonged administration of tamoxifen

correlated with slightly reduced fidelity (Figure 5K-L). We further sought to assess the

specificity of tdTomato expression using flow cytometry. Among a total of 1690

CD11b$^+$CD45int/hi cells in the CNS, 33 were positive for tdTomato, indicating minimal, yet

nonzero, expression in non-microglial myeloid cells.

The flow cytometry data revealed that recombination was largely, but not 100 percent specific. Furthermore, when examining the sections, we noticed some tdTomato fluorescence that did not appear to be associated with parenchymal microglia. To explore the nature of this ectopic fluorescence in this dosing paradigm, we imaged cells in the choroid plexus (Figure 5N-P) as well as cells in the cortex, which were immunostained for the endothelial cell marker CD31 (Figure 5Q-S). In the choroid plexus, there was very weak tdTomato fluorescence and most IBA1-positive macrophages did not express tdTomato (Figure 5N-P). In the cortex, high power confocal micrographs showed that the tdTomato and the CD31-immunolabeling signal were largely separate, with some tdTomato overlap with a larger CD31-positive blood vessel (Figure 5Q-S). Furthermore, tdTomato fluorescence was present on the pia in IBA1-negative cells (Figure 5T-V) and absent from CD163-expressing perivascular macrophages (Figure 5W-Z). Together, these data indicate that the *Tmem119-CreERT2* line can be used to conditionally control gene expression in adult and early postnatal microglia with good specificity.

*Characterization of transgene expression in monocytes from Tmem119 knock-in mice*

Monocytes infiltrate the brain and contribute to pathology in a number of brain disorders. Currently widely used transgenic microglia lines also target monocytes in addition to microglia and cannot readily distinguish the two cell types, thus hampering unequivocal attribution of microglia function in these disorders. To assess the utility of *Tmem119* knock-in mice for investigating microglia in models of CNS disorders, we isolated blood monocytes and examined transgene expression (Figure 6A). Using flow cytometry on white blood cells isolated from control or *Tmem119-EGFP*$^{+/-}$ mice, we did not detect any EGFP-expression (Figure 6B-C). To further determine whether *Tmem119-CreERT2*$^{+/-}$ mice display recombination in monocytes, we administered tamoxifen and sacrificed the mice for analysis 7 days after the last dose (Figure 6D). Using flow cytometry on white blood cells, we found that about 3 percent of CD45$^+$ cells expressed tdTomato (Figure 6E-F). This suggests that there is a low level of CreERT2 activity in *Tmem119-CreERT* mice. Together, these data indicate high applicability of *Tmem119-EGFP* mice and good applicability of *Tmem119-CreERT2* mice to the study of brain disorders with potential monocyte contribution.

**Statistical table**

|   | Data Structure | Type of test | 95% Confidence interval of difference between medians |
|---|---|---|---|
| a | N too small to test normality | Mann-Whitney test | -0.3393 to 0.2262 |
| b | N too small to test normality | Mann-Whitney test | -0.1601 to 0.3359 |
| c | N too small to test normality | Mann-Whitney test | -11.59 to 7.300 |
| d | N too small to test normality | Mann-Whitney test | -2.000 to 1.000 |
| e | N too small to test normality | Mann-Whitneåy test | -963.9 to 560.5 |
| f | N too small to test normality | Mann-Whitney test | -109.0 to 120.0 |

**Discussion**

In this study, we addressed the critical need for transgenic mouse lines that specifically

label or control microglia, but not other closely related cell types with overlapping signature

gene expression (Bennett et al., 2016; Wieghofer and Prinz, 2016; Haimon et al., 2018). We

targeted the recently identified microglia-specific *Tmem119* gene to knock in *EGFP* or

*CreERT2*. Multiple analyses confirm correct insertion of the transgenes, intact expression of

endogenous *Tmem119*, and normal basic morphology of microglia in the knock-in mice . Using

immunofluorescence and flow cytometry approaches, we demonstrate that EGFP is expressed in

microglia, but not other cells of the monophagocytic system defined anatomically and by

signature protein expression. Flow cytometry further shows that EGFP is almost exclusively

expressed in CD11b$^+$CD45lo cells corresponding to microglia. We further provide evidence for

early postnatal expression of EGFP and activity of CreERT2 in microglia upon tamoxifen

administration. In addition, we show that blood monocytes do not express EGFP and display

minimal activity of CreERT2. Together, our findings indicate proper generation of the two

mouse lines, which are now readily available through a public repository (JAX#031823 for

*Tmem119-EGFP* and JAX#031820 for *Tmem119-CreERT2*).

TMEM119 is a single transmembrane protein that is highly expressed in microglia in the

brains of humans and mice (Bennett et al., 2016; Satoh et al., 2016). In this study, we opted to

refrain from knocking *Tmem119* out through insertion of *EGFP* or *CreERT2* into the *Tmem119*

ORF and instead chose to preserve expression by using a polycistronic knock-in approach with

ribosomal skipping peptide P2A (Figure 1A). This was primarily motivated by the high

expression levels of *Tmem119*, conservation of expression in microglia across species, and

potential developmental regulation patterns, all of which suggest the importance of this gene.

Disruption of important genes in microglia could lead to confounding haploinsufficiency

phenotypes, as previously shown for other transgenic microglia lines (Lee et al., 2010; Rogers et

al., 2011). Our data rule out such potential confounding effects by showing intact *Tmem119*

mRNA and protein expression (Figure 1E-G), as well as normal basic morphology of microglia

in the knock-in mice (Figure 1H-M).

Examination of EGFP expression and co-expression analysis with TMEM119, IBA1, and

CD163 revealed that EGFP labels only parenchymal microglia (Figures 2 and 3). Identification

of these cells was possible by direct observation of EGFP fluorescence as well as by analysis of

immunostaining-enhanced tissue sections, supporting the conclusion that EGFP expression levels

are robust. Flow cytometry analysis further showed expression of EGFP in virtually all

CD11b[+]CD45lo microglia (Figure 2N) and a minor fraction of 6.9 percent (153/2212) of

CD11b[+]CD45int/hi cells (Figure 3G-I). This reflects that *Tmem119* expression is highly

selective, but not completely restricted to microglia, which has been observed in recent single-

cell sequencing studies (Li et al., 2019). An interesting observation was the expression of EGFP

at early postnatal time points in microglia (P1, P3, and P6), but also endothelial cells of blood

vessels at P1 (Figure 4). This is interesting for two reasons. First, it suggests that *Tmem119* is

initially expressed in microglia and endothelial cells and only becomes a specific microglia

marker during early postnatal development, presumably by selective up- and downregulation of

expression in microglia and endothelial cells, respectively. Second, translational processing of

*Tmem119* transcript appears to stand in opposition to a previous report showing that, while RNA

sequencing detects *Tmem119* mRNA throughout development, immunostaining only identifies

TMEM119 protein at later postnatal stages (Bennett et al., 2016). Although our data also do not

directly show TMEM119 protein expression at early stages, ribosomal processing of the

transcript is at least indirectly implied by the presence of EGFP protein, suggesting TMEM119

protein might be expressed but undetected due to expression level, protein stability, differential

posttranslational modification, or methodological limitations in immunostaining and flow

cytometry. Together, these data indicate that *Tmem119* knock-in mice may enable the study of

microglia and endothelial cells until postnatal day (P1) and microglia specifically after P3 and

later.

690           Our experiments using *Tmem119-CreERT2*$^{+/-}$ Ai14(RCL-tdT)$^{+/-}$ mice show tdTomato

expression in IBA1-expressing microglia, indicating that this line is suitable for Cre-dependent

manipulation of genes and transgene expression in microglia without affecting other major cell

types in the CNS (Figure 5). In this particular experiment, we observed very high, dose-

dependent recombination efficiencies close to 100 percent with a trend for slight reduction of the

generally high specificity with increasing tamoxifen dosage (Figure 5K, L). The recombination

efficiency observed across all doses compares well with other published models, such as the

*Cx3cr1* mouse line (Goldmann et al., 2013), and might even permit shorter dosing paradigms

dependent on the assay and relevant conditional allele. We further examined tissues other than

parenchymal microglia and observed that recombination was largely if not completely absent in

choroid plexus macrophages (5N-P). This is consistent with reported endogenous TMEM119

expression and other reports in the literature, suggesting other brain macrophages may express

microglia-specific genes at very low levels (Bennett et al., 2016; Goldmann et al., 2016; Mildner

et al., 2017), which might be sufficient to cause occasional all-or-none recombination events. In

addition, our data showed tdTomato expression in a few larger blood vessels and IBA1-negative

cells in the pia (Figure 5Q-V). While we did not determine the precise cellular origin of this

signal, our data from CD31-immunostaining exclude endothelial cells of the intracerebral

microvasculature (Figure 5Q-S). Together with the spatial extent of tdTomato expression and

recent single cell RNA sequencing data of vascular cells, our observations point to

leptomeningeal cells and their projections that penetrate deep into the brain ensheathing large

blood vessels and the choroid plexus stroma (Decimo et al., 2012, Vanlandewijck et al., 2018).

Together, these data render the *Tmem119-CreERT2* line a highly useful tool for the conditional

study of microglia, while also suggesting potential application for leptomeningeal cells.

Examination of EGFP and tdTomato expression in blood monocytes of *Tmem119-EGFP*

and *Tmem119-CreERT2; Ai14* mice revealed complete absence and minimal expression,

respectively (Figure 6). While, the lack of EGFP expression in blood monocytes of the

*Tmem119-EGFP* knock-in mice was expected based on previous observations, tdTomato

expression in 3% blood monocytes of the *Tmem119-CreERT2; Ai14* mice was rather surprising.

Similar to other brain macrophages discussed above, blood monocytes may express *Tmem119* at

very low, almost undetectable levels, which may nonetheless be sufficient to cause occasional

all-or-none Cre recombination events. Together, these data render the *Tmem119* knock-in lines

powerful tools to investigate microglia in diseases with monocyte contribution, while also

providing indication to allow time for monocyte turnover in the *Tmem-CreERT2* line after

tamoxifen dosing.

During the past decade, available microglia lines have been instrumental in advancing

our understanding of microglia. These lines remain powerful tools to address aspects of

microglia function. At the same time, it is widely acknowledged that ontogenetic differences, as

well as differences in localization and microenvironment between parenchymal microglia and

their closely related cells from the monophagocytic system warrant more discerning observation

and manipulation (Bennett et al., 2016; Wieghofer and Prinz, 2016; Haimon et al., 2018).

Recently developed antibodies for TMEM119 immunostaining are one tool that has been

adopted quickly to identify parenchymal microglia. As TMEM119 is a transmembrane protein

localizing to processes, immunostaining for this protein is particularly powerful to study

morphology of microglia. To identify discrete microglia, however, relatively diffuse staining

patterns of membrane-bound TMEM119 present a challenge (Figure 2B). In this case, the more

discretely identifiable cytosolic distribution of EGFP in the *Tmem119-EGFP* line might be of

great value. In addition, the robustness of the EGFP signal in the *Tmem119-EGFP* line may

enable cell sorting without the need for antibodies, possibly even at early developmental stages

before the onset of TMEM119 appearance on the membrane, as well as *in vivo* imaging

approaches. Furthermore, the absence of EGFP fluorescence in blood monocytes renders this line

uniquely useful for the study of microglia in brain disorders that involve monocyte infiltration.

*Tmem119-CreERT2* line will significantly facilitate conditional control of gene expression

selectively in microglia. Compared to constitutive Cre lines, which cumulatively recombine

throughout development and thus deny temporal control, the *CreERT2* line will allow precise

control of gene expression to examine the role of microglia in temporally restricted

developmental processes. In this capacity, dose-dependency of CREERT2 and potentially

resulting mosaic expression of recombined alleles using low doses may allow the study of

candidate genes in adjacent microglia populations sharing the same microenvironment. Finally,

as recently shown in an important study by *Jung and colleagues* using RiboTag, microglia gene

expression profiles strongly depend on the isolation method and defining marker genes or Cre

lines used for molecular identification or enrichment (Haimon et al., 2018). The *Tmem119-*

*CreERT2* line provided here, possibly in conjunction with approaches such as RiboTag and

removal of the pia, might enable the generation of the purest microglia gene expression profiles

to date.

**Legends**

**Figure 1.**

**Generation and validation of *Tmem119-EGFP and Tmem119-CreERT2* knock-in lines. A**,
Experimental strategy using a guide RNA (blue bar) to introduce a double-strand break at the
*Tmem119* stop codon in mouse zygotes and an injection mixture containing *EGFP* ssDNA
template with short homology arms of 55-300 bp or *CreERT2* dsDNA template with long 1.5 kb
homology arms. Each knock-in is designed to cause in-frame insertion of P2A peptide followed
by *EGFP* or *CreERT2* and a stop codon. Not drawn to scale. **B**, Confirmation of insertion by
amplification of fragments spanning the 5' and 3' junctions using sets of primers binding inside
the template and outside the homology arms  (open and closed arrows in A with arrowheads
indicating corresponding products). **C and D**, Sanger sequencing across the 5' and 3' junction of
the CreERT2 line and comparison to wildtype (WT) sequence indicate seamless insertion in the
transgenic mice (Tg). **E**, qPCR analysis of gene expression for *Tmem119* in wildtype controls
and *Tmem119-EGFP$^{+/-}$* knock-in animals. N=4 WT, N=6 *Tmem119-EGFP$^{+/-}$* mice. **F,**
Representative TMEM119 immunoreactivity in wildtype controls and *Tmem119-EGFP$^{+/-}$* knock-
in animals. **G,** Relative intensity of TMEM119 immunostaining signal. N=WT, N=4 *Tmem119-
EGFP$^{+/-}$* mice. **H and I,** Representative IBA-immunostained microglia and corresponding
neurolucida traces in WT and *Tmem119-EGFP$^{+/-}$* knock-in mice. **J-M,** Quantified cell body area,
number of processes, convex hull area, and total process length of microglia in WT and
*Tmem119-EGFP$^{+/-}$* knock-in mice. N= 12 microglia from 4 mice per genotype. ssDNA, single
strand DNA; dsDNA, double strand DNA; TAA (red bar), stop codon; Kb, kilo bases.

**Figure 2**

**EGFP labels microglia in all regions across the brain**. **A**, Representative epifluorescence
image of a sagittal section of a P25 *Tmem119-EGFP$^{+/-}$* mouse stained with an anti-GFP antibody
for improved signal-to-noise ratio (data for one of three mice shown). **B-E**, High-power confocal
micrographs of native EGFP fluorescence (B, green) in TMEM119 (C, red) and IBA1 (D, white)
microglia in the cortical region outlined in A. FIJI-calculated composite image showing all three
labels (E). Boxes 3A and 3B cross-reference areas shown in figure 3.**F-J,** Flow cytometry
analysis showing gating for of single, live, CD45-positive cells (one of three independent

experiments shown). Numbers in or adjacent to outlined areas indicate percent cells in each gate.

**K and L,** Representative density blots showing EGFP expression in WT and *Tmem119-EGFP*$^{+/-}$

mice (pre-gated on CD45$^+$ cells). **M,** Representative density plot showing CD11b$^+$CD45lo cells

corresponding to microglia (pre-gated on CD45$^+$ cells). **N,** Histogram showing fraction of

CD11b$^+$CD45lo microglia expressing EGFP. **O-R,** Representative immunostaining for

oligodendrocytes (P, Olig2, red), neurons (P, NeuN, red), and astrocytes (Q, GFAP, red) in

EGFP-expressing *Tmem119-EGFP*$^{+/-}$ mice. One of three independent experiments shown.

**Figure 3**

**EGFP discerns parenchymal microglia from other brain macrophages. A**, IBA1 labels

meningeal macrophages and parenchymal microglia from the region indicate with a dashed box

in figure 2A. **B**, EGFP labels only parenchymal macrophages (open arrowheads), but not

meningeal IBA1-positive macrophages (closed arrowheads). Diffuse fluorescence is associated

with pia. ECIC, external cortex of inferior colliculus; Cer, cerebellum. Scale bar 100 µm. **C and**

**D**, Choroid plexus IBA1$^+$ macrophages (closed arrowheads) from the region boxed in figure 2A

are not labeled with EGFP, while adjacent parenchymal microglia are (open arrowheads). Hpc,

hippocampus. Scale bar 100 µm. **E and F**, Perivascular CD163-expressing macrophages (E, red,

closed arrowheads) are not labeled with EGFP (F, green, open arrowheads). Scale bar 100 µm.

**G-I**, Flow cytometry analysis of CD11b and CD45 expression in pre-gated single live CD45$^+$

cells and EGFP-expressing CD45$^+$ cells of *Tmem119-EGFP*$^{+/-}$ mice shown as density plots (G,

CD45$^+$ and H CD45$^+$ EGFP$^+$) and overlay (I, black CD45$^+$, green CD45$^+$ EGFP$^+$). One of three

independent experiments (N ≥ 3 mice per group) shown. Numbers in or adjacent to outlined

areas indicate percent cells in each gate.

**Figure 4**

*Tmem119-EGFP* **mice label microglia at early postnatal stages**. Sagittal slices from postnatal

day 1, 3 and 6 (P1, P3, P6) mice were stained against IBA1. **A-C**, High power confocal images

of the somatosensory cortex show EGFP expression in IBA-positive cells and blood vessels at P1

(solid arrowheads in magnified panel). **D-F**, Confocal micrographs around the forceps major of

corpus callosum (CC) and hippocampus (Hpc) show EGFP-labeled, IBA1-positive microglia at

P3. **G-I**, Confocal micrographs of the striatum (CPu) and corpus callosum (CC) show EGFP

labeling of IBA-positive cells at P3. **J-L**, Confocal micrographs of the P6 cortex reveal EGFP-

expression in IBA-immunostained cells. Error bars 100 µm.

**Figure 5**

*Tmem119-CreERT2* **mice effectively recombine a conditional allele in microglia. A**,

*Tmem119-CreERT2* mice were crossed to *Ai14(RCL-tdT)-D* mice and tamoxifen (TAM) was

administered to induce tdTomato expression. Different sets of adult mice received tamoxifen per

os (p.o) for 3, 5 or 10 days and a set of neonatal mice received TAM for 3 days intraperitoneally

(i.p). The mice were sacrificed for flowcytometry (FC) 7 days after the 3-day dosing paradigm or

for immunofluorescence (IF) 9 or more days after dosing. **B-D**, Representative confocal

micrograph showing tdTomato expression in IBA-positive microglia upon tamoxifen

administration. **E-G**, Absence of tdTomato expression in untreated animals. **H-J**, Representative

immunostaining for oligodendrocytes (H, Olig2, green), neurons (I, NeuN, green), and astrocytes

(J, GFAP, green) in tdTomato-expressing microglia of *Tmem119-CreERT2$^{+/-}$; Ai14(RCL-tdT)-*

*D$^{+/-}$* mice. One of three independent experiments shown. **K and L,** Percentage completion **(K)**

and fidelity **(L)** of the labeling in different brain regions for the different dosing schemes. Hpc =

Hippocampus. CPu = Caudate-Putamen. Ctx = Cortex. N=2 mice for 3 days, 3 mice each for 5

and 10 days of TAM in adults, and 3 mice for 3 days administration at P2 = postnatal day 2.

Floating bars = min to max, line at mean. **M,** Flow cytometry analysis of CD11b and CD45

expression in pre-gated single live CD45$^{+}$ cells and tdTomato-expressing CD45$^{+}$ cells of

*Tmem119-CreERT2$^{+/-}$; Ai14(RCL-tdT)-D$^{+/-}$* mice shown as overlay (I, black CD45$^{+}$, red CD45$^{+}$

tdTomato$^{+}$). One of two representative experiments (N=3 mice). **N-O**, Confocal micrograph of

the choroid plexus and adjacent parenchyma. **Q-S**, Confocal micrograph of CD31-

immunostained blood vessels (green) shows tdTomato$^{+}$ microglia (arrowheads) and tdTomato

expression in a large blood vessel (open arrowhead). **T-V**, Confocal micrograph showing

tdTomato expression in cortical microglia and cells of the pia (open arrowheads). **W-Z**, Confocal

micrograph showing CD163- (white) and IBA1-expressing (green) macrophages. The open

arrowhead indicates a tdTomato$^{-}$CD163$^{+}$IBA1$^{+}$ perivascular macrophage. Closed arrowheads

parenchymal microglia. One of three independent experiments shown (N=3 mice).

**Figure 6**

**Blood monocyte profiling in Tmem119 knock-in mice**

**A,** Experimental approach for flow cytometry of monocytes isolated from whole blood. **B and**

**C,** Representative flow cytometry analysis of EGFP expression in pre-gated single live CD45$^+$

cells in wildtype control (B) and *Tmem119-EGFP$^{+/-}$* mice (C). One of three experiments is

shown (N=3 mice). Numbers in or adjacent to outlined areas indicate percent cells in each gate.

**D,** Adult mice of *Tmem119-CreERT2$^{+/-}$; Ai14(RCL-tdT)-D$^{+/-}$* mice received tamoxifen (TAM)

per os (p.o) for 3 days. The mice were sacrificed 7 days after the last dose for flow cytometry

(FC). **E and F,** Representative flow cytometry analysis of tdTomato expression in pre-gated

single live CD45$^+$ cells in *Ai14(RCL-tdT)-D$^{+/-}$* (E) and *Tmem119-CreERT2$^{+/-}$; Ai14(RCL-tdT)-*

*D$^{+/-}$* mice (F). One of two independent experiments is shown (N=3 mice).

**Tables**

**Table 1: Quantification of EGFP labeled microglia in Tmem119-EGFP mice.**

| Brain Region | Completeness (EGFP$^+$Iba$^+$/Iba$^+$) | Fidelity (EGFP$^+$Iba$^+$/EGFP$^+$) |
|---|---|---|
| Somatosensory cortex | 98.2 ± 3.3 % | 96.1 ± 2.0 % |
| Striatum | 97.2 ± 1.8 % | 99.2 ± 2.1 % |
| Thalamus | 96.9 ± 2.3 % | 98.5 ± 0.8 % |
| Hippocampus | 98.5 ± 3.5 % | 98.9 ± 1.9 % |

**References**

Aguzzi A, Zhu C (2017) Microglia in prion diseases. The Journal of clinical investigation
127:3230-3239.

Bennett ML, Bennett FC, Liddelow SA, Ajami B, Zamanian JL, Fernhoff NB, Mulinyawe SB,
Bohlen CJ, Adil A, Tucker A, Weissman IL, Chang EF, Li G, Grant GA, Hayden Gephart MG,
Barres BA (2016) New tools for studying microglia in the mouse and human CNS.
Proceedings of the National Academy of Sciences.

Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, Koeglsperger T, Dake B,
Wu PM, Doykan CE, Fanek Z, Liu L, Chen Z, Rothstein JD, Ransohoff RM, Gygi SP, Antel
JP, Weiner HL (2013) Identification of a unique TGF-β–dependent molecular and
functional signature in microglia. Nature Neuroscience 17:131.

Buttgereit A, Lelios I, Yu X, Vrohlings M, Krakoski NR, Gautier EL, Nishinakamura R, Becher B,
Greter M (2016) Sall1 is a transcriptional regulator defining microglia identity and
function. Nature immunology 17:1397-1406.

Clausen B, Burkhardt C, Reith W, Renkawitz R, Förster I (1999) Conditional gene targeting in
macrophages and granulocytes using LysMcre mice. Transgenic research 8:265-277.

Davalos D, Grutzendler J, Yang G, Kim JV, Zuo Y, Jung S, Littman DR, Dustin ML, Gan W-B (2005)
ATP mediates rapid microglial response to local brain injury in vivo. Nature neuroscience
8:752.

Decimo I, Fumagalli G, Berton V, Krampera M, Bifari F (2012) Meninges: from protective
membrane to stem cell niche. American Journal of Stem Cells 1:92-105.

Ferron M, Vacher J (2005) Targeted expression of Cre recombinase in macrophages and
osteoclasts in transgenic mice. Genesis 41:138-145.

Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, Mehler MF, Conway SJ, Ng LG,

Stanley ER (2010) Fate mapping analysis reveals that adult microglia derive from

primitive macrophages. Science (New York, NY):1194637.

Goldmann T, Wieghofer P, Müller PF, Wolf Y, Varol D, Yona S, Brendecke SM, Kierdorf K,

Staszewski O, Datta M, Luedde T, Heikenwalder M, Jung S, Prinz M (2013) A new type of

microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation.

Nature Neuroscience 16:1618.

Goldmann T et al. (2016) Origin, fate and dynamics of macrophages at central nervous system

interfaces. Nature immunology 17:797.

Haimon Z, Volaski A, Orthgiess J, Boura-Halfon S, Varol D, Shemer A, Yona S, Zuckerman B,

David E, Chappell-Maor L, Bechmann I, Gericke M, Ulitsky I, Jung S (2018) Re-evaluating

microglia expression profiles using RiboTag and cell isolation strategies. Nature

immunology 19:636-644.

Hirasawa T, Ohsawa K, Imai Y, Ondo Y, Akazawa C, Uchino S, Kohsaka S (2005) Visualization of

microglia in living tissues using Iba1-EGFP transgenic mice. Journal of Neuroscience

Research 81:357-362.

Itagaki S, McGeer P, Akiyama H, Zhu S, Selkoe D (1989) Relationship of microglia and astrocytes

to amyloid deposits of Alzheimer disease. Journal of neuroimmunology 24:173-182.

Keren-Shaul H, Spinrad A, Weiner A, Matcovitch-Natan O, Dvir-Szternfeld R, Ulland TK, David E,

Baruch K, Lara-Astaiso D, Toth B, Itzkovitz S, Colonna M, Schwartz M, Amit I (2017) A

Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease.

Cell 169:1276-1290.e1217.

Kim W-K, Alvarez X, Fisher J, Bronfin B, Westmoreland S, McLaurin J, Williams K (2006) CD163

Identifies Perivascular Macrophages in Normal and Viral Encephalitic Brains and

Potential Precursors to Perivascular Macrophages in Blood. The American Journal of

Pathology 168:822-834.

Lawson LJ, Perry VH, Dri P, Gordon S (1990) Heterogeneity in the distribution and morphology
of microglia in the normal adult mouse brain. Neuroscience 39:151-170.

Lee S, Varvel NH, Konerth ME, Xu G, Cardona AE, Ransohoff RM, Lamb BT (2010) CX3CR1
Deficiency Alters Microglial Activation and Reduces Beta-Amyloid Deposition in Two
Alzheimer's Disease Mouse Models. The American Journal of Pathology 177:2549-2562.

Li Q, Cheng Z, Zhou L, Darmanis S, Neff NF, Okamoto J, Gulati G, Bennett ML, Sun LO, Clarke LE et al.
2019. Developmental Heterogeneity of Microglia and Brain Myeloid Cells Revealed by Deep
Single-Cell RNA Sequencing. Neuron 101: 207-223.e210.

Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD, Hawrylycz
MJ, Jones AR, Lein ES, Zeng H (2010) A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. Nat Neurosci 13:133-140.

Mathys H, Adaikkan C, Gao F, Young JZ, Manet E, Hemberg M, De Jager PL, Ransohoff RM,
Regev A, Tsai LH (2017) Temporal Tracking of Microglia Activation in Neurodegeneration
at Single-Cell Resolution. Cell reports 21:366-380.

Mildner A, Huang H, Radke J, Stenzel W, Priller J (2017) P2Y12 receptor is expressed on human
microglia under physiological conditions throughout development and is sensitive to
neuroinflammatory diseases. Glia 65:375-387.

O'Loughlin E, Madore C, Lassmann H, Butovsky O (2018) Microglial Phenotypes and Functions in
Multiple Sclerosis. Cold Spring Harbor Perspectives in Medicine 8.

Paolicelli RC, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, Giustetto M, Ferreira TA,
Guiducci E, Dumas L, Ragozzino D, Gross CT (2011) Synaptic pruning by microglia is
necessary for normal brain development. Science (New York, NY) 333:1456-1458.

Parkhurst CN, Yang G, Ninan I, Savas JN, Yates JR, Lafaille JJ, Hempstead BL, Littman DR, Gan W-
B (2013) Microglia promote learning-dependent synapse formation through brain-
derived neurotrophic factor. Cell 155:1596-1609.

Prinz M, Priller J (2014) Microglia and brain macrophages in the molecular age: from origin to
neuropsychiatric disease. Nature Reviews Neuroscience 15:300.

Ransohoff RM (2016) How neuroinflammation contributes to neurodegeneration. Science (New
York, NY) 353:777-783.

Rogers JT, Morganti JM, Bachstetter AD, Hudson CE, Peters MM, Grimmig BA, Weeber EJ,
Bickford PC, Gemma C (2011) CX3CR1 Deficiency Leads to Impairment of Hippocampal
Cognitive Function and Synaptic Plasticity. The Journal of Neuroscience 31:16241-16250.

Samokhvalov IM, Samokhvalova NI, Nishikawa S-i (2007) Cell tracing shows the contribution of
the yolk sac to adult haematopoiesis. Nature 446:1056.

Sasmono RT, Oceandy D, Pollard JW, Tong W, Pavli P, Wainwright BJ, Ostrowski MC, Himes SR,
Hume DA (2003) A macrophage colony-stimulating factor receptor–green fluorescent
protein transgene is expressed throughout the mononuclear phagocyte system of the
mouse. Blood 101:1155-1163.

Satoh J-i, Kino Y, Asahina N, Takitani M, Miyoshi J, Ishida T, Saito Y (2016) TMEM119 marks a
subset of microglia in the human brain. Neuropathology 36:39-49.

Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, Ransohoff RM,
Greenberg ME, Barres BA, Stevens B (2012) Microglia sculpt postnatal neural circuits in
an activity and complement-dependent manner. Neuron 74:691-705.

Sierra A, Encinas JM, Deudero JJ, Chancey JH, Enikolopov G, Overstreet-Wadiche LS, Tsirka SE,
Maletic-Savatic M (2010) Microglia shape adult hippocampal neurogenesis through
apoptosis-coupled phagocytosis. Cell stem cell 7:483-495.

Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N, Micheva KD, Mehalow
AK, Huberman AD, Stafford B, Sher A, Litke Alan M, Lambris JD, Smith SJ, John SWM,
Barres BA (2007) The Classical Complement Cascade Mediates CNS Synapse Elimination.
Cell 131:1164-1178.

Ueno M, Fujita Y, Tanaka T, Nakamura Y, Kikuta J, Ishii M, Yamashita T (2013) Layer V cortical
neurons require microglial support for survival during postnatal development. Nat
Neurosci 16:543-551.

Vanlandewijck M, He L, Mäe MA, Andrae J, Ando K, Del Gaudio F, Nahar K, Lebouvier T, Laviña B,
Gouveia L et al. 2018. A molecular atlas of cell types and zonation in the brain vasculature.
Nature 554: 475.

Weinhard L, di Bartolomei G, Bolasco G, Machado P, Schieber NL, Neniskyte U, Exiga M,
Vadisiute A, Raggioli A, Schertel A, Schwab Y, Gross CT (2018) Microglia remodel
synapses by presynaptic trogocytosis and spine head filopodia induction. Nature
Communications 9:1228.

Wieghofer P, Prinz M (2016) Genetic manipulation of microglia during brain development and
disease. Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease 1862:299-309.

Yona S, Kim K-W, Wolf Y, Mildner A, Varol D, Breker M, Strauss-Ayali D, Viukov S, Guilliams M,
Misharin A, Hume David A, Perlman H, Malissen B, Zelzer E, Jung S (2013) Fate Mapping
Reveals Origins and Dynamics of Monocytes and Tissue Macrophages under
Homeostasis. Immunity 38:79-91.

# Figure 1

# Figure 2

# Figure 3

# Figure 4

# Figure 5

# Figure 6