MHC genotyping from rhesus macaque exome sequences

John R. Caskey[1], Roger W. Wiseman[1,2], Julie A. Karl[2], David A. Baker[1], Taylor Lee[1], Muthuswamy

Raveendran[3], R. Alan Harris[3], Jianhong Hu[3], Donna M. Muzny[3],Jeffrey Rogers[3], David H. O'Connor[1,2]

[1]Wisconsin National Primate Research Center, University of Wisconsin-Madison, Madison, WI 53715,

USA

[2]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison, Madison, WI

53705, USA

[3]Human Genome Sequencing Center, Baylor College of Medicine, Houston, TX 77030, USA

Corresponding author: Dr. David H. O'Connor (doconnor@primate.wisc.edu; +1 608 890 0845)

Author	ORCID
Caskey	0000-0001-5665-524X
Wiseman	0000-0002-7682-7085
Karl	0000-0002-4447-4721
Baker	0000-0002-6811-8238
Lee	0000-0002-8768-4358
Raveendran	0000-0001-6185-4059
Jianghong	0000-0002-6318-8595
Harris	0000-0002-7333-4752
Rogers	0000-0002-7374-6490
O'Connor	0000-0003-2139-470X

Keywords: Macaca mulatta, exome, major histocompatibility complex

Abstract

Indian rhesus macaque major histocompatibility complex (MHC) variation can influence the outcomes of transplantation and infectious disease studies. Frequently, rhesus macaques are MHC genotyped to identify variants that could account for unexpected results. Since the MHC is only one region in the genome where variation could impact experimental outcomes, strategies for simultaneously profiling variation in the macaque MHC and the remainder of the protein coding genome would be useful. Here we introduce macaque exome sequence (MES) genotyping, in which MHC class I and class II genotypes are determined with high confidence using target-enrichment probes that are enriched for MHC sequences. For a cohort of 27 Indian rhesus macaques, we describe two methods for obtaining MHC genotypes from MES data and demonstrate that the MHC class I and class II genotyping results obtained with these methods are 98.1% and 98.7% concordant, respectively, with expected MHC genotypes. In contrast, conventional MHC genotyping results obtained by deep sequencing of short multiplex PCR amplicons were only 92.6% concordant with expectations for this cohort.

Introduction

The major histocompatibility complex (MHC) is an intensively studied set of genes in macaques (Shiina et al. 2017; Wiseman et al. 2013). The genomic MHC region contains clusters of genes that encode the MHC class I complex and the MHC class II complex. Cells use MHC class I molecules to present intracellular peptides to immune cells like the CD8+ T cell or natural killer cells (Garcia and Adams 2005). MHC class I molecules accommodate intracellular peptides of varying specificity by having diverse amino acid sequences in the α1 and α2 subunits, which form the peptide-binding cleft (Silver and Watkins 2017; Loffredo et al. 2009). These α 1 and α 2 subunits correspond to exons 2 and 3, respectively, of an MHC class I gene (Malissen et al. 1982). Most of the polymorphisms that distinguish MHC class I alleles are concentrated in exons 2 and 3 (Williams 2001). Thousands of individual MHC allelic variants have been identified in the three most widely used macaque species for biomedical research: rhesus (*Macaca mulatta*), cynomolgus (*Macaca fascicularis)*, and pig-tailed macaques (*Macaca nemestrina*) (Semler et al. 2018; Karl et al. 2017; Maccari et al. 2017).

In humans, the MHC is also termed the human leukocyte antigen complex (HLA). HLA class I has a single copy of the *HLA-A*, *HLA-B*, and *HLA-C* genes on each chromosome (Daza-Vamenta et al. 2004; Shiina et al. 2017). In contrast, macaques have a variable number of genes on each chromosome that encode MHC class I MHC-A and MHC-B proteins, and macaques lack an *HLA-C* orthologue (Daza-Vamenta et al. 2004; Shiina et al. 2017; Wiseman et al. 2013). Initial approaches to genotype macaque MHC class I relied on using sequence-specific PCR oligonucleotides to test for the presence or absence of individual alleles (Kaizu et al. 2007). More recently, deep sequencing genomic DNA or complementary DNA PCR amplicons spanning a highly variable region of exon 2 has become commonplace (Wiseman et al. 2009; Karl et al. 2013). Amplicon sequences can be used to genotype groups of closely related MHC class I alleles, which are denoted as lineages. For example, an amplicon deep sequence that corresponds to the rhesus macaque *Mamu-A1*001* lineage demonstrates that an animal possesses *Mamu-A1*001:01*, *Mamu-A1*001:02*, or another closely related variant that has not yet been identified. This lineage-level reporting of MHC class I genotypes can be sufficient for designing experiments where certain MHC class I genotypes need to be matched between animals, balanced among experimental groups, or excluded entirely from a study (Loffredo et al.

2007; Loffredo et al. 2009; Muhl et al. 2002; Karl et al. 2013; Wiseman et al. 2009; Nomura et al. 2012; Mothe et al.

2003).

Amplicon deep sequencing can also be used for MHC class II genotyping. The human HLA class II genes *DQA1*,

DQB1, *DPA1*, and *DPB1* have direct orthologues in macaques (Otting et al. 2017). Both macaques and humans have

a variable number of MHC class II DRB genes on a single chromosome, while the DRA gene is oligomorphic

(Daza-Vamenta et al. 2004; Shiina et al. 2017), and typically is not used for genotyping purposes. MHC class II

molecules are members of the immunoglobulin superfamily, but they differ from MHC class I in several ways. The

MHC class II complex is comprised of an α and β subunit heterodimer, and separate genes encode each MHC class

II α and β subunit (Brown et al. 1993). Highly polymorphic regions that are diagnostic for MHC class II allele

lineages can be PCR amplified in a manner that is similar to the MHC class I genotyping. The most extensive

polymorphism among alleles of the MHC class II genes is found in exon 2 (Williams 2001). Each of the *DRB*,

DQA1, *DQB1*, *DPA1* and *DPB1* polymorphic MHC class II genes are sufficiently divergent that separate PCR

amplicons are required for deep sequencing (Karl et al. 2014).

Comprehensive MHC class I and class II genotyping of macaques by amplicon deep sequencing requires preparation

of six separate PCR amplicons for MHC class I and class II DRB, DQA1, DQB1, DPA1, and DPB1 (Karl et al.

2014; Karl et al. 2017). Despite this complexity, a major advantage to amplicon deep sequencing has been its cost

effectiveness. The output from a single MiSeq sequencing run can be used to determine the MHC class I and MHC

class II genotypes for up to 192 macaque samples. In recent years, improved sequencing hardware and software

have prompted new approaches to MHC genotyping. Instead of a utilizing PCR amplicons of variable gene regions

to genotype samples, researchers can use human whole exome sequencing (WES) and whole genome sequencing

(WGS) datasets to determine HLA genotypes (Xie et al. 2017; Kishikawa et al. 2019; Yang et a.l 2014; Posey et al.

2016). Likewise, target capture approaches have been described for HLA genotyping with next-generation

sequencing (Cao et al. 2013; Wittig et al. 2015). In contrast, MHC genotyping of macaques from WGS (Xue et al.

2016; Bimber et al. 2017; de Manuel et al. 2018) or whole exome sequence (Vallender 2011; Cornish et al. 2016)

datasets have not been reported to date. This reflects the challenges presented by mapping short sequence reads

against complex, duplicated gene families like the MHC class I genes of macaques where genomic reference

sequences and reference databases of known alleles are incomplete (Zimin et al. 2014; Maccari et al. 2017). In an

era where the per-base cost of sequencing is dropping rapidly, we explored the feasibility of obtaining whole exome

sequencing data while maintaining parity of results with the traditional MHC PCR amplicon approach.

Here we introduce MHC genotyping via macaque exome sequencing (MES), which is an exome sequencing-based

workflow for comprehensive MHC class I and class II genotyping in macaques. This workflow uses a commercially

available human exome target-capture enrichment kit in conjunction with specialized spike-in target-capture probes

to specifically cope with the high copy number of macaque MHC genes. We show that accuracy of Indian rhesus

macaque MHC class I and class II results from this workflow are comparable to conventional MiSeq genotyping

when using exon 2 reference sequences.

Methods

Animals

Twenty-seven whole blood samples were collected from Indian rhesus macaques (*Macaca mulatta*). Five of these samples came from a breeding group of animals living at the Wisconsin National Primate Research Center (WNPRC). The remaining 22 samples were provided by Dr. Michele Di Mascio from the National Institutes of Health's National Institutes of Allergy and Infectious Diseases. Blood sampling was performed under anesthesia and in accordance with the regulations and guidelines outlined in the Animal Welfare Act, the Guide for the Care and Use of Laboratory Animals, and the Weatherall report (Animal Welfare Act 1966; Weatherall 2006).

Data

Exome sequence datasets have been deposited in the sequence read archive (SRA) under BioProjects PRJNA529708 and PRJNA527214. Fasta reference sequences used for MHC genotyping, and sequence analysis scripts are available from https://go.wisc.edu/jb0926.

MHC class I and class II genotyping by amplicon deep sequencing

Genomic DNA was isolated from $250\mu L$ of whole blood using a Maxwell® 48 LEV Blood DNA Kit (Promega Corporation, Fitchburg, WI). Following isolation, DNA concentrations were determined with a Nanodrop 2000 and samples were normalized to 60 ng/ul. MHC class I and class II PCR amplicons were generated using exon 2-specific primers with adapters (CS1 and CS2) necessary for 4-primer amplicon tagging with the Fluidigm Access Array™ System (Fluidigm, San Francisco, CA, USA) by previously described methods (Karl et al. 2017; Karl et al. 2014). Pooled PCR products were purified using the AMPure XP beads (Agencourt Bioscience Corporation, Beverly, MA, USA) and quantified using the Quant-iT dsDNA HS Assay kit with a Qubit fluorometer (Invitrogen, Carlsbad, CA, USA), following the manufacturer's protocols. The MHC exon 2 genotyping amplicon pools were sequenced on an Illumina MiSeq instrument (San Diego, CA, USA) as previously described (Karl et al. 2017).

Analysis of the MiSeq exon 2 genotyping amplicon sequences was performed using a custom Python workflow. The workflow contained a step to remove oligonucleotide primers and sequencing adapters with bbduk, merge a step to

reads bbmerge, a step to identify unique sequences/remove chimeras with USEARCH, and finally mapping unique

reads against a deduplicated reference database of rhesus macaque partial MHC class I and class II exon 2 sequences

with bbmap (Bushnell et al. 2017; Edgar 2010). The database was derived from sequences in the IPD-MHC NHP

database downloaded from the European Bioinformatics Institute website (Maccari et al. 2017). For this publication,

we define "IPD exon 2" as this database of reference sequences. SAM output files from bbmap were parsed with the

Python package pandas to enumerate the reads from each animal that were identical to IPD exon 2 reference

sequences. *Mamu-A, -B, -DRB, -DQA1, -DQB1, -DPA1* and *-DPB1* lineage-level haplotypes were inferred for each

of the samples with a semi-automated custom workflow that identifies diagnostic alleles associated with previously

defined rhesus macaque haplotypes (Karl et al. 2013; Otting et al. 2017).

MHC class I and class II genotyping by exome sequencing

Genomic DNA was isolated as described above and shipped to the Human Genome Sequencing Center at the Baylor

College of Medicine. MHC and exon-containing genomic DNA fragments were selectively enriched using a custom

target-capture probeset. Genome-wide exons were captured using SeqCap EZ HGSC VCRome2.1, an optimized

human clinical exome probeset (Clark et al. 2013, Yang et al. 2013). SeqCap EZ HGSC VCRome2.1 contains

probes designed to enrich 23,585 human genes and 189,028 non-overlapping exons. A low coverage audit was

performed to identify rhesus macaque exons inferred from the reference genome rheMac2 that were not sufficiently

enriched (<20x coverage) with these human probes, and an additional 22,884 rhesus macaque exons were

incorporated into the genotyping probe design (Prall et al. 2017). Finally, and most importantly for MHC analyses,

we modified the SeqCap EZ Design: Human MHC Design to selectively enrich MHC class I and class II sequences.

This previous design was prepared by the Beijing Genome Institute in collaboration with Roche/Nimblegen and it

targeted the complete 4.97Mb HLA region with non-redundant probes designed against 8 fully sequenced HLA

haplotypes (Horton et al. 2008; Cao et al. 2013). For our macaque studies, we prepared a minimal MHC target

capture design using a subset of these probes that are based on all functional HLA class I (*HLA-A*, *-B*, *-C*, and *-E*)

and class II (*HLA-DRA*, *-DRB1*, *-DRB3*, *-DRB4*, *-DRB5*, *-DQA1*, *-DQB1*, *-DPA1* and *-DPB1*) genes. Probes were

included to capture complete gene sequences (exons + introns + 3' UTR) as well as ~1 kb of 5' upstream flanking

sequence. The BED file of rhesus rheMac2 target coordinates lifted over to rheMac8 was used to prepare this

combined minimal MHC and supplemental rhesus spike-in probe design. Because derivation of MHC results from

genotyping is paramount, we used a ratio of 2.5x spike-in probes to 1x VCRom2.1 probes. The supplemental probes

for MHC and rhesus macaque were a single reagent, and the MHC-specific probes only constituted 609 kb of the

37.9 Mb probes (Prall et al. 2017).

An Illumina paired-end pre-capture library was constructed with 750 nanograms of DNA, as described by the Baylor

College of Medicine Human Genome Sequencing Center. Pre-capture libraries were pooled into 10-plex library

pools for target capture according to the manufacturer's protocol. Samples were pooled in 10-plex sequence capture

library pools for 151 bp paired-end sequencing in a single lane of an S4 flow cell on an Illumina Novaseq 6000 at

the Baylor College of Medicine Human Genome Sequencing Center.

Enumeration of MHC reads in exome data

The effective enrichment of MHC reads in the exome datasets was calculated by mapping the reads of each sample's

exome dataset against an individual genomic reference file for each individual locus: *HLA-A* exons 2-3 (NCBI Gene

ID: 3105) and *HLA-E* exons 2-3 (Gene ID: 3133); *HLA-DPA1* exons 2-4 (Gene ID: 3113); *HLA-DPB1* exons 2-4

(Gene ID: 3115); *HLA-DQA1* exons 2-4 (Gene ID: 3117); *HLA-DQA2* exons 2-4 (Gene ID: 3118); *HLA-DQB1*

exons 2-4 (Gene ID: 3119); *HLA-DQB2* exons 2-4 (Gene ID: 3120); *HLA-DRB1* exons 2-4 (Gene ID: 3123), *HLA-*

DRB3 exons 2-4 (Gene ID: 3125), *HLA-DRB4* exons 2-4 (Gene ID: 3126) and *HLA-DRB5* exons 2-4 (Gene ID:

3127). This mapping was done by using bbmap with default parameters, which corresponds to a minimum

alignment identity of approximately 76% (Bushnell et al. 2017). Empirically, these mapping parameters are

sufficient to map macaque MHC reads to their human orthologues. Mapped reads were written to a new fastq file

using bbmap's outm= parameter. To quantify the total number of reads in a sample and the number of reads

extracted with our reference file, we created a custom Python script, which is available to download.

MHC genotyping from exome data

Two complementary data analysis strategies were employed to analyze the exome sequence data, and to verify

reproducibility and confidence in the MHC genotyping results. For accuracy and quantification purposes, the

expected MHC genotypes for each animal were established based on concordance among at least two out of the

three described strategies and biological plausibility, e.g., no more than two alleles per *Mamu-DQA1, -DQB1, -*

DPA1, or *-DPB1* locus.

Strategy 1: MHC genotyping using Diagnostic Sub-Region (DSR)

The Diagnostic Sub-Region (DSR) was an intra-allelic region that encompassed polymorphisms, and these

polymorphisms were distinguishable from alleles with similar sequences. Therefore, this method ensured the DSR

was captured in at least one read for each called allele. MHC class I and class II reads initially were extracted from

each animal by mapping the FASTQ reads to HLA class I and class II reference sequences containing exons 2-3 and

exons 2-4, respectively, plus the intervening intron(s) as described above in 'Enumeration of MHC reads in exome

data'. Reads were mapped to these reference sequences using bbmap with default parameters and the parameter

(qtrim=lr) (Bushnell et al. 2017).

Following extraction of MHC reads from the total exome sequences, the MHC reads were prepared for assembly

using a modified version of a data pre-processing pipeline, which included tools from the BBTools package

(Bushnell et al. 2017). Briefly, optical duplicates and reads from low-quality regions of the sequencing run were

removed. Next, Illumina sequencing adapters were trimmed from the ends of sequencing reads. Any residual spike-

in or PhiX sequences that inadvertently survived mapping to HLA class I and class II were then removed. Three

rounds of error-correction and read merging were performed to create high-confidence merged reads that were well-

supported by common kmers found in the extracted MHC reads. The error-corrected reads were not merged with a

minimum overlap, but instead were separately mapped against the IPD exon 2 sequences using bbmapskimmer. The

default settings for the software tool bbmapskimmer were used with the following modified parameters

(semiperfectmode=t ambiguous=all ssa=t maxsites=50000 maxsites2=50000 expectedsites=50000) (Bushnell et al.

2017). The semiperfectmode setting accepted reads with perfect matches, as well as reads that extended off the end

of contigs for no more than half of the length of the mapped read segment. The ambiguous setting and the

'expectedsites' setting reported the first 50,000 matched read segments that met the 'semiperfectmode' filtering,

which was set exceedingly high in order to exhaustively map our IPD exon 2 reference file of ~1600 alleles. Using

semiperfect mode, these were the segment sequences with the longest matching length. This output file included all

mapped reads to all IPD exon 2 reference allele sequences.

We then used samtools mpileup with the settings (-A -a --ff UNMAP -x -B -q 0 -Q 0) on the bbmapskimmer output

to calculate a depth of coverage at each position for each IPD exon 2 sequence. Next, we removed any aligned reads

to the IPD exon 2 database sequences that contained less than a minimum depth of coverage of two across the entire

reference sequence. For ambiguously-mapped reads, the alignments with the longest-matching region were selected

for further analysis; ties among alignments were counted multiple times. To reduce the number of false positives, we

used Python pandas to only report database sequence matches that had at least one unambiguously mapped read.

Based on these mapping parameters, unambiguously-mapped reads must span the DSR. The MHC genotypes from

this method were reported as the minimum depth of coverage for each IPD exon 2 database sequence per animal.

Strategy 2: De novo reconstruction of MHC sequences from exome reads

Most *de novo* sequence assemblers have been optimized for resolving long contigs, and are tolerant of small

sequence mismatches that can otherwise fragment assemblies. In the case of MHC alleles, however, such closely

related sequences were often biologically distinct. Because of the gene duplications in the macaque MHC, there

were many valid MHC contigs that could be assembled from a single sample within the exome data. This is

conceptually similar to the computational challenge of assembling viral haplotypes, where rapidly evolving viruses

such as human immunodeficiency virus accumulate variants that frequently co-segregate as minor populations

within an infected person (Baaijens et al. 2017). Therefore, we utilized the overlap assembly algorithm SAVAGE,

originally designed to reconstruct viral haplotypes, to reconstruct MHC allele sequences from exome reads (Baaijens

et al. 2017). Similar to Strategy 1 above, HLA-mapped reads were pre-processed using the BBTools package to

remove low quality reads and optical duplicates. Next, adapters were trimmed, residual spike-in and PhiX sequences

were removed, three steps of error-correction took place, and reads were merged. These merged reads, as well as

high-confidence unmerged paired-end reads that could not be grouped into an overlapping merged read, were used

for SAVAGE assembly. The following workflow was implemented in a reproducible snakemake workflow that fully

documents parameter selection (Koster and Rahman 2012), and is available upon request.

SAVAGE is designed to construct individual haplotypes from the overlap graph of individual reads. We processed

the totality of sequence data in a single patch in order to maximize the sensitivity of the contiguous reconstruction

with parameter '--split 1', as well as the parameter '--revcomp' to handle reverse complement reads. The IPD exon 2

reference database that was used for MiSeq and DSR genotyping was also used to assess the quality of SAVAGE

genotypes. The IPD sequences were mapped to contigs produced by SAVAGE using sensitive parameters in bbmap

(minlength=100 vslow=t subfilter=0 indelfilter=0 lengthtag=t ignorefrequentkmers=t kfilter=100) designed to

identify sequences that perfectly match SAVAGE contigs. A post-processing script refined these mappings, and

only retained the mappings where the length of the mapped region was the same as the length of the IPD exon 2

sequence. These mappings indicated where the reference database sequence was fully and exactly contained within a

SAVAGE contig.

Results

MHC reads are efficiently enriched using target-capture probes

We designed a custom target enrichment probeset that accounts for the extensive duplication of macaque MHC genes (Prall et al. 2017). A tripartite target capture system was used in this study. The first component is SeqCap EZ HGSC VCRome2.1, an optimized human clinical exome probeset. Since macaques are closely related to humans, SeqCap EZ HGSC VCRome2.1 can also be used in macaques, though some sequences that are most divergent between macaques and humans are not efficiently captured. The second component of the target capture system is an additional 22,884 rhesus macaque exon sequences that were not effectively captured with the SeqCap EZ HGSC VCRome2.1 reagent. The third component is a collection of probes designed to specifically enrich macaque MHC class I and class II sequences. These probes span the full length of HLA class I and class II genes including introns, 3' UTRs, and approximately 1,000 bp of flanking 5' sequence.

We obtained a median coverage of 100X for the target exon sequences across the genome with >20X coverage for 94.97% of bases that were targeted in this study. As illustrated in **Table 1**, an average of 70,549,789 Illumina sequence reads per sample were obtained for the 27 animals evaluated in this study. These reads were mapped against reference files containing representative genomic HLA exons 2-3 for *HLA-A,* and *HLA-E* exons 2 - 4 for *HLA-DRA*, *-DRB*1, *-DRB3*, *-DRB4*, *-DRB5*, *-DQA1*, *-DQB1*, *-DPA1* and *-DPB1* sequences. We identified an average of 269,057 MHC class I and class II sequence reads per sample which corresponds to an average of 0.37% of the total sequence reads evaluated per sample (**Table 1**). In a previous study by Ericsen *et al*. (Ericsen et al. 2014), we found that MHC sequences only accounted for an average of 0.13% of the total Illumina sequence reads that were evaluated per animal when the standard HGSC VCRome2.1 panel was used alone for target capture (**Supplementary Table 1**). Thus, we achieved an almost three-fold increase of MHC genomic sequences after inclusion of the spike-in probes for target capture compared to use of the VCRome2.1 probeset alone.

MHC genotypes defined from target-enriched genomic DNA are comparable to those obtained by amplicon deep sequencing

We hypothesized that MHC genotypes derived from target-enriched genomic sequence would be comparable in

accuracy to MHC genotypes derived from conventional amplicon deep sequencing. MHC class I and class II PCR

amplicons were generated from the same 27 animals and deep sequenced on an Illumina MiSeq. Representative

MHC class I and class II genotypes supported by the amplicon data are shown in **Figure 1** and **Figure 2**,

respectively. These figures illustrate genotypes from a pedigreed family of Indian rhesus macaques: a Sire, a Dam,

and three Progeny that are paternal half-siblings. Comprehensive genotypes for all 27 Indian rhesus macaques

evaluated in this study are shown in **Supplementary Figure 1**.

The superset of alleles supported by at least two of the methods described in this manuscript (MiSeq amplicon

genotyping, DSR genotyping of individual exome sequencing reads, or SAVAGE genotyping of contigs derived

from exome data), was considered to be the expected MHC genotypes for these animals. A complication for this

assertion was certain genotypes were difficult for MiSeq analysis to report correctly. As we have shown previously

(Karl et al. 2017), the number of reads supporting each genotyping call was highly variable, ranging from tens of

reads to thousands of reads per allele. A small subset of allelic variants contained nucleotide substitutions in their

sequences within the binding sites of the PCR oligonucleotides that interfere with efficient PCR amplification. This

result is exemplified when members of the *Mamu-B11L*01* allele lineage were closely examined. Members of the

*Mamu-B11L*01* allele lineage have two nucleotide substitutions relative to the 5' oligonucleotide that was used to

generate MHC class I amplicons, so these *Mamu-B11L*01* sequences were routinely absent in MiSeq genotypes

(**Figure 1**). Likewise, the *Mamu-DPA1*11:01* allele has four nucleotide substitutions relative to the oligonucleotide

pair used to generate *DPA1* amplicons for MiSeq genotyping (**Figure 2**).

These nucleotide substitutions do not fully account for all differences in the abundance of sequence reads for each

allele. False negatives were also noted for certain allele lineages, such as *Mamu-B*074* and *Mamu-B*098*, which

exhibited significantly diminished PCR efficiency despite being completely matched with the oligonucleotides used

for amplification (**Figure 1**). False positive genotyping calls were also noted in the MiSeq assay that resulted from

intermolecular recombination during the PCR process. This was exemplified by read support for the presence of

*Mamu-B*007:07* in Sire r07010, and Progeny 2 r17041 (**Figure 1**). *Mamu-B*007:07* only differs from the *Mamu-*

*B*007g1* allele group that was determined to be present in this pair of animals by a single nucleotide variant at the

extreme 5' end of the class I genotyping amplicon. Chimeric PCR products equivalent to the *Mamu-B*007:07*

sequence were formed between the 3' portion of the *Mamu-B*007g1* sequence and other allelic variants in these

animals with this 5' SNP. Taken together, these results illustrate that the MiSeq amplicon genotyping, while

generally reflective of an animal's MHC genotype, can yield both false positive and false negative results. When

compared to the expected genotypes for this dataset, MiSeq amplicon genotyping has 90.4% MHC class I and

97.8% MHC class II concordance (**Table 2**).

Two separate strategies were used to derive MHC genotypes from target-capture data. The DSR analysis was a

straightforward extension of the methodology used for deriving genotypes from MiSeq amplicons. MHC sequence

reads were extracted from the total exome dataset. A simplified workflow that mapped those MHC reads to

reference sequences, and then assigned the genotypes based on the presence of exome reads overlapping each

position of the reference sequence, was problematic. This simplified workflow was extremely vulnerable to false

positive genotypes, and the similarities among different MHC sequences often enabled those reads to map to

multiple different alleles (Wiseman et al. 2013). Two or more reads each could partially match a portion of a

sequence in the IPD exon 2 database, which could complement each other to provide support for an allele that was

not biologically relevant. As discussed in the Methods, the DSR encompassed polymorphisms that discriminated

among closely related alleles by requiring at least one mapped read to unambiguously map to the corresponding

allele within the IPD exon 2 sequences. This strategy can identify specific polymorphisms of interest among the IPD

exon 2 sequences to produce genotypes for MHC class I and MHC class II (**Figures 1**, **2** and **Supplementary**

Figure 1). For the 27 animals evaluated in this study, DSR was 97.3% and 100% concordant with expected MHC

class I and class II genotypes, respectively (**Table 2**).

The DSR strategy did not consider sequences that were not among the IPD exon 2 sequences and this lack of

consideration contributes to overcalled alleles. The apparent *Mamu-A1*059:01* allele in Dam r05029 was

erroneously derived from reads that were from *Mamu-A1*004g1* and an unknown allele that was not among the

sequences in the IPD exon 2 database. As a result of this unknown sequence not being among the known IPD exon

2 sequences, reads unambiguously mapped to *Mamu-A1*059:01*, which caused this allele to be overcalled. This type

of overcalling will be mitigated with the discovery of additional allelic variants, and their inclusion in future

iterations of the IPD database.

The second approach for determining genotypes performed *de novo* assembly on MHC class I and class II reads

from each sample. The resulting assembled contigs were then mapped against IPD exon 2 reference sequences to

define perfectly matching contigs. Because most assemblers were not tuned for the challenge of assembling large

numbers of contigs that differ from one another by as little as 1 bp, we relied on an assembler, SAVAGE, originally

designed to reconstruct viral sequencing haplotypes. As shown in **Figures 1, 2 and Supplementary Figure 1**,

contigs produced by SAVAGE matched the expected genotypes. Unlike the DSR method, SAVAGE assembled

contigs for downstream analyses. False positives and false negatives among closely-related variants were mitigated

by the inclusion of the filtering steps described under Strategy 2. Across all 27 samples, the SAVAGE contigs are

98.3% and 99.7% accurate with respect to the expected MHC genotypes (**Table 2**).

While this analysis focused on genotyping using the same exon 2 reference sequences that are commonly utilized for

MiSeq amplicon analyses, the SAVAGE contigs are frequently much longer than these reference sequences (**Figure**

3). These extended contigs frequently contain exons 2 through 4, plus the intervening introns, and could be used to

provide higher resolution genotyping than is possible using exon 2 sequence alone. Moreover, contigs that contain

complete sequences for exons 2-3 of MHC class I and exon 2 of MHC class II alleles meet the minimum criteria for

obtaining formal allele nomenclature for non-human primates from the IPD-MHC (Robinson et al. 2013).

Discussion

Here we describe MES genotyping of Indian rhesus macaques. In our estimation, this method will supersede MiSeq PCR amplicon genotyping as the most widely used macaque MHC genotyping assay in the future. This methodology has a number of compelling advantages. Most importantly, exome sequencing dramatically improves the overall quantity and quality of genomic information obtained from each sample. The same datasets used for MHC analyses may be used to evaluate protein-coding genetic variation throughout the genome. The loss of start and stop codons exome-wide can be obtained from the same datasets by modifying the workflows *in silico*, as opposed to requiring the sequencing of multiple new sets of target genes (Yang et al. 2013). Exome-wide datasets offer the promise of retrospectively identifying candidate DNA sequence variants that may be responsible for unexpected experimental outcomes in studies with macaques and other nonhuman primates. The availability of exome-wide sequences in conjunction with MHC genotypes may also increase the rigor of prospective macaque experiments by enabling more sophisticated balancing of experimental groups, and exclusion of animals whose genetics are likely to strongly bias experimental results (Loffredo et al. 2007; Reynolds et al. 2011; Haus et al. 2014).

These same exome datasets also offer the potential to improve the quality of MHC genotyping. Full-length long-read MHC transcript sequencing offers exquisite resolution, but this technology can be labor intensive and difficult to scale (Karl et al. 2017; Semler et al. 2018). The MiSeq exon 2 amplicon approach is limited in its allelic resolution, but it is the current standard deep sequencing approach for high-throughput MHC genotyping in macaques. In this report, we compare two novel strategies for MHC genotyping from MES datasets to this standard MiSeq amplicon genotyping method. The results that we obtained with all three approaches show strong concordance with expected MHC genotypes for the Indian rhesus macaques that were evaluated. As illustrated in **Figure 3a**, MHC class I genomic contigs with an average length of approximately 1.1 kb can be assembled from sequence reads that were initially extracted from the whole exome datasets.

Following the initial enrichment step, MHC class II genomic contigs averaging approximately 1.9 kb in length could be assembled from whole exome sequence reads that mapped to *HLA-DRB1*,*-DRB4*, *-DRB5*, *-DQA1*, *-DQB1*, *-DPA1* and *-DPB1* reference sequences, which contained exons 2 - 4, and the pairs of introns (**Figure 3b**). The

addition of HLA spike-in probes containing both exon and intron sequences for the target capture step, and the

introduction of 151 bp paired end reads for the Illumina NovaSeq platform greatly facilitated our ability to assemble

these extended MHC genomic contigs. Additional technological advances, including even longer sequence reads and

more efficient assembly algorithms, will undoubtedly increase genomic contig lengths as well as MHC allelic

resolution in future studies.

Both MHC genotyping approaches described here depend upon mapping exome sequence reads against a reference

database of MHC class I and class II allele sequences. Currently, the IPD database essentially is restricted to coding

regions for macaque MHC sequences, and many IPD entries are only partial transcript sequences that lack complete

coding regions. In addition, it is very challenging to correctly phase short Illumina sequence read that map to

different exons of a specific allelic variant and are separated by intronic sequences that span hundreds to thousands

of base pairs in genomic DNA. Our results with the SAVAGE workflow (**Figure 3**) demonstrate that exome

sequence reads that have been enriched with the enhanced MHC probe design described here can be assembled into

genomic contigs that span multiple exons and introns. These contigs, therefore, could also be used to improve MHC

reference databases, though this requires a major effort with more animals, and is beyond the scope of this

manuscript.

The current cost of genotyping is relatively high compared to amplicon deep sequencing, largely due to two

expenses. First, the amount of sequence data needed for a single sample can be high. Compared to conventional

MHC genotyping, where 192 macaques' data can be collected on a single instrument run, exome data acquisition is

much more expensive. On an Illumina Novaseq instrument, which has a much higher run cost than the MiSeq, only

70 exome samples can be sequenced simultaneously per lane of a S4 flow cell. However, this cost of sequencing is

rapidly decreasing, and as of early 2019, commercial providers have advertised sequencing for $9 USD per Gb of

whole genome sequence data[1]. Second, a major expense in MHC genotyping is the production, validation, and use

of target-capture arrays, as well as the development of *in silico* data analysis workflows. The approaches described

here, in particular MHC genotyping from SAVAGE contigs produced by *de novo* assembly of exome reads, is

flexible and should be adaptable as sequencing approaches evolve and improve.

---

[1] Comment via Twitter 03/18/2019 @albertvilella https://twitter.com/AlbertVilella/status/1107524501645000705

A major goal for future studies will be to attempt to extend these contigs to encompass full length genomic MHC

sequences using SAVAGE or other assembly software tools. The relatively compact genomic structure and

consistent length of MHC class I genes increase the attainability of this goal. Establishment of comprehensive

macaque MHC allele databases of extended genomic sequences will greatly facilitate mapping of exome sequence

reads since they will be contiguous with the reference sequences instead of being interrupted by intervening

sequences between each exon that are not included in current non-human primate IPD-MHC databases (Maccari et

al. 2017).

These results demonstrate that MHC genotypes can be obtained by analyzing genomic DNA selectively enriched for

MHC and protein-coding gene sequences. This represents an important advance for characterizing MHC genetics in

macaques, and this suggests that analyses of whole exome and whole genome data will become the predominant

method for studying macaque genetics in the coming decade.

Acknowledgements

We gratefully acknowledge Michele Di Mascio and his group at the National Institute of Allergy and Infectious Diseases of the National Institutes of Health for providing rhesus macaque samples used in this study. We also gratefully thank Brian Bushnell for assistance with the BBTools software, and the WNPRC for providing samples from five related macaques.

This research was supported by contracts HHSN272201600007C from the National Institute of Allergy and Infectious Diseases of the National Institutes of Health. This work was also supported in part by the Office of Research Infrastructure Programs/OD (P51OD011106) awarded to the Wisconsin National Primate Research Center at the University of Wisconsin-Madison. This research was also supported in part by grant R24-OD011173 from the National Institutes of Health. This research was conducted in part at a facility constructed with support from Research Facilities Improvement Program grants RR15459-01 and RR020141-01.

Figure Legends

Table 1. Fraction of total exome sequence reads corresponding to exons 2 - 3 of MHC class I and exons 2 - 4 of MHC class II genes.

Table 2. Summary of discordant results for each MHC genotyping method by animal.

Figure 1. Comparison of MHC class I results from MiSeq PCR amplicon versus whole exome genotyping strategies for a representative breeding group of rhesus macaques. Results for each of the three methods are provided side-by-side in the columns for each macaque. Each row indicates the detection of a specific MHC class I allele or lineage group of closely related sequences that are ambiguous because they are identical over the IPD exon 2 database sequence. Values in the body of this figure indicate the number of sequence reads supporting each allele call for the MiSeq and DSR methods while alleles supported by a SAVAGE contig are reported with a "1". Discrepancies between the MiSeq (pink), DSR (yellow) or SAVAGE (blue) methods are highlighted by filled cells with borders.

Figure 2. Comparison of MHC class II results from MiSeq PCR amplicon versus whole exome genotyping strategies for a representative breeding group of rhesus macaques. Results for each of the three methods are provided side-by-side in the columns for these five related macaques. Each row indicates detection of a specific MHC class II allele or lineage group of closely related sequences that are ambiguous because they are identical over the IPD exon 2 database sequence. Values in the body of this figure indicate the number of sequence reads supporting each allele call for the MiSeq and DSR methods while alleles supported by a SAVAGE contig are reported with a "1". The *Mamu-DPA1*11:01* allele missed by the MiSeq assay due to multiple mismatches versus the amplification primers is highlighted in pink.

Figure 3. Lengths of MHC sequence contigs assembled by the SAVAGE method for a representative breeding group of rhesus macaques. Values in the body of this figure indicate contig lengths generated for these MHC sequences in each animal. (**A**) MHC class I genotyping results are illustrated for the five animals in this breeding group. Three false negative allele calls for SAVAGE versus the expected genotypes for these animals are highlighted in magenta, e.g., *Mamu-B*030g1* in Sire r07010. Sequences highlighted in red are associated with the maternal MHC Haplotype a that was inherited by Progeny 1 from its Dam. Sequences in Progeny 1 and Progeny 3

for their MHC Haplotype b that was inherited from Sire r07010 are highlighted in dark blue while the light blue

sequences represent the alternate paternal Haplotype c that was inherited by Progeny 2. Three unique extended

MHC Haplotypes in this breeding group are indicated with shades of grey (Haplotypes d - f). MHC allele groups

that are shared by both parental haplotypes are indicated by colored borders around filled cells. (**B**) MHC class II

genotyping results are illustrated for this same breeding group. (**C**) Abbreviated Mamu haplotype designations (Karl

et al. 2013) are summarized for the six extended MHC haplotypes identified by segregation in this breeding group.

For example, the twenty MHC sequences (red) that are associated with extended Haplotype a can be summarized by

the following string of abbreviated Mamu haplotype designations: Mamu-

A004/B048/DR04a/DQA01g1/DQB06:01/DPA02g1/DPB15g.

Supplementary Materials

Supplementary Table 1. Fraction of total exome sequence reads corresponding to MHC class I and class II genes after target capture with the VCRome2.1 probe design alone. This exome sequence dataset was described previously by Ericsen and coworkers (Ericsen et al. 2014).

Supplementary Figure 1. Comparison of MHC class I results from MiSeq PCR amplicon versus whole exome genotyping assays with the DSR and SAVAGE strategies for all 27 animals. Results for each of the three method are provided side-by-side in the columns for each macaque. Each row indicates the detection of a specific MHC class I allele or lineage group of closely related sequences that are ambiguous because they are identical over the IPD exon 2 database sequence. Values in the body of this figure indicate the number of sequence reads supporting each allele call for the MiSeq and DSR methods while alleles supported by a SAVAGE contig are reported with a "1". Discrepancies between the MiSeq (pink), DSR (yellow) or SAVAGE (blue) methods are highlighted by filled cells with borders.

References

Animal Welfare Act. 'The Animal Welfare Act - Public Law 89-544 Act of August 24, 1966. 1966.

https://www.nal.usda.gov/awic/animal-welfare-act-public-law-89-544-act-august-24-1966 . Accessed

05/01/2019

Baaijens, J. A., A. Z. E. Aabidine, E. Rivals, and A. Schonhuth. 2017. De novo assembly of viral quasispecies using

overlap graphs. *Genome Res* 27: 835-848. https://www.ncbi.nlm.nih.gov/pubmed/28396522

Bimber, B. N., R. Ramakrishnan, R. Cervera-Juanes, R. Madhira, S. M. Peterson, R. B. J. Norgren, and B. Ferguson.

2017. Whole genome sequencing predicts novel human disease models in rhesus macaques. *Genomics* 109:

214-220. https://www.ncbi.nlm.nih.gov/pubmed/28438488

Brown, J. H., T. S. Jardetzky, J. C. Gorga, L. J. Stern, R. G. Urban, J. L. Strominger, and D. C. Wiley. 1993. Three-

dimensional structure of the human class II histocompatibility antigen HLA-DR1. *Nature* 364: 33-39.

https://www.ncbi.nlm.nih.gov/pubmed/8316295

Bushnell, B., J. Rood, and E. Singer. 2017. BBMerge - Accurate paired shotgun read merging via overlap. *PLoS*

One 12: e0185056. https://www.ncbi.nlm.nih.gov/pubmed/29073143

Cao, H., J. Wu, Y. Wang, H. Jiang, T. Zhang, X. Liu, Y. Xu, D. Liang, P. Gao, Y. Sun, B. Gifford, M. D'Ascenzo,

X. Liu, L. C. Tellier, F. Yang, X. Tong, D. Chen, J. Zheng, W. Li, T. Richmond, X. Xu, J. Wang, and Y. Li.

2013. An integrated tool to study MHC region: accurate SNV detection and HLA genes typing in human

MHC region using targeted high-throughput sequencing. *PLoS One* 8: e69388.

https://www.ncbi.nlm.nih.gov/pubmed/23894464

Clark, M. J., R. Chen, and M. Snyder. 2013. Exome sequencing by targeted enrichment. *Curr Protoc Mol Biol*

Chapter 7: Unit7.12. https://www.ncbi.nlm.nih.gov/pubmed/23547016

Cornish, A. S., R. M. Gibbs, and R. B. J. Norgren. 2016. Exome screening to identify loss-of-function mutations in

the rhesus macaque for development of preclinical models of human disease. *BMC Genomics* 17: 170.

https://www.ncbi.nlm.nih.gov/pubmed/26935327

Daza-Vamenta, R., G. Glusman, L. Rowen, B. Guthrie, and D. E. Geraghty. 2004. Genetic divergence of the rhesus

macaque major histocompatibility complex. *Genome Res* 14: 1501-1515.

https://www.ncbi.nlm.nih.gov/pubmed/15289473

de Manuel, M., T. Shiina, S. Suzuki, N. Dereuddre-Bosquet, H. J. Garchon, M. Tanaka, N. Congy-Jolivet, A.

Aarnink, R. Le Grand, T. Marques-Bonet, and A. Blancher. 2018. Whole genome sequencing in the search

for genes associated with the control of SIV infection in the Mauritian macaque model. *Sci Rep* 8: 7131.

https://www.ncbi.nlm.nih.gov/pubmed/29739964

Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26: 2460-2461.

https://www.ncbi.nlm.nih.gov/pubmed/20709691

Ericsen, A. J., G. J. Starrett, J. M. Greene, M. Lauck, M. Raveendran, D. R. Deiros, M. S. Mohns, N. Vince, B. T.

Cain, N. H. Pham, J. T. Weinfurter, A. L. Bailey, M. L. Budde, R. W. Wiseman, R. Gibbs, D. Muzny, T. C.

Friedrich, J. Rogers, and D. H. O'Connor. 2014. Whole genome sequencing of SIV-infected macaques

identifies candidate loci that may contribute to host control of virus replication. *Genome Biol* 15: 478.

https://www.ncbi.nlm.nih.gov/pubmed/25418588

Garcia, K. C., and E. J. Adams. 2005. How the T cell receptor sees antigen--a structural view. *Cell* 122: 333-336.

https://www.ncbi.nlm.nih.gov/pubmed/16096054

Haus, T., B. Ferguson, J. Rogers, G. Doxiadis, U. Certa, N. J. Rose, R. Teepe, G. F. Weinbauer, and C. Roos. 2014.

Genome typing of nonhuman primate models: implications for biomedical research. *Trends Genet* 30: 482-

487. https://www.ncbi.nlm.nih.gov/pubmed/24954183

Horton, R., R. Gibson, P. Coggill, M. Miretti, R. J. Allcock, J. Almeida, S. Forbes, J. G. Gilbert, K. Halls, J. L.

Harrow, E. Hart, K. Howe, D. K. Jackson, S. Palmer, A. N. Roberts, S. Sims, C. A. Stewart, J. A. Traherne,

S. Trevanion, L. Wilming, J. Rogers, P. J. de Jong, J. F. Elliott, S. Sawcer, J. A. Todd, J. Trowsdale, and S.

Beck. 2008. Variation analysis and gene annotation of eight MHC haplotypes: the MHC Haplotype Project.

Immunogenetics 60: 1-18. https://www.ncbi.nlm.nih.gov/pubmed/18193213

Kaizu, M., G. J. Borchardt, C. E. Glidden, D. L. Fisk, J. T. Loffredo, D. I. Watkins, and W. M. Rehrauer. 2007.

Molecular typing of major histocompatibility complex class I alleles in the Indian rhesus macaque which

restrict SIV CD8+ T cell epitopes. *Immunogenetics* 59: 693-703.

https://www.ncbi.nlm.nih.gov/pubmed/17641886

Karl, J. A., P. S. Bohn, R. W. Wiseman, F. A. Nimityongskul, S. M. Lank, G. J. Starrett, and D. H. O'Connor. 2013.

Major histocompatibility complex class I haplotype diversity in Chinese rhesus macaques. *G3 (Bethesda)* 3:

1195-1201. https://www.ncbi.nlm.nih.gov/pubmed/23696100

Karl, J. A., M. E. Graham, R. W. Wiseman, K. E. Heimbruch, S. M. Gieger, G. G. Doxiadis, R. E. Bontrop, and D.

H. O'Connor. 2017. Major histocompatibility complex haplotyping and long-amplicon allele discovery in

cynomolgus macaques from Chinese breeding facilities. *Immunogenetics* 69: 211-229.

https://www.ncbi.nlm.nih.gov/pubmed/28078358

Karl, J. A., K. E. Heimbruch, C. E. Vriezen, C. J. Mironczuk, D. M. Dudley, R. W. Wiseman, and D. H. O'Connor.

2014. Survey of major histocompatibility complex class II diversity in pig-tailed macaques. *Immunogenetics*

66: 613-623. https://www.ncbi.nlm.nih.gov/pubmed/25129472

Kishikawa, T., Y. Momozawa, T. Ozeki, T. Mushiroda, H. Inohara, Y. Kamatani, M. Kubo, and Y. Okada. 2019.

Empirical evaluation of variant calling accuracy using ultra-deep whole-genome sequencing data. *Sci Rep* 9:

1784. https://www.ncbi.nlm.nih.gov/pubmed/30741997

Koster, J., and S. Rahmann. 2012. Snakemake--a scalable bioinformatics workflow engine. *Bioinformatics* 28: 2520-

2522. https://www.ncbi.nlm.nih.gov/pubmed/22908215

Loffredo, J. T., J. Maxwell, Y. Qi, C. E. Glidden, G. J. Borchardt, T. Soma, A. T. Bean, D. R. Beal, N. A. Wilson,

W. M. Rehrauer, J. D. Lifson, M. Carrington, and D. I. Watkins. 2007. Mamu-B*08-positive macaques

control simian immunodeficiency virus replication. *J Virol* 81: 8827-8832.

https://www.ncbi.nlm.nih.gov/pubmed/17537848

Loffredo, J. T., J. Sidney, A. T. Bean, D. R. Beal, W. Bardet, A. Wahl, O. E. Hawkins, S. Piaskowski, N. A. Wilson,

W. H. Hildebrand, D. I. Watkins, and A. Sette. 2009. Two MHC class I molecules associated with elite

control of immunodeficiency virus replication, Mamu-B*08 and HLA-B*2705, bind peptides with sequence

similarity. *J Immunol* 182: 7763-7775. https://www.ncbi.nlm.nih.gov/pubmed/19494300

Maccari, G., J. Robinson, K. Ballingall, L. A. Guethlein, U. Grimholt, J. Kaufman, C. S. Ho, N. G. de Groot, P.

Flicek, R. E. Bontrop, J. A. Hammond, and S. G. Marsh. 2017. IPD-MHC 2.0: an improved inter-species

database for the study of the major histocompatibility complex. *Nucleic Acids Res* 45: D860-D864.

https://www.ncbi.nlm.nih.gov/pubmed/27899604

Malissen, M., B. Malissen, and B. R. Jordan. 1982. Exon/intron organization and complete nucleotide sequence of

an HLA gene. *Proc Natl Acad Sci U S A* 79: 893-897. https://www.ncbi.nlm.nih.gov/pubmed/6461010

Mothe, B. R., J. Weinfurter, C. Wang, W. Rehrauer, N. Wilson, T. M. Allen, D. B. Allison, and D. I. Watkins. 2003.

Expression of the major histocompatibility complex class I molecule Mamu-A*01 is associated with control

of simian immunodeficiency virus SIVmac239 replication. *J Virol* 77: 2736-2740.

https://www.ncbi.nlm.nih.gov/pubmed/12552014

Muhl, T., M. Krawczak, P. Ten Haaft, G. Hunsmann, and U. Sauermann. 2002. MHC class I alleles influence set-

point viral load and survival time in simian immunodeficiency virus-infected rhesus monkeys. *J Immunol*

169: 3438-3446. https://www.ncbi.nlm.nih.gov/pubmed/12218167

Nomura, T., H. Yamamoto, T. Shiino, N. Takahashi, T. Nakane, N. Iwamoto, H. Ishii, T. Tsukamoto, M. Kawada,

S. Matsuoka, A. Takeda, K. Terahara, Y. Tsunetsugu-Yokota, N. Iwata-Yoshikawa, H. Hasegawa, T. Sata, T.

K. Naruse, A. Kimura, and T. Matano. 2012. Association of major histocompatibility complex class I

haplotypes with disease progression after simian immunodeficiency virus challenge in burmese rhesus

macaques. *J Virol* 86: 6481-6490. https://www.ncbi.nlm.nih.gov/pubmed/22491464

Otting, N., M. K. van der Wiel, N. de Groot, A. J. de Vos-Rouweler, N. G. de Groot, G. G. Doxiadis, R. W.

Wiseman, D. H. O'Connor, and R. E. Bontrop. 2017. The orthologs of HLA-DQ and -DP genes display

abundant levels of variability in macaque species. *Immunogenetics* 69: 87-99.

https://www.ncbi.nlm.nih.gov/pubmed/27771735

Posey, J. E., J. A. Rosenfeld, R. A. James, M. Bainbridge, Z. Niu, X. Wang, S. Dhar, W. Wiszniewski, Z. H.

Akdemir, T. Gambin, F. Xia, R. E. Person, M. Walkiewicz, C. A. Shaw, V. R. Sutton, A. L. Beaudet, D.

Muzny, C. M. Eng, Y. Yang, R. A. Gibbs, J. R. Lupski, E. Boerwinkle, and S. E. Plon. 2016. Molecular

diagnostic experience of whole-exome sequencing in adult patients. *Genet Med* 18: 678-685.

https://www.ncbi.nlm.nih.gov/pubmed/26633545

Prall, T. M., M. E. Graham, J. A. Karl, R. W. Wiseman, A. J. Ericsen, M. Raveendran, R. Alan Harris, D. M.

Muzny, R. A. Gibbs, J. Rogers, and D. H. O'Connor. 2017. Improved full-length killer cell immunoglobulin-

like receptor transcript discovery in Mauritian cynomolgus macaques. *Immunogenetics* 69: 325-339.

https://www.ncbi.nlm.nih.gov/pubmed/28343239

Reynolds, M. R., J. B. Sacha, A. M. Weiler, G. J. Borchardt, C. E. Glidden, N. C. Sheppard, F. A. Norante, P. A.

Castrovinci, J. J. Harris, H. T. Robertson, T. C. Friedrich, A. B. McDermott, N. A. Wilson, D. B. Allison, W.

C. Koff, W. E. Johnson, and D. I. Watkins. 2011. The TRIM5{alpha} genotype of rhesus macaques affects

acquisition of simian immunodeficiency virus SIVsmE660 infection after repeated limiting-dose intrarectal

challenge. *J Virol* 85: 9637-9640. https://www.ncbi.nlm.nih.gov/pubmed/21734037

Robinson, J., J. A. Halliwell, H. McWilliam, R. Lopez, and S. G. Marsh. 2013. IPD--the Immuno Polymorphism

Database. *Nucleic Acids Res* 41: D1234-40. https://www.ncbi.nlm.nih.gov/pubmed/23180793

Semler, M. R., R. W. Wiseman, J. A. Karl, M. E. Graham, S. M. Gieger, and D. H. O'Connor. 2018. Novel full-

length major histocompatibility complex class I allele discovery and haplotype definition in pig-tailed

macaques. *Immunogenetics* 70: 381-399. https://www.ncbi.nlm.nih.gov/pubmed/29134258

Shiina, T., A. Blancher, H. Inoko, and J. K. Kulski. 2017. Comparative genomics of the human, macaque and mouse

major histocompatibility complex. *Immunology* 150: 127-138.

https://www.ncbi.nlm.nih.gov/pubmed/27395034

Silver, Z. A., and D. I. Watkins. 2017. The role of MHC class I gene products in SIV infection of macaques.

Immunogenetics 69: 511-519. https://www.ncbi.nlm.nih.gov/pubmed/28695289

Vallender, E. J. 2011. Expanding whole exome resequencing into non-human primates. *Genome Biol* 12: R87.

https://www.ncbi.nlm.nih.gov/pubmed/21917143

Weatherall, D. 2006. The use of non-human primates in research. 147. https://mrc.ukri.org/documents/pdf/the-use-

of-non-human-primates-in-research/ . Accessed 04/30/2019

Williams, T. M. 2001. Human leukocyte antigen gene polymorphism and the histocompatibility laboratory. *J Mol*

Diagn 3: 98-104. https://www.ncbi.nlm.nih.gov/pubmed/11486048

Wiseman, R. W., J. A. Karl, B. N. Bimber, C. E. O'Leary, S. M. Lank, J. J. Tuscher, A. M. Detmer, P. Bouffard, N.

Levenkova, C. L. Turcotte, E. J. Szekeres, C. Wright, T. Harkins, and D. H. O'Connor. 2009. Major

histocompatibility complex genotyping with massively parallel pyrosequencing. *Nat Med* 15: 1322-1326.

https://www.ncbi.nlm.nih.gov/pubmed/19820716

Wiseman, R. W., J. A. Karl, P. S. Bohn, F. A. Nimityongskul, G. J. Starrett, and D. H. O'Connor. 2013. Haplessly

hoping: macaque major histocompatibility complex made easy. *ILAR J* 54: 196-210.

https://www.ncbi.nlm.nih.gov/pubmed/24174442

Wittig, M., J. A. Anmarkrud, J. C. Kassens, S. Koch, M. Forster, E. Ellinghaus, J. R. Hov, S. Sauer, M. Schimmler,

M. Ziemann, S. Gorg, F. Jacob, T. H. Karlsen, and A. Franke. 2015. Development of a high-resolution NGS-

based HLA-typing and analysis pipeline. *Nucleic Acids Res* 43: e70.

https://www.ncbi.nlm.nih.gov/pubmed/25753671

Xie, C., Z. X. Yeo, M. Wong, J. Piper, T. Long, E. F. Kirkness, W. H. Biggs, K. Bloom, S. Spellman, C. Vierra-

Green, C. Brady, R. H. Scheuermann, A. Telenti, S. Howard, S. Brewerton, Y. Turpaz, and J. C. Venter.

2017. Fast and accurate HLA typing from short-read next-generation sequence data with xHLA. *Proc Natl*

Acad Sci U S A 114: 8059-8064. https://www.ncbi.nlm.nih.gov/pubmed/28674023

Xue, C., M. Raveendran, R. A. Harris, G. L. Fawcett, X. Liu, S. White, M. Dahdouli, D. Rio Deiros, J. E. Below, W.

Salerno, L. Cox, G. Fan, B. Ferguson, J. Horvath, Z. Johnson, S. Kanthaswamy, H. M. Kubisch, D. Liu, M.

Platt, D. G. Smith, B. Sun, E. J. Vallender, F. Wang, R. W. Wiseman, R. Chen, D. M. Muzny, R. A. Gibbs, F.

Yu, and J. Rogers. 2016. The population genomics of rhesus macaques (Macaca mulatta) based on whole-

genome sequences. *Genome Res* 26: 1651-1662. https://www.ncbi.nlm.nih.gov/pubmed/27934697

Yang, Y., D. M. Muzny, J. G. Reid, M. N. Bainbridge, A. Willis, P. A. Ward, A. Braxton, J. Beuten, F. Xia, Z. Niu,

M. Hardison, R. Person, M. R. Bekheirnia, M. S. Leduc, A. Kirby, P. Pham, J. Scull, M. Wang, Y. Ding, S. E.

Plon, J. R. Lupski, A. L. Beaudet, R. A. Gibbs, and C. M. Eng. 2013. Clinical whole-exome sequencing for

the diagnosis of mendelian disorders. *N Engl J Med* 369: 1502-1511.

https://www.ncbi.nlm.nih.gov/pubmed/24088041

Yang, Y., D. M. Muzny, F. Xia, Z. Niu, R. Person, Y. Ding, P. Ward, A. Braxton, M. Wang, C. Buhay, N.

Veeraraghavan, A. Hawes, T. Chiang, M. Leduc, J. Beuten, J. Zhang, W. He, J. Scull, A. Willis, M.

Landsverk, W. J. Craigen, M. R. Bekheirnia, A. Stray-Pedersen, P. Liu, S. Wen, W. Alcaraz, H. Cui, M.

Walkiewicz, J. Reid, M. Bainbridge, A. Patel, E. Boerwinkle, A. L. Beaudet, J. R. Lupski, S. E. Plon, R. A.

Gibbs, and C. M. Eng. 2014. Molecular findings among patients referred for clinical whole-exome

sequencing. *JAMA* 312: 1870-1879. https://www.ncbi.nlm.nih.gov/pubmed/25326635

Zimin, A. V., A. S. Cornish, M. D. Maudhoo, R. M. Gibbs, X. Zhang, S. Pandey, D. T. Meehan, K. Wipfler, S. E.

Bosinger, Z. P. Johnson, G. K. Tharp, G. Marcais, M. Roberts, B. Ferguson, H. S. Fox, T. Treangen, S. L.

Salzberg, J. A. Yorke, and R. B. J. Norgren. 2014. A new rhesus macaque assembly and annotation for next-

generation sequencing analyses. *Biol Direct* 9: 20. https://www.ncbi.nlm.nih.gov/pubmed/25319552

Table 1. Fraction of total exome sequence reads corresponding to exons 2 - 3 of MHC class I and exons 2 - 4 of MHC class II genes

Animal ID	SRA Accession	Total Raw Reads	MHC-I Reads Extracted	DRB1/3/4/5 Reads Extracted	DQA1 Reads Extracted	DQB1 Reads Extracted	DPA1 Reads Extracted	DPB1 Reads Extracted	MHC Reads Extracted	MHC % of total
r05029	SAMN11131055	87,345,430	79,712	215,654	24,266	39,188	29,604	112,376	500,800	0.57
r17099	SAMN11131056	67,632,236	32,406	89,946	9,694	16,782	11,214	47,974	208,016	0.31
r07010	SAMN11173514	81,062,856	71,694	204,352	22,528	38,382	25,392	111,626	473,974	0.58
r17041	SAMN11131058	80,004,430	30,852	244,326	9,746	16,310	12,644	151,376	465,254	0.58
r17061	SAMN11131059	78,428,820	70,082	204,694	20,776	41,018	19,440	95,538	451,548	0.58
J8R	SAMN11282383	80,868,782	45,164	87,864	11,878	19,736	13,810	57,146	235,598	0.29
J01	SAMN11282382	61,009,650	34,342	87,124	9,050	16,628	10,958	46,964	205,066	0.34
ZC08	SAMN11282384	60,693,394	31,168	81,586	10,392	15,660	10,172	45,748	194,726	0.32
DGKG	SAMN11282378	80,530,772	48,658	130,256	12,788	22,682	13,372	62,990	290,746	0.36
CF18	SAMN11282368	68,593,710	37,792	96,842	10,864	18,226	13,118	51,556	228,398	0.33
DE1AA	SAMN11282369	65,712,570	39,334	108,002	10,060	17,124	11,840	51,134	237,494	0.36
DEG8	SAMN11282370	68,490,344	35,566	116,824	10,404	19,278	12,294	55,086	249,452	0.36
DEXX	SAMN11282371	67,451,626	33,900	128,128	10,442	18,898	9,778	53,258	254,404	0.38
DF24	SAMN11282372	67,768,724	41,362	96,172	10,606	18,176	10,976	50,612	227,904	0.34
DF64	SAMN11282373	65,762,818	37,426	107,974	11,222	15,632	11,446	51,214	234,914	0.36
DF6T	SAMN11282374	71,472,222	41,160	101,102	11,052	18,794	13,492	51,626	237,226	0.33
DFET	SAMN11282375	77,040,660	49,068	128,862	12,588	20,952	12,586	58,238	282,294	0.37
DFV0	SAMN11282377	73,877,184	41,924	98,866	11,630	17,690	13,952	52,394	236,456	0.32
DJ94	SAMN11282379	66,051,412	39,616	88,196	9,910	15,892	10,978	45,972	210,564	0.32
HIH	SAMN11282380	76,491,578	41,308	85,732	11,148	18,366	14,102	50,746	221,402	0.29
DFJT	SAMN11282376	62,249,064	37,728	78,860	9,666	15,268	11,426	43,086	196,034	0.31
HLJ	SAMN11282381	68,288,558	41,748	94,570	9,960	17,240	12,756	48,356	224,630	0.33
37360	SAMN11282363	62,310,824	38,200	97,452	9,866	15,024	10,320	46,140	217,002	0.35
0L7	SAMN11282364	64,145,834	35,302	128,778	9,310	19,992	10,686	54,196	258,264	0.40
0R1	SAMN11282365	64,789,208	36,462	86,772	10,656	17,258	11,982	46,124	209,254	0.32
0R8	SAMN11282366	72,836,490	38,286	118,184	10,500	22,248	13,170	56,142	258,530	0.35
0TI	SAMN11282367	80,730,750	41,552	106,356	13,364	20,714	14,860	57,744	254,590	0.32
	Average	**70,549,789**	**42,660**	**119,018**	**12,014**	**20,487**	**13,569**	**61,310**	**269,057**	**0.37**

Table 2. Summary of discordant results for each MHC genotyping method by animal

Animal ID	MHC class I				MHC class II			
	Total Alleles Expected	# MiSeq Discordance	# DSR Discordance	# SAVAGE Discordance	Total Alleles Expected	# MiSeq Discordance	# DSR Discordance	# SAVAGE Discordance
r05029	28	2	1		10			
r17099	25	3		1	12			
r07010	26	3	1	1	12			
r17041	31	3		1	10			
r17061	29	1	2		12	1		
J8R	29	4	1		12	2		
J01	30	3	1	1	11	1		
ZC08	24	2		1	12			
DGKG	28	4	1		13			
CF18	28	3			12			
DE1AA	30	5	1		13			
DEG8	27	3	1		13			
DEXX	24	3	1	1	14			
DF24	32	4		1	12			
DF64	30	3	1	1	13			
DF6T	31	2		1	10			
DFET	30	3	1	1	13	1		1
DFV0	28	2			6			
DJ94	34	3	2	1	12			
HIH	30	3			10			
DFJT	28	2	3		11			
HLJ	29	2	1	1	11			
37360	29	2	1	1	13	1		
0L7	20	2			14			
0R1	17	1			12			
0R8	25	2			13	1		
0TI	25	2	1		12			
Total	747	72	20	13	318	7	0	1
Discordance (%)		9.6	2.7	1.7		2.2	0.0	0.3
Concordance (%)		90.4	97.3	98.3		97.8	100.0	99.7

Assay	Dam r05029 MiSeq	DSR	SAVAGE	Progeny 1 r17099 MiSeq	DSR	SAVAGE	Sire r07010 MiSeq	DSR	SAVAGE	Progeny 2 r17041 MiSeq	DSR	SAVAGE	Progeny 3 r17061 MiSeq	DSR	SAVAGE	Ambiguous Allele Group
Mamu-A Major Alleles																
Mamu-A1*001g2				168	27	1	158	62	1				210	74	1	A1*001:01,A1*001:02
Mamu-A1*002:01	335	136	1										447	127	1	
Mamu-A1*004g1	151	137	1	126	51	1										A1*004_6_alleles
Mamu-A1*008g1							306	79	1	352	36	1				A1*008_5_alleles
Mamu-A1*023_A1*106g1										228	36	1				A1*023:01,A1*023:02,A1*106:01
Mamu-A1*059:01		45														
Mamu-A Minor Alleles																
Mamu-A2*05g1				319	25	1	712	162	1	423	15	1	438	56	1	A2*05_32_alleles
Mamu-A3*13_A4*14g1	77	79	1										91	88	1	A3*13_7_alleles,A4*14_1_allele
Mamu-A3*13g1							93	85	1	97	31	1				A3*13:03,A3*13:11
Mamu-A4*14g1	145	104	1	164	23	1				199	29	1				A4*14_14_alleles
Mamu-E Alleles																
Mamu-E*02g1	775	185	1	1923	169	1	2041	303	1	3240	125	1	914	120	1	E*02_9_alleles
Mamu-E*02g4	307	106	1										382	95	1	E*02:13,E*02:21,E*02:29
Mamu-E*02g5	1104	134	1										1583	69	1	E*02:04,E*02:06
Mamu-E*02g7							467	152	1	634	50	1				E*02:12:01,E*02:28
Mamu-E*02g8	689	116	1	1914	92	1	781	85	1				710	85	1	E*02:20,E*02:30
Mamu-B Major Alleles																
Mamu-B*001g1							161	60	1	143	28	1				B*001_alleles
Mamu-B*007:07							25			27						
Mamu-B*007g1							161	33	1	132	12	1				B*007_6_alleles
Mamu-B*012:01													284	60	1	
Mamu-B_022g1													171	46	1	B*022:01,B*022:02
Mamu-B*027g1										93	32	1				B*027:03,B*027:04
Mamu-B*030g1				270	12	1	303	73	-	318	11	1	671	103	1	B*030_6_alleles
Mamu-B*030g2							306	79	1	306	29	1				B*030:02,B*030:05
Mamu-B*031:03				105	12	1	139	26	1				198	35	1	
Mamu-B*031g1													127	40	1	B*031:01,B*031:02
Mamu-B*041g1	128	47	1	57	24	1										B*041:01,B*041:02
Mamu-B*043:01				223	18	1	224	49	1	263	16	1	220	41	1	
Mamu-B*048:01	275	91	1	318	26	1										
Mamu-B*052:01	54	63	1													
Mamu-B*055:01	180	65	1													
Mamu-B*058:02	177	43	1													
Mamu-B*064:01	179	56	1	44	31	1										
Mamu-B*073g1				129	19	1	140	60	1				149	32	1	B*073:01,B*073:02
Mamu-B*074g1	-	82	1							-	18	1	16	48	1	B*074:01,B*074:02,B*074:03
Mamu-B Minor Alleles																
Mamu-B*035_B*049g1	161	92	1							293	37	1	261	80	1	B*035:01,B*035:02,B*049:01
Mamu-B*053g1													114	59	1	B*053:02,B*053:03
Mamu-B*054:01	194	95	1													
Mamu-B*054:04										321	26	1				
Mamu-B*054:05													169	59	1	
Mamu-B*057:01													104	41	1	
Mamu-B*057:06				59	24	1	113	61	1				51	40	1	
Mamu-B*057g1										152	10	1				B*057_alleles
Mamu-B*063:03	198	74	1	122	29	1	184	73	1	243	16	1				
Mamu-B*063:05														16		
Mamu-B*063g1	121	74	1													B*063:01_alleles,B*063:02:02_allele
Mamu-B*070g2	28	100	1										89	68	1	B*070:01,B*070:06
Mamu-B*072g1				218	41	1	447	161	1	365	25	1	330	74	1	B*072_8_alleles
Mamu-B*092g1				-	35	1	14	88	1	23	25	1	82	84	1	B*092:01,B*092:02,B*092:03
Mamu-B*098g1	198	108	1							324	32	1	238	60	1	B*098_9_alleles
Mamu-B*109:06	479	92	1	479	49	1										
Mamu-B*134g2	257	82	1	276	24	1								19		B*134:03,B*134:04,B*134:06
Mamu-B*188g1							336	42	1	535	11	1				B*188:01,B*188:02
Mamu-B11L*01:12				-	29	-	-	89	1				-	74	1	
Mamu-B11L*01g1	-	58	1	-	36	1	-	79	1	-	49	1				B11L*01_alleles,B11L*01:04,B11L*01:05
Mamu-B16*01:01				44	34	1	72	61	1				110	29	1	
Mamu-B16*01g1	181	74	1							176	23	1				B16*01_alleles
Mamu-B16*01g2							51	47	1	77	14	-				B16*01:02,B16*01:09
Mamu-B17*01g1										1221	12	1	1227	17	1	B17*01_14_alleles
Mamu-B17*01g2	1305	74	1	773	18	1	593	24	1	971	9	1				B17*01_9_alleles
Mamu-B17*01g4								6								B17*01:09,B17*01:15
Mamu-B20*01:02							304	43	1							
Mamu-B20*01g1	176	66	1	181	21	1	272	82	1	365	33	1				B20*01:01,B20*01:04
Mamu-I Alleles																
Mamu-I*01g1	130	63	1	129	31	1	485	151	1	532	44	1	252	68	1	I_01_66_alleles
Mamu-I*01g3										81	28	1				I*01:19,I*01:20:01

Fig. 1 Comparison of MHC class I results from MiSeq PCR amplicon versus whole exome genotyping strategies for a representative breeding group of rhesus macaques

Relationship Animal_ID Assay	Dam r05029 MiSeq	DSR	SAVAGE	Progeny 1 r17099 MiSeq	DSR	SAVAGE	Sire r07010 MiSeq	DSR	SAVAGE	Progeny 2 r17041 MiSeq	DSR	SAVAGE	Progeny 3 r17061 MiSeq	DSR	SAVAGE	Ambiguous Allele Group
Mamu-DRB Alleles																
Mamu-DRB1*03:09	198	370	1	72	131	1	170	305	1	272	107	1	156	200	1	
Mamu-DRB1*03g1				2789	217	1	2714	497	1				3052	499	1	DRB1*03:06,DRB1*03:26
Mamu-DRB1*10:03				1761	168	1	1952	358	1				2711	310	1	
Mamu-DRB*W2g1	154	451	1	72	183	1	132	410	1	206	176	1	107	398	1	DRB*W2:01,DRB*W2:05
Mamu-DRB*W3g1	1936	546	1							3097	217	1				DRB*W3:03:01,DRB*W3:03:02
Mamu-DRB*W4:01	1777	414	1							2631	158	1				
Mamu-DQA/DQB Alleles																
Mamu-DQA1*01:02				2384	98	1	2444	256	1				2964	154	1	
Mamu-DQA1*01g1	2298	318	1	2046	96	1	2276	288	1	4317	107	1	3360	210	1	DQA1*01:04:01,DQA1*01:04:02
Mamu-DQA1*23:01	2883	334	1							5079	108	1				
Mamu-DQB1*06:01	3803	160	1	3980	41	1	3892	152	1	5757	33	1	3892	122	1	
Mamu-DQB1*06g2				5229	173	1	4784	345	1				6400	282	1	DQB1*06:05,DQB1*06:18
Mamu-DQB1*18g4	2656	329	1							4504	108	1				DQB1*18:02,DQB1*18:27
Mamu-DPA/DPB Alleles																
Mamu-DPA1*02g1	5450	1765	1	5301	316	1	5153	840	1	8727	758	1				DPA1*02_9_alleles
Mamu-DPA1*04g1				435	181	1	489	385	1				1075	350	1	DPA1*04:03:01,DPA1*04:03:02,DPA1*04:05
Mamu-DPA1*11:01													-	225	1	
Mamu-DPB1*02g1	106	25	1	87	73	1							342	47	1	DPB1*02:01,DPB1*02:04:01,DPB1*02:04:02
Mamu-DPB1*15g1	254	217	1	286	31	1	207	129	1	3084	60	1				DPB1*15:01,DPB1*15:02
Mamu-DPB1*16:01													1769	98	1	

Fig. 2 Comparison of MHC class II results from MiSeq PCR amplicon versus whole exome genotyping strategies for a representative breeding group of rhesus macaques

A

B

Relationship / GS ID	Ave Length	Dam r05029	Progeny 1 r17099	Sire r07010	Progeny 2 r17041	Progeny 3 r17061	Ambiguous Allele Groups
MHC Haplotype 1		a	a	b	c	b	
MHC Haplotype 2		d	b	c	e	f	
Mamu-DRB Alleles							
Mamu-DRB1*03:09	1,763	3384	761	1053	559	3057	
Mamu-DRB1*03g1	866		1131	768		700	DRB1*03:06,DRB1*03:26
Mamu-DRB1*10:03	653		631	710		618	
Mamu-DRB*W2g1	591	655	520	556	613	612	DRB*W2:01,DRB*W2:05
Mamu-DRB*W3g1	2,213	1430			2996		DRB*W3:03:01,DRB*W3:03:02
Mamu-DRB*W4:01	691	644			737		
Mamu-DQA/DQB Alleles							
Mamu-DQA1*01:02	1,942		2087	1475		2264	
Mamu-DQA1*01g1	1,677	1467	1988	1429	1980	1522	DQA1*01:04:01,DQA1*01:04:02
Mamu-DQA1*23:01	2,073	2070			2076		
Mamu-DQB1*06:01	2,820	2726	1334	3199	4063	2780	
Mamu-DQB1*06g2	2,436		1076	3257		2975	
Mamu-DQB1*18g4	4,332	4336			4328		DQB1*18:02,DQB1*18:27
Mamu-DPA/DPB Alleles							
Mamu-DPA1*02g1	1,897	1989	1795	1929	1873		DPA1*02_9_alleles
Mamu-DPA1*04g1	1,985		1894	2051		2011	DPA1*04:03:01,DPA1*04:03:02,DPA1*04:05
Mamu-DPA1*11:01	1,842					1842	
Mamu-DPB1*02g1	3,693		3946	3073		4059	DPB1*02:01,DPB1*02:04:01,DPB1*02:04:02
Mamu-DPB1*15g1	2,687	2470	2506	3190	2581		DPB1*15:01,DPB1*15:02
Mamu-DPB1*16:01	827					827	

C

MHC Haplotype	a	b	c	d	e	f
Mamu-A Haplotype	A004	A001	A008	A002a	A023	A002a
Mamu-B Haplotype	B048	B043b	B001a	B055	B043a	B012b
Mamu-DRB Haplotype	DR04a	DR03f	DR04a	DR06	DR06	DR04a
Mamu-DQA	01g1	01:02	01g1	23:01	23:01	01g1
Mamu-DQB	06:01	06g2	06:01	18g4	18g4	06:01
Mamu-DPA	02g1	04g	02g1	02g1	02g1	11:01
Mamu-DPB	15g	02g	15g	15g	15g	16:01

Fig. 3 Lengths of MHC sequence contigs assembled by the SAVAGE method for a representative breeding group of rhesus macaques. Values in the body of this figure indicate contig lengths generated for these MHC sequences in each animal.

(A) MHC class I genotyping results are illustrated for the five animals in this breeding group. Three false negative allele calls for SAVAGE versus the expected genotypes for these animals are highlighted in magenta, e.g., Mamu-B*030g1 in sire r07010. Sequences highlighted in red are associated with the maternal MHC Haplotype a that was inherited by Progeny 1 from its dam. Sequences in Progeny 1 and Progeny 3 for their MHC Haplotype b that was inherited from sire r07010 are highlighted in dark blue while the light blue sequences represent the alternate paternal Haplotype c that was inherited by Progeny 2. Three unique extended Haplotypes in this breeding group are indicated with shades of grey (d-f). MHC allele groups that are shared by both parental haplotypes are indicated by colored borders around filled cells.

(B) MHC class II genotyping results are illustrated for this same breeding group.

(C) Abbreviated Mamu haplotype designations are summarized for the sx extended MHC haplotypes identified by segregation in this breeding group. For example, the twenty MHC sequences (red) that are associated with extended Haplotype a can be summarized by the following string of abbreviated Mamu haplotype designations: Mamu-A004/B048/DR04a/DQA01g1/DQB06:01/DPA02g1/DPB15g   (Karl et al 2013).

Supplementary Table 1. Fraction of exome sequence reads corresponding to MHC class I and class II genes after target capture with the VCRome2.1 probe design alone

GS ID	SRA Accession	Total Raw Reads	MHC-I Reads Extracted	DRB1/3/4/5 Reads Extracted	DQA1 Reads Extracted	DQB1 Reads Extracted	DPA1 Reads Extracted	DPB1 Reads Extracted	MHC Reads Extracted	MHC % of total
cy0642	SAMN11281734	79,795,404	4,848	26,822	2,568	2,274	900	81,340	118,752	0.15
cy0643	SAMN11281735	86,265,844	4,408	26,930	2,542	2,320	936	74,370	111,506	0.13
cy0646	SAMN11281736	109,909,444	4,724	25,476	2,382	2,196	910	72,206	107,894	0.1
cy0648	SAMN11281737	106,315,304	5,558	34,442	2,966	2,714	1,092	97,254	144,026	0.14
cy0649	SAMN11281738	78,575,038	6,022	33,742	3,176	2,970	1,134	90,556	137,600	0.18
cy0651	SAMN11281739	102,890,862	6,686	33,194	3,388	3,094	1,294	94,532	142,188	0.14
cy0652	SAMN11281740	89,074,544	4,652	25,314	2,660	2,478	976	69,976	106,056	0.12
cy0654	SAMN11281741	99,169,798	5,294	30,328	2,924	2,756	1,042	79,664	122,008	0.12
	Average	**93,999,530**	**5,274**	**29,531**	**2,826**	**2,600**	**1,036**	**82,487**	**123,754**	**0.13**