Establishing reference samples for detection of somatic mutations and germline variants with NGS technologies

Li Tai Fang[1*], Bin Zhu[2*], Yongmei Zhao[3*], Wanqiu Chen[4], Zhaowei Yang[4,30], Liz Kerrigan[5], Kurt Langenbach[5], Maryellen de Mars[5], Charles Lu[6], Kenneth Idler[6], Howard Jacob[6], Ying Yu[7], Luyao Ren[7], Yuanting Zheng[7], Erich Jaeger[8], Gary Schroth[8], Ogan D. Abaan[8], Justin Lack[3], Tsai-Wei Shen[3], Keyur Talsania[3], Zhong Chen[4], Seta Stanbouly[4], Jyoti Shetty[9], Parimal Kumar[9], John Bettridge[9], Bao Tran[9], Daoud Meerzaman[10], Cu Nguyen[10], Virginie Petitjean[11], Marc Sultan[11], Margaret Cam[12], Tiffany Hung[13], Eric Peters[13], Rasika Kalamegham[13], Sayed Mohammad Ebrahim Sahraeian[1], Marghoob Mohiyuddin[1], Yunfei Guo[1], Lijing Yao[1], Lei Song[2], Hugo YK Lam[1], Jiri Drabek[14,15], Roberta Maestro[15,16], Daniela Gasparotto[15,16], Sulev Kõks[15,17,18], Ene Reimann[15,18], Andreas Scherer[19,15], Jessica Nordlund[20,15], Ulrika Liljedahl[20,15], Roderick V Jensen[21], Mehdi Pirooznia[22], Zhipan Li[23], Chunlin Xiao[24], Stephen Sherry[24], Rebecca Kusko[25], Malcolm Moos[26], Eric Donaldson[27], Zivana Tezak[28], Baitang Ning[29], Weida Tong[29], Jing Li[30], Penelope Duerken-Hughes[31], Huixiao Hong[29#], Leming Shi[7#], Charles Wang[4,31#], Wenming Xiao[28,29#], and The Somatic Working Group of SEQC-II Consortium

[1]Bioinformatics Research & Early Development, Roche Sequencing Solutions Inc., 1301 Shoreway Road, Suite #300, Belmont, CA 94002; [2]Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, 9609 Medical Center Drive, Bethesda, Maryland 20892, USA.; [3]Advanced Biomedical and Computational Sciences, Biomedical Informatics and Data Science Directorate, Frederick National Laboratory for Cancer Research, 8560 Progress Drive, Frederick, MD21701; [4]Center for Genomics, Loma Linda University School of Medicine, 11021 Campus St., Loma Linda, CA 92350; [5]ATCC (American Type Culture Collection), 10801 University Blvd, Manassas, VA 20110; [6]Computational Genomics, Genomics Research Center (GRC), AbbVie, 1 North Waukegan Road, North Chicago, IL 60064; [7]State Key Laboratory of Genetic Engineering, School of Life Sciences and Shanghai Cancer Center, Fudan University, 2005 SongHu Road, Shanghai, China 200438; [8]Core applications group, Product development, Illumina Inc , 200 Lincoln Centre Dr. Foster City, CA 94404; [9]Genomics Laboratory, Cancer Research Technology Program, Frederick National Laboratory for Cancer Research, 8560 Progress Drive, Frederick, MD21701; [10]Computational Genomics Research, Center for Biomedical Informatics and Information Technology (CBIIT), National Cancer Institute, 9609 Medical Center Drive, Rockville, MD 20850; [11]Biomarker development, Novartis Institutes for Biomedical Research, Fabrikstrasse 10, CH-4056 Basel, Switzerland; [12]CCR Collaborative Bioinformatics Resource (CCBR), Office of Science and Technology Resources, Center for Cancer Research, NCI, Bldg 37, Rm 3041C, 37 Convent Drive, Bethesda, MD 20892; [13]Companion Diagnostics Development, Oncology Biomarker Development, Genentech, 1 DNA Way, South San Francisco, CA 94080; [14]IMTM, Faculty of Medicine and Dentistry, Palacky University Olomouc, Hnevotinska 5, 77900 Olomouc, the Czech Republic; [15]member of EATRIS ERIC- European Infrastructure for Translational Medicine; [16]Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, National Cancer Institute, Unit of Oncogenetics and Functional Oncogenomics, Via Gallini 2, 33081 Aviano (PN), Italy; [17]Perron Institute for Neurological and Translational Science, Verdun St, Nedlands, Western Australia, 6009, Australia; [18] the Centre for Molecular Medicine and Innovative Therapeutics, Murdoch

University, Murdoch, 6150, Western Australia; [19]Institute for Molecular Medicine Finland (FIMM), HiLIFE, P.O.Box 20, FI-00014 University of Helsinki, Finland; [20]Department of Medical Sciences, Molecular Medicine and Science for Life Laboratory, Uppsala University, Molekylär Medicin, Box 1432, BMC, Uppsala, 75144 Sweden; [21]Department of Biological Sciences, Virginia Tech, Life Sciences 1, 970 Washington St., Blacksburg, VA 24061; [22]Bioinformatics and Computational Biology Core, National Heart Lung and Blood Institute, National Institutes of Health, 12 SOUTH DR. Bethesda MD 20892; [23]Sentieon Inc., 465 Fairchild Drive, Suite 135, Mountain View CA 94043; [24]National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health,45 Center Drive, Bethesda, Maryland 20894; [25]Immuneering Corporation, One Broadway 14th Fl Cambridge MA 02142 USA; [26]The Center for Biologics Evaluation and Research, U.S. Food and Drug Administration, FDA, Silver Spring, Maryland; [27]Division of Antiviral Products, Office of Antimicrobial, Center for Drug Evaluation and Research, FDA, Silver Spring, Maryland; [28]The Center for Devices and Radiological Health, U.S. Food and Drug Administration, FDA, Silver Spring, Maryland; [29]Bioinformatics branch, Division of Bioinformatics and Biostatistics, National Center for Toxicological Research, Food and Drug Administration, 3900 NCTR Road, Jefferson, AR 72079; [30]Departemnt of Allergy and Clinical Immunology, State Key Laboratory of Respiratory Disease, Guangzhou Institute of Respiratory Health, the First Affiliated Hospital of Guangzhou Medical University, Guangzhou, Guangdong, 510182, P. R. China; [31]Department of Basic Science, Loma Linda University School of Medicine, Loma Linda, CA 92350

 * Contributed equally
To whome correspondence should be addressed to: Wenming.Xiao@fda.hhs.gov, chwang@llu.edu, lemingshi@fudan.edu.cn and Huixiao.Hong@fda.hhs.gov

Abstract

We characterized two reference samples for NGS technologies: a human triple-negative breast cancer cell line and a matched normal cell line. Leveraging several whole-genome sequencing (WGS) platforms, multiple sequencing replicates, and orthogonal mutation detection bioinformatics pipelines, we minimized the potential biases from sequencing technologies, assays, and informatics. Thus, our "truth sets" were defined using evidence from 21 repeats of WGS runs with coverages ranging from 50X to 100X (a total of 140 billion reads). These "truth sets" present many relevant variants/mutations including 193 COSMIC mutations and 9,016 germline variants from the ClinVar database, nonsense mutations in *BRCA1/2* and missense mutations in *TP53* and *FGFR1.* Independent validation in three orthogonal experiments demonstrated a successful stress test of the truth set. We expect these reference materials and "truth sets" to facilitate assay development, qualification, validation, and proficiency testing. In addition, our methods can be extended to establish new fully characterized reference samples for the community.

Introduction

87   In oncology, accurate somatic mutation detection is essential to diagnose cancer, pinpoint
88 targeted therapies, predict survival, and identify resistance mutations. Despite the recent
89 explosion of technological advancements, many studies have reported difficulties in obtaining
90 consistent and concordant somatic mutation calls from individual platforms or pipelines[1-3], which
91 hampers clinical validation and advancement of these biomarkers.
92
As more sequencing technologies can detect clinically actionable somatic mutations for
oncology, the need grows stronger for benchmark samples with known "ground-truth" variants.
Such a publicly available sample set would allow platform and pipeline developers to quantify
accuracy of somatic mutation calls, study reproducibility across platforms or pipelines, perform
validation usig orthogonal techniques, and calibrate best practices of protocols and methods. The
FDA has released a guidance on the use of NGS technologies for *in vitro* diagnosis of suspected
germline diseases[4], in which well-characterized reference materials are recommended to
establish NGS test performance.
101
In the absence of well-characterized samples with somatic mutations, normal samples
such as the Platinum Genome[5], HapMap[6] cell lines, or Genome in a Bottle (GiaB) consortium
materials[7,8] are often used in clinical test development and validation of somatic applications.
Also there are some gene-specific reference samples available, such as KRAS in the WHO 1st
International Reference Panel[9], or from synthetic materials[10]. Such samples do not adequately
address cancer-specific quality metrics such as somatic mutation variant allele frequency (VAF),
heterogeneity, tumor mutation burden (TMB), etc. Therefore, cancer reference samples with an
abundance of well-defined genetic alterations characterized across the whole genome are highly
desirable and urgent needed.
111
Previous attempt has characterized a cancer cell line (from metastatic melanoma) that
inquired somatic mutations (SNV/indels) in exon regions only. Germline variants and somatic
mutations across the rest of the genome were not defined[11]. In addition, this dataset is
distributed under dbGAP-controlled access, limiting its accesibility and utility. In fact, a recent
landscape analysis of currently available somatic variant reference samples published by the
Medical Devices Innovation Consortium (MDIC) did not identify any reference mutation sets that
can be used to evaluate the somatic mutation calling accuracy on a whole-genome basis[12].
119
To fulfill this unmet need, we chose a pair of cell lines, HCC1395 (triple-negative breast
cancer) and HCC395BL (B lymphocytes) from the same donor, supplied by the American Type
Culture Collection (ATCC). These two specific cell lines were chosen because they are rich in
testable features (CNVs, SNVs, indels, SVs, and genome rearrangements[13]), and may have a
potential to serve as a long-term, publicly available, and renewable reference samples with
appropriate consent from donor. Using multiple next generation sequencing (NGS) platforms,
sequencing centers, and various bioinformatics analysis pipelines we profiled these tumor-
normal matching cell lines. Thus, we minimized biases that were specific to any platform,
sequencing center, or bioinformatic algorithm, to create a list of high-confidence mutation calls
across the whole genome, here called the "truth set." A subset of these calls was further
confirmed with orthogonal targeted sequencing and Whole Exome Sequencing (WES). We also

sequenced a series of titrations between HCC1395 and HCC1395BL genomic DNA (gDNA) to
confirm candidate somatic SNV/indels.
133
We defined truth sets containing somatic mutations and germline variants in a paired cell
lines, HCC1395/HCC1395BL, with methods that minimized potential bias from library preparation,
sequencing center, or bioinformatics pipeline. While the "truth set" germline variants in
HCC1395BL can be used for benchmarking germline variant detection, the "truth set" somatic
mutations in HCC1395 can be used for benchmarking cancer mutation detection with VAF as low
as 5%. Many of variants and mutations have clinical implications. In the coding regions, a total of
193 somatic mutations are documented in the COSMIC database and 8 germline variants are
annotated as pathogenic in the ClinVar database. Interestingly, there is a nonsense somatic
mutation in the *BRCA2* gene and a nonsense germline variant in the *BRAC1* gene. Other hotspot
somatic mutations are also observed in the *TP53* and *FGFR1* genes. Thus, we believe these paired
cell lines may be highly valuable for those looking for reference samples to benchmark products
in detection of mutations in these four genes.
146

Results

148
149 **Massive data generated to characterize the reference samples**
150
To provide reference samples for the community well into the future, a matched pair,
HCC1395 and HCC1395BL was selected for profiling[14]. Previous studies of this triple negative
breast cancer cell line have revealed the existence of many somatic structural and ploidy
changes[13], which are confirmed by our cell karyotype and cytogenetic analysis (Suppl. Fig S1, S2).
Several attempts have been made to identify SNVs and small indels[15–17]. Given that appropriate
consent from the donor has been obtained for tumor HCC1395 and normal HCC1395BL for the
purposes of genomic research, we sought to characterize this pair of cell lines as publicly available
reference samples for the NGS community. In this manuscript, we focued our efforts on germline
and somatic SNVs and indels. By performing numerous sequencing experiments with multiple
platforms at different sequencing centers, we obtained high-confidence call sets of both somatic
and germline SNVs and indels (Table 1). Larger structural variants and copy number analysis will
be included in a separate manuscript that will discuss these fundings in greater detail.
163
164 **Initial Determination of Somatic Mutation Call Set**
165
High-confidence somatic SNVs and indels were obtained based primarily on 21 pairs of
tumor-normal Whole Genome Sequencing (WGS) replicates from six sequencing centers;
sequencing depth ranged from 50X to 100X (see manuscript NBT-RA46164). Each of the 21
tumor-normal sequencing replicates was aligned with BWA MEM[18], Bowtie2[19], and NovoAlign[20]
to create 63 pairs of tumor-normal Binary Sequence Alignment/Map (BAM) files. Six mutation
callers (MuTect2[21], SomaticSniper[22], VarDict[23], MuSE[24], Strelka2[25], and TNscope[26]) were applied
to discover somatic mutation candidates for each pair of tumor-normal BAM files (Fig. 1).
SomaticSeq[27] was then utilized to combine the call sets and classify the candidate mutation calls
into "PASS", "REJECT", or "LowQual". Four confidence levels (HighConf, MedConf, LowConf, and

Unclassified) were determined based on the cross-aligner and cross-sequencing center
reproducibility of each mutation call. HighConf and MedConf calls were grouped together as the
"truth set" (also known as high-confidence somatic mutations). The call set in its entirety is
referred to as the "super set" which includes low-confidence (LowConf) and likely false positive
(Unclassified) calls. For low-VAF (Variant Allele Frequency) calls, a HiSeq data set with 300×
coverage and a NovaSeq data set with 380× coverage were employed to rescue initial LowConf
and Unclassified calls into the truth set. The details are described in the Methods.
182
A breakdown of the four confidence levels is displayed in Fig. 2a. In the truth set, HighConf
calls consist of 94% of the SNVs and 79% of the indels. LowConf calls typically do not have enough
"PASS" classifications across the 63 data sets to be included in the truth set. Variants calls labeled
as Unclassified are not reproducible and likely false positives, with more "REJECT" classifications
than "PASS". The vast majority of the calls in the super set are either HighConf or Unclassified. In
other words, super set calls tend to be either highly reproducible or not at all reproducible.
189
In general, HighConf calls were classified as "PASS" in the vast majority of the data sets,
with no variant read in the matched normal and high mapping quality scores. MedConf calls
tended to be low-VAF (VAF $\lesssim$ 0.10) variants. Due to stochastic sampling of low frequency variants,
MedConf calls were not reproduced as highly across different sequencing replicates as HighConf
calls. LowConf calls (not a part of truth set) tended to have VAF near or below our detection limits
(VAF $\lesssim$ 0.05). Distinguishing the LowConf calls with sequencing noise is challenging because they
were not reproduced enough to be high-confidence calls (Fig. 2b).
197
198 **Independent AmpliSeq confirmation of Call Set**
199
We randomly selected 450 SNV and 21 indel calls of different confidence levels from the
super set and performed PCR-based AmpliSeq with approximately 2000× depth for tumor and
normal cells on an Illumina MiSeq sequencer. As we treated the AmpliSeq data set as a
confirmatory experiment, simple rules were devised to determine whether a variant call was
deemed positively confirmed, not confirmed, or uninterpretable based on the presence or
absense of somatic mutation evidence in the AmpliSeq data. Overall, positively confirmed calls
had at least 100 variant-supporting reads in the tumor but had no variant read in the normal
sample, despite sequencing depths of 600× or more in the normal. Not confirmed calls either
had no more variant-supporting read than the expected from base call errors, and/or had
VAF≥10% in the normal cells. Uninterpretable calls did not satisfy the criteria for either positive
or no validation, either because they did not have enough read depth (<50) or had fewer than
10 variant-supporting reads. (See Methods for details).
212
Both HighConf and MedConf SNV calls had very high coverage in validation and thus had
impressive validation rates (99% and 92%) (Table 2). There were only three HighConf SNV calls
that were not confirmed by AmpliSeq. Two of them had germline signals below the detection
limit of 50× in the WGS, and the third one was likely an actual somatic mutation missed by
AmpliSeq. There were only seven "positively confirmed" Unclassified SNV calls. Four of those
seven were either a part of di-nucleotide change or had deletions within 1 bp of the call. The

other three had low mapping quality scores (MQ), which drove the categorization of
"Unclassified". This result suggests that some of the "positively confirmed" Unclassified calls
might be false positives after all, but it also exposes the limitations of our truth set with regard
to complex variants and low mappability regions. LowConf and Unclassified calls (not part of the
truth set) also had higher fractions of uninterpretable calls, which consist of low-coverage
genomic positions or ambiguous variant signals. In addition, there were also 17 HighConf, 2
MedConf, 1 LowConf, and 1 Unclassified indel calls re-sequenced by AmpliSeq. The only not
confirmed HighConf indel call was caused by a germline signal. The lone Unclassified indel call
was not confirmed (we expect Unclassified calls to be not confirmed). For the inquisitive reader,
these discrepant calls (i.e., not confirmed HighConf calls and confirmed Unclassified calls) are
discussed in greater detail in the Supplementary Material.
230
The VAF calculated from the truth set correlated highly with the VAF calculated from
AmpliSeq data set, especially for HighConf calls (Fig. 2c). On the other hand, almost all the data
points at the bottom of the graph (i.e., VAF = 0 by AmpliSeq) are Unclassified calls (red). It
suggests that despite high VAFs (from 21 WGS replicates) for some of the calls, they were
categorized correctly as Unclassified (implying likely false positives). In addition, a large number
of uninterpretable Unclassified calls (red X's) lying at the bottom suggest those were correctly
labeled as Unclassified in addition to the not confirmed ones (open red circles). Moreover, some
of the seven "positively confirmed" Unclassified calls had dubious supporting evidence. Taken
together, these results suggest that the actual true positive rate for the Unclassified calls may be
even lower than the validation rate (11%) we reported here. The indel equivalent is portrayed in
Suppl. Fig S7a.
242
243 **Orthogonal Confirmation of Call Set with WES on Ion Torrent**
244
We have also sequenced the tumor-normal pair with Whole Exome Sequencing (WES) on
the Ion Torrent S5 XL sequencer with the Agilent SureSelect All Exon + UTR v6 hybrid capture.
The sequencing depths for the HCC1395 and HCC1395BL were $34\times$ and $47\times$, respectively.
Results from this Ion Torrent sequencing were leveraged to evaluate high-VAF SNV calls (Table 1
and 2). HighConf and MedConf SNV calls had high positive validation rates (99% and 89%).
However, because the Ion Torrent sequencing was performed at much lower depth, nearly 50%
of the calls were deemed uninterpretable (compared with 16% for AmpliSeq, despite having
AmpliSeq custom target enriched for low-confidence calls vs. WES). The trend of higher
uninterpretable fraction with lower confidence level calls was even more pronounced in this data
set because the coverage was too low to confirm or invalidate many low-VAF calls. The validation
rate for MedConf calls (predominantly low-VAF calls) may have suffered due to low coverage.
256
257   The VAF correlation between truth set and Ion Torrent WES (R=0.928) is lower than that
258 between truth set and AmpliSeq (R=0.958), although the vast majority of the HighConf SNV calls
259 in Ion Torrent data still stay within the 95% confidence interval area (Fig. 2d).
260
261   There are uninterpretable Unclassified calls (red X's) at the bottom for high-VAF calls,
262 which is again highly suggestive that the true positive rate for Unclassified calls may be lower

263   than the reported validation rate (25%) for Ion Torrent data as well. The indel equivalent is
264   included in Suppl. Fig. S7b.
265
266   **Independent Confirmation of Call Set with WES on HiSeq**
267
We used 14 HiSeq WES replicates from six sequencing centers to evaluate the
concordance between these data sets and the WGS data sets employed to construct the truth
set. While the WES data sets were not sequenced from orthogonal platforms, they provide
insights in terms of the reproducibility of our call sets in different library preparations. The scatter
plot between the super set derived VAF and medium HiSeq WES-derived VAF is presented in Fig.
2e. Almost all truth set (HighConf and MedConf calls) variants had consistent VAFs calculated
from both sources.
275
Again, simple rules were implemented for validation with the WES data as well (Table 2).
The validation rate for HighConf, MedConf, LowConf, and Unclassified SNV calls by WES were
100%, 98.4%, 93.1%, and 42.4%. These validation rates are higher than other methods because
these WES data were sequenced on the same platform and sequencing centers as those used to
build the truth set. Thus, the truth set variant calls are reproducible in WES, though these data
sets do not eliminate sequencing center or platform specific artifacts that may exist in both WGS
and WES data sets. The indel equivalent is the subject of Suppl. Fig. S7c.
283
284   **Validation with tumor content titration series**
285
To evaluate the effects of tumor purity, we pooled HCC1395 DNA with HCC1395BL DNA
at different ratios to create a range of admixtures representing tumor purity levels of 100%, 75%,
50%, 20%, 10%, 5%, and 0%. For each tumor DNA dilution point, we performed WGS on a HiSeq
4000 with 300× total coverage by combining three repeated runs (manuscript NBT-RA46164).
We plotted the VAF fitting score between the expected values based on the super set vs. the
observed values at each tumor fraction (Fig. 2f). For real somatic mutations, their observed VAF
should scale linearly with tumor fraction in the tumor-normal titration series. In contrast, the
observed VAF for sequencing artifacts or germline variants will not scale in this fashion. Fig. 2f
shows that the fitting scores for HighConf and MedConf calls are much higher than LowConf and
Unclassified calls across all VAF brackets, indicating that the HighConf and MedConf calls contain
far more real somatic mutations than LowConf and Unclassified calls. The formula [Eq. 2] for the
fitting score is described in the Methods.
298
299   **Definition and Confirmation of Germline SNVs/Indels in matched normal**
300
For the 21 WGS sequencing replicates of HCC1395BL (aligned with BWA MEM, Bowtie2,
and NovoAlign to create 63 BAM files) we employed four germline variant callers, i.e.,
FreeBayes[28], Real Time Genomics (RTG)[29], DeepVariant[30], and HaplotypeCaller[31], to discover
germline variants (SNV/indels). To consolidate all the calls, a generalized linear mixed model
(GLMM) was fit for each set of SNV calls which are sequenced at different centers on various
replicates, aligned by the three aligners, and discovered by the four callers. We estimated the

SNVs/indel call probability (SCP) averaged across four factors (sequencing center, sequencing replicate, aligner, and caller), and examined the variance of SCP across these factors. The SNV candidates considered were called at least four times (out of a maximum of 21x3x4=252 times) by various combination of the four factors. The frequency histogram of the averaged SCPs demonstrates a bimodal pattern (Fig. 3a). The vast majority of SNV calls (97%) had SCPs either below 0.1 (57%) or above 0.9 (40%). Only a small minority of calls (3%) lie between 0.1 and 0.9. This indicates when SNVs were repeatedly sequenced and called, only a small proportion of them would be recurrently called as SNVs, and those recurrent calls were in fact highly recurrent.

Each of our germline SNV or indel calls had annotated SCP. See the Methods and Eq. 2 for details. Suppl. Table S7 demonstrates that, of the highest-confidence calls (SCP=1, i.e, they were called everywhere), the validation rates were approximately 99% for SNV and 98% for indels by Illumina MiSeq, and 98% and 97% for Ion Torrent. Of the 11 SNV with SCP below 0.5, all were not confirmed by MiSeq. Other calls had intermediate validation rates.

Figs 3b and 3c display that the vast majority of confirmed germline VAF was around 50% and 100%. A considerable number of lower-confidence germline SNV calls clustered around 20% VAF in non-exonic regions (Fig. 3b), with a large proportion of them being uninterpretable during validation. Scatter plots for indels are qualitatively similar (Suppl. Fig. S13).

SNV Functional Relevance and TMB Benchmarks

Among the truth set somatic mutations, 186 COSMIC SNVs and 7 COSMIC indels are in the coding region. One hotspot somatic mutation of particular biological significance is a *TP53 c.128G>A* (COSMIC99023, chr17:7675088 C>T, VAF>99%), which causes an amino acid change *p.Arg43His* that leads to the inactivation of *TP53* tumor suppressive function[32]. In addition, there is also a stop gain mutation in *BRCA2 c.4777G>T* (COSMIC13843, chr13:32339132 G>A), which causes a nonsense at *p.Glu1593**, though it is only a heterozygous variant with VAF of 37.5%. Furthermore, there is a missense mutation in *FGFR1 c.473C>T* (COSM1456963, chr8: 38428420 G>A, VAF>99%).

Of the over 3.5 million high-confidence germline variants discovered in HCC1395BL, 9,016 of them are in the ClinVar database. Most of them were annotated as "benign" or "like benign"; however, 8 SNVs were annotated as "pathogenic" (Suppl. Table S9). One germline variant likely to substantially increase the risk of an affected patient to develop breast cancer is a premature stop gain in *BRCA1* (chr17:43057078, *c.5251C>A*, *p.Arg1751**, ClinVar #55480, OMIM Entry #604370). The lifetime risk of breast cancer for carriers of this variant is 80 to 90%[33]. The premature stop codon deactivates *BRCA1*'s function to repair DNA double-strand breaks. It is one of the most common germline variants among breast cancer patients. HCC1395 has both *BRCA1* and *TP53* completely inactivated, one from germline and one acquired somatically. The loss of two critical tumor suppressor genes likely contributed to tumorigenesis. A full list of COSMIC somatic mutations and ClinVar germline variants in the coding region is provided in Supplemental File 2.

Tumor mutational burden (TMB) is defined as the number of non-synonymous somatic
variants per unit area of the genome, i.e., typically the number of non-synonymous mutations
per Mbps[34]. Recent literature increasingly has reported correlations between TMB and response
to anti-PD(L)-1 immunotherapy treatment[35]. The "gold standard" to measure TMB is to perform
tumor-normal WES and find the total number of non-synonymous mutations (all in the coding
regions). Due to the high cost and time required for WES, researchers are trying to infer TMB
with much smaller and less expensive targeted oncology panels. One way to increase the
statistical power of a much smaller panel is to measure all somatic mutations, including
synonymous mutations, which is expected to correlate highly with frequency of non-synonymous
mutations if we believe most somatic mutations, especially in high-TMB patients, occur more-or-
less randomly. We inferred TMB with various commercially available target panels. The
uncertainties of mutation rate (calculated as the 95% binomial confidence interval) inferred by
smaller oncology panels are quite large, so we advise caution when attempting to infer TMB from
targeted oncology panels (Suppl. Table S10).
365
366 **Defining Genome Callable Regions**
367
Accurate variant calling requires an abundance of high-quality reads aligned accurately to
the genomic coordinates in question. False positives are overwhelmingly enriched in genomic
regions where the alignments are challenging, base call qualities are low, and/or reported
coverage is far from the mean or median[36]. There are parts of the human genome that cannot be
covered by current technologies (Fig. 4a). To obtain the callable regions, we ran GATK CallableLoci
on each of the 63 HCC1395 and HCC1395BL BAM files to identify regions of low coverage (<10),
ultra-high coverage (8× the mean coverage of the sample), difficult to map (MQ<20), poor
reads (Base Quality Score BQ<20), or with N in the reference genome. We then created
consensus callable regions that we deemed callable for our truth set. A limitation of our callable
regions and our truth set is that they were defined and relied on short-read sequencing
technologies (i.e., Illumina sequencers), because currently only high-accuracy short-read
technologies are fit for somatic variant detection due to their low VAF. Variant calls outside the
consensus callable regions were labeled NonCallable in the super set and truth set to warn users
of these potential problems (details in Methods). NonCallable regions consisted of approximately
8% of the genome but contained over 34% of all Unclassified calls and 23% of all LowConf calls in
the super set (Suppl. Table S6).
384
The consensus callable regions consist of a total of 2.73 billion bps (Fig. 4b). In comparison
with GiaB NA12878 genome's more strictly defined high-confidence (HC) regions[7], 88% of our
consensus callable regions are in common with GiaB's HC regions. On the other hand, 98% of
GiaB's HC regions are a part of our conensus callable regions. Unlike GiaB's HC which exclude
regions with structural variations as well as regions where variant calls are inconsistent with
pidigree or regions with unexplained pipeline inconsnstencies, when there were disagreements
in a variant call from various sequencing data, we did not exclude the region. Instead, we
attempted to resolve these discrepancies. When there were nearby structural changes, we relied
on machine learning algorithms to resolve these challenging events. As a result, our conensus
callable regions included some difficult genomic regions, such as human leukocyte antigen (HLA)

and olfactory receptor genes which contain high homologus sequences. The confidence (or the
lack thereof) we hold for each variant call is annotated on a per call basis. We have demonstrated
some benchmarking results with different regions in the Supplementary, section 1.10.
398
399

Discussion

401
Through a community effort, we generated a high confidence somatic mutation call set
with limit of detection (LOD) at 5% VAF (Fig. 2b). To employ as an accuracy benchmark, we
recommend considering the variant calls labeled with both HighConf and MedConf as true
positives. These true positive variants can be used to assess sensitivity, i.e., the fraction of those
variants detected by a pipeline. On the other hand, variant calls labeled as Unclassified plus any
unspecified genomic coordinates are likely false positives. LowConf calls could not be confidently
determined here and should be blacklisted for current accuracy evaluation. LowConf calls had
validation rates around 50%, and often had VAF below our $50\times$ depth detection limit. They
represent opportunities for future work to ascertain their actual somatic status.
411
The confidence level of each variant call was determined by the "PASS" classifications
provided by SomaticSeq across different sequencing centers with different aligners (see
Methods). If a variant was not detected by any caller in a data set, it was considered "Missing" in
that data set, which is common for low-VAF calls due to stochastic sampling. For most calls,
however, they either had "PASS" classifications or "REJECT or Missing" classifications, but not
both. Few variant candidates had a large number of "PASS *and* REJECT" classifications (Suppl. Fig.
S6a). HighConf calls had many "PASS" classifications, very few "REJECT" classifications, and a full
range of VAFs. MedConf calls had fewer "PASS" calls (still high), still very few "REJECT"
classifications, but were mostly low-VAF, which explains the lower number of "PASS" calls.
LowConf calls had even fewer "PASS" calls than MedConf though they overlaped significantly,
and also a low number of "REJECT" classifications. LowConf calls tended to have even lower VAF
than MedConf, around or below our detection limit (Fig. 2b). Only Unclassified calls suffered a
significant number of "REJECT" classifications, and they also displayed a full range of VAF. The
performance of Unclassified calls indicated that SomaticSeq labeled them "REJECT" due to poor
mapping, poor alignment, germline risk, or causes other than lack of variant reads. HighConf and
Unclassified calls are far apart in all of the metrics describedabove.
428
429   Variant re-sequencing with AmpliSeq (Suppl. Fig. S6c) pointed to a high validation rate for
430 HighConf and MedConf calls. Suppl. Fig. S6c also contains a cluster of Unclassified and LowConf
431 calls in the middle of the XY plane, representing calls with some conflicts (i.e., large number of
432 "PASS" and "REJECT" calls).
433
Each time a human cell divides, somatic mutations could be introduced by replication
errors. Somatic mutations can occur much more frequently in cancer cells with malfunctioning
DNA repair systems. It is not feasible to detect extremely low-VAF somatic mutations because
they may appear in few tumor cells. Our "truth set" for somatic mutation was built upon WGS
with $50\times$-100X coverages, and thus it was designed to detect somatic mutations limited to 5% of

VAF. Variants with low-VAF (≤12%) were cross-referenced with two data sets with depths over
300× to ascertain their presence. While we do not expect our truth set to be 100% accurate or
100% comprehensive, the AmpliSeq and Ion Torrent data sets demonstrated combined 99% and
91% validation rates for HighConf and MedConf SNV calls, respectively. AmpliSeq also showed a
94% validation rate for HighConf indel calls. VAF of 5% represents the lower detection limit of the
first release of the somatic mutation truth set, even though there are many true mutations with
VAF under that threshold. We recommend that if using this truth set as a benchmark, novel
variant calls (i.e., variants calls not present in our super set) with VAF<5% should be blacklisted
from the accuracy calculations because we cannot confidently determine their status. Due to
losses of chr6p, chr16q, and chrX in HCC1395BL (Suppl. Fig S1, S2), somatic mutations in these
regions were excluded.
450
For the first time, tumor-normal paired "reference samples" with a whole-genome
characterized somatic mutation and germline "truth sets" are available to the community. Our
samples, data sets, and the list of known somatic mutations can serve as a public resource for
evaluating NGS platforms and pipelines. The massive and diverse amount of sequencing data
generated from multiple platforms at multiple sequencing centers can help tool developers to
create and validate new algorithms and to build more accurate artificial intelligence (AI) models
for somatic mutation detection. The reference samples and call set presented here can help in
assay development, qualification, validation, and proficiency testing. Such community defined
tumor-normal paired reference samples can be helpful in quality assessment by clinical
laboratories engaged in NGS, data exchange between laboratories, characterization of gene
therapy products, and premarket review of NGS-based products. Furthermore, the methodology
used in this study can be extended to establish truth sets for additional cancer reference samples.
Other reference sample efforts may be able to build on the data sets we established or consider
using these samples as a genomic background for other reference samples.
465
466

467 ## Methods
468 See Online Methods
469

Acknowledgements

We thank Justin Zook of the National Institute of Standard Technology for advice in establishing
reference samples and truth set; Sivakumar Gowrisankar of Novartis, Susan Chacko of the Center
for Information Technology, the National Institute of Health for their assistance with data
transfer; Dr. Jun Ye of Sentieon for providing the Sentieon software package. We also appreciate
Dr. Laufey Amundadottir of the Division of Cancer Epidemiology and Genetics, National Cancer
Institute (NCI), National Institutes of Health (NIH), for the sponsorship and the usage of the NIH
Biowulf cluster; Drs Reena Phillip, Yun-Fu Hu, Sharon Liang, and You Li of the Center for Devices
and Radiological Health, U.S. Food and Drug Administration, for their advices on study design and
manuscript writing; and Seven Bridges for providing storage and computational support on the
Cancer Genomic Cloud (CGC). The CGC has been funded in whole or in part with Federal funds
from the National Cancer Institute, National Institutes of Health, Contract No.

HHSN261201400008C and ID/IQ Agreement No. 17X146 under Contract No.
HHSN261201500003I. Chunlin Xiao and Steve Sherry were supported by the Intramural Research
Program of the National Library of Medicine, National Institutes of Health. This work also used
the computational resources of the NIH Biowulf cluster (http://hpc.nih.gov). Original data was
also backed up on the servers provided by Center for Biomedical Informatics and Information
Technology (CBIIT), NCI. We would also like to thank the partially support from the Ardmore
Institute of Health (AIH) grant (2150141) and Dr. Charles A. Sims' gift to Loma Linda University
(LLU) Center for Genomics. The LLU Center for Genomics is partially supported by AIH grant
(2150141) and Charles A. Sims' gift.
491

Disclaimer

493 This is a research study, not intended to guide clinical applications. The views presented in this
494 article do not necessarily reflect current or future opinion or policy of the US Food and Drug
495 Administration. Any mention of commercial products is for clarification and not intended as
496 endorsement.
497

Data availability

499 All raw data (FASTQ files) are available on NCBI's SRA database (SRP162370). The truth set for
500 somatic mutations in HCC1395, VCF files derived from individual WES and WGS runs, and source
501 codes are available on NCBI's ftp site (ftp://ftp-
502 trace.ncbi.nlm.nih.gov/seqc/ftp/Somatic_Mutation_WG/). Some alignment files (BAM) are also
503 available on Seven Bridges' s Cancer Genomics Cloud (CGC) platform.

Author contributions

506 Study conceived and designed by: W.X., C.W., L.S., H.H., E.D., Z.T., B.N., W.T., R.J.
507
508 Biosample preparation: L.K., K.L., M. M., T.H., W.C.
509
510 NGS library preparation and sequencing: W.C., Z.C., S.S., K.I., H.J., E.J., G.S., S.S., J.S., P.K., J.B., B.T.,
511 V.P., M.S., T.H., E.P., R.K., J.D., P.V., R.M., D.G., S.K., E.R., A.S., J.N., U.L., J.W., J.L., P.PH.
512
513 Data analysis: L.T.F., W.X., B.Z., Y.Z., Z.Y., C.L., O.A., L.S., J.L., T.S., K.T., D.M., C.N., M.C., S.M.S.,
514 M.M., Y.G., L.Y., H.L., M.P., Z.L.
515
516 Data management: W.X., C.X., S. S.
517
518 Manuscript writing: L.T.F., W.X., R.K., M.M., C.X., S.S.
519
520 Project management: W.X.
521

References

1. Hofmann, A. L. *et al.* Detailed simulation of cancer exome sequencing data reveals differences
and common limitations of variant callers. *BMC Bioinformatics* **18**, 8 (2017).

2. Krøigård, A. B., Thomassen, M., Lænkholm, A.-V., Kruse, T. A. & Larsen, M. J. Evaluation
of Nine Somatic Variant Callers for Detection of Somatic Mutations in Exome and Targeted
Deep Sequencing Data. *PLOS ONE* **11**, e0151664 (2016).

3. Shi, W. *et al.* Reliability of Whole-Exome Sequencing for Assessing Intratumor Genetic
Heterogeneity. *Cell Rep.* **25**, 1446–1457 (2018).

4. Tezak, Z. & Berger, A. Considerations for Design, Development, and Analytical Validation
of Next Generation Sequencing (NGS) – Based In Vitro Diagnostics (IVDs) Intended to Aid
in the Diagnosis of Suspected Germline Diseases. (2018).

5. Eberle, M. A. *et al.* A reference data set of 5.4 million phased human variants validated by
genetic inheritance from sequencing a three-generation 17-member pedigree. *Genome Res.* **27**,
157–164 (2017).

6. The International HapMap Project. *Nature* **426**, 789 (2003).

7. Zook, J. M. *et al.* Integrating human sequence data sets provides a resource of benchmark SNP
and indel genotype calls. *Nat. Biotechnol.* **32**, 246–251 (2014).

8. Zook, J. M. *et al.* An open resource for accurately benchmarking small variant and reference
calls. *Nat. Biotechnol.* 1 (2019). doi:10.1038/s41587-019-0074-6

9. National Institute for Biological Standards and Control. *WHO Reference Panel 1st*
International Reference Panel for genomic KRAS codons 12 and 13 mutations NIBSC code:
16/250. (World Health Organization).

10. Oncospan   Reference   Standard   HD827.   Available   at:
https://www.horizondiscovery.com/reference-standards/type/oncospan. (Accessed: 17th April

2019)

11. Craig, D. W. *et al*. A somatic reference standard for cancer genome sequencing. *Sci. Rep.* **6**,
549   24607 (2016).

12. Zehnbauer, B. *et al*. MDIC SRS Report: Somatic Variant Reference Samples for NGS.
551   (2019).

13. Popova, T. *et al*. Ploidy and Large-Scale Genomic Instability Consistently Identify Basal-like
553   Breast Carcinomas with BRCA1/2 Inactivation. *Cancer Res.* **72**, 5454–5462 (2012).

14. Gazdar, A. F. *et al*. Characterization of paired tumor and non-tumor cell lines established from
555   patients with breast cancer. *Int. J. Cancer* **78**, 766–774 (1998).

15. Kalyana-Sundaram, S. *et al*. Gene Fusions Associated with Recurrent Amplicons Represent a
557   Class of Passenger Aberrations in Breast Cancer. *Neoplasia* **14(8)**, 702–708 (2012).

16. Zhang, J. *et al*. INTEGRATE: gene fusion discovery using whole genome and transcriptome
559   data. *Genome Res.* **26**, 108–118 (2016).

17. Robinson, D. R. *et al*. Functionally recurrent rearrangements of the MAST kinase and Notch
561   gene families in breast cancer. *Nat. Med.* **17**, 1646–1651 (2011).

18. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
563   *ArXiv13033997 Q-Bio* (2013).

19. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
565   357–359 (2012).

20. NovoCraft   Technologies   Sdn   Bhd.   NovoAlign.   Available   at:
567   http://www.novocraft.com/products/novoalign/.

21. Cibulskis, K. *et al*. Sensitive detection of somatic point mutations in impure and heterogeneous
cancer samples. *Nat. Biotechnol.* **31**, 213–219 (2013).

22. Larson, D. E. *et al*. SomaticSniper: identification of somatic point mutations in whole genome

sequencing data. *Bioinformatics* **28**, 311–317 (2012).

23. Lai, Z. *et al*. VarDict: a novel and versatile variant caller for next-generation sequencing in

cancer research. *Nucleic Acids Res*. **44**, e108 (2016).

24. Fan, Y. *et al*. MuSE: accounting for tumor heterogeneity using a sample-specific error model

improves sensitivity and specificity in mutation calling from sequencing data. *Genome Biol*.

17, 178 (2016).

25. Kim, S. *et al*. Strelka2: fast and accurate calling of germline and somatic variants. *Nat*.

Methods **15**, 591 (2018).

26. Freed, D., Pan, R. & Aldana, R. TNscope: Accurate Detection of Somatic Mutations with

Haplotype-based Variant Candidate Detection and Machine Learning Filtering. *bioRxiv*

250647 (2018). doi:10.1101/250647

27. Fang, L. T. *et al*. An ensemble approach to accurately detect somatic mutations using

SomaticSeq. *Genome Biol*. **16**, 197 (2015).

28. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

ArXiv12073907 Q-Bio (2012).

29. Real Time Genomics (RTG) Variant Caller. Available at: https://www.realtimegenomics.com/.

30. Poplin, R. *et al*. A universal SNP and small-indel variant caller using deep neural networks.

Nat. *Biotechnol*. **36**, 983–987 (2018).

31. Poplin, R. *et al*. Scaling accurate genetic variant discovery to tens of thousands of samples.

bioRxiv 201178 (2017). doi:10.1101/201178

32. Soussi, T., Leroy, B. & Taschner, P. E. M. Recommendations for Analyzing and Reporting

TP53 Gene Variants in the High-Throughput Sequencing Era. *Hum*. *Mutat*. **35**, 766–778

(2014).

33. OMIM Clinical Synopsis - #604370 - BREAST-OVARIAN CANCER, FAMILIAL,

SUSCEPTIBILITY   TO,   1;   BROVCA1.   Available   at:

https://www.omim.org/clinicalSynopsis/604370. (Accessed: 22nd March 2019)

34. Meléndez, B. *et al*. Methods of measurement for tumor mutational burden in tumor tissue.

Transl. Lung Cancer Res. **7**, 661–667 (2018).

35. Goodman, A. M. *et al*. Tumor Mutational Burden as an Independent Predictor of Response to

Immunotherapy in Diverse Cancers. *Mol. Cancer Ther*. **16**, 2598–2608 (2017).

36. Li, H. Toward better understanding of artifacts in variant calling from high-coverage samples.

Bioinformatics **30**, 2843–2851 (2014).

603
604

NGS technologies		platforms	# of reads (coverage)	
			HCC1395	HCC1395BL
Initial	WGS	HiSeq	57 billion (2,800X)	57 billion (2,800X)
		NovaSeq	13 billion (650X)	13 billion (650X)
		10X Genomics	20 billion (1,000X)	20 billion (1,000X)
		PacBio	20 million (50X)	20 million (50X)
Validation	WGS-tumor content	HiSeq	7.6 billion (380X)	7.6 billion (380X)
	WES	HiSeq	5 billion (12,500X)	5 billion (12,500X)
		Ion Torrent	67 million (34X)	82 million (47X)
	AmpliSeq	MiSeq	3.3 million (2000X)	3.3 million (2000X)

605   **Table 1**. Massive data from multiple NGS platforms was obtained to derive and confirm germline
606   and somatic variants in HCC1395 and HCC1395BL

Validation Platform	Variant Type	Category	Total Number	Fraction Interpretable	Validation Rate (Interpretable)	Validation Rate (Total)
AmpliSeq Deep Sequencing	SNV	HighConf	247	(237/247) 96.0%	**(234/237) 98.7%**	94.7%
		MedConf	40	(37/40) 92.5%	**(34/37) 91.9%**	85.0%
		LowConf	58	(41/58) 70.7%	**(22/41) 53.7%**	37.9%
		Unclassified	105	(62/105) 59.0%	**(7/62) 11.3%**	6.7%
	INDEL	HighConf	17	(17/17) 100.0%	**(16/17) 94.1%**	94.1%

Platform	Type	Confidence	Count			
		MedConf	2	(2/2) 100.0%	**(2/2) 100.0%**	100.0%
		LowConf	1	(0/1) 0.0%	**(0/0) NA**	nan%
		Unclassified	1	(1/1) 100.0%	**(0/1) 0.0%**	0.0%
Ion Torrent WES	SNV	HighConf	703	(629/703) 89.5%	**(623/629) 99.0%**	88.6%
		MedConf	43	(27/43) 62.8%	**(24/27) 88.9%**	55.8%
		LowConf	134	(39/134) 29.1%	**(28/39) 71.8%**	20.9%
		Unclassified	802	(155/802) 19.3%	**(39/155) 25.2%**	4.9%
	INDEL	HighConf	31	(25/31) 80.6%	**(22/25) 88.0%**	71.0%
		MedConf	15	(7/15) 46.7%	**(6/7) 85.7%**	40.0%
		LowConf	8	(0/8) 0.0%	**(0/0) NA**	nan%
		Unclassified	36	(8/36) 22.2%	**(6/8) 75.0%**	16.7%
WES	SNV	HighConf	1074	(1068/1074) 99.4%	**(1068/1068) 100%**	99.4%
		MedConf	64	(63/64) 98.4%	**(62/63) 98.4%**	96.9%
		LowConf	197	(144/197) 73.1%	**(134/144) 93.1%**	68.0%
		Unclassified	1218	(436/1218) 35.8%	**(184/436) 42.4%**	15.1%
	INDEL	HighConf	45	(43/45) 95.6%	**(43/43) 100%**	95.6%
		MedConf	17	(17/17) 100.0%	**(17/17) 100%**	100.0%
		LowConf	13	(10/13) 76.9%	**(9/10) 90%**	69.2%
		Unclassified	54	(19/54) 35.2%	**(14/19) 73.7%**	25.9%

Table 2: Validation of SNVs of different confidence levels by three different methods

Figure legends

Figure 1: Schematic of the bioinformatics pipeline used to define the confidence levels of the super set and truth set (see Online Methods for detail)

Figure 2: Initial definition of somatic mutation truth set and subsequent validation. (a) A breakdown of the four confidence levels in the super set. (b) Histograms of VAF for SNVs (top) and Indels (bottom) calls. (c) Validation of initial definition of somatic mutation truth set with AmpliSeq. Solid circles are variant calls that were positively confirmed. Open circles are variants that were not confirmed. X's are when validation data were deemed uninterpretable due to low depth or unclear signal. The dashed lines at the diagonal represent the 95% binomial confidence-interval of observed VAF given the actual VAF, calculated based on 2000× depth for AmpliSeq. The figure shows very high correlation between VAF estimated from super set data and validation data for HighConf calls (R=0.958). Many Unclassified data points lie at the bottom, implying that those calls were not real mutations despite the large number of apparent variant-supporting reads in the super set data. X-axis: VAF calculated from the super set. Y-axis: VAF calculated from AmpliSeq data. (d) Validation of the initial definition of the somatic mutation truth set with Ion Torrent WES. The 95% binomial confidence-interval dash lines were calculated based on 34× depth for Ion Torrent. R=0.928 for HighConf calls. (e) Validation of initial definition of somatic mutation truth set with 12 repeats of WES on the HiSeq platform. Y-axis: median VAF calculated based on 12 HiSeq WES replicates. The 95% binomial confidence-interval dashed lines were calculated based on 150× depth for HiSeq WES. R=0.992 for HighConf calls. (f) Average tumor

purity fitting scores for the VAF of each SNV across the four different confidence levels vs. the
observed VAF in the tumor-normal titration series. The formula for fitting scores is described in
Eq. 1 in the Online Methods.

Figure 3: Initial definition of germline variants and validation. (a) Histogram of SNV call probability
for germline variants identified by four callers from 63 BAM files. (b) VAF scatter plot of germline
SNVs by the truth set and AmpliSeq. R=0.986 for SCP=1 calls. (c) VAF scatter plot of germline SNVs
by the truth set and Ion Torrent WES. R=0.758 for SCP=1 calls.

Figure 4: Genome coverage and high-confidence regions on reference genome GRCh38. a)
Genome coverage comparison between three technologies. Inner track: PacBio. Middle track:
10X Genomics. Outer track: Illumina. Red line: HCC1395. Green line: HCC1395BL. b) Genome
regions coverage by Illumina short reads in comparison to NA12878. Inner track: NA12878.
Middle track: the callable regions in HCC1395 and HCC1395BL. Outer track: gene density