Knocking out non-muscle myosin II in retinal ganglion cells promotes long-distance optic nerve regeneration

Xue-Wei Wang[1, #], Shu-Guang Yang[1, #], Chi Zhang[1, #], Jin-Jin Ma[3], Yingchi Zhang[1], Bin-Bin Yang[1], Yi-Lan Weng[5], Guo-Li Ming[6], Anish R. Kosanam[1], Saijilafu [3,4, *], and Feng-Quan Zhou[1, 2, *]

[1]Department of Orthopaedic Surgery, The Johns Hopkins University School of Medicine, Baltimore, MD 21287

[2]The Solomon H. Snyder Department of Neuroscience, The Johns Hopkins University School of Medicine, Baltimore, MD 21287

[3]Orthopaedic Institute, Medical College, Soochow University, Suzhou, Jiangsu, China

[4]Department of Orthopaedic Surgery, the First Affiliated Hospital, Soochow University, Suzhou, China

[5]Center for Neuroregeneration, Department of Neurosurgery, Houston Methodist Research Institute, Houston, TX 77030

[6]Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

[#]These authors contributed equally to this study

*Correspondence to:

Feng-Quan Zhou, Ph.D., Room 291, The John G. Rangos Sr. Building, 855 North Wolfe Street, Baltimore, MD 21287, Email: fzhou4@jhmi.edu

25   Saijilafu, Ph.D. Orthopaedic Institute, Medical College, Soochow University, Suzhou, Jiangsu, China

26   Email: saijilafu@suda.edu.cn

Summary

In addition to changed gene expression, pathological cytoskeletal dynamics in the axon is another key intrinsic barrier for axon regeneration in the central nervous system (CNS). Here we showed that knocking out myosin IIA/B in retinal ganglion cells alone was sufficient to induce marked and sustained optic nerve regeneration. Combined Lin28 overexpression and myosin IIA/B knockout led to remarkable synergistic promoting effect and long-distance axon regeneration. Immunostaining, RNA-seq and western blot analyses revealed that myosin II deletion did not affect known axon regeneration signaling pathways or the expression of regeneration associated genes. Instead, it abolished the retraction bulb formation and significantly enhanced the axon extension efficiency. The study provided clear and strong evidence that directly targeting neuronal cytoskeleton was sufficient to induce strong CNS axon regeneration, and combining gene expression in the soma and modified cytoskeletal dynamics in the axon was an optimal approach for long-distance CNS axon regeneration.

Keywords

Axon regeneration, optic nerve regeneration, non-muscle myosin II, Lin28, growth cone, cytoskeleton

Introduction

Axon regeneration in the mammalian central nervous system (CNS) has been a long standing and highly challenging issue in the biomedical research field. The current consensus is that there are two major reasons that neurons in the mature mammalian CNS do not regenerate their axons after the injury. One is the hostile environment caused by inhibitors in the scar tissues and degenerating myelin, and the other is the diminished intrinsic neural regeneration ability of mature CNS neurons[1,2]. Therefore, the widely accepted view is that combination strategies that target both intrinsic growth ability and inhibitory environment are likely the best option for successful CNS axon regeneration and function recovery. Early studies[3-6] using peripheral nerve graft transplants have shown that some mature CNS neurons, such as spinal cord neurons and retinal ganglion cells (RGCs), could regenerate their axons into the permissive nerve grafts, indicating clearly that these neurons still retain limited intrinsic regeneration ability. However, to date many studies targeting selected inhibitory molecules resulted in no or very modest CNS regeneration[7,8]. A likely reason is that there are multiple classes of inhibitory molecules, potentially including unidentified ones, which inhibit axon regeneration via distinct cellular and molecular mechanisms. Thus, targeting a few inhibitory signals while leaving the others intact may not result in a permissive environment similar to that in the peripheral nerve grafts.

In contrast, several studies targeting the intrinsic axon growth ability (e.g. Pten, SOCS3, KLF4, Lin28)[9-13] have produced very promising results. However, in the corticospinal tract (CST) regeneration model, although modulation of the intrinsic regeneration ability substantially enhanced axon regeneration, most regenerating axons still cannot pass the lesion site, likely due to the effects of inhibitory molecules at the injury site, especially glial scar-based inhibitors. For instance, Pten deletion has been shown to induce by far the strongest promoting effect on CST axon regeneration[9]. However, the most robust promoting effect can only be achieved in young mice (< 1 month). A recent study[14] showed that Pten

deletion-induced regeneration of CST axons beyond the injury site was greatly diminished in aged mice. Specifically, in 12-18-month-old mice, Pten deletion led to little, if any, CST regeneration beyond the injury site. One likely reason for the diminished effect in older animals was the increased response to the inhibitory CNS environment. Thus, developing a successful strategy for stimulating regeneration of injured CST remains a challenge, especially in older animals. In the optic nerve regeneration model, boosting the intrinsic regeneration ability of RGCs have produced strong optic nerve regeneration. However, tissue clearing and 3D imaging studies have revealed that many of the regenerating RGC axons make U-turns in the optic nerve, at the optic chiasm, or make wrong guidance decisions after the chiasm[15,16]. Together, a new strategy is needed to enable neurons with increased intrinsic axon growth ability to grow axons in the inhibitory environment more efficiently with less U-turns and unnecessary branching, and can cross the inhibitory boundary more efficiently.

Neuronal cytoskeleton is not only the major machinery that drives axon growth[17-19], but also the converging targets of most, if not all, inhibitory signaling pathways[17,19]. In other words, by directly manipulating growth cone cytoskeletal motility it is possible to interfere with how the growth cones respond to multiple inhibitory signals, regardless if these signals are from different inhibitors or downstream pathways. Indeed, our previous study[20] showed that knocking down or pharmacologically inhibiting non-muscle myosin IIA/B could allow regenerating sensory axons to grow straight, make less branches, and completely ignore chondroitin sulfate proteoglycans (CSPGs) and myelin-based inhibitors. The effects were much stronger than that of Rho kinase inhibitor. Here we examined if knocking out non-muscle myosin IIA/B in RGCs could promote optic nerve regeneration in vivo. The results showed that deleting myosin IIA/B alone was sufficient to induce robust and sustained optic nerve regeneration. Moreover, combination of myosin IIA/B knockout and Lin28 overexpression, which enhances the intrinsic axon regeneration ability of RGCs[13], led to synergistic promoting effect and surprisingly long-

distance optic nerve regeneration. Importantly, the promoting effect was independent of well-known

signaling mediators of optic nerve regeneration, such as increased mTOR activity and GSK3β

inactivation[2]. RNA-seq and western blot analyses comparing wild type and myosin IIA/B knockout

neurons showed no significant difference in the expression of known regeneration associated genes,

indicating local effects in the axons. In support, detailed analyses of growth cone morphologies and axon

trajectories revealed that myosin II deletion almost abolished the formation of retraction bulbs, a hallmark

of failed axon regeneration, and significantly enhanced the axon extension efficiency in the optic nerve.

Collectively, our study demonstrated clearly that manipulation of neuronal cytoskeleton alone was

sufficient to promote significant CNS axon regeneration in vivo. The study also provided strong evidence

that combining enhanced intrinsic regeneration ability in the neuronal soma with local manipulation of

axonal cytoskeleton was an optimal approach to induce long-distance CNS axon regeneration in vivo.

Results

Double knockout of non-muscle myosin IIA/B in RGCs led to significant and sustained optic nerve regeneration in vivo

Non-muscle myosin II consists of two essential light chains, two regulatory light chains, and a myosin II heavy chain (MHC). There are three different isoforms of MHCs in mammalian cells, IIA, IIB, and IIC, encoded by *Myh 9,10*, and *14* genes, respectively. Myosin IIA and IIB with MHC IIA/B (myosin IIA/B) are the major isoforms in neurons[20]. In nerve growth cones, both myosin IIA/B are localized near the transition zone, where microtubules and actin filaments interact[20]. Our previous study[20] has shown that pharmacological inhibition or double knockdown of myosin IIA/B in developing or regenerating sensory neurons drastically promoted sensory axon growth over two major inhibitory substrates, myelin extracts or CSPGs. Here we tested if myosin IIA/B loss of function could also induce axon regeneration in RGCs after optic nerve injury. To knock out both myosin IIA/B in RGCs, we crossed *Myh9^{f/f}* and *Myh10^{f/f}* mice to generate *Myh9^{f/f}: Myh10^{f/f}* mice (hereafter *myosin IIA/B^{f/f}*), and injected AAV2-Cre into the vitreous humors of these mice. Wild type mice injected with AAV2-Cre were used in the control group. To examine viral vector infection rate, immunostaining of Cre recombinase in whole-mount retina was performed two weeks after the injection. The results showed that the infection rate in RGCs was about 90% (Supplementary Figure 1A, C). Immunostaining of retinal sections also showed a nice colocalization of Cre staining with Tuj1-positive RGCs (Supplementary Figure 1B). We also injected AAV2-Cre into tdTomato reporter mice to examine the efficiency of Cre-mediated gene recombination. Strong expression of tdTomato in RGCs was observed two weeks after AAV2-Cre injection (Supplementary Figure 2A, B), indicating successful gene recombination. Lastly, we examined if myosin IIA/B were indeed deleted in RGCs after AAV2-Cre injection. By western blot analysis of the whole retina tissue, we found that the protein level of myosin IIA was markedly reduced (Supplementary Figure 3A). Immunostaining of retinal

sections with anti-myosin IIB antibody showed significantly reduced level of myosin IIB in RGCs (Supplementary Figure 3B, C). Together, these results demonstrated clearly that myosin IIA/B were successfully deleted in RGCs.

To determine how myosin IIA/B double knockout (dKO) in RGCs affected optic nerve regeneration, we performed optic nerve crush (ONC) 2 weeks after the viral injection. We first assessed optic nerve regeneration 2 weeks after the ONC. The regenerating RGC axons were labeled with anterogradely transported cholera toxin subunit B (CTB) conjugated with Alexa Fluor 594, which was injected into the vitreous humor 2 days prior to tissue harvest (Figure 1A). The fixed optic nerves were first tissue cleared to be transparent as previously described[13] and axon regeneration was imaged with confocal microscopy. The results showed that very limited optic nerve regeneration occurred in wild type mice infected with AAV2-Cre. In contrast, there was greatly enhanced optic nerve regeneration in myosin IIA/B dKO mice (Figure 1A, B). In the majority of optic nerves, regenerating axons reached 750 μm from the crush site. To further determine if myosin IIA/B dKO led to sustained promoting effect on optic nerve regeneration, we assessed optic nerve regeneration 4 weeks after the ONC. We found that at 4 weeks post ONC myosin IIA/B dKO significantly increased not only the lengths but also the number of regenerating axons, indicating continued axon regeneration along time. Specifically, most optic nerves had regenerating axons reaching 2000 μm from the crush site (Figure 1B). In addition to counting axon numbers at different distances from the crush site, we also quantified the lengths of the top 5 longest regenerating axons in each condition. The result showed that in control condition the lengths of the top 5 axons remained unchanged at 2 and 4 weeks, whereas the top 5 axons of myosin IIA/B dKO RGCs continued to grow from 2 to 4 weeks (Figure 1C). Both quantification results demonstrated that knocking out myosin IIA/B in RGCs led to sustained optic nerve regeneration at the same speed up to 4 weeks. When RGC survival rates were assessed, myosin IIA/B dKO showed no effect (Figure 1D, E), indicating that it promoted optic

nerve regeneration by enhancing axon extension rather than protecting RGCs from ONC-induced cell death. Collectively, these results demonstrated clearly that knocking out myosin IIA/B was sufficient to induce robust and sustained optic nerve regeneration without affecting cell survival.

Knocking out non-muscle myosin IIA/B acted synergistically with Lin28a overexpression to promote long-distance optic nerve regeneration

Our recent study showed that overexpression of Lin28 in RGCs induced robust and sustained optic nerve regeneration via regulation of gene expression and enhanced intrinsic axon regeneration ability[13]. Therefore, we tested if combining myosin IIA/B dKO, which alters axonal cytoskeletal dynamics, with Lin28 overexpression, which controls gene expression, could have combinatory promoting effect on optic nerve regeneration. The results showed that either myosin IIA/B dKO or Lin28a overexpression alone could lead to robust optic nerve regeneration 2 weeks after ONC (Figure 2A, B). When both treatments were combined, optic nerve regeneration was greatly enhanced, with the longest length reaching 3.5mm from the crush site (Figure 2A, B). In particular, either myosin IIA/B dKO or Lin28a overexpression RGCs had almost no regenerating axons growing beyond 1.75 mm from the crush site. In contrast, in the combinatory treatment group there were significant number of regenerating axons at 3 mm and the longest axons reached up to 3.5 mm (Figure 2A, B). Because the dehydration process in the tissue clearing approach results in 18% of shrinkage in nerve lengths[13], the real lengths of the longest regenerating axons were more than 4 mm. Indeed, in about half of nerves in the combinatory group the longest regenerating axons almost reached the optic chiasm (Figure 2A and Supplementary Figure 4) 2 weeks after ONC. This result suggested that myosin IIA/B dKO and Lin28a acted synergistically to promote long-distance optic nerve regeneration. Similarly, we quantified the average distances of the top 5 longest regenerating axons in each condition. The results showed that the longest regenerating axons in the combinatory group were

markedly longer than either the myosin IIA/B dKO or Lin28a overexpression group (Figure 2C). To better

show the regenerating axons in the tissue cleared transparent optic nerves, we created a 3D animation of

an optic nerve (Supplementary Video 1).

Knocking out non-muscle myosin IIA/B in RGCs did not significantly affect known optic nerve regeneration signaling pathways

In our previous study[20], we showed that treating adult sensory neurons cultured on CSPGs with

the myosin II inhibitor, blebbistatin, could induce halted axon to regrow within minutes. Conversely,

washing out the blebbistatin stopped axon growth in a very short time. Such rapid response to blebbistatin

in both ways indicated that inhibition of myosin II promoted axon growth by its direct effects on growth

cone cytoskeleton without affecting signaling events in the neuronal soma. To test this idea, we examined

how myosin IIA/B dKO affected two well-known pathways governing the intrinsic axon regeneration

ability, the activation of mTOR, marked by increased level of phospho-S6 (pS6), and the inactivation of

GSK3β, marked by phosphorylation of its serine 9 residue (pGSK3β), at 2 weeks after ONC. Previous

studies have shown that most identified molecules promoting optic nerve regeneration act via either of

these pathways, including Pten knockout[10], Akt overexpression[21,22], Lin28 overexpression[13], osteopontin

overexpression[23], SOCS3 deletion[12], melanopsin overexpression[24], HDAC5 manipulation[25], and direct

modulation of mTOR[26] or GSK3β signaling[21]. For mTOR activation, we examined the level of pS6 in

RGCs under different conditions. The results showed that knocking out myosin IIA/B had no effect on

pS6 level, whereas Lin28a overexpression markedly increased the level of pS6 in RGCs (Figure 3A).

Quantification demonstrated that the percentage of pS6 positive (pS6$^+$) RGCs increased by nearly 8 folds

in the Lin28a overexpression group compared to that in the wild type group, whereas myosin IIA/B dKO

had no effect (Fig. 3B). To provide a more objective measurement of pS6 level in RGCs, we also

quantified the average fluorescence intensity of pS6 staining in all Tuj1 positive RGCs. The results showed that myosin IIA/B dKO actually slightly but significantly reduced the pS6 level compared to that of the wild type group. In contrast, Lin28a overexpression greatly increased the pS6 level (Figure 3C). Lastly, we quantified the average fluorescence intensity of pS6 staining only in pS6$^+$ RGCs under different conditions. Similarly, there was no significant difference between wild type and myosin IIA/B dKO RGCs, whereas the Lin28a overexpression group had a much higher value (Figure 3D).

For GSK3β inactivation, we found that there was very few wild type RGCs showing positive staining of pGSK3β, and knocking out myosin IIA/B had no impact on it (Figure 3E, F). In contrast, Lin28a overexpression increased the percentage of pGSK3β positive (pGSK3β$^+$) RGCs by about 6 folds (Figure 3F). The average fluorescence intensity of pGSK3β staining in all Tuj1 positive RGCs was not affected by myosin IIA/B dKO, whereas Lin28a overexpression greatly increased the pGSK3β level (Figure 3G). When the average fluorescence intensity of pGSK3β were quantified only in pGSK3β$^+$ RGCs, the level of pGSK3β was significantly decreased in myosin IIA/B dKO group compared with that in the wild type group. The level of pGSK3β in the Lin28a overexpression group was still the highest (Figure 3H).

Taken together, these results provided clear and strong evidence that knocking out myosin IIA/B had no effects on two well-known signaling pathways occurred in the neuronal soma supporting intrinsic regenerative ability of optic nerves.

Knocking out non-muscle myosin IIA/B in sensory neurons did not significantly affect known regeneration associated genes and pathways

To further explore the potential molecular mechanisms underlying myosin IIA/B dKO induced axon regeneration, we performed sham surgery or bilateral sciatic nerve injury (SNI) on *myosin IIA/B$^{f/f}$*

and *Advillin-Cre: myosin IIA/B^f/f* mice, in which myosin IIA/B were conditionally knocked out in sensory

neurons. Three days later, we collected lumbar 4 and 5 dorsal root ganglia (L4/5 DRGs) and isolated

mRNA and protein for RNA-seq and western blot analyses. As expected, mRNA levels of myosin IIA/B

in L4/5 DRGs of *Advillin-Cre: myosin IIA/B^f/f* mice were lowered (Figure 4A). Pearson correlation

coefficient of gene expression indicated by FPKM (Fragments Per Kilobase of exon model per Million

mapped reads) among different samples showed that under same treatment (sham or SNI), the two wild

type replicates were quite similar to the two dKO replicates (Figure 4B). When the numbers of

differentially expressed genes (DEGs) were compared, we found SNI significantly altered the mRNA

levels of nearly 2000 genes, whereas myosin IIA/B dKO caused little change. Specifically, the number of

DEGs between wild type neurons and myosin IIA/B dKO neurons was actually equivalent to that between

the two replicates within each condition (Figure 4C). These results indicated that myosin IIA/B dKO *per*

se did not significantly change the transcriptome in sensory neurons. In addition, in wild type and myosin

IIA/B dKO neurons we closely examined and compared the FPKMs of many classic regeneration-

associated genes (RAGs) and genes well-known to control axon regeneration, such as *Atf3*, *Sox11*, *Lin28a*,

Gap43, *Pten*, *Klf9*, and *Rab27*, etc[13,27-31]. The results showed that the mRNA levels of these genes were

up- or downregulated by SNI as expected, but were not largely affected by myosin IIA/B dKO (Figure

4D, E). Gene ontology analysis of DEGs between different conditions revealed that SNI stimulated many

neuron-related programs (Figure 4G), whereas myosin IIA/B dKO induced gene expression change had

no specific connection with neurons or axon regeneration (Figure 4H). Lastly, we directly examined the

protein levels of several genes and pathways regulating axon regeneration by western blot and found that

in either uninjured or injured condition, myosin IIA/B dKO did not change the protein levels of Atf3, c-

Jun, Gap43 or c-Myc (Figure 4F). Moreover, consistent with our RGC immunostaining results, myosin

IIA/B dKO had no impact on mTOR or GSK3β pathway (Figure 4F) in sensory neurons. Taken together,

these results further supported that myosin IIA/B dKO induced optic nerve regeneration was unlikely to

be caused by enhanced intrinsic axon regeneration ability.

Knocking out non-muscle myosin IIA/B in RGCs changed axon tip morphology and regenerating

axon trajectory

To better understand the cellular mechanisms by which myosin IIA/B knockout promoted optic

nerve regeneration, we first performed detailed analysis of axonal tip morphologies in wild type and

myosin IIA/B dKO optic nerves 2 and 4 weeks after ONC. Based on a previous study[32], there are mainly

three types of axonal tip morphologies in vivo. One is the retraction bulb (Figure 5A, C), which is the

hallmark structure of dystrophic axons that failed to regenerate[17,19]. The other two are growth-competent

growth cones with two different end shapes (Figure 5A, C). We found that in wild type optic nerves, a

significant percentage of axons had retraction bulbs at their ends, whereas knocking out myosin IIA/B in

RGCs almost abolished the formation of retraction bulbs (Figure 5A, B). Most regenerating axons in the

myosin IIA/B knockout nerves had growth-competent growth cones, indicating that deleting myosin IIA/B

efficiently transformed dystrophic axon tips into growth cones and rendered subsequent axon regeneration.

The tissue clearing and confocal imaging of whole-mount optic nerves allowed us to visualize the

bona fide morphology of regenerating axons. Thus, we next examined how myosin IIA/B knockout

influenced the axon extension trajectories 4 weeks after ONC. In wild type nerves, the majority of axons

followed a wandering path with many curves and backward turns (U-turns), which resulted in very

inefficient axon regeneration towards the distal optic nerve. In contrast, in myosin IIA/B dKO nerves,

most regenerating axons were straight with significantly reduced U-turns (Figure 5D-F and

Supplementary Figures 5, 6), indicating a higher efficiency of axon regeneration. Together, we think that

enhanced optic nerve regeneration induced by myosin IIA/B dKO was achieved through 1) switching

retraction bulbs into growth-competent growth cones, and 2) more efficient axon regeneration with

straighter axon growth and less U-turns.

Discussion

266
In addition to diminished intrinsic axon regeneration capacity regulated by changes in gene

expression during neuronal maturation, dystrophic growth cone with disruptive cytoskeletal dynamics is

another key intrinsic barrier for successful CNS axon regeneration[17]. Although it has been well recognized

that modulation of axonal cytoskeleton would be a plausible approach to enhance CNS axon regeneration,

very few studies have shown direct and convincing results. Two previous elegant studies[33,34] have shown

that moderate stabilization of microtubules with taxol or epothilone B could promote axon regeneration

after the spinal cord injury. The promoting effects were achieved through decreased glial scar formation,

which rendered the lesion site more permissive, and improved microtubule protrusion in the growth cones

of the injured axons. However, the promoting effects were moderate with regenerating axons only entering

the injury site. Similarly, after optic nerve injury low dose taxol treatment alone had little promoting effect

on axon regeneration [35]. In this study we provided clear and strong evidence that knocking out myosin

IIA/B in RGCs alone was sufficient to induce robust and sustained optic nerve regeneration. Based on our

previous in vitro study[20] we think that deleting myosin IIA/B acted locally at the growth cone without

affecting signaling events at the neuronal soma and gene transcription. Thus, we first examined two well-

known signaling pathways in RGCs, mTOR activation and GSK3β inactivation, which occur in the

neuronal soma to support the intrinsic axon regeneration ability. The results showed that deleting myosin

IIA/B had no effects on these two pathways, suggesting that the intrinsic axon regeneration ability of

RGCs might not be elevated. In contrast, overexpression of Lin28a, which promotes optic nerve

regeneration by enhancing the intrinsic axon regeneration ability, led to marked activation of mTOR and

inactivation of GSK3β in RGCs. Next, by using sensory neuron specific myosin IIA/B conditional

knockout mice, we explored how deleting myosin IIA/B affected sensory neuron transcriptome. The

results showed that deleting myosin IIA/B did not significantly change the transcription profile in sensory

neurons. More importantly, when the transcription levels of many known RAGs were examined, we found that peripheral axotomy significantly changed their mRNA levels, whereas knocking out myosin IIA/B had little effects. Lastly, we directly examined the protein levels of selected RAGs and signaling mediators with western blot. Similarly, knocking out myosin IIA/B had little effects, whereas peripheral axotomy showed significant impact. Collectively, these results provided strong and clear evidence that myosin IIA/B knockout promoted optic nerve regeneration without affecting gene transcription and the intrinsic axon regeneration ability.

To better understand how knocking out myosin IIA/B promoted optic nerve regeneration, we carefully examined how RGC axonal morphology and trajectory were affected. The inability of mature CNS axon to form a growth-competent growth cone in the inhibitory environment after injuries is a hallmark of regeneration failure[17,19]. In most cases, after CNS injuries, a bulb-like structure is formed at the tip of the injured axon, called the retraction bulb[32]. Indeed, very limited optic nerve regeneration was observed in wild type mice, and retraction bulbs were often observed at the tips of the axons. In contrast, in myosin IIA/B dKO nerves, very few retraction bulbs were identified, while growth-competent growth cones could be found at almost all axonal tips, indicating that deleting myosin IIA/B was an efficient strategy to transform retraction bulbs into growth cones. Based on our previous in vitro study[20], it is likely achieved through slowed retrograde flow of actin filaments and the subsequent protrusion of microtubules in the dystrophic growth cones. In addition, confocal microscopy of cleared whole-mount optic nerves allowed us to trace the trajectories of regenerating axons. We found that in the wild type group, the majority of axons showed wandering trajectories with many kinks and U-turns, likely due to the inhibitory substrates in the optic nerve. As a result, such axon extension resulted in very inefficient axon regeneration. When myosin IIA/B were knocked out, most axons followed a straighter path with reduced U-turns, indicating overcoming inhibitory cues upon myosin IIA/B deletion and greatly enhanced regeneration

efficiency. Collectively, these results suggested that deleting myosin IIA/B promoted optic nerve regeneration via efficiently transforming retraction bulbs into active growth cones and more efficient axon extension within the inhibitory CNS environment.

Long-distance axon regeneration is one of the most important aspects and a prerequisite for successful functional recovery after neural injuries. Several previous studies[36,37], including ours[13], have shown that combined manipulation of multiple genes/pathways usually have additive promoting effects on optic nerve regeneration. Here we showed that combining myosin IIA/B knockout with Lin28a overexpression resulted in surprisingly long-distance optic nerve regeneration. Two weeks after ONC, regenerating axon reached up to 3.25 mm from the crush site (nearly 4 mm in real distance considering 18% shrinkage due to tissue clearing process[13]) without enhancing the RGC survival rate. The longest regenerating axons were about 4.3 mm and close to the optic chiasm. To our knowledge, such distance was longer than most previous published studies using combinatory approaches. For instance, we recently showed that combining Lin28a overexpression with Pten knocking down in RGCs resulted in optic nerve regeneration up to 2.4 mm from the crush site 2 weeks after ONC[13]. In an earlier study[36], combined knockout of Pten and SOCS3, together with CNTF, led to optic nerve regeneration up to 3 mm from the crush site 2 weeks after ONC. Moreover, combining Zymosan, c-AMP, and Pten deletion could promote optic nerve regeneration to 3 mm from the crush site[37]. The effectiveness of the combinatory approach will be optimal if each gene/pathway acts independently. Indeed, Lin28a and myosin II acted in distinct neuronal compartments with distinct cellular mechanisms, resulting in an optimized combination. Although the number of long-distance regenerating axons observed in this study was relatively low, it was likely due to the poor survival of RGCs as neither myosin IIA/B knockout (Figure 1D, E) nor Lin28a overexpression[13] protects RGCs from dying after injury. Thus, future studies combining myosin II knockout with other regeneration approaches, as well as enhanced RGC survival, would very likely lead

335   to large number of axons regenerating back to their original targets in the brain and gain recovery of lost

336   visual function. Moreover, the availability of water soluble and stable pharmacological inhibitor of myosin

337   II[38] would make future translational applications possible to repair axonal injuries induced by glaucoma,

338   spinal cord injury, traumatic brain injury, and neurodegenerative diseases.

Methods

Mice

All animal experiments were conducted in accordance with the protocol approved by the Institutional Animal Care and Use Committee of the Johns Hopkins University. The *Myh9^f/f* (stock#032096-UNC) and *Myh10^f/f* (stock#016981-UNC) mouse strains were obtained from Mutant Mouse Resource and Research Center (MMRRC) at University of North Carolina at Chapel Hill, an NIH-funded strain repository, and were donated to the MMRRC by Robert S. Adelstein, M.D., National Heart, Lung, and Blood Institute (NHLBI). The two lines were crossed to generate *Myh9^f/f: Myh10^f/f* mice. *Advillin-Cre* mouse line was a kind gift from Dr. Fan Wang's laboratory at Duke University, and was crossed with *Myh9^f/f: Myh10^f/f* to get *Advillin-Cre: Myh9^f/f: Myh10^f/f* conditional knockout mice. The tdTomato reporter line (stock#007909) was purchased from The Jackson Laboratory. Adult mice (6 weeks) of both sexes were used. Genotypes of the mice were determined by PCR using primers provided by MMRRC and The Jackson Laboratory. All animal surgeries were performed under anesthesia induced by intraperitoneal injection of ketamine (100 mg/kg) and xylazine (10 mg/kg) diluted in sterile saline. Details of the surgeries are described below.

Construct

The pAAV-Ef1a-Lin28a-FLAG plasmid was constructed in a previous study[13]. Briefly, the Lin28a-FLAG open reading frame with a 5' BamHI and a 3' EcoRV restriction sites was synthesized (codon optimized, gBlocks of Integrated DNA Technologies) and used for replacing the EYFP open reading frame in pAAV-Ef1a-EYFP, to obtain the pAAV-Ef1a-Lin28a-FLAG plasmid. pAAV-Ef1a-EYFP was a kind gift from Dr. Hongjun Song's laboratory at University of Pennsylvania. All restriction

361 enzymes and T4 DNA ligase were purchased from New England Biolabs. Plasmids were amplified using

362 DH5α competent cells (Thermo Fisher Scientific) and purified with Endofree plasmid maxi kit (Qiagen).

Optic nerve regeneration model

Intravitreal viral injection, optic nerve crush and RGC axon labeling were performed as previously

described[10]. Briefly, under anesthesia, 1.5 μl of AAV2 virus was injected into the right vitreous humor of

a mouse with a Hamilton syringe (32-gauge needle). The position and direction of the injection were well-

controlled to avoid injury to the lens. Two weeks later, the right optic nerve of the mouse was exposed

intraorbitally and crushed with Dumont #5 fine forceps (Fine Science Tools) for 5 s at approximately 1

mm behind the optic disc. To label RGC axons in the optic nerve, 1.5 μl of Alexa Fluor 594-conjugated

CTB (2 μg/μl, Thermo Fisher Scientific) was injected into the right vitreous humor with a Hamilton

syringe (32-gauge needle) 2 days before the mouse was sacrificed by transcardial perfusion under

anesthesia. The right optic nerve and bilateral retinas were dissected out and post-fixed in 4% PFA

overnight at 4°C. AAV2-Cre (SL100813) was purchased from SignaGen Laboratories. AAV2-Lin28a-

FLAG was also packaged by SignaGen Laboratories. All viruses used had titers over 1 x 10^{13} gc/ml.

Optic nerve dehydration and clearing

378   Dehydration and clearing of optic nerves were done based on previous studies[15,39]. Briefly, fixed

379 optic nerves were first dehydrated in increasing concentrations of tetrahydrofuran (50%, 70%, 80%, 100%

380 and 100%, v/v % in distilled water, 20 min each, Sigma-Aldrich) and then cleared in a solution of benzyl

381 alcohol and benzyl benzoate (BABB, 1:2, Sigma-Aldrich). Incubations were done on an orbital shaker at

382 room temperature. The nerves were stored in BABB in the dark at room temperature.

Analysis of RGC axon regeneration

Tissue cleared whole-mount optic nerves were imaged with a 20x objective on a Zeiss 800 confocal microscope. For each optic nerve, Z-stack and tiling (10% overlap) functions were used to acquire stacked 2-μm-thick planes of the whole area of interest and the tiles were stitched.

To quantify the number of regenerating axons in each optic nerve, every 8 consecutive planes were Z-projected (maximum intensity) to generate a series of Z-projection images of 16-μm-thick optical sections. At each 250-μm interval from the crush site, the number of CTB-labeled axons was counted in each Z-projection image and summed over all optical sections.

To quantify the average length of top 5 longest axons of each optic nerve, all stitched 2-μm-thick planes were Z-projected (maximum intensity) to obtain a single Z-projection image of the nerve. Top 5 longest regenerating axons were manually traced from the axonal tips to the crush site using the Fiji software (NIH) to acquire the lengths of the axons.

Immunohistochemistry of whole-mount retinas

Fixed retinas were first radially cut into a petal shape (4 incisions) and blocked with PBST (1%) containing 10% goat serum for 1 hr at room temperature. The retinas were then sequentially stained with primary antibodies overnight at 4°C, and corresponding Alexa Fluor-conjugated secondary antibodies (1:500, Thermo Fisher Scientific) for 2 hr at room temperature. All antibodies were diluted with the blocking buffer. Following each antibody incubation, the retinas were washed with PBST (0.3%) for 4 times (15 min each). After the last wash, the retinas were mounted onto slides with Fluoroshield (Sigma-Aldrich). Fluorescent images of the flat-mounted retinas were acquired with a 20x objective on a Zeiss 800 confocal microscope.

Analysis of RGC transduction rate

To quantify RGC transduction rate, uninjured right retinas (no optic nerve crush) were taken from transcardially perfused *Myh9^{f/f}: Myh10^{f/f}* mice 2 weeks after intravitreal AAV2-Cre injection. The retinas were stained with mouse anti-tubulin β3 (Tuj1, 1:500, BioLegend) and rabbit anti-Cre recombinase (1:100, Cell Signaling Technology) antibodies following the steps mentioned above (see Immnnunohistochemistry of whole-mount retinas). Five to eight fields under 20x objective were randomly obtained from the peripheral regions of each flat-mounted retina. For each mouse, RGC transduction rate was calculated by dividing the total number of Cre$^+$/Tuj1$^+$ cells in all fields by the total number of Tuj1$^+$ cells in all fields. Only cells in the ganglion cell layer were counted.

Analysis of RGC survival rate

To quantify RGC survival rate, C57Bl6/J and *Myh9^{f/f}: Myh10^{f/f}* mice injected with AAV2-Cre were transcardially perfused 2 weeks after optic nerve crush and both retinas of each mouse were collected. The retinas were stained with mouse anti-tubulin β3 antibody (Tuj1, 1:500, BioLegend) following the steps mentioned above (see Immnnunohistochemistry of whole-mount retinas). Seven or eight fields under 20x objective were randomly taken from the peripheral regions of each flat-mounted retina. For each mouse, RGC survival rate was calculated by dividing the average number of Tuj1$^+$ cells in one field in the injured retina (right) by that in the uninjured retina (left). Only cells in the ganglion cell layer were counted.

Immunohistochemistry of retinal sections

Fixed retinas were sectioned with a cryostat (10 μm) and the retinal sections were warmed on a slide warmer at 37°C for 1 hr. Sections were rinsed once in PBS, soaked in 100°C citrate buffer (pH 6) for 15 min, let to cool in the buffer to room temperature and then washed twice (5 min each) in PBS. After

being blocked with PBST (0.3%) containing 10% goat serum at room temperature for 1 hr, the sections were stained with primary antibodies against target molecules overnight at 4 °C, followed by corresponding Alexa Fluor-conjugated secondary antibodies (1:500, Thermo Fisher Scientific) at room temperature for 1 hr. All antibodies were diluted with the blocking buffer. The sections were washed for 4 times (5, 5, 10, 10 min) with PBST (0.3%) following each antibody incubation and finally mounted with DAPI Fluoromount-G (SouthernBiotech). Fluorescent images of the retinal sections were taken with a CCD camera connected to a Zeiss inverted fluorescence microscope controlled by AxioVision software.

Analysis of myosin IIB level in RGCs

To analyze myosin IIB level in RGCs, both retinas (uninjured) of each mouse were taken from transcardially perfused $Myh9^{f/f}$: $Myh10^{f/f}$ mice 2 weeks after intravitreal AAV2-Cre injection and sectioned. The retinal sections were stained with mouse anti-tubulin β3 (Tuj1, 1:500, BioLegend) and rabbit anti-myosin IIB (1:100, Thermo Fisher Scientific) antibodies following the steps mentioned above (see Immnunohistochemistry of retinal sections).

To quantify the fluorescence intensity of myosin IIB in all RGCs, at least 7 non-adjacent retinal sections acquired with identical imaging configurations were analyzed for each retina. Fluorescence intensity was measured using the "outline spline" function of AxioVision and the background fluorescence intensity was subtracted.

Analysis of S6 and GSK3β phosphorylation

To analyze S6 and GSK3β phosphorylation in RGCs, C57Bl6/J and $Myh9^{f/f}$: $Myh10^{f/f}$ mice injected with AAV2-Cre, and $Myh9^{f/f}$: $Myh10^{f/f}$ mice injected with AAV2-Lin28a-FLAG were transcardially perfused 2 weeks after optic nerve crush and the right retina of each mouse was collected and sectioned.

The retinal sections were stained with mouse anti-tubulin β3 antibody (Tuj1, 1:500, BioLegend), and rabbit anti-pS6 Ser235/236 (1:200, Cell Signaling Technology) or rabbit anti-pGSK3β Ser9 (1:200, Cell Signaling Technology) antibody following the steps mentioned above (see Immnunohistochemistry of retinal sections).

To quantify the percentage of pS6$^+$ or pGSK3β$^+$ RGCs, at least 363 or 434 RGCs from at least 7 non-adjacent retinal sections from each mouse were analyzed. For each mouse, the percentage of pS6$^+$ or pGSK3β$^+$ RGCs was calculated by dividing the number of pS6$^+$/Tuj1$^+$ or pGSK3β$^+$/Tuj1$^+$ cells by the number of Tuj1$^+$ cells. Only cells in the ganglion cell layer were counted.

The relative fluorescence intensity of each RGC was calculated by dividing the fluorescence intensity of the RGC by that of its adjacent tissue. To quantify the fluorescence intensity of pS6 or pGSK3β in all RGCs, 20 retinal sections acquired with identical imaging configurations from at least 2 mice were analyzed for each group. To quantify the fluorescence intensity of pS6 or pGSK3β in S6-activated or GSK3β-inactivated RGCs, at least 40 RGCs with identical imaging configurations from at least 2 mice were analyzed for each group. Fluorescence intensity of RGCs were measured using the "outline spline" function of AxioVision.

Sciatic nerve injury model

Under anesthesia, bilateral sciatic nerves of a mouse were exposed and axotomized right below pelvis. Nerves were only exposed but not axotomized for a sham surgery. Three days after the surgery, the mouse was euthanized and bilateral L4/5 DRGs were collected and used for total RNA or protein extraction.

mRNA sequencing and data analysis

Total RNA was isolated with RNeasy mini kit (Qiagen) and RNA integrity was determined by

Agilent 2100 bioanalyzer and RNA 6000 Nano kit (Agilent Technologies). Paired-end libraries were

synthesized using the TruSeq RNA library preparation kit (Illumina). Briefly, mRNA molecules were

purified using oligo dT-attached magnetic beads. Following purification, the mRNA molecules were

fragmented into small pieces using divalent cations at 94°C for 8 min. The cleaved RNA fragments were

reversely transcribed into first strand cDNA using reverse transcriptase and random primers. This was

followed by second strand cDNA synthesis using DNA Polymerase I and RNase H. These cDNA

fragments then went through an end repair process, the addition of a single base, and the ligation of the

adapters. The products were then purified and enriched with PCR to obtain the final cDNA library.

Purified libraries were quantified by Qubit 2.0 fluorometer (Thermo Fisher Scientific) and validated by

Agilent 2100 bioanalyzer (Agilent Technologies) to confirm the insert size and calculate the mole

concentration. Cluster was generated by cBot with the library diluted to 10 pM and sequencing was done

on HiSeq 2500 (Illumina).

Sequencing raw reads were preprocessed by filtering out rRNA reads, sequencing adapters, short-

fragment reads and other low-quality reads using Seqtk (github.com/lh3/seqtk). Hisat2 (version 2.0.4)[40]

was used to map the cleaned reads to the mouse GRCm38.p4 (mm10) reference genome with two

mismatches. After genome mapping, Stringtie (version 1.3.0)[41,42] was run with a reference annotation to

generate FPKM (Fragments Per Kilobase of exon model per Million mapped reads) values for known

gene models[43]. Differentially expressed genes were identified using edgeR[44]. The P value significance

threshold in multiple tests was set by the false discovery rate (FDR)[45,46]. The cut-off for differentially

expressed genes was set as $P < 0.05$ and |log2fold change| > 1. For genes used for gene ontology (GO)

analyses, an additional cut-off of average FPKM > 1 in at least one condition was applied. GO analysis

was done using DAVID Bioinformatics Resources 6.8[47,48]. GO terms came from biological process, cellular component and molecular function categories.

Western blot analysis

Total protein was extracted from L4/5 DRGs or retinas using the RIPA buffer containing protease inhibitor cocktail (Sigma-Aldrich) and phosphatase inhibitor cocktail (Sigma-Aldrich). Identical amount of total protein from each condition was then separated by 4-12% gradient SDS-PAGE gel electrophoresis and transferred onto polyvinylidene fluoride membranes. After being blocked with TBST (1%) containing 5% blotting-grade blocker (Bio-Rad), the membranes were incubated overnight with primary antibodies against target molecules at 4 °C, followed by corresponding HRP-linked secondary antibodies (1:2000, Cell Signaling Technology) for 1 hr at room temperature. All antibodies were diluted with blocking buffer. The membranes were washed with TBST (1%) for four times (5, 5, 10, 10 min) after each antibody incubation. Rabbit primary antibodies against myosin IIA (1:1000), Gap43 (1:1000), c-Jun (1:1000), c-Myc (1:1000), pAkt Ser473 (1:2000), pGSK3β Ser9 (1:1000), pS6 Ser235/236 (1:2000) were purchased from Cell Signaling Technology. Mouse anti-Atf3 primary antibody (1:100) was from Santa Cruz Biotechnology. Mouse anti-β-actin primary antibody (1:5000) was from Sigma-Aldrich.

Analysis of axonal tip morphology

The method was derived from a previous study[32]. For each optic nerve, top 10 longest axons were first identified in the Z-projection image of the nerve. Then the maximum diameter of each axonal tip and the diameter of the cylindrical shaft of the corresponding axon were measured using Fiji software (NIH), and a tip/shaft ratio was calculated. An axonal tip was defined as a retraction bulb if its tip/shaft ratio was over 4. Otherwise, it was defined as a growth cone.

Analysis of axon extension efficiency

For each optic nerve, a 250-μm-long region with equivalent number of axons in the Z-projection image of the nerve was used for analysis. All traceable axons within this region were manually traced. For each axon, the length (covered distance on its trajectory) and the displacement (distance along the longitudinal axis of the nerve, sometimes could be zero or negative) between the start point and the end point were measured using Fiji software (NIH). The extension efficiency of each nerve was calculated by dividing the summed displacement by the summed length of all axons.

Analysis of U-turn rate

For each optic nerve, top 15 longest axons were identified in the Z-projection image of the nerve and their trajectories near the axonal tips were traced. A U-turn was defined when the angle between the final direction of the axonal tip and the positive longitudinal axis of the optic nerve was wider than 90 degrees. U-turn rate of each nerve was calculated by dividing the number of axons that made U-turns by 15.

Quantification and statistical analysis

Statistical analyses were done with GraphPad Prism 7 and the significance level was set as $P < 0.05$. Data are represented as mean ± SEM unless specifically stated. For comparisons between two groups, two-tailed unpaired or paired t test was used. For comparisons among three or more groups, one-way ANOVA followed by Tukey's multiple comparisons test was used to determine the statistical significance. Fisher's exact test was used to test contingency tables. All details regarding statistical analyses, including the tests used, P values, exact values of n, definitions of n, are described in figure legends.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request. RNA-seq raw data will be deposited and accession code will be available before publication.

Acknowledgements: We appreciate Dr. Michele Pucak's help in confocal imaging experiments and analyses. The study was supported by grants (to F.Q.Z.) from NIH (R01NS064288, R01NS085176, R01GM111514, R01EY027347), the Craig H. Neilsen Foundation (259450), and the BrightFocus Foundation (G2017037). Saijilafu was supported by the grants from the National Natural Science Foundation of China (81571189, 81772353), the National Key Research and Development Program (2016YFC1100203), and Innovation and Entrepreneurship Program of Jiangsu Province.

Author Contributions: X.-W.W., S.-G.Y., S., and F.-Q.Z. conceived the study and designed the project; X.-W.W. and S.-G.Y. performed most of the experiments; C.Z. performed and analyzed the immunostaining experiments; Y.Z. analyzed optic nerve regeneration; C.Z. and Y.Z. analyzed the axon morphologies and trajectories; J.-J.M. performed the sensory neuron RNA-seq experiment; Y.-L.W. and G.-L.M. helped with the optic nerve regeneration experiments; B.-B.Y., and A.R.K. helped with the data analyses; X.-W.W. and F.-Q.Z. wrote the manuscript with contribution from all authors.

Competing interests: The authors declare no competing interests.

Reference:

He, Z. & Jin, Y. Intrinsic Control of Axon Regeneration. *Neuron* **90**, 437-451, doi:10.1016/j.neuron.2016.04.022 (2016).

Curcio, M. & Bradke, F. Axon Regeneration in the Central Nervous System: Facing the Challenges from the Inside. *Annu Rev Cell Dev Biol* **34**, 495-521, doi:10.1146/annurev-cellbio-100617-062508 (2018).

David, S. & Aguayo, A. J. Axonal elongation into peripheral nervous system "bridges" after central nervous system injury in adult rats. *Science* **214**, 931-933 (1981).

Fawcett, J. W. The Paper that Restarted Modern Central Nervous System Axon Regeneration Research. *Trends Neurosci* **41**, 239-242, doi:10.1016/j.tins.2018.02.012 (2018).

Richardson, P. M., Mcguinness, U. M. & Aguayo, A. J. Axons from Cns Neurons Regenerate into Pns Grafts. *Nature* **284**, 264-265, doi:DOI 10.1038/284264a0 (1980).

So, K. F. & Aguayo, A. J. Lengthy regrowth of cut axons from ganglion cells after peripheral nerve transplantation into the retina of adult rats. *Brain Res* **328**, 349-354 (1985).

Geoffroy, C. G. & Zheng, B. Myelin-associated inhibitors in axonal growth after CNS injury. *Current opinion in neurobiology* **27**, 31-38, doi:10.1016/j.conb.2014.02.012 (2014).

Lee, J. K. & Zheng, B. Role of myelin-associated inhibitors in axonal repair after spinal cord injury. *Exp Neurol* **235**, 33-42, doi:10.1016/j.expneurol.2011.05.001 (2012).

Liu, K. *et al.* PTEN deletion enhances the regenerative ability of adult corticospinal neurons. *Nat Neurosci* **13**, 1075-1081, doi:10.1038/nn.2603 (2010).

Park, K. K. *et al.* Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway. *Science* **322**, 963-966, doi:10.1126/science.1161566 (2008).

Moore, D. L. *et al.* KLF family members regulate intrinsic axon regeneration ability. *Science* **326**, 298-301, doi:10.1126/science.1175737 (2009).

Smith, P. D. *et al.* SOCS3 Deletion Promotes Optic Nerve Regeneration In Vivo. *Neuron* **64**, 617-623, doi:10.1016/j.neuron.2009.11.021 (2009).

Wang, X. W. *et al.* Lin28 Signaling Supports Mammalian PNS and CNS Axon Regeneration. *Cell Rep* **24**, 2540-2552 e2546, doi:10.1016/j.celrep.2018.07.105 (2018).

Geoffroy, C. G., Hilton, B. J., Tetzlaff, W. & Zheng, B. Evidence for an Age-Dependent Decline in Axon Regeneration in the Adult Mammalian Central Nervous System. *Cell Rep* **15**, 238-246, doi:10.1016/j.celrep.2016.03.028 (2016).

Luo, X. *et al.* Three-dimensional evaluation of retinal ganglion cell axon regeneration and pathfinding in whole mouse tissue after injury. *Exp Neurol* **247**, 653-662, doi:10.1016/j.expneurol.2013.03.001 (2013).

Pernet, V. *et al.* Misguidance and modulation of axonal regeneration by Stat3 and Rho/ROCK signaling in the transparent optic nerve. *Cell Death Dis* **4**, e734, doi:10.1038/cddis.2013.266 (2013).

17 Blanquie, O. & Bradke, F. Cytoskeleton dynamics in axon regeneration. *Current opinion in neurobiology*
51, 60-69, doi:10.1016/j.conb.2018.02.024 (2018).

18 Hur, E. M. *et al.* GSK3 controls axon growth via CLASP-mediated regulation of growth cone microtubules.
Genes Dev **25**, 1968-1981, doi:10.1101/gad.17015911 (2011).

19 Hur, E. M., Saijilafu & Zhou, F. Q. Growing the growth cone: remodeling the cytoskeleton to promote axon
regeneration. *Trends in Neurosciences* **35**, 164-174, doi:10.1016/j.tins.2011.11.002 (2012).

20 Hur, E. M. *et al.* Engineering neuronal growth cones to promote axon regeneration over inhibitory
molecules. *Proc Natl Acad Sci U S A* **108**, 5057-5062, doi:10.1073/pnas.1011258108 (2011).

21 Guo, X., Snider, W. D. & Chen, B. GSK3beta regulates AKT-induced central nervous system axon
regeneration via an eIF2Bepsilon-dependent, mTORC1-independent pathway. *Elife* **5**, e11903,
doi:10.7554/eLife.11903 (2016).

22 Miao, L. *et al.* mTORC1 is necessary but mTORC2 and GSK3beta are inhibitory for AKT3-induced axon
regeneration in the central nervous system. *Elife* **5**, doi:10.7554/eLife.14908 (2016).

23 Duan, X. *et al.* Subtype-specific regeneration of retinal ganglion cells following axotomy: effects of
osteopontin and mTOR signaling. *Neuron* **85**, 1244-1256, doi:10.1016/j.neuron.2015.02.017 (2015).

24 Li, S. *et al.* Promoting axon regeneration in the adult CNS by modulation of the melanopsin/GPCR
signaling. *Proc Natl Acad Sci U S A* **113**, 1937-1942, doi:10.1073/pnas.1523645113 (2016).

25 Pita-Thomas, W., Mahar, M., Joshi, A., Gan, D. & Cavalli, V. HDAC5 promotes optic nerve regeneration
by activating the mTOR pathway. *Exp Neurol* **317**, 271-283, doi:10.1016/j.expneurol.2019.03.011 (2019).

26 Lim, J. H. *et al.* Neural activity promotes long-distance, target-specific regeneration of adult retinal axons.
Nat Neurosci **19**, 1073-1084, doi:10.1038/nn.4340 (2016).

27 Tsujino, H. *et al.* Activating transcription factor 3 (ATF3) induction by axotomy in sensory and
motoneurons: A novel neuronal marker of nerve injury. *Mol Cell Neurosci* **15**, 170-182, doi:DOI
10.1006/mcne.1999.0814 (2000).

28 Tanabe, K., Bonilla, I., Winkles, J. A. & Strittmatter, S. M. Fibroblast growth factor-inducible-14 is induced
in axotomized neurons and promotes neurite outgrowth. *Journal of Neuroscience* **23**, 9675-9686 (2003).

29 Christie, K. J., Webber, C. A., Martinez, J. A., Singh, B. & Zochodne, D. W. PTEN Inhibition to Facilitate
Intrinsic Regenerative Outgrowth of Adult Peripheral Axons. *Journal of Neuroscience* **30**, 9306-9315,
doi:10.1523/Jneurosci.6271-09.2010 (2010).

30 Apara, A. *et al.* KLF9 and JNK3 Interact to Suppress Axon Regeneration in the Adult CNS. *Journal of
Neuroscience* **37**, 9632-9644, doi:10.1523/Jneurosci.0643-16.2017 (2017).

31 Sekine, Y. *et al.* Functional Genome-wide Screen Identifies Pathways Restricting Central Nervous System
Axonal Regeneration. *Cell Reports* **23**, 415-428, doi:10.1016/j.celrep.2018.03.058 (2018).

32 Erturk, A., Hellal, F., Enes, J. & Bradke, F. Disorganized microtubules underlie the formation of retraction
bulbs and the failure of axonal regeneration. *J Neurosci* **27**, 9169-9180, doi:10.1523/JNEUROSCI.0612-
07.2007 (2007).

33   Hellal, F. *et al.* Microtubule Stabilization Reduces Scarring and Causes Axon Regeneration After Spinal
Cord Injury. *Science* **331**, 928-931, doi:10.1126/science.1201148 (2011).

34   Ruschel, J. *et al.* Axonal regeneration. Systemic administration of epothilone B promotes axon regeneration
after spinal cord injury. *Science* **348**, 347-352, doi:10.1126/science.aaa2958 (2015).

35   Sengottuvel, V., Leibinger, M., Pfreimer, M., Andreadaki, A. & Fischer, D. Taxol facilitates axon
regeneration in the mature CNS. *J Neurosci* **31**, 2688-2699, doi:10.1523/JNEUROSCI.4885-10.2011
(2011).

36   Sun, F. *et al.* Sustained axon regeneration induced by co-deletion of PTEN and SOCS3. *Nature* **480**, 372-
375, doi:10.1038/nature10594 (2011).

37   Kurimoto, T. *et al.* Long-distance axon regeneration in the mature optic nerve: contributions of
oncomodulin, cAMP, and pten gene deletion. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **30**, 15654-15663, doi:10.1523/JNEUROSCI.4340-10.2010 (2010).

38   Varkuti, B. H. *et al.* A highly soluble, non-phototoxic, non-fluorescent blebbistatin derivative. *Sci Rep* **6**,
26141, doi:10.1038/srep26141 (2016).

39   Erturk, A. *et al.* Three-dimensional imaging of solvent-cleared organs using 3DISCO. *Nat Protoc* **7**, 1983-
1995, doi:10.1038/nprot.2012.119 (2012).

40   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory requirements.
Nat Methods **12**, 357-360, doi:10.1038/nmeth.3317 (2015).

41   Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat*
Biotechnol **33**, 290-295, doi:10.1038/nbt.3122 (2015).

42   Pertea, M., Kim, D., Pertea, G. M., Leek, J. T. & Salzberg, S. L. Transcript-level expression analysis of
RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc* **11**, 1650-1667,
doi:10.1038/nprot.2016.095 (2016).

43   Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying mammalian
transcriptomes by RNA-Seq. *Nat Methods* **5**, 621-628, doi:10.1038/nmeth.1226 (2008).

44   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
expression   analysis   of   digital   gene   expression   data.   *Bioinformatics*   **26**,   139-140,
doi:10.1093/bioinformatics/btp616 (2010).

45   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate - a Practical and Powerful Approach
to Multiple Testing. *J R Stat Soc B* **57**, 289-300 (1995).

46   Benjamini, Y. & Yekutieli, D. The control of the false discovery rate in multiple testing under dependency.
Ann Stat **29**, 1165-1188 (2001).

47   Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene lists
using DAVID bioinformatics resources. *Nature Protocols* **4**, 44-57, doi:10.1038/nprot.2008.211 (2009).

48   Huang, D. W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths toward the
comprehensive functional analysis of large gene lists. *Nucleic Acids Research* **37**, 1-13,
doi:10.1093/nar/gkn923 (2009).

Figure 1

Figure 1. Deletion of myosin IIA/B in RGCs induced robust and sustained optic nerve regeneration.

(A) Top: experimental timeline. Bottom: representative images of optic nerves showing that deletion of myosin IIA/B in RGCs produced robust and persistent axon regeneration 2 and 4 weeks after optic nerve crush. The columns on the right display magnified images of the areas in white dashed boxes on the left, showing axons at 250, 500, 750, 1250 and 2000 μm distal to the crush sites. The yellow line indicates the crush sites. Yellow arrows indicate top 3 longest axons of each nerve. Scale bar, 100 μm (50 μm for the magnified images).

(B) Quantification of optic nerve regeneration in (A) (one-way ANOVA followed by Tukey's multiple comparisons test, $P < 0.0001$ at 250, 500, 750, 1000, 1250 and 1500 μm, $P = 0.0002$, 0.0071, 0.3875 and 0.3875 at 1750, 2000, 2250 and 2500 μm, respectively; n = 10 mice in 4-week WT group, n = 9 mice in other groups).

(C) Quantification of average length of top 5 longest axons of each nerve in (A) (one-way ANOVA followed by Tukey's multiple comparisons test, $P < 0.0001$; n = 10 mice in 4-week WT group, n = 9 mice in other groups).

(D) Representative images of flat-mounted retinas showing that deletion of myosin IIA/B had no effect on RGC survival rate 2 weeks after optic nerve crush. Flat-mounted retinas were stained with anti-tubulin β3 antibody (Tuj1, green). Scale bar, 50 μm.

(E) Quantification of RGC survival rate in (D) (unpaired t test, $P = 0.9092$, n = 4 and 3 mice in WT and dKO groups, respectively, 7-8 fields were analyzed for each retina).

Data are represented as mean ± SEM. n.s., not significant, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$. WT, wild type; dKO, double knockout of myosin IIA/B.

Figure 2

Figure 2. Myosin IIA/B deletion and Lin28a overexpression had synergistic effect on optic nerve regeneration.

(A) Top: experimental timeline. Bottom: representative images of optic nerves showing that combined myosin IIA/B deletion and Lin28a overexpression in RGCs produced much stronger axon regeneration 2 weeks after the optic nerve crush. The columns on the right display magnified images of the areas in white dashed boxes on the left, showing axons at 250, 750, 1000, 1500, 2000 and 3000 μm distal to the crush sites. The yellow line indicates the crush sites. Yellow arrows indicate top 3 longest axons of each nerve. Scale bar, 100 μm (50 μm for the magnified images).

(B) Quantification of optic nerve regeneration in (A) (one-way ANOVA followed by Tukey's multiple comparisons test, $P < 0.0001$ at 250, 500, 750, 1000, 1250, 1500, 1750 and 2000 μm, $P = 0.0004$, 0.0206, 0.0092, 0.0042, 0.0026, 0.0844 at 2250, 2500, 2750, 3000, 3250 and 3500 μm, respectively; n = 9 mice in Lin28a O/E group, n = 8 mice in other groups).

(C) Quantification of average length of top 5 longest axons of each nerve in (A) (one-way ANOVA followed by Tukey's multiple comparisons test, $P < 0.0001$; n = 9 mice in Lin28a O/E group, n = 8 mice in other groups).

Data are represented as mean ± SEM. n.s., not significant, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. WT, wild type; dKO, double knockout of myosin IIA/B; O/E, overexpression.

Figure 3

Figure 3. Myosin IIA/B deletion-induced optic nerve regeneration was independent of mTOR or GSK3β pathway.

(A) Representative images of retinal sections showing that deletion of myosin IIA/B did not activate mTOR (marked by pS6) in RGCs, whereas Lin28a overexpression markedly activated mTOR in RGCs 2 weeks after optic nerve crush. The two columns on the right display magnified images of the RGCs marked in white dashed boxes on the left. Retinal sections were stained with anti-pS6 (green) and anti-tubulin β3 (magenta) antibodies. Scale bar, 50 μm (12.5 μm for the magnified images).

(B) Quantification of the percentage of pS6$^+$ RGCs in (A) (one-way ANOVA followed by Tukey's multiple comparisons, $P < 0.0001$, n = 3 mice in WT group, n = 5 mice in dKO group, n = 4 mice in Lin28a O/E group, at least 363 RGCs from at least 7 non-adjacent retinal sections were analyzed for each mouse).

(C) Quantification of average fluorescence intensity of pS6 in all RGCs (one-way ANOVA followed by Tukey's multiple comparisons, $P < 0.0001$, n = 20 retinal sections with identical imaging configurations from at least 2 mice were analyzed for each group).

(D) Quantification of average fluorescence intensity of pS6 in pS6$^+$ RGCs (one-way ANOVA followed by Tukey's multiple comparisons, $P < 0.0001$, n = 42, 49 and 53 RGCs with identical imaging configurations from at least 2 mice were analyzed for WT, dKO and Lin28a O/E groups, respectively).

(E) Representative images of retinal sections showing that deletion of myosin IIA/B did not inactivate GSK3β (marked by pGSK3β) in RGCs, whereas Lin28a overexpression markedly inactivated GSK3β in RGCs 2 weeks after optic nerve crush. The right two columns display magnified images of the RGCs marked in white dashed boxes on the left. Retinal sections were stained with anti-pGSK3β (green) and anti-tubulin β3 (magenta) antibodies. Scale bar, 50 μm (12.5 μm for the magnified images).

(F) Quantification of the percentage of pGSK3β$^+$ RGCs in (E) (one-way ANOVA followed by Tukey's multiple comparisons, $P < 0.0001$, n = 3 mice in WT group, n = 4 mice in other groups, at least 434 RGCs from at least 7 non-adjacent retinal sections were analyzed for each mouse).

(G) Quantification of average fluorescence intensity of pGSK3β in all RGCs (one-way ANOVA followed by Tukey's multiple comparisons, $P < 0.0001$, n = 20 retinal sections with identical imaging configurations from at least 2 mice were analyzed for each group).

(H) Quantification of average fluorescence intensity of pGSK3β in pGSK3β$^+$ RGCs (one-way ANOVA followed by Tukey's multiple comparisons, $P < 0.0001$, n = 54, 40 and 65 RGCs with identical imaging configurations from at least 2 mice were analyzed for WT, dKO and Lin28a O/E groups, respectively).

Data are represented as mean ± SEM. n.s., not significant, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. WT, wild type; dKO, double knockout of myosin IIA/B; O/E, overexpression.

Figure 4

Figure 4. Conditional knockout of myosin IIA/B in sensory neurons did not significantly affect known regeneration associated genes and pathways

(A) Decreased mRNA levels of myosin IIA/B in L4/5 DRGs of *Advillin-Cre: myosin IIA/B^{f/f}* mice (n = 2 mice in each group, data are represented as mean ± SEM).

(B) Pearson correlation heatmap showing the high degree of similarity in gene transcription between wild type and myosin IIA/B deleted neurons under same condition (sham or SNI).

(C) Number of differentially expressed genes between replicates or conditions showing that myosin IIA/B deletion did not significantly change gene expression in neurons.

(D, E) Expression heatmap of genes known to be upregulated (D) or downregulated (E) following SNI. Results showed that myosin IIA/B deletion did not affect the transcription of these genes in neurons.

(F) Representative western blot results showing that neuron specific myosin IIA/B conditional knockout did not affect classic regeneration associated genes or pathways (n = 2 independent experiments).

(G, H) Gene ontology analyses of differentially expressed genes between uninjured (sham) and injured (SNI) condition in wild type neurons (G), and between wild type neurons and myosin IIA/B deleted neurons under uninjured (sham) condition (H).

WT, wild type. dKO, double knockout of myosin IIA/B. SNI, sciatic nerve injury.

Figure 5

A

WT 2 weeks | dKO 2 weeks

WT 4 weeks | dKO 4 weeks

B

■ Growth cone ■ Retraction bulb

19 (15.8%) 2 (2%) 8 (13.3%) 1 (1.7%)

% of axonal tips

WT 2 weeks · dKO 2 weeks · WT 4 weeks · dKO 4 weeks

*** *

C

Retraction bulb

Growth cone I

Growth cone II

D

Regeneration efficiency (%)

WT 4 weeks · dKO 4 weeks

**

E

U turn rate (%)

WT 4 weeks · dKO 4 weeks

*

F

WT

dKO

$$\text{extention efficiency} = \frac{\sum d}{\sum s}$$

Figure 5. Myosin IIA/B deletion modified cytoskeletal dynamics to promote axon regeneration.

(A) Representative images of optic nerves showing that deletion of myosin IIA/B in RGCs abolished formation of retraction bulbs in optic nerves 2 and 4 weeks after optic nerve crush. Yellow arrows indicate retraction bulbs. Scale bar, 50 μm.

(B) Quantification of retraction bulbs in (A) (Fisher's exact test, $P = 0.0004$ and 0.0322 for 2 weeks and 4 weeks after optic nerve crush, respectively; n = 120 and 100 axonal tips from 12 nerves in 2-week WT and 10 nerves in dKO groups, respectively, n = 60 axonal tips from 6 nerves in each 4-week group).

(C) Representative images of retraction bulbs and growth cones found in different optic nerves. Scale bar, 5 μm.

(D) Quantification of axon extension efficiency in (F) and Figure S5 (unpaired t test, $P = 0.0032$, n = 6 mice in each group, at least 35 axons were analyzed for each mouse).

(E) Quantification of U-turn rate in Figure S6 (unpaired t test, $P = 0.0210$, n = 6 mice in each group, top 15 longest axons were analyzed for each mouse).

(F) Left: representative images of optic nerves showing that deletion of myosin IIA/B in RGCs improved axon extension efficiency 4 weeks after optic nerve crush. Middle: sketches of all axon traces in the left column. Right: detailed trajectories of a few axons (each color represents a single axon) in the left column. As illustrated, the extension efficiency of each nerve was calculated by dividing the summed displacement by the summed length of all traced axons. Scale bar, 50 μm.

Data are represented as mean ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. WT, wild type; dKO, double knockout of myosin IIA/B.

Supplementary Figure 1. Transduction rate of AAV2-Cre. Related to Figure 1.

(A) Representative images of flat-mounted retinas showing the high transduction rate of AAV2-Cre in RGCs. Flat-mounted retinas were stained with anti-tubulin β3 (Tuj1, green) and anti-Cre recombinase (red) antibodies. Scale bar, 50 μm.

(B) Representative images of retinal sections verifying the specificity of the anti-Cre antibody and the high transduction rate of AAV2-Cre in RGCs. Retinal sections were stained with anti-tubulin β3 (Tuj1, green) and anti-Cre (red) antibodies. Scale bar, 50 μm.

(C) Quantification of the transduction rate of AAV2-Cre in RGCs in (A). The average transduction rate is $90.59 \pm 1.114\%$ (n = 3 mice, 5-8 fields were analyzed for each mouse, data are represented as mean $\pm$ SD).

Supplementary Figure 2

A

B

Tuj1   tdTomato   Merge

Supplementary Figure 2. Verification of the Cre-mediated gene recombination. Related to Figure 1.

(A, B) Representative images of flat-mounted retinas (A) and retinal sections (B) showing the expression of tdTomato in RGCs of tdTomato reporter mice, indicating successful Cre-mediated gene recombination 2 weeks after intravitreal AAV2-Cre injection. Flat-mounted retinas and retinal sections were stained with anti-tubulin β3 (Tuj1, green). Scale bar, 50 μm.

Supplementary Figure 3. Verification of myosin IIA/B deletion. Related to Figure 1.

(A) Western blot result showing the decreased protein level of myosin IIA in retina of a *myosin IIA/B^{f/f}* mouse 2 weeks after intraviteal AAV2-Cre injection.

(B) Quantification of average fluorescence intensity of myosin IIB in (C) (paired t test, $P = 0.0057$, n = 3 mice, data are represented as mean ± SEM, **$P < 0.01$, all images were taken with identical configurations).

(C) Representative images of retinal sections from the naïve (top) or the AAV2-Cre injected retina (bottom) of a *myosin IIA/B^{f/f}* mouse showing the successful deletion of myosin IIB in RGCs 2 weeks after intravitreal AAV2-Cre injection. The insets display magnified images of the RGCs marked in white dashed boxes. Retinal sections were stained with anti-tubulin β3 (Tuj1, red) and anti-myosin IIB (green) antibodies. Scale bar, 50 μm (25 μm for the magnified images).

Supplementary Figure 4. Combination of myosin IIA/B deletion and Lin28a overexpression greatly enhanced optic nerve regeneration up to the optic chiasm. Related to Figure 2.

Images of two optic nerves in the combinatory treatment group with longest axons reaching optic chiasm. Magnified images show detailed morphologies of axons in white dashed boxes. Scale bar, 200 μm (66.7 μm for the magnified images).

WT

dKO

Supplementary Figure 5. Deletion of myosin IIA/B in RGCs improved axon extension efficiency. Related to Figure 5.

Images of the regions used for the quantification of axon extension efficiency (one region from each optic nerve). Scale bar, 50 μm. WT, wild type; dKO, double knockout of myosin IIA/B.

Supplementary Figure 6. Deletion of myosin IIA/B reduced U-turn rate. Related to Figure 5.

Top: representative images of the leading-edge area of optic nerves showing that myosin IIA/B deletion reduced U-turn rate. Middle: axon trajectories near axonal tips, U-turns are labeled in yellow. Bottom: overlay of top and middle rows. Scale bar, 50 μm. WT, wild type; dKO, double knockout of myosin IIA/B.