Juvenile hormone interacts with multiple factors to modulate aggression and dominance in groups of orphan bumble bee (*Bombus terrestris*) workers

Atul Pandey[1], Uzi Motro[1], Guy Bloch[1]

[1] - Department of Ecology, Evolution and Behavior, The Hebrew University of Jerusalem, Israel

Corresponding author

Guy Bloch: guy.bloch@mail.huji.ac.il, Phone: +972-(0)26584320

Short title:
Juvenile hormone and dominance in bumble bee workers

Abstract:

Juvenile hormone (JH) is a key regulator of insect development and reproduction. Given that JH commonly affects adult insect fertility, it has been hypothesized to also regulate behaviors such as dominance and aggression that are associated with reproduction. We tested this hypothesis in the bumble bee *Bombus terrestris* for which JH has been shown to be the major gonadotropin. We used the allatoxin Precocene-I (P-I) to reduce hemolymph JH titers and replacement therapy with the natural JH to revert this effect. In small orphan groups of workers with similar body size but mixed treatment, P-I treated bees showed lower aggressiveness, oogenesis, and dominance rank compared with control and replacement therapy treated bees. In similar groups in which all bees were treated similarly, there was a clear dominance hierarchy, even in P-I and replacement therapy treatment groups in which the bees showed similar levels of ovarian activation. In a similar experiment in which bees differed in body size, larger bees were more likely to be dominant despite their similar JH treatment and ovarian state. In the last experiment, we show that JH manipulation does not affect dominance rank in groups that had already established a stable dominance hierarchy. These findings solve previous ambiguities concerning whether or not JH affects dominance in bumble bees. JH positively affects dominance, but bees with similar levels of JH can nevertheless establish dominance hierarchies. Thus, multiple factors including JH, body size, and previous experience affect dominance and aggression in social bumble bees.

1. Introduction

Juvenile hormone (JH) is a well-studied hormone named after its morphogenic functions in the regulation of insect metamorphosis. In adults of many, but not all, insects studied so far, JH functions as a gonadotropic hormone regulating fertility (Adams, 2009; De Loof et al., 2001; Riddiford, 2012). One of the best studied examples for a non-gonadotropin function in adult insects is the Western honeybee (*Apis mellifera*), in which JH does not affect adult female oogenesis but rather regulates age-related division of labor (Reviewed in Bloch et al., 2002; Hartfelder, 2000; Robinson and Vargo, 1997; Wegener et al., 2013). These findings for the honeybee contrast with evidence that JH retains its ancestral gonadotropic functions in species of bees and wasps that live in solitary lifestyle or in small and simple societies (Bell, 1973; Bloch et al., 2000a, 1996; Chen et al., 1979; Pan and Wyatt, 1971; Shorter and Tibbetts, 2009; Shpigler et al., 2014; Smith et al., 2013; Wasielewski et al., 2011). These observations led to hypotheses stating that the evolution of advanced sociality in bees was associated with modification in JH signaling pathways (Bloch et. al., 2002; Hartfelder K, 1998; Robinson and Vargo, 1997; West-Eberhard, 1996). There is also evidence consistent with the premise that JH lost its ancestral gonadotrophic functions along the evolution of advanced sociality in additional lineages such as ants and wasps (Giray et al., 2005; Lengyel et al., 2007; Norman and Hughes, 2016; O'Donnell and Jeanne, 1993; Penick et al., 2011; Shorter and Tibbetts, 2009). This association between modifications in JH functions and the evolution of sociality is commonly attributed to assumed fitness costs related to high JH titers, similar to the well-established costs of testosterone in vertebrates (Flatt and Kawecki, 2007; Rantala et al., 2003; Rodrigues and Flatt, 2016). Accordingly, the loss of JH gonadotropic functions in complex insect societies

is assumed to allow highly social insect queens to be exceptionally fertile over long

periods while escaping the hypothesized fitness cost of high JH levels (Pamminger

et al., 2016; Rodrigues and Flatt, 2016).

Bumble bees provide an excellent model system for studying proximate and

ultimate aspects of the relationships between JH signaling and sociality because

they are social, but their societies are in many aspects simpler than that of the highly

eusocial honeybees and ants. By contrast to honeybees, JH does not influence task

performance (Shpigler et al., 2016), and there is good evidence consistent with the

hypothesis that JH is the major gonadotropin in bumble bees. JH titers are positively

correlated with ovarian activity (Bloch et al., 2000a, 1996), and manipulations

increasing or decreasing JH levels, result in ovarian activation or inactivation,

respectively (Amsalem et al., 2014b; Röseler, 1977; Shpigler et al., 2014, 2010,

2016; Van Doorn, 1989). JH was also shown to regulate additional processes that

are associated with reproduction such as wax secretion and comb construction

(Shpigler et al., 2014). As expected for a gonadotropic hormone, JH treatment

upregulates the expression of the major yolk protein vitellogenin (Vg) in the fat body

and its protein levels in the hemolymph (Shpigler et al., 2014).

In vertebrates such as mammals, birds, and fish, the gonadotropic steroid

hormones commonly regulate behaviors related to reproduction such as courtship,

mating, aggression, and dominance (Brain, 1977; Nelson, 2005; Norris and Lopez,

2011). The relationships between gonadotropic hormones and behavior have

received significantly less attention in insects and other invertebrates. The current

study aims at testing the hypothesis that JH regulates agonistic behaviors underlying

dominance hierarchy establishment in the social bumble bee *Bombus terrestris*.

Dominance rank is typically achieved by means of overt aggression and agonistic interactions and is an important factor determining reproduction in many eusocial insects, including bumble bees (Andrade-Silva and Nascimento, 2015; Bloch et al., 1996; Duchateau and Velthuis, 1989; Geva et al., 2005; Monnin and Peeters, 1999; Roseler, 1991; Sasaki et al., 2016; Van Doorn, 1989; Van Doorn and Heringa, 1986). Earlier studies on JH, dominance, and aggression in bumble bees led to somewhat conflicting conclusions. In *B. terrestris*, the best-studied bumble bee, JH titers, and in-vitro biosynthesis rates are typically correlated with dominance rank; dominant queens and workers have active ovaries, large corpora allata (the JH producing glands) and high JH titers; dominant workers also typically show more overt aggression and threatening displays (Amsalem et al., 2014b; Amsalem and Hefetz, 2011; Bloch et al., 2010, 2000a; Larrere and Couillaud, 1993; Röseler, 1977; Van Doorn, 1989). However, in small groups of callow workers, a dominance hierarchy is established and aggression is typically highest during the first few days post pupal emergence, a period when hemolymph JH titers are still low. At later ages when the dominance hierarchy is already clear, some subordinate individuals, which show little aggression or dominance displays, may nevertheless have developed ovaries or high JH titers (Bloch et al., 2000a, 1996; Röseler, 1977; Van Doorn, 1989; Van Doorn and Heringa, 1986). A recent study in which circulating JH titers were reduced using the allatotoxin precocene-I (P-I), reported reduced aggressive behavior, but the interpretation of this finding is difficult because the effect could not be recovered by replacement therapy with JH-III (Amsalem et al., 2014b).

To thoroughly test the hypothesis that JH influences aggression and dominance, we manipulated circulating JH titers in callow orphan ("queenless") workers using a combination of JH reduction with P-I, and replacement therapy with JH-III, the

natural JH of bumble bees (Bloch et al., 2000, 1996). We tested the hypothesis that JH influence dominance and aggression in bumble bee in 4 complementary experiments: In Experiment 1, we studied queenless groups composed of same body size, same age workers, each subjected to a different JH treatment. This experiment directly tested whether a decrease or an increase in JH levels affect dominance rank. In Experiment 2, we studied similar groups composed of workers of different body size, but of similar age, and subjected to the same treatment. This experiment tested whether body size influences dominance hierarchy in bees with similar JH levels. In Experiment 3, we studied groups of workers with a similar body size, age and subjected to the same JH treatment. This experiment tested whether bees with similar JH levels and size nevertheless establish dominance hierarchies. In the last experiment (Experiment 4), we studied queenless groups which had already established a dominance hierarchy and tested whether treatment with P-I or JH can reduce or increase, the dominance rank of the most dominant or most subordinate individual, respectively. This experiment tested the interaction between JH levels and previous experience (i.e., dominance rank). Taken together, our results suggest that JH is one of several factors affecting aggressiveness and dominance behavior in bumble bees and can explain the inconsistent results in previous studies on JH and dominance in bumble bees.

2. Materials and Methods:

2.1. Bees

Bombus terrestris colonies with a queen, 5-10 workers and brood at various stages of development (typically 2–4 days post first worker emergence) were obtained from Polyam Pollination Services, Yad-Mordechai, Israel. Each colony was housed in a wooden nesting box (21 × 21 × 12 cm) and placed in an environmental

chamber (29 ± 1 °C; 55% ± 10% RH) in constant darkness at the Bee Research

Facility at the Edmond J. Safra campus of the Hebrew University of Jerusalem, Givat

Ram, Jerusalem. The top and front walls of the nest boxes were made of transparent

Plexiglass panels, and the other parts were made of wood. Commercial sugar syrup

(Polyam Pollination Services) and fresh pollen (collected by honey bees; mixed with

sugar syrup) were provided *ad libitum*. We painted each individual bee within the

treatment with xylene free silver color paint (Pilot-PL01735) on the thorax creating a

similar pattern across treatment groups. All treatments and behavioral observations

were performed under dim red light. Given that bumble bees are very sensitive to

substrate-borne vibrations, we paid special attention to minimize touching the cages

or substrate during treatments and observations.

2.2. Manipulating JH titers

Our protocols are based on those developed and validated by Shpigler et al.,

(2016) with the following modifications: (A) We kept the concentration of P-I constant

at 50µg/µl and varied the volume in which it was dissolved in order to adjust the total

amount applied to a bee to her body-size (see below). (B) In the second treatment,

we applied the JH-III or vehicle to the abdomen rather than to the thorax. This was

done in order to minimize possible mixing of the P-I and JH treatments, and reduce

the number of times the bee is anesthetized by chilling. (C) We used a higher and

somewhat different range of doses (200-260µg/bee compared to 160-210µg/bee). In

brief, we collected callow workers (<24 h after emergence from the pupa) and cold

anesthetized them in 50 ml tubes immersed in ice (~2 °C) until being immobile for 5–

10 min. The bees in the control groups *("Control")* in experiments 1, 2 & 3 were

handled and chilled on ice (20-25 minutes) similarly to bees from the other

experimental groups, but not treated with a drug or vehicle. **Sham-treated bees**

("Sham") were similarly chilled and then treated on successive days with the castor

oil and dimethylformamide (DMF) vehicles: On day 1 post-emergence, we applied

the castor oil with an amount (4.0-5.2µl/bee) corrected for body size as detailed in

Supplementary Table S1. On Day 2, in parallel to the JH-III treatments (see below),

we treated the Sham treatment bees with DMF (3.5µl/ bee, irrespective of body size).

All other treatments and handling conditions were kept similar to the other treatment

groups. **Precocene-I treatment ("P-I"):** Given that the CA is the only known source

of JH in insects (Riddiford, 2008), and that Precocnes cause atrophy of the corpora

allata glands (Haunerland and Bowers, 1985), we used Precocene-I (P-I; Sigma-

Aldrich, cat # 195855) to reduce circulating JH titers (for details see Shpigler et al.

2016). We suspended the P-I in castor oil (Sigma-Aldrich, cat # 259853) and applied

the solution to the dorsal part of the thorax. The P-I + castor oil mixture, which is

highly viscous, was thoroughly mixed with a pipette and vortexed at high speed for 2

minutes to better disperse the drug. The precise P-I dose was selected based on our

previous studies (Shpigler et al., 2016) and further ad hoc validations for each new

P-I batch that we used. These validations helped to assure a high survival rate and

effective manipulation of JH levels (Fig.1A). Based on these considerations, we used

200-260µg per bee which was delivered in 4.0-5.2µl vehicle volume (Table S1). The

volume was adjusted according to the bee body size such that the drug

concentration remains similar at 50µg/µl/bee to minimize density-dependent

adsorption of the drug. As an index for body size, we used the length of the front

wing marginal cell (Shpigler et al., 2013; Yerushalmi, 2006). The cold-anesthetized

and treated bees were placed on ice-chilled glass plates immobilized for ~10 minutes

after P-I treatments to allow the drugs to be absorbed, and minimize the incidents of

wiping-off, as this can lead to variation in treatment affectivity. **Replacement**

therapy (*"P-I+JH-III"*): we applied JH-III (Sigma-Aldrich, cat # J2000) dissolved in

3.5µl of Dimethylformamide (DMF, J.T Backers, cat # 7032) to bees that were

already treated with P-I on the day before. This one day gap between the treatments

was important for stress relief and efficacy of the two tandem treatments. We treated

each bee with an amount of 50µg/bee JH-III, which we selected based on

preliminary dose-response JH-III application studies and assessments of ovarian

state (Fig.1B). This dose was as effective as higher doses used in previous studies

with bumble bees in which 70 or 100µg/bee were applied on the thorax (Amsalem et

al., 2014; Shpigler et al., 2014, 2016). P-I treated bees were collected and

anesthetized by chilling on ice for 5-10 minutes. When the bee was motionless, we

topically applied JH-III dissolved in DMF (replacement therapy). The treated bees

were left immobilized on ice for additional ~10 min for efficient penetration and for

minimizing drug wiping off.

Figure 1: Dose-dependent effect of topical treatments with Precocene-I and Juvenile

hormone on worker ovarian development: (A) Treatment with Precocene-I (P-I). The

allatotoxin P-I was mixed with castor oil and amounts were adjusted according to the bee

body size (4.0-5.2µl/ bee; see Table S1 for details). (**B**) Replacement therapy with juvenile

hormone III (JH-III), the natural JH of bumble bees. In this experiment bees were first treated with P-I and then on the following day, with increasing amounts of JH-III (1.0-100µg/bee/3.5µl DMF). The sample size was 2-3 groups, each consisting of 4 worker bees (n=8-12 bees/ treatment). The box plots show the median (line) and mean (+), the box frame spans over the first to the third quartile. The whiskers depict the 5th/95th percentile, outliers are depicted with dots. The number above the bars show the percentage survival.

2.3. Behavioral observations

Focal orphan groups were observed twice a day during days 3-5 after the first treatment (a total of 120 min observations per group). Each observation day included a morning observation session between 09:00–11:00h, and an evening observation session between 16:00–18:00h. Each session lasted 20 minutes. Orphan ("*queenless*") *B. terrestris* workers at this age typically show a high level of agonistic behaviors and establish clear dominance hierarchies (Amsalem and Hefetz, 2010; Bloch et al., 1996; Geva et al., 2005; Van Doorn, 1989). Given the importance of agonistic interactions for both the establishment and maintenance of dominance hierarchies in bumble bees, we recorded threatening displays that include *"buzzing"*, and *"pumping"* as described in Bloch et al., (1996). Buzzing displays are characterized by fast, short wing vibrations of a worker facing another bee; "pumping" is performed by a bee standing and facing a nestmate bee while performing distinct dorso-ventral pumping movement with her abdomen (Bloch et al., 1996; Duchateau, 1989; Geva et al., 2005). Dominant bees typically switched between buzzing and pumping behaviors. "Overt aggression", which commonly follows threatening displays, includes darting and attacks directed towards another bee. To estimate the dominance of individual bees we used the Dominance Index (DI) that was calculated as in previous studies (Bloch et al., 1996; Geva et al., 2005; Van Doorn and Heringa, 1986). Briefly, the DI is the proportion of encounters

between each pair of bees in which the focal bee did not retract, out of the total encounters [1-(Retractions/Total encounters)]. Thus, we used the amount of overt aggression and threatening displays as indices for aggressiveness, and the DI as an index for dominance. The dominance rank of a bee depicts its DI relative to that of others in the same queenless group. In each group, the worker with the highest DI value was termed "*alpha*" (α), following by "*beta" (β)*, "*gamma" (γ)* and "*delta" (δ)* in descending DI order. Observers were blind to treatments. To ensure similar recording of focal behaviors across observers, all the observers were trained individually. The similarity in the behavioral recording was ensured by training sessions in which the instructor and trainee recorded behavioral observations in parallel. The observations were conducted under dim red light that bees do not see well. The observers entered the room carefully and avoided touching the tables on which the colonies were placed in order to avoid any movements or vibrations that can startle bumble bees. Observations were conducted silently and there were no visible signs that the bees were disturbed by the observations. The reported indices for each of these behaviors are the sum of total events recorded over 120 min of observations across 3 days (i.e. morning and evening observations on days 3, 4 and 5) and are presented as event per hour.

2.4. Assessment of ovarian state

At the end of each experiment, we stored the focal bees at -20^0C. To assess ovarian state, we fixed the bee on a wax filled dissecting plate under a stereomicroscope (Nikon SMZ645) and immersed it in honey bee saline (Huang et al., 1991). We cut three incisions through the lateral and ventral abdominal cuticle using fine scissors and then immersed the internal organs in saline. We gently removed the ovaries into a drop of saline on a microscope slide, and use the ocular

ruler to measure the length of the four largest terminal oocytes from both ovarioles.

We used the average oocyte length of the four largest terminal oocytes as an index

for ovarian state.

2.5. Experiment 1: The influence of JH on dominance and aggression in

groups of similar size, differently-treated queenless worker bees

We collected callow bees (<24hrs of age) from various source colonies,

determined their respective sizes, and assigned them randomly to one of the four

manipulations described in Section 2.2. Following the treatments on days 1 & 2, we

established groups on Day 3. Each orphan ("queenless") group of four included one

untreated (handling control) bee, one sham control bee, one P-I treated bee, and

one replacement therapy treated bee. The four bees in the same group were

selected to be of a similar body size. Bees were kept in isolation in a small Petri dish

(35 × 12 mm) during the first and second day of the experiment (Section 2.2). The

isolated bees were fed ad libitum with pollen cakes and sugar syrup and kept under

controlled environmental conditions (29 ± 1 °C; 55% ± 10% RH). This procedure

minimized the possible lateral transfer of treatment solutions (i.e., JH or P-I) across

differentially treated groupmate bees. We made this adjustment in protocol following

a preliminary experiment in which the ovaries of P-I treated bees were larger than

expected in mixed groups that were constituted immediately after the first treatment

(compare Fig. S1 and Fig. 2A). Each individual bee in this experiment was tagged

with a different numbered colored disk allowing individual identification. The number

tag (Opalith tags, Germany) was attached at the time of the second treatment. On

Day 3 bees were assigned in groups of four per cage. Behavioral observations and

oocyte measurements were performed as described in sections 2.3 & 2.4.

2.6. Experiment 2: The influence of JH on dominance and aggression in groups of different size, similarly-treated queenless worker bees

We treated callow bees and assigned them to queenless groups within 1hr of collection from the source colony. Worker bees in each group differed in body size – each bee was chosen from a different size class (as denoted in Table S1; Two way ANOVA, $F_{(Size\ range)\ 3,\ 76} = 546.8$, $p < 0.05$; $F_{(Treatment\ groups)\ 3,\ 76} = 0.54$, $p > 0.05$, partial $\eta2 = 0.99$). The experimental outline was different in several ways from Experiment 1: (1) All the bees within a treatment group were treated similarly; (2) groups were constituted of bees having four different body sizes (Table S1). Each focal bee was paint marked with a different color to allow individual identification. The individual paint marks were applied to the dorsal part of the thorax soon after applying JH-III to the bee abdomen on Day 2. At this day, the castor oil on the thorax was already absorbed. After drying up, the bees of all the groups were reassembled into their respective wooden cages. Behavioral observations and oocyte measurements were performed as described in sections 2.3 & 2.4. This experimental design allowed us to test the effects of both body size and JH manipulation on ovarian development and behavior.

2.7. Experiment 3: The influence of JH on dominance and aggression in groups of similar size, similarly-treated queenless worker bees

The experimental outline is similar to Exp. 2 other than that the bees in each group were of similar body size (one way ANOVA, $F_{3,\ 100} = 0.40$, $p > 0.05$, partial $\eta2 = 0.01$). This experiment allows us to test the effect of JH on aggression and dominance while controlling for the possibly confounding factors of size and age.

2.8. Experiment 4: The influence of JH on dominance and aggression in groups of same size bees in which dominance hierarchy has been already established

In contrast to the three previous experiments, here we treated bees that have already established a stable dominance hierarchy during the first four days following group constitution. We performed the first set of behavioral observations on days 3 & 4, before treating the bees. On the evening of Day 4, after determining the dominance rank within each group, we divided the groups into two sets. In one set, the α ranked individual was treated with P-I (500µg/5µl/ bee, henceforth termed "Experiment 4A"). In the other set of groups, we treated the δ rank individuals with JH-III (100µg/5µl/bee, henceforth termed "Experiment 4B"). We used here higher doses compared to the previous three experiments because the treated bees were older. Following treatment, we kept the bees on ice for 10 minutes for absorbing the treatment solution, followed with about 60 minutes of isolation in closed petri-dish with pollen and syrup to avoid the lateral transfer of the treatment solution. Each focal bee within a group was number tagged during group formation. The tag was attached to the dorsal part of the abdomen for the P-I treated bees, and on the thorax for JH-III treated bees. We performed additional sessions of behavioral observations on days 5 & 6 as described in section 2.3. Bees within the same group were of similar body size (One-way ANOVA, $F_{3, 132}= 0.06$, $p>0.05$, partial $\eta2=0.00$). The bees were collected from several donor colonies and assigned randomly to treatment groups.

2.9. Statistical analyses

Experiment 1: We studied 22 Groups across three Trials with a total number of 88 bees. For each bee, we measured three variables: oocyte size, the number of

threatening displays she performed, and their calculated dominance Index. For the statistical analyses, we used two-factorial General Linear Model, considering the Groups as independent units of sampling: One factor being the Trial, and another factor is the Treatment, having 4 repeated measures (Control, Sham, P-I, and P-I+JH). We thus have $3 \times 4 = 12$ random variables, and normality was ascertained for each after subjecting them to the $\ln(x+1)$ transformation.

Experiment 2: Here we studied 23 groups (5 Control, 5 Sham, 8 P-I, and 5 P-I+JH) with a total of 92 bees. The Groups are nested within the trials, which in turn are nested within Treatment. We analyzed three variables: oocyte length, the number of threatening displays, and the Dominance Index. Each was considered as the dependent variable in a two-way nested ANOVA. Normality was ascertained after subjecting each dependent variable to the square-root transformation. We also tested the effect of Treatment on the within-group dominance hierarchy, measured by the Dominance Range (i.e., the difference in Dominance Indices between α and δ bees in the group), using a two-way nested ANOVA.

Experiment 3: We studied 26 Groups (5 Control, 5 Sham, 8 P-I, and 8 P-I+JH) consisted of 104 bees. The statistical analysis is similar to that described for Experiment 2.

Experiment 4: In Experiment 4A we had 20 groups, 10 in which the α bee was treated with PI, and 10 in which the α bee was treated with only the vehicle (altogether 80 bees). In Experiment 4B we had 16 groups, 8 in which the δ bee was treated with JH-III, and 8 in which the δ bee was treated with only the vehicle (altogether 64 bees). For each bee we measured the terminal oocyte size, the number of threatening displays, and the DI at the first and at the second sets of behavioral observations. After applying the square-root transformation (for achieving

normality), we considered for each bee three variables: (1) the size of terminal

oocyte, (2) the difference between the number of threatening displays during the

second and the first sets of behavioral observations, and (3) the difference between

the dominance rank at the second and the first sets of behavioral observations.

For each dominance rank, we then compared in Experiment 4A the means of

these three variables between the bees of the treated (P-I) and Control (vehicle)

groups. Similarly, in Experiment 4B we compared for each dominance rank, the

means of the abovementioned three variables between the bees of the treated (JH-

III) and Control (vehicle) groups (all comparisons were performed using independent

samples t-tests, assuming equal or unequal variances, as determined by the

corresponding F-tests).

Dominance hierarchy as a function of JH level: We used three measures for

assessing the linearity of the dominance hierarchy within a group: (1) The DI range,

(2) the DI standard deviation, and (3) the inverse of the Shannon's Diversity Index,

which we calculated using the DI data. We used the ovarian state as a proxy for JH

level (see above) and considered two measures: (1) The mean length of the four

largest oocytes and (2) the length of only the largest oocyte in the ovaries. For

testing the relationship between the extent of the dominance hierarchy and ovarian

state, we performed six linear regressions, one for each of the 3 × 2 possible

combinations. These were done twice – once on the data of Experiment 2 (different

size bees) and once on the data of Experiment 3 (same size bees). We used partial

η^2 for effect size in ANOVA, whereas Cohen's d (Cd) measure of effect size was

calculated for pairwise comparisons.

3. Results

3.1. Experiment 1: The influence of JH on dominance and aggression in groups of similar size, differently-treated queenless worker bees

The assessment of ovarian state confirm that our JH manipulations were effective; P-I treated bees had ovaries containing smaller oocytes compared to control bees, an effect that was reverted by replacement therapy (Fig. 2A; Wilks' Lambda = 0.04, $F_{3,\ 17}$ = 130.45, P < 0.001, partial η^2 = 0.96; Bonferroni adjusted pairwise comparisons, Control vs. Sham: P > 0.999, Cd=0.31; Control vs. PI: P < 0.001, Cd=3.76; Control vs. JH: P = 0.180, Cd=0.71; Sham vs. PI: P < 0.001, Cd=4.41; Sham vs. JH: P = 0.006, Cd=1.16; PI vs. JH: P < 0.001, Cd=4.64). Trial had no significant effect on oocyte size ($F_{2,\ 19}$ = 0.54, P = 0.59, partial η^2 =0.01). The P-I treated bees also showed reduced amounts of threatening displays, an effect that was recovered by the replacement therapy, showing that it is regulated by JH (Fig. 2B; Wilks' Lambda = 0.11, $F_{3,17}$ = 45.46, P < 0.001, partial η^2 = 0.89; Bonferroni adjusted pairwise comparisons, Control vs. Sham: P > 0.999, Cd=0.38; Control vs. PI: P < 0.001, Cd=2.46; Control vs. JH: P = 0.061, Cd=0.86; Sham vs. PI: P < 0.001, Cd=2.37; Sham vs. JH: P < 0.001, Cd=1.54; PI vs. JH: P < 0.001, Cd=3.51). There was a small but statistically significant effect of Trial ($F_{2,\ 19}$ = 6.68, P = 0.006, partial η^2=0.07). JH manipulation also had a significant influence on the Dominance Index (Fig. 2C; Wilk's Lambda = 0.03, $F_{3,\ 17}$ = 166.90, P < 0.001, partial η^2 = 0.96; Bonferroni adjusted pairwise comparisons, Control vs. Sham: P = 0.944, Cd=0.44; Control vs. PI: P < 0.001, Cd=3.43; Control vs. JH: P = 0.054, Cd=0.88; Sham vs. PI: P < 0.001, Cd =2.12; Sham vs. JH: P = 0.003, Cd=1.26; PI vs. JH: P < 0.001, Cd=5.75) with no effect of Trial ($F_{2,\ 19}$ = 2.78, P = 0.087, partial η^2=0.01). The Bonferroni adjusted pairwise comparisons (Fig. 2C) show that the P-I treated bees had a significantly lower Dominance Index. Thus, the rank in the dominance

hierarchy was influenced by treatment (Chi-Square Test $_{(n=22)}$ =114.55, df=9,

p<0.001). In all groups except one, the P-I treated bee showed the lowest

dominance rank (δ; 95.45%), and in the remaining one group it had the second

lowest (γ) rank. The majority of the replacement therapy treated bees were the most

dominant in their group (73%; Fig. 2D).

The replacement therapy not only recovered the P-I treatment effect, but actually

resulted in better-developed ovaries and higher levels of threatening displays and

dominance index compared to the sham-treated (but not the control bees, Fig. 2).

This experiment shows that JH can influence the amount of threatening displays and

dominance rank in groups composed of workers bees with similar body size.

Figure 2: JH affects ovarian development, aggression and dominance status in

groups of four orphan workers of similar body size and mixed treatments. (A) Ovarian

state; (B) threatening displays; (C) Dominance Index. In A – C: Each colored symbol depicts

the value of an individual worker bee. The color and shape of the symbols correspond to the

bee's dominance rank within its group with α, β, γ & δ describing descending dominance

ranks. Data were [Ln(X+1)] transformed to normalize the distribution for statistical analyses.

Treatments marked with different small letters are statistically different in a Wilk's Lambda

test followed by Bonferroni's correction. N=22 bees treated and tested per treatment; (D)

Dominance rank distribution. The number in each stack box depicts the percentage of bees

of each dominance rank. Treatment had a significant effect on the propensity to acquire a

certain dominance rank (Chi-square test, p<0.001). Other details as in Fig.1.

3.2. Experiment 2: The influence of JH on dominance and aggression in

groups of different size, similarly-treated queenless worker bees

Ovarian state differed between treatment groups, confirming the affectivity of JH

manipulations in this experiment (Fig.3A, Treatment: $F_{3,\ 69}$ = 21.91, P < 0.001, partial

η^2 = 0.49; Group: $F_{19,\ 69}$ = 0.71, P = 0.793, partial η^2 = 0.16; Bonferroni adjusted

pairwise comparisons, Control vs. Sham: P > 0.999, Cd=0.12; Control vs. PI: P <

0.001, Cd=1.69; Control vs. PI+JH: P > 0.999, Cd=0.11; Sham vs. PI: P < 0.001,

Cd=1.82; Sham vs. PI+JH: P > 0.999, Cd=0.01; PI vs. PI+JH: P < 0.001, Cd=1.81).

All the bees in the P-I treatment groups, including the most dominant individuals, had

undeveloped ovaries, similar in size to those of the low ranked bees (γ & δ) in the

Control and Sham treatment groups (Fig. 3A). In the Replacement Therapy groups,

all the bees, including the low ranked individuals, had similarly active ovaries (One

way ANOVA $_{(replacement\ group)}$, $F_{3,16}$=1.17, p=0.35, partial $\eta2$=0.18). The average oocyte

length of the γ and δ ranked individuals in the replacement therapy groups appeared

larger than for bees of similar dominance rank in the Control, Sham and P-I

treatment groups (Fig. 3A).

Threatening displays did not differ between treatment groups (Fig. 3B, $F_{3,\,69}$ =

0.99, P = 0.403, partial η^2 = 0.04) and there was no significant effect for group factor

(Fig. 3B, $F_{19,\,69}$ = 0.57, P = 0.916, partial η^2 = 0.14). In all groups, the most dominant

individual performed most of the threatening displays (Fig. 3B). All the treatment

groups established clear dominance hierarchy (Fig. 3C; Two way Nested ANOVA

$_{(Treatment)}$, $F_{3,\,69}$=0.30, p > 0.05, partial $\eta2$=0.01). There was also no significant effect

of treatment on the Dominance Index (Fig.3C, $F_{3,\,69}$ = 0.30, P = 0.823, partial η^2 =

0.01) and no significant effect of the Group factor (Fig.3C, $F_{19,\,69}$ = 0.16, P > 0.999,

partial η^2 = 0.04).

Figure 3. JH effects on ovarian development, dominance status and threatening displays in groups of four orphan workers of different body size and similar treatment. (A) Ovarian state (B) Threatening behavior, (C) Dominant Index, and (D) Dominance rank distribution. Data were square-root transformed to normalize the distribution for statistical analysis. Letters (L, LM, SM, and S) refer to the body size of the bees (Table S1). Other details as in Fig. 1 & 2.

To compare the strength of the linear dominance hierarchies across treatments, we first estimated the range of dominance indexes by subtracting the dominance index of δ from that of the α individual in each group. This value is expected to be larger when the dominance hierarchy is clear. We found no significant effect of treatment on Dominance Range (Fig.S3, $F_{3, 12}$ = 1.02, P = 0.418, partial η^2=0.20) and no effect of the Trial ($F_{7, 12}$ = 1.96, P = 0.147, partial η^2 = 0.53). We next tested the correlation between ovary state (length of the largest oocyte, or mean of the four largest oocyte) as a proxy for JH titer and three different indices for the strength of the dominance hierarchy: The range of the Dominance Index (see above), the Standard Deviation of the Dominance Index and the inverse of the Shannon's Diversity Index. We found that none of the six combinations produced a statistically significant correlation between the measure of the dominance hierarchy and that of ovarian state (Table S2).

Taken together, these analyses show that bees subjected to similar JH manipulation and thus, have similar ovarian state, can nevertheless establish clear dominance hierarchies, similar to that observed in groups of workers in which JH levels were not manipulated (i.e., the Control and Sham groups). Dominance rank in all treatment groups was strongly influenced by body size with the largest individual most likely to acquire the top tank in the dominance hierarchy, and the smaller individuals typically occupying the lower ranks. Importantly, these include the P-I and

replacement therapy treatment in which all the bees in a group are assumed to have

overall similar JH titers. In 7 out of 8 P-I treated groups (87.5%, Fig. 3D, Chi-Square

Test $_{(P-I)}$ =448.5, df=9, p<0.001) and in 3 out of 5 replacement therapy treated groups

the largest individual was the most dominant in the group (60%, Fig. 3D, Chi-Square

Test $_{(P-I+JH)}$ =432.00, df=9, p<0.001). This proportion is comparable to the Control and

sham groups (3 out of 5 groups = 60%, Fig. 3D, Chi-Square Test $_{(Control)}$ =240.00,

df=9, p<0.001, Chi-Square Test $_{(Sham)}$ =304.00, df=9, p<0.001). This experiment

shows that body size has a strong influence on agonistic behavior and dominance

rank in small groups of queenless workers, and this effect of body size is not

compromised in groups in which all bees are manipulated to have similar JH titers

and level of ovarian activation.

3.3. Experiment 3: The influence of JH on dominance and aggression in

groups of similar size, similarly-treated orphan worker bees

Given that Exp.1 and Exp.2 indicate that both JH and body size (respectively)

influence dominance rank in small queenless groups, we next studied dominance

hierarchy formation in groups of bees of similar body size and JH levels. The strong

influence of treatment on ovarian state confirmed the effectiveness of our JH

manipulations (Fig. 4A; Treatment: $F_{3,\ 78} = 43.78$, $P < 0.001$, partial $\eta^2 = 0.63$; Group:

$F_{22,78} = 0.35$, $P = 0.996$, partial $\eta^2 = 0.09$; Bonferroni adjusted pairwise comparisons,

Control vs. Sham: $P > 0.999$, Cd=0.15; Control vs. PI: $P < 0.001$, Cd=2.07; Control

vs. PI+JH: $P = 0.114$, Cd=0.68; Sham vs. PI: $P < 0.001$, Cd=1.93; Sham vs. PI+JH:

$P = 0.028$, Cd=0.83; PI vs. PI+JH: $P < 0.001$, Cd=2.76). Even the most dominant P-I

treated individuals had undeveloped ovaries, with the largest oocytes appearing

similar in size to those of subordinate bees in the Control and Sham treatment

groups. Similarly, all the bees in the Replacement Therapy treatment groups,

including those with low dominance rank, had active ovaries appearing similar to

dominant bees in the Control and Sham treatment groups (Fig. 4A). These ovarian

analyses suggest that in both the P-I and Replacement Therapy groups all four

individuals had comparable, either low or high, JH levels, respectively (Fig.4A; One

way ANOVA within groups: P-I -- F=1.70, df=3, p>0.05, partial $\eta 2$=0.15;

Replacement Therapy -- F=29.90, df=3, p<0.0001, partial $\eta 2$=0.76).

JH manipulation also affected threatening behavior. P-I treated bees performed

overall less, and replacement therapy treated bees more, threatening displays

compared to the Sham and Control groups (Fig. 4B, Treatment: $F_{3,\ 78} = 26.81$, $P <$

0.001, partial $\eta^2 = 0.51$; Group: $F_{22,\ 78} = 0.10$, $P > 0.999$, partial $\eta^2 = 0.03$; Bonferroni

adjusted pairwise comparisons, Control vs. Sham: $P > 0.999$, Cd=0.19; Control vs.

PI: $P < 0.001$, Cd=1.73; Control vs. PI+JH: $P > 0.999$, Cd=0.38; Sham vs. PI: $P <$

0.001, Cd=1.55; Sham vs. PI+JH: $P = 0.301$, Cd=0.57; PI vs. PI+JH: $P < 0.001$,

Cd=2.12). These effects of JH treatment were closely linked to dominance rank: The

frequency of threatening displays was severely reduced in high-rank individuals in

the P-I treated groups (Fig. 4B) and elevated in low ranked individuals in the

Replacement Therapy treatment groups. JH manipulation did not affect Dominance

Index (Fig. 4C; $F_{3,\ 78} = 0.67$, $P = 0.574$, partial $\eta^2 = 0.03$; Group effect - $F_{22,\ 78} = 0.07$,

$P > 0.999$, partial $\eta^2 = 0.02$). It is remarkable that despite their similar JH levels,

ovary state, and elevated threatening displays by subordinate individuals, bees in the

Replacement Therapy groups nevertheless differ in their Dominance Index (Fig. 4C;

One way ANOVA within group, F=122.77, df=3, p<0.001, partial $\eta 2$=0.93). Although

the variation was lower in groups of P-I treated bees, the Dominance Index was

nevertheless significantly different among bees differing in their dominance rank (Fig.

4C; $F_{3, 28}$=13.41, p<0.001, partial η^2 = 0.50). The Dominance Range was influenced

by treatment ($F_{3, 14}$ = 150.82, P < 0.001, partial η^2 = 0.97; Trial effect − $F_{8, 14}$ = 2.38,

P = 0.074, partial η^2 = 0.58), and was significantly lower in the P-I and Replacement

Therapy groups compared with the Control and the Sham groups (Fig. 4C, D;

Bonferroni adjusted pairwise comparisons, Control vs. Sham: P > 0.999, Cd=0.38;

Control vs. PI: P < 0.001, Cd=10.02; Control vs. PI+JH: P < 0.001, Cd=5.94; Sham

vs. PI: P < 0.001, Cd=10.39; Sham vs. PI+JH: P < 0.001, Cd=6.32; PI vs. PI+JH: P <

0.001, Cd=4.07). By contrast to Exp. 2, here the strength of the dominance hierarchy

was positively correlated with the ovarian state when estimated using the range of

dominance index, Shannon's diversity index, or the standard deviation of dominance

index (Fig. 4D; Table S3). These results suggest that even in groups composed of

workers with similar body size, age, and JH levels, there is a weak dominance

hierarchy.

Figure 4: JH effects on ovarian development, dominance rank and threatening displays in groups of four orphan workers of similar body size and treatment. (A) ovarian state, (B) threatening displays, (C) Dominance Index and (D) Correlation between mean oocyte length and range of dominance index. Data were square-root transformed to normalize the distribution for statistical analysis. Other details as in Fig. 1&2.

3.4. Experiment 4: The influence of JH on dominance and aggression in groups of same size bees in which dominance hierarchy has been already established

Here, we tested whether P-I treatment can decrease the rank of dominant individuals (Exp. 4A), and whether JH treatment can increase, the dominance of subordinate (Exp. 4B), in groups that had already established stable dominance

hierarchies. Dominant (α) individuals that were treated with P-I had less developed

ovaries compared to same age dominant bees that were treated with only the vehicle

(Fig. 5A, left; Table 1Ai). It is notable, however, that the P-I treated dominant bees

still have developed ovaries when assessed at 7 days of age, suggesting that

reducing JH levels at this stage only slows or prevents further growth, but does not

lead to a significant decrease in oocyte size. The ovarian state was similar for the β,

γ & δ ranked bees (that were not treated) in the Sham and P-I treatment groups (Fig.

5A, left; Table 1Ai). The P-I treated dominant bees did not show a decrease in the

amount of threatening displays (Fig. 5B, left; Table 1Aii) or DI (Fig.5C, left; Table

1Aiii). None of the α-ranked individuals lost their high dominance rank after the P-I

treatment. Treating α individual with P-I also did not affect the level of threatening

displays or the Dominance Index of the other bees in the group.

In Experiment 4B, we found that δ-ranked individuals that were treated with JH

had significantly better-developed ovaries compared to sham-treated δ individuals in

control groups suggesting that our manipulation successfully elevated circulating JH

titers (Fig. 5A, right column; Table 1Bi). JH treatment also caused a significant

increase in the frequency of threatening displays performed by the treated individual

but not by her three groupmates (Fig. 5B, right column; Table 1Bii). This change in

behavior was associated with a higher DI for the treated individuals when compared

with the values before the treatment (Fig. 5C right column; Table 1Biii). Dominance

rank comparison showed that in 1 out of the 8 (12.5%) groups the JH treated δ

showed an increase in dominance rank, but this is not significantly different than the

probability to change rank in the control groups in which the δ-ranked bee was

treated with vehicle alone (0 out of the 8 groups, Fisher's exact test $_{(δ\text{-treated})}$, p=0.54).

Figure 5: Effect of JH on dominance rank in groups that have already established dominance hierarchies. Left column – Exp. 4A in which we reduced JH levels for the most dominant individual; right column – Exp. 4B in which we elevated JH for the most subordinate individual. (**A**) ovarian state, (**B**) threatening displays. and (**C**) Dominance Index. The letters on the x-axis show the treatment groups: S=Sham, T=Treatment (P-I or JH), B=before the treatment and A=after the treatment). Data were square-root transformed to normalize the distribution for statistical analysis. Other details as in Figs. 1 and 2.

610 **Table 1A.** The influence of treating the α ranked individual with P-I on (i) the terminal

611 oocyte length, (ii) the *change* in the number of threatening displays and (iii) the

612 *change* in the dominance index for all bees in the group as function of their

613 dominance rank (variables were square-root transformed for the statistical analyses).

Dominance Rank	α	B	γ	δ
(i). Terminal Oocyte Length	$t_{11} = -4.66$ $P < 0.001$ $Cd=2.08$	$t_{12} = 0.80$ $P = 0.440$ $Cd=0.36$	$t_{14} = -0.52$ $P = 0.607$ $Cd=0.23$	$t_{18} = -1.82$ $P = 0.086$ $Cd=0.81$
(ii). Threatening Displays	$t_{12} = -0.69$ $P = 0.503$ $Cd=0.31$	$t_{18} = 0.94$ $P = 0.359$ $Cd=0.42$	$t_{18} = 1.62$ $P = 0.122$ $Cd=0.73$	$t_{18} = -0.44$ $P = 0.663$ $Cd=0.20$
(iii). Dominance Index	$t_{12} = -0.35$ $P = 0.729$ $Cd=0.16$	$t_{10} = -1.28$ $P = 0.228$ $Cd=0.57$	$t_{14} = 0.61$ $P = 0.548$ $Cd=0.27$	$t_{10} = -0.73$ $P = 0.480$ $Cd=0.33$

614 **Cd*=Cohen's d*

615 **Table 2B.** The influence of treating the δ ranked individual with JH-III on (i) the

616 terminal oocyte length, (ii) the *change* in the number of threatening displays and (iii)

617 the *change* in the dominance index for all bees in the group as function of their

618 dominance rank (variables were square-root transformed for the statistical analyses).

Dominance Rank	α	B	γ	δ
(i). Terminal Oocyte Length	t14 = −1.44 P = 0.172 *Cd*=0.72	t14 = −0.15 P = 0.883 *Cd*=0.08	t14 = 1.57 P = 0.139 *Cd*=0.78	t14 = 8.21 $P = 10^{-6}$ *Cd*=4.11
(ii). Threatening Displays	t14 = 0.50 P = 0.629 *Cd*=0.25	t14 = 1.36 P = 0.196 *Cd*=0.68	t14 = −0.68 P = 0.508 *Cd*=0.34	t14 = 5.25 $P = 10^{-4}$ *Cd*=2.62
(iii). Dominance Index	t14 = 0.05 P = 0.958 *Cd*=0.03	t14 = −1.12 P = 0.281 *Cd*=0.56	t10 = 0.63 P = 0.543 *Cd*=0.31	t14 = 3.98 P = 0.001 *Cd*=1.99

619 **cd*=Cohen's d*

620 The correlation between the DI standard deviation before and after the treatment did

621 not show any significant effect for α and δ treated individuals. This experiment

622 suggests that JH has little, or only limited, effect on dominance rank after a stable

623 dominance hierarchy had already been established.

4. Discussion

We combined treatments with the allatoxin P-I and replacement therapy with the

natural JH to manipulate circulating JH levels in orphan groups of bumble bee

workers. Our measurements of ovarian activity (terminal oocyte length) confirm that

our JH manipulations were effective. Using this powerful system we show that in

addition to its gonadotrophic effects on physiology, JH also affects social behaviors

that are associated with reproduction such as aggressiveness and dominance. Our

results suggest that JH interacts with additional factors such as body size and

previous experience that also influence aggressiveness and dominance. These

findings explain previous ambiguities concerning whether or not JH influences

dominance in bumble bees.

In orphan quartets in which each callow worker was subjected to a different JH

manipulation (Exp. 1), the P-I and replacement therapy treated bees typically show

the lowest and highest dominance rank, respectively. These findings are consistent

with the premise that JH influences dominance in this species. However, in groups in

which callow bees of a similar size were subjected to the same treatment (Exp. 3),

dominance rank was apparent even in the P-I and replacement therapy quartets in

which all four bees were subjected to similar JH manipulation and had comparable

ovarian activity, although dominance hierarchy was not as pronounced as in the

Control and Sham treatment groups (Fig. 4C). These findings are consistent with the

hypothesis that variation in JH levels is not necessary for the establishment of

dominance hierarchies. This notion received further support in Exp. 2 in which the

bees in the same group differ in body size, but all of which were subjected to the

same treatment. Here, the dominance hierarchies in the P-I and replacement therapy

treated groups were overall comparable to that in the Control and Sham treatment

groups (Fig. 3C). Large body size increased and small size decreased, the

propensity to secure the top dominance rank. These findings show that variation in

body size is sufficient for establishing stable dominance hierarchies among bees that

do not differ in JH titers and ovarian state. In the last experiment, we treated

somewhat older (4-day-old) bees that have already established dominance

hierarchies. We downregulated JH titers for the most dominant individual (α) and

elevated JH titers for the most subordinate (γ) individual. These treatments affected

ovarian state but had only limited effects on the level of threatening behavior and on

the dominance rank of the treated individual. This experiment exemplifies the

importance of previous experiment by showing that the influence of JH on

dominance and aggression is small at best in bees that have already established a

stable dominance hierarchy (Fig. 5). Taken together, our results show that JH is

sufficient but not necessary for the regulation of aggressiveness, and the

establishment and maintenance of a dominance hierarchy in *B. terrestris*. Body size,

previous experience, and conceivably additional factors, also influence dominance

status.

Our findings are consistent with the notion that JH modulates aggressiveness in

bumble bees. In the first three experiments, bees treated with P-I to reduce JH titers

showed less threatening displays relative to control and sham-treated bee's (this

trend was not statistically significant in Exp. 2), an effect that was reverted by

replacement therapy with JH-III. JH manipulation however, did not affect

aggressiveness in the fourth experiment, in which bees were treated after the

dominance hierarchy had been already established. These manipulation

experiments are consistent with the typical correlation between dominance rank and

JH hemolymph titer (Bloch et al., 2000a; Bloch and Hefetz, 1999), in vitro JH

biosynthesis rates (Bloch et al., 1996; Larrere and Couillaud, 1993), CA volume

(Röseler et al., 1984; Van Doorn, 1989), and brain transcript abundance of the

transcription factor *Kr-h1* which is considered a major JH readout gene in insects

(Shpigler et al., 2010). Our findings are also consistent with studies showing that JH

modulates aggression in other species of insects (Barth et al., 1975; Hunt, 2007; Kou

et al., 2009, 2008; Pearce et al., 2001; Robinson et al., 1987; Röseler et al., 1984;

Scott, 2006a, 2006b; Tibbetts and Huang, 2010), including other bees (Breed, 1983;

Huang et al., 1994). It should be noted however, that some previous findings are not

consistent with the hypothesis that JH affects dominance or aggression in bumble

bees. Amsalem et al., (2014a, b) reported that oral treatment with JH, P-I, or both did

not affect Vg mRNA levels and aggression, and suggested an alternative hypothesis

stating that Vg regulates aggression independent of JH. Some of the inconsistencies

between our results and that of Amsalem et al., (2014a, b), may be explained by

methodological differences. For example, Amsalem et al., (2014a, b) used oral P-I

treatment (3 mg/bee) and a higher JH-III dose (100 µg/bee compared to 50µg/bee in

our study), that may affect the response to JH treatment (Pinto et al., 2000; Rutz et

al., 1976). In addition, the analyses of ovarian activity in Amsalem et al., (2014b)

suggest that their replacement therapy did not fully recover the effect of the P-I

treatment in some of their experiments (e.g., Figs 2 and 3 in this paper). Our

procedures also differ from that of Van Doorn (1989) who injected JH-I into one

individual in a queenless group. He reported only a small effect on the propensity of

the injected bee to acquire the top dominance rank. However, given that control bees were not disturbed in this experiment, it is possible that the stress of injection interfered with the hormonal influence; it is also not clear at what age the bees were injected. Similarly, JH was not found necessary for the expression of dominance behavior in the wasp *Synoeca surinama* (Kelstrup et al., 2014).

Our study points to factors other than JH that also affect dominance and aggression. Bees manipulated to have similar JH levels (i.e., all workers in the same group are similarly treated with P-I or replacement therapy) nevertheless established dominance hierarchies (Exp. 2 and 3). Exp. 2 further indicates that one important factor is body size. Bees in the P-I and Replacement Therapy groups showed agnostic behaviors and formed clear dominance hierarchy comparable to those in the control and sham-treated groups, despite having similarly developed ovaries (which is a proxy for JH titers; Fig. 3). Dominance rank in this experiment was not correlated with the ovarian state but rather with body size. These findings are consistent with earlier studies showing that body size affects dominance in untreated *B. terrestris* (Röseler, 1977; Van Doorn, 1989) and *B. atratus* worker bees (Matos and Garofalo, 1995). Body size is also correlated with dominance rank in other social bees (e.g., carpenter bees, Withee and Rehan, (2016), Lawson et al., (2017); halictine bee, Smith and Weller, (1989)) as well as other social insects such as paper wasps (Cervo et al., 2008; Chandrashekara and Gadagkar, 1991; Zanette and Field, 2009) and queenless ants (Heinze and Oberstadt, 1999; Nowbahari et al., 1999; Trible and Kronauer, 2017). The common correlation between body size and dominance rank can be explained by the importance of body size in determining the outcome of agonistic interactions and the establishment of dominance hierarchies. However, it should be noted that size is not always correlated with dominance rank

in studies with bumble bees (Amsalem and Hefetz, 2010; Foster et al., 2004; Free, 1955) and other insects such as the paper wasp, *Polybia occidentalis* (O'Donnell and Jeanne, 1995)*.* Size and aggression do not seem to be important for reproductive dominance in the primitively eusocial wasp *Ropalidia marginata* (Gadagkar, 2009; Bang and Gadagkar, 2012).

Previous experience can also influence the outcome of agonistic interactions and may explain the little effect of JH manipulation in Exp. 4. In many animals, older (but not too old) individuals are more likely to dominate younger opponents (Hughes and Strassmann, 1988; Khazraïe and Campan, 1999; Stevenson and Schildberger, 2013; Tsuji and Tsuji, 2005). More specific effects of previous agonistic interactions, known as "winner" and "loser" effects, reflect the increasing tendency of previous winners to win again, and of previous losers to be beaten in a conflict following a defeat (Goubault et al., 2019; reviewed in Hsu et al., 2006; Kim et al., 2018), and were recently suggested to be also important in a social wasp (Bang and Gadagkar, 2016). In most groups in Experiment 4, reducing JH and ovarian activity in the already established α-ranked bee did not cause her to lose her top position. Given that the bees in each group were of a similar body size, it seems unlikely that the α bee kept her top rank after JH was reduced, and the δ bee remained the most subordinate after her JH titers were increased because they were the largest or smallest in their groups, respectively. We suggest that a more likely explanation is that previous experience as the dominant or subordinate individual, or both, have a stronger influence on subsequent dominance rank than our manipulation of circulating JH titers. The prior experience was found to affect dominance rank in many animals, including species of insects such as crickets and beetles (Arneson and Wcislo, 2003; Goubault and Decuignière, 2012; Khazraïe and Campan, 1999;

Lee et al., 2014). To the best of our knowledge, loser and winner effects have not been explicitly tested in bumble bees, but there is evidence lending credence to this notion. For example, in queenless groups composed of both callow and older dominant workers, the later ones typically attain the top rank and inhibit ovarian development in the callows, but in groups composed of only callows, one of the youngsters acquires the top dominance rank (Bloch and Hefetz, 1999).

Our finding that JH is one of several factors affecting dominance and aggression in bumble bees can explain the apparently conflicting findings in previous studies (Amsalem et al., 2014b, 2014a; Röseler, 1977; Van Doorn, 1989). The effects of JH manipulation is influenced by the context, including JH titers of other bees in the group. This multiplicity of factors raises questions concerning the ways these factors interact during dominance hierarchy formation. We suggest that in groups of callow workers, hemolymph JH titers are less important during the initial stages of hierarchy formation. This premise is consistent with observations that the highest aggression and the initial establishment of a clear dominance hierarchy in small groups of callow workers typically occur during the first few days in which both JH biosynthesis rates (as determined in vitro) and JH titers are low (Bloch et al., 2000a, 1996; Van Doorn, 1989). Body size, aggressiveness, age, and previous experience may all play a role during this initial stage. We speculate that the burst of intensive agonistic interactions leads to the activation of the CA in the more aggressive and dominant individuals. This idea is consistent with the "challenge hypothesis", which was developed to explain the social modulation of testosterone to increase aggression in the context of courtship and reproduction in vertebrates (Wingfield et al., 1990), and later extended to JH in insects (Scott, 2006a; Tibbetts and Huang, 2010; Bernstein et al., 1979; Kou et al., 2019, 2009; Tibbetts and Huang, 2010) . Indeed, as predicted by the challenge

hypothesis, JH titers and CA activity are higher during conflicts associated with
unstable social situations under both queenless and queenright conditions (Bloch et
al., 2002; Bloch et al., 2000a, 1996; Röseler, 1977; Van Doorn, 1989). The JH –
ovary axis is not activated in workers that do not experience intensive agonistic
interactions of this kind. For example, JH and oogenesis are low in workers that are
placed individually (Cnaani et al., 2002; Duchateau and Velthuis, 1989; Röseler PF.,
1990; Sibbald and Plowright, 2015) or in relatively stable queenright conditions. The
dominant individuals further inhibit CA activity and JH signaling in subordinate
workers (Bloch et al., 2000a, 1996; Bloch and Hefetz, 1999; Shpigler et al., 2010).
The consequential elevated JH titers in the dominant bees coordinate many tissues
involved in reproduction. These include modulation of the fat body, exocrine glands,
and ovaries, and stimulating processes such as vitellogenesis and oogenesis
(Amsalem et al., 2013; Shpigler et al., 2014, 2010). Our unpublished RNA
sequencing analyses further show that JH regulates the expression of hundreds of
genes in the brain, which is consistent with the hypothesis that JH influences
behaviors that are associated with reproduction. Thus, JH may stabilize the
dominance hierarchy by enhancing the aggressiveness in the most dominant
individual, which in turn inhibits its levels in the subordinate ones. It should be noted,
however, that currently there is no evidence that JH acts directly on the neuronal and
molecular underpinnings of aggression, and its effects may be mediated by
additional neuroendocrine signals. For example, dominance in bumble bees is also
correlated with neuroendocrine signals such as octopamine (Bloch et al., 2000c),
ecdysteroids (Bloch et al., 2000; Geva et al., 2005), and Vg (Amsalem et al., 2014b,
2014a). As for ecdysteroids, although the activated ovaries of the dominant
individuals produce high amounts of ecdysteroids (Geva et al., 2005), these probably

have little influence (if at all) on aggression and dominance hierarchy formation

because ovariectomized workers can attain high dominance rank even if their

ovaries are removed at an early age (van Doorn, 1989). Factors other than JH are

also important after the initial elevation of JH titers in dominant bees, as evident by

the limited effect of JH manipulation in bees that have already established

dominance hierarchy in Exp. 4. Additional studies in which the different factors are

manipulated are needed for clarifying how and when they interact to regulate

aggression and dominance.

Conclusions

We used multiple experimental designs to manipulate JH in quartets of orphan

bumble bee workers to clarify some of the ambiguity concerning the behavioral

functions of JH in conflicts over reproductive dominance. We unequivocally show

that JH affects aggressiveness and dominance rank in *B. terrestris*, and under some

conditions can even be the pivotal factor determining rank during the establishment

of a dominance hierarchy. We also show that JH is not the only player in the system.

Body size, age, and previous experience are also important and the interplay

between these multiple factors is just starting to be unveiled. Our study highlights

many open questions pertaining to the mechanisms by which JH affects

aggressiveness and social behavior to shape a dominance hierarchy. Although our

results show that JH manipulations affect these behaviors, it is yet not clear if JH

acts directly on the brain or that its effect is mediated by some yet unknown

intermediary neuroendocrine signals.

Acknowledgements

We thank Michael Schneider and David Zlotkin for helping with behavioral

observations. This research was supported by grants from the US-Israel Binational

Agricultural Research and Development Fund (BARD, number IS-4418-11, and IS-

5077-18, to G.B.) and The Planning and Budgeting Committee (PBC) Fellowship

Program for Outstanding Chinese and Indian Post-Doctoral Fellows (to A.P).

Conflict of interest: There is no conflict of interest for any of the authors.

Supplementary figures:

Supplementary Figure S1: Possible lateral transfer of drugs used to

manipulate JH titers. The graph shows ovarian state for bees in a preliminary

experiment in which individuals subjected to different treatments were housed

together shortly after treatment on Day-1. The ovaries of the P-I treated bees in this

experiment are better developed compared with Experiment 1 in which the treated

bees where kept individually isolated for the 48hrs after treatment. Treatments

marked with different small letters are statistically different in a Kruskal-Wallis test

followed by Dunn's posthoc analysis. Other details as in Figure 1.

Supplementary Figure S2: Effect of sham treatments with vehicle only on aggressiveness and dominance rank in groups which had already established stable hierarchies. Left column – Exp. 4A, groups in which we treated the dominant individual with castor oil (5µl/bee, applied to the thorax); right column – Exp. 4B, groups in which we treated the most subordinate individual with DMF (5µl/ bee, applied to the abdomen). (**A**) Threatening behavior and (**B**) Dominance Index B- *Before* and **A-** *After* the treatment. Other details as in Fig. 1 & 5.

Supplementary Figure S3: Effect of JH on the linearity of the dominance

hierarchy in Experiment 2. The range of Dominance Index is the difference

between the Dominance Index of the α and the δ in each group. Treatment groups

consisted of different size, similarly-treated queenless worker bees. Other details as

in Fig. 4D.

Supplementary Table S1. Size-adjusted amounts of Precocene-I and castor oil

topically applied to *B. terrestris* workers in Experiments 1-3. The final concentration

applied to all the bees was 50µg P-I/ 1µl castor oil.

Serial Number	Wing size (mm)	Body size denotation	P-I amount (µg/bee)	Castor oil Volume (µl/bee)

1	2.5-2.6	Small (S)	200	4.0
2	2.7-2.8	Small medium (SM)	220	4.4
3	2.9-3.0	Large medium (LM)	240	4.8
4	3.1-3.2	Large (L)	260	5.2

853 **Supplementary Table S2.** Correlations between three measures of dominance

854 hierarchy and two measures of oocyte size in the Experiment 2.

Dependent Variable	Independent Variable	Correlation Coefficient	P value
Dominance Range	Mean oocyte size	0.04	0.852
	Maximal oocyte	0.12	0.576
Dominance SD	Mean oocyte size	0.03	0.877
	Maximal oocyte	0.14	0.515
(Shannon)-1	Mean oocyte size	−0.1	0.653
	Maximal oocyte	0.025	0.911

856 **Supplementary Table S3.** Correlations between three estimates of dominance

857 hierarchy and two measures of oocyte size in the Experiment 3.

Dependent Variable	Independent Variable	Correlation Coefficient	P value
Dominance Range	Mean oocyte size	0.63	< 0.001
	Maximal oocyte	0.78	$<10^{-5}$
Dominance SD	Mean oocyte size	0.66	< 0.001
	Maximal oocyte	0.80	$<10^{-6}$
$(Shannon)^{-1}$	Mean oocyte size	0.39	0.051
	Maximal oocyte	0.63	< 0.001

References:

Adams, M.E., 2009. Juvenile Hormones, in: Encyclopedia of Insects.

https://doi.org/10.1016/B978-0-12-374144-8.00155-7

Amsalem, E., Hefetz, A., 2011. The effect of group size on the interplay between

dominance and reproduction in Bombus terrestris. PLoS One 6.

https://doi.org/10.1371/journal.pone.0018238

Amsalem, E., Hefetz, A., 2010. The appeasement effect of sterility signaling in

dominance contests among Bombus terrestris workers. Behav. Ecol. Sociobiol.

64, 1685–1694. https://doi.org/10.1007/s00265-010-0982-4

Amsalem, E., Malka, O., Grozinger, C., Hefetz, A., 2014a. Exploring the role of

juvenile hormone and vitellogenin in reproduction and social behavior in bumble

bees. BMC Evol. Biol. 14, 45. https://doi.org/10.1186/1471-2148-14-45

Amsalem, E., Shamia, D., Hefetz, A., 2013. Aggression or ovarian development as

determinants of reproductive dominance in Bombus terrestris: Interpretation

using a simulation model. Insectes Soc. 60, 213–222.

https://doi.org/10.1007/s00040-013-0285-7

Amsalem, E., Teal, P., Grozinger, C.M., Hefetz, A., 2014b. Precocene-I inhibits

juvenile hormone biosynthesis, ovarian activation, aggression and alters sterility

signal production in bumble bee (Bombus terrestris) workers. J. Exp. Biol. 217,

3178–3185. https://doi.org/10.1242/jeb.107250

Andrade-Silva, A.C.R., Nascimento, F.S., 2015. Reproductive regulation in an orchid

bee: Social context, fertility and chemical signalling. Anim. Behav. 106, 43–49.

https://doi.org/10.1016/j.anbehav.2015.05.004

Arneson, L., Wcislo, W.T., 2003. Dominant-Subordinate Relationships in a

Facultatively Social, Nocturnal Bee, Megalopta genalis (Hymenoptera:

Halictidae). Source J. Kansas Entomol. Soc. https://doi.org/10.2307/25086104

Bang, A., Gadagkar, R., 2016. Winner–loser effects in a eusocial wasp. Insectes

Soc. 63, 349–352. https://doi.org/10.1007/s00040-015-0455-x

Bang, A., Gadagkar, R., 2012. Reproductive queue without overt conflict in the

primitively eusocial wasp Ropalidia marginata. Proc. Natl. Acad. Sci.

https://doi.org/10.1073/pnas.1212698109

890 Barth, R.H., Lester, L.J., Sroka, P., Kessler, T., Hearn, R., 1975. Juvenile hormone

891   promotes dominance behavior and ovarian development in social wasps

892   (Polistes annularis). Experientia 31, 691–692.

893   https://doi.org/10.1007/BF01944632

894 Bell, W.J., 1973. Factors controlling initiation of vitellogenesis in a primitively social

895   bee, Lasioglossum zephyrum (Hymenoptera: Halictidae). Insectes Soc.

896   https://doi.org/10.1007/BF02223194

897 Bernstein, I.S., Rose, R.M., Gordon, T.P., Grady, C.L., 1979. Agonistic rank,

898   aggression, social context, and testosterone in male pigtail monkeys. Aggress.

899   Behav. https://doi.org/10.1002/1098-2337(1979)5:4<329::AID-

900   AB2480050402>3.0.CO;2-Z

901 Bloch, G., Borst, D.W., Huang, Z.-Y., Robinson, G.E., Cnaani, J., Hefetz, A., 2000a.

902   Juvenile hormone titers, juvenile hormone biosynthesis, ovarian development

903   and social environment in Bombus terrestris. J. Insect Physiol. 46, 47–57.

904   https://doi.org/10.1016/S0022-1910(99)00101-8

905 Bloch, G., Borst, D.W., Huang, Z.-Y., Robinson, G.E., Hefetz, A., 1996. Effects of

906   social conditions on Juvenile Hormone mediated reproductive development in

907   Bombus terrestris workers. Physiol. Entomol. 21, 257–267.

908   https://doi.org/10.1111/j.1365-3032.1996.tb00863.x

909 Bloch, G., Hefetz, A., 1999. Regulation of reproduction by dominant workers in

910   bumblebee (Bombus terrestris) queenright colonies. Behav. Ecol. Sociobiol. 45,

911   125–135. https://doi.org/10.1007/s002650050546

912 Bloch, G., Hefetz, A., Hartfelder, K., 2000b. Ecdysteroid titer, ovary status, and

913   dominance in adult worker and queen bumble bees (Bombus terrestris). J.

914   Insect Physiol. 46, 1033–1040. https://doi.org/10.1016/S0022-1910(99)00214-0

Bloch, G., Shpigler, H., Wheeler, D.E., Robinson, G.E., 2010. Endocrine influences

on the organization of insect societies, in: Hormones, Brain and Behavior

Online. pp. 1027–1070. https://doi.org/10.1016/B978-008088783-8.00030-9

Bloch, G., Simon, T., Robinson, G.E., Hefetz, A., 2000c. Brain biogenic amines and

reproductive dominance in bumble bees (Bombus terrestris). J. Comp. Physiol. -

A Sensory, Neural, Behav. Physiol. 186, 261–268.

https://doi.org/10.1007/s003590050426

Bloch, G., Wheeler, D., Robinson, G., 2002a. Endocrine Influences on the

Organization of Insect Societies, in: Hormones, Brain and Behavior.

https://doi.org/10.1016/B978-012532104-4/50042-1

Bloch, G., Wheeler, D., Robinson, G., 2002b. Endocrine Influences on the

Organization of Insect Societies, in: Donald W. Pfaff, Arthur P. Arnold, ... Robert

T. Rubin (Ed.), Hormones, Brain and Behavior. Academic Press, pp. 195–235.

Brain, P.F., 1977. Hormones and aggression. Annu. Res. Rev. Horm. Aggress. 1,

126.

Breed, M.D., 1983. Correlations between aggressiveness and corpora allata volume,

social isolation, age and dietary protein in worker honeybees. Insectes Soc. 30,

482–495. https://doi.org/10.1007/BF02223979

Cervo, R., Dapporto, L., Beani, L., Strassmann, J.E., Turillazzi, S., 2008. On status

badges and quality signals in the paper wasp Polistes dominulus: Body size,

facial colour patterns and hierarchical rank. Proc. R. Soc. B Biol. Sci. 275,

1189–1196. https://doi.org/10.1098/rspb.2007.1779

Chandrashekara, K., Gadagkar, R., 1991. Behavioural Castes, Dominance and

Division of Labour in a Primitively Eusocial Wasp. Ethology.

https://doi.org/10.1111/j.1439-0310.1991.tb00252.x

Chen, T.T., Couble, P., Abu-Hakima, R., Wyatt, G.R., 1979. Juvenile hormone-

controlled vitellogenin synthesis in Locusta migratoria fat body. Hormonal

induction in vivo. Dev. Biol. 69, 59–72. https://doi.org/10.1016/0012-

1606(79)90274-4

Cnaani, J., Schmid-Hempel, R., Schmidt, J.O., 2002. Colony development, larval

development and worker reproduction in Bombus impatiens Cresson. Insectes

Soc. 49, 164–170. https://doi.org/10.1007/s00040-002-8297-8

De Loof, A., Baggerman, G., Breuer, M., Claeys, U., Cerstiaens, A., Clynen, E.,

Janssen, T., Schoofs, L., Broeck, J. Vanden, 2001. Gonadotropins in insects: An

overview. Arch. Insect Biochem. Physiol. https://doi.org/10.1002/arch.1044

Duchateau, M.J., 1989. Agonistic behaviours in colonies of the bumblebee Bombus

terrestris. J. Ethol. 7, 141–151. https://doi.org/10.1007/BF02350036

Duchateau, M.J., Velthuis, H.H.W., 1989. Ovarian development and egg laying in

workers of Bombus terrestris. Entomol. Exp. Appl. 51, 199–213.

https://doi.org/10.1111/j.1570-7458.1989.tb01231.x

Flatt, T., Kawecki, T.J., 2007. Juvenile hormone as a regulator of the trade-off

between reproduction and life span in Drosophila melanogaster. Evolution (N.

Y). https://doi.org/10.1111/j.1558-5646.2007.00151.x

Foster, R.L., Brunskill, A., Verdirame, D., O'Donnell, S., 2004. Reproductive

physiology, dominance interactions, and division of labour among bumble bee

workers. Physiol. Entomol. 29, 327–334. https://doi.org/10.1111/j.0307-

6962.2004.00388.x

Free, J.B., 1955. The behaviour of egg-laying workers of bumblebee colonies. Br. J.

Anim. Behav.

Gadagkar, R., 2009. Interrogating an insect society. Proc. Natl. Acad. Sci.

https://doi.org/10.1073/pnas.0904317106

Geva, S., Hartfelder, K., Bloch, G., 2005. Reproductive division of labor, dominance,

and ecdysteroid levels in hemolymph and ovary of the bumble bee Bombus

terrestris. J. Insect Physiol. 51, 811–823.

https://doi.org/10.1016/j.jinsphys.2005.03.009

Giray, T., Giovanetti, M., West-Eberhard, M.J., 2005. Juvenile hormone,

reproduction, and worker behavior in the neotropical social wasp Polistes

canadensis. Proc. Natl. Acad. Sci. 102, 3330–3335.

https://doi.org/10.1073/pnas.0409560102

Goubault, M., Decuignière, M., 2012. Previous Experience and Contest Outcome:

Winner Effects Persist in Absence of Evident Loser Effects in a Parasitoid

Wasp. Am. Nat. https://doi.org/10.1086/667192

Goubault, M., Exbrayat, M., Earley, R.L., 2019. Fixed or flexible? Winner/loser

effects vary with habitat quality in a parasitoid wasp. Behav. Ecol. Sociobiol.

https://doi.org/10.1007/s00265-019-2688-6

Hartfelder, K., 2000. Insect juvenile hormone: From "status quo" to high society.

Brazilian J. Med. Biol. Res. 33, 157–177. https://doi.org/10.1590/S0100-

879X2000000200003

Hartfelder K, E.W., 1998. Social Insect Polymorphism: Hormonal Regulation of

Plasticity in Development and Reproduction in the Honeybee, in: Roger A.

Pedersen, G.P.S. (Ed.), Current Topics in Developmental Biology. pp. 45–77.

https://doi.org/https://doi.org/10.1016/S0070-2153(08)60364-6

Haunerland, N.H., Bowers, W.S., 1985. Comparative studies on pharmacokinetics

and metabolism of the anti-juvenile hormone precocene II. Arch. Insect

Biochem. Physiol. https://doi.org/10.1002/arch.940020106

Heinze, J., Oberstadt, B., 1999. Worker age, size and social status in queenless

colonies of the ant Leptothorax gredleri. Anim. Behav. 58, 751–759.

https://doi.org/10.1006/anbe.1999.1188

Hsu, Y., Earley, R.L., Wolf, L.L., 2006. Modulation of aggressive behaviour by

fighting experience: Mechanisms and contest outcomes. Biol. Rev. Camb.

Philos. Soc. https://doi.org/10.1017/S146479310500686X

Huang, Z.Y., Robinson, G.E., Tobe, S.S., Yagi, K.J., Strambi, C., Strambi, A., Stay,

B., 1991. Hormonal regulation of behavioural development in the honey bee is

based on changes in the rate of juvenile hormone biosynthesis. J. Insect

Physiol. 37, 733–741. https://doi.org/10.1016/0022-1910(91)90107-B

Huang, Z.Y., Robinson1, G.E., Borst, D.W., 1994. Physiological correlates of division

of labor among similarly aged honey bees. J. Comp. Physiol. A 174, 731–739.

https://doi.org/10.1007/BF00192722

Hughes, C.R., Strassmann, J.E., 1988. Age Is More Important Than Size in

Determining Dominance Among Workers in the Primitively Eusocial Wasp,

Polistes Instabilis. Behaviour 107, 1–14.

https://doi.org/10.1163/156853988X00151

Hunt, G.J., 2007. Flight and fight: A comparative view of the neurophysiology and

genetics of honey bee defensive behavior. J. Insect Physiol.

https://doi.org/10.1016/j.jinsphys.2007.01.010

Kelstrup, H.C., Hartfelder, K., Nascimento, F.S., Riddiford, L.M., 2014. The role of

juvenile hormone in dominance behavior, reproduction and cuticular pheromone

signaling in the caste-flexible epiponine wasp, Synoeca surinama. Front. Zool.

11. https://doi.org/10.1186/s12983-014-0078-5

Khazraïe, K., Campan, M., 1999. The role of prior agonistic experience in dominance

relationships in male crickets Gryllus bimaculatus (Orthoptera: Gryllidae).

Behav. Processes. https://doi.org/10.1016/S0376-6357(98)00058-8

Kim, Y.-K., Saver, M., Simon, J., Kent, C.F., Shao, L., Eddison, M., Agrawal, P., Texada, M., Truman, J.W., Heberlein, U., 2018. Repetitive aggressive encounters generate a long-lasting internal state in *Drosophila melanogaster* males. Proc. Natl. Acad. Sci. 201716612. https://doi.org/10.1073/pnas.1716612115

Kou, R., Chou, S.Y., Chen, S.C., Huang, Z.Y., 2009. Juvenile hormone and the ontogeny of cockroach aggression. Horm. Behav. 56, 332–338. https://doi.org/10.1016/j.yhbeh.2009.06.011

Kou, R., Chou, S.Y., Huang, Z.Y., Yang, R.L., 2008. Juvenile hormone levels are increased in winners of cockroach fights. Horm. Behav. 54, 521–527. https://doi.org/10.1016/j.yhbeh.2008.05.011

Kou, R., Hsu, C.C., Chen, S.C., Chang, P.Y., Fang, S., 2019. Winner and loser effects in lobster cockroach contests for social dominance. Horm. Behav. https://doi.org/10.1016/j.yhbeh.2018.12.002

Larrere, M., Couillaud, F., 1993. Role of juvenile hormone biosynthesis in dominance status and reproduction of the bumblebee, Bombus terrestris. Behav. Ecol. Sociobiol. 33, 335–338. https://doi.org/10.1007/BF00172932

Lawson, S.P., Helmreich, S.L., Rehan, S.M., 2017. Effects of nutritional deprivation on development and behavior in the subsocial bee *Ceratina calcarata* (Hymenoptera: Xylocopinae). J. Exp. Biol. jeb.160531. https://doi.org/10.1242/jeb.160531

Lee, V.E., Head, M.L., Carter, M.J., Royle, N.J., 2014. Effects of age and experience on contest behavior in the burying beetle, Nicrophorus vespilloides. Behav. Ecol.

https://doi.org/10.1093/beheco/art101

Lengyel, F., Westerlund, S.A., Kaib, M., 2007. Juvenile hormone III influences task-

specific cuticular hydrocarbon profile changes in the ant Myrmicaria

eumenoides. J. Chem. Ecol. 33, 167–181. https://doi.org/10.1007/s10886-006-

9185-x

Matos, E.V.S., Garofalo, C.A., 1995. Observations on the development of queenless

colonies of Bombus atratus (Hymenoptera, Apidae). J. Apic. Res. 34, 177–185.

https://doi.org/10.1080/00218839.1995.11100903

Monnin, T., Peeters, C., 1999. Dominance hierarchy and reproductive conflicts

among subordinates in a monogynous queenless ant. Behav. Ecol. 10, 323–

332. https://doi.org/10.1093/beheco/10.3.323

Nelson, R.J., 2005. Biology of Aggression, Biology of Aggression.

https://doi.org/10.1093/acprof:oso/9780195168761.001.0001

Norman, V.C., Hughes, W.O.H., 2016. Behavioural effects of juvenile hormone and

their influence on division of labour in leaf-cutting ant societies. J. Exp. Biol. 219,

8–11. https://doi.org/10.1242/jeb.132803

Norris, D.O., Lopez, K.H., 2011. Hormones and Reproduction of Vertebrates -

Volume 4, Hormones and Reproduction of Vertebrates - Volume 4.

https://doi.org/10.1016/C2009-0-01697-3

Nowbahari, E., Fénéron, R., Malherbe, M.C., 1999. Effect of body size on aggression

in the ant, Cataglyphis niger (Hymenoptera; Formicidae). Aggress. Behav. 25,

369–379. https://doi.org/10.1002/(SICI)1098-2337(1999)25:5<369::AID-

AB5>3.0.CO;2-C

O'Donnell, S., Jeanne, R.L., 1995. The Roles of Body-Size and Dominance in

Division-of-Labor among Workers of the Eusocial Wasp Polybia-Occidentalis

(Olivier) (Hymenoptera, Vespidae). J. Kansas Entomol. Soc.

O'Donnell, S., Jeanne, R.L., 1993. Methoprene accelerates age polyethism in

workers of a social wasp (Polybia occidentalis). Physiol. Entomol.

https://doi.org/10.1111/j.1365-3032.1993.tb00467.x

Pamminger, T., Treanor, D., Hughes, W.O.H., 2016. Pleiotropic effects of juvenile

hormone in ant queens and the escape from the reproduction–

immunocompetence trade-off. Proc. R. Soc. B Biol. Sci. 283, 20152409.

https://doi.org/10.1098/rspb.2015.2409

Pan, M.L., Wyatt, G.R., 1971. Juvenile hormone induces vitellogenin synthesis in the

monarch butterfly. Science (80-.). 174, 503–505.

https://doi.org/10.1126/science.174.4008.503

Pearce, A.N., Huang, Z.Y., Breed, M.D., 2001. Juvenile hormone and aggression in

honey bees. J. Insect Physiol. 47, 1243–1247. https://doi.org/10.1016/S0022-

1910(01)00109-3

Penick, C.A., Liebig, J., Brent, C.S., 2011. Reproduction, dominance, and caste:

Endocrine profiles of queens and workers of the ant Harpegnathos saltator. J.

Comp. Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol. 197, 1063–

1071. https://doi.org/10.1007/s00359-011-0667-0

Pinto, L.Z., Bitondi, M.M.G., Simões, Z.L.P., 2000. Inhibition of vitellogenin synthesis

in Apis mellifera workers by a juvenile hormone analogue, pyriproxyfen. J. Insect

Physiol. 46, 153–160. https://doi.org/10.1016/S0022-1910(99)00111-0

Rantala, M.J., Vainikka, A., Kortet, R., 2003. The role of juvenile hormone in immune

function and pheromone production trade-offs: A test of the immunocompetence

handicap principle. Proc. R. Soc. B Biol. Sci.

https://doi.org/10.1098/rspb.2003.2472

Riddiford, L.M., 2012. How does juvenile hormone control insect metamorphosis and

reproduction? Gen. Comp. Endocrinol. 179, 477–484.

https://doi.org/10.1016/j.ygcen.2012.06.001

Riddiford, L.M., 2008. Juvenile hormone action: A 2007 perspective. J. Insect

Physiol. https://doi.org/10.1016/j.jinsphys.2008.01.014

Robinson, G.E., 1992. Regulation of Division of Labor in Insect Societies. Annu. Rev.

Entomol. 37, 637–665. https://doi.org/10.1146/annurev.en.37.010192.003225

Robinson, G.E., Strambi, A., Strambi, C., Paulino-Simões, Z.L., Tozeto, S.O.,

Negraes Barbosa, J.M., 1987. Juvenile hormone titers in European and

Africanized honey bees in Brazil. Gen. Comp. Endocrinol. 66, 457–459.

https://doi.org/10.1016/0016-6480(87)90258-9

Robinson, G.E., Vargo, E.L., 1997. Juvenile hormone in adult eusocial hymenoptera:

Gonadotropin and behavioral pacemaker. Arch. Insect Biochem. Physiol. 35,

559–583. https://doi.org/10.1002/(SICI)1520-6327(1997)35:4<559::AID-

ARCH13>3.0.CO;2-9

Rodrigues, M.A., Flatt, T., 2016. Endocrine uncoupling of the trade-off between

reproduction and somatic maintenance in eusocial insects. Curr. Opin. Insect

Sci. 16, 1–8. https://doi.org/10.1016/j.cois.2016.04.013

Roseler, P., 1991. Reproductive competition during colony establishment, in: RW,

R.K. and M. (Ed.), The Social Biology of Wasps,. Cornell University Press, pp.

309–335.

Röseler, P.F., 1977. Juvenile Hormone Control of Oogenesis in Bumblebee Workers,

Bombus terrestris. J. Insect Physiol. 23, 985–992. https://doi.org/10.1016/0022-

1910(77)90126-3

Röseler, P.F., Röseler, I., Strambi, A., Augier, R., 1984. Influence of insect hormones

on the establishment of dominance hierarchies among foundresses of the paper

wasp, Polistes gallicus. Behav. Ecol. Sociobiol. 15, 133–142.

https://doi.org/10.1007/BF00299381

Röseler PF., van H.C.G.J., 1990. Castes and Reproduction in Bumblebees. In:

Engels W. (eds) Social Insects. Springer, Berlin, Heidelberg.

Rutz, W., Gerig, L., Wille, H., Lüscher, M., 1976. The function of juvenile hormone in

adult worker honeybees, Apis mellifera. J. Insect Physiol. 22, 1485–1491.

https://doi.org/10.1016/0022-1910(76)90214-6

Sasaki, T., Penick, C.A., Shaffer, Z., Haight, K.L., Pratt, S.C., Liebig, J., 2016. A

Simple Behavioral Model Predicts the Emergence of Complex Animal

Hierarchies. Am. Nat. 187, 765–775. https://doi.org/10.1086/686259

Scott, M.P., 2006a. Resource defense and juvenile hormone: The "challenge

hypothesis" extended to insects. Horm. Behav. 49, 276–281.

https://doi.org/10.1016/j.yhbeh.2005.07.003

Scott, M.P., 2006b. The role of juvenile hormone in competition and cooperation by

burying beetles. J. Insect Physiol. 52, 1005–1011.

https://doi.org/10.1016/j.jinsphys.2006.04.006

Shorter, J.R., Tibbetts, E.A., 2009. The effect of juvenile hormone on temporal

polyethism in the paper wasp Polistes dominulus. Insectes Soc. 56, 7–13.

https://doi.org/10.1007/s00040-008-1026-1

Shpigler, H., Amsalem, E., Huang, Z.Y., Cohen, M., Siegel, A.J., Hefetz, A., Bloch,

G., 2014. Gonadotropic and physiological functions of juvenile hormone in

bumblebee (Bombus terrestris) workers. PLoS One 9.

https://doi.org/10.1371/journal.pone.0100650

Shpigler, H., Patch, H.M., Cohen, M., Fan, Y., Grozinger, C.M., Bloch, G., 2010. The

transcription factor Krüppel homolog 1 is linked to hormone mediated social

organization in bees. BMC Evol. Biol. 10. https://doi.org/10.1186/1471-2148-10-

Shpigler, H., Tamarkin, M., Bloch, G., Poleg, M., Siegel, A.J., Gruber, Y., 2013.

Social influences on body size and developmental time in the bumblebee

Bombus terrestris. Behav. Ecol. Sociobiol. 1–12.

Shpigler, H.Y., Siegel, A.J., Huang, Z.Y., Bloch, G., 2016. No effect of juvenile

hormone on task performance in a bumblebee (Bombus terrestris) supports an

evolutionary link between endocrine signaling and social complexity. Horm.

Behav. 85, 67–75. https://doi.org/10.1016/j.yhbeh.2016.08.004

Sibbald, E.D., Plowright, C.M.S., 2015. Reproductive potential and its behavioural

consequences in orphaned bumblebee workers (Bombus impatiens). Apidologie

46, 618–627. https://doi.org/10.1007/s13592-015-0351-4

Smith, A.R., Kapheim, K.M., Pérez-Ortega, B., Brent, C.S., Wcislo, W.T., 2013.

Juvenile hormone levels reflect social opportunities in the facultatively eusocial

sweat bee Megalopta genalis (Hymenoptera: Halictidae). Horm. Behav. 63, 1–4.

https://doi.org/10.1016/j.yhbeh.2012.08.012

Smith, B.H., Weller, C., 1989. Social competition among gynes in halictine bees: The

influence of bee size and pheromones on behavior. J. Insect Behav. 2, 397–

411. https://doi.org/10.1007/BF01068064

Stevenson, P.A., Schildberger, K., 2013. Mechanisms of experience dependent

control of aggression in crickets. Curr. Opin. Neurobiol.

https://doi.org/10.1016/j.conb.2013.03.002

Tibbetts, E.A., Huang, Z.Y., 2010. The Challenge Hypothesis in an Insect: Juvenile

Hormone Increases during Reproductive Conflict following Queen Loss in

Polistes Wasps. Am. Nat. 176, 123–130. https://doi.org/10.1086/653664

Trible, W., Kronauer, D.J.C., 2017. Caste development and evolution in ants: it's all

    about size. J. Exp. Biol. 220, 53–62. https://doi.org/10.1242/jeb.145292

Tsuji, K., Tsuji, N., 2005. Why is dominance hierarchy age-related in social insects?

    The relative longevity hypothesis. Behav. Ecol. Sociobiol. 58, 517–526.

    https://doi.org/10.1007/s00265-005-0929-3

Van Doorn, A., 1989. Factors influencing dominance behaviour in queenless

    bumblebee workers (*Bombus terrestris*). Physiol. Entomol. 14, 211–221.

    https://doi.org/10.1111/j.1365-3032.1989.tb00954.x

Van Doorn, A., Heringa, J., 1986. The ontogeny of a dominance hierarchy in

    colonies of the Bumblebee Bombus terrestris (Hymenoptera, Apidae). Insectes

    Soc. 33, 3–25. https://doi.org/10.1007/BF02224031

Wasielewski, O., Wojciechowicz, T., Giejdasz, K., Krishnan, N., 2011. Influence of

    methoprene and temperature on diapause termination in adult females of the

    over-wintering solitary bee, Osmia rufa L. J. Insect Physiol. 57, 1682–1688.

    https://doi.org/10.1016/j.jinsphys.2011.09.002

Wegener, J., Huang, Z.Y., Lorenz, M.W., Lorenz, J.I., Bienefeld, K., 2013. New

    insights into the roles of juvenile hormone and ecdysteroids in honey bee

    reproduction. J. Insect Physiol. 59, 655–661.

    https://doi.org/10.1016/j.jinsphys.2013.04.006

West-Eberhard, M.J., 1996. Wasp societies as micrcosms for the study of

    development and evolution., in: Natural History and Evolution of Paper-Wasps.

    pp. 290–317.

Wingfield, J.C., Hegner, R.E., Dufty, A.M., Ball, G.F., 1990. The "Challenge

    Hypothesis": Theoretical Implications for Patterns of Testosterone Secretion,

Mating Systems, and Breeding Strategies. Am. Nat. 136, 829–846.

https://doi.org/10.1086/285134

Withee, J.R., Rehan, S.M., 2016. Cumulative effects of body size and social

experience on aggressive behaviour in a subsocial bee. Behaviour 153, 1365–

1385. https://doi.org/10.1163/1568539X-00003382

Yerushalmi, S., 2006. Developmentally determined attenuation in circadian rhythms

links chronobiology to social organization in bees. J. Exp. Biol. 209, 1044–1051.

https://doi.org/10.1242/jeb.02125

Zanette, L., Field, J., 2009. Cues, concessions, and inheritance: Dominance

hierarchies in the paper wasp Polistes dominulus. Behav. Ecol.

https://doi.org/10.1093/beheco/arp060

1201