Title: Extensive and diverse patterns of cell death sculpt neural networks in insects.

Authors: Sinziana Pop[1], Chin-Lin Chen[2], Connor J Sproston[1], Shu Kondo[3], Pavan Ramdya[2,*] and Darren W Williams[1,4,*]

[1]Centre for Developmental Neurobiology, King's College London, New Hunt's House 4th Floor, Guy's Hospital Campus, London SE1 1UL, UK.

[2]Neuroengineering Laboratory, Brain Mind Institute and Institute of Bioengineering, École Polytechnique Fédérale de Lausanne, Lausanne CH-1015, Switzerland.

[3]Genetic Strains Research Center, National Institute of Genetics, Shizuoka 411-8540, Japan.

[4]Lead Contact

[*]Correspondence: darren.williams@kcl.ac.uk, pavan.ramdya@epfl.ch.

ABSTRACT

Changes to the structure and function of neural networks are thought to underlie the

evolutionary adaptation of animal behaviours. Among the many developmental phenomena

that generate change programmed cell death appears to play a key role. We show that cell

death occurs continuously throughout insect neurogenesis and happens soon after neurons are

born. Focusing on two dipterans which have lost flight during evolution we reveal that

reductions in populations of flight interneurons are likely caused by increased cell death

during development.

Mimicking an evolutionary role for increasing cell numbers, we artificially block

programmed cell death in the medial neuroblast lineage in *Drosophila melanogaster*, which

results in the production of 'undead' neurons with complex arborisations and distinct

neurotransmitter identities. Activation of these 'undead' neurons and recordings of neural

activity in behaving animals demonstrate that they are functional. Our findings suggest that

the evolutionary modulation of death-based patterning could generate novel network

configurations.

KEYWORDS

Programmed cell death, postembryonic neurogenesis, hemilineage, medial neuroblast,

octopamine, evo-devo, ventral nerve cord, *Drosophila melanogaster*, *Braula coeca*,

Crataerina pallida, *Thermobia domestica*

INTRODUCTION

Nervous systems are exquisitely adapted to the biomechanical and ecological environments

in which they operate. How they evolve to be this way is largely unknown. Such changes can

occur through modifications in receptor tuning, transmitter/receptor repertoires, neuronal

excitability, neuromodulation, structural connectivity, or in the number of neurons within

specific regions of the central nervous system (CNS). The differences seen in networks, over

an evolutionary timescale, ultimately result from heritable changes in developmental

processes (Horder, 1989). Advancing our knowledge of the mechanisms of neural

development using comparative approaches will help us understand how specific elements

can be modified, how new 'circuits' and behaviours evolve, and will ultimately lead to a

better understanding of how nervous systems function (Ramdya and Benton, 2010). Studies

comparing the nervous systems of mammalian species that occupy diverse ecological niches

reveal clear differences in the number of cells within homologous brain regions (Herculano-

Houzel et al., 2014). Such differences have occurred either through expansion or reduction of

specific cell populations, through changes in proliferation or apoptotic programmed cell

death (PCD) during development (Charvet et al., 2011). Most studies of nervous system

evolution have focused on stem cell identity and the role of differential proliferation

dynamics (Truman and Ball, 1998; Rakic, 2009; Biffar and Stollewerk, 2014). While one

recent study has elegantly shown a role for PCD in the evolution of peripheral olfactory

sensory neurons in drosophilids and mosquitoes (Prieto-Godino et al., 2020), how changes in

cell death can modify central circuits still remains an open question.

In insects the number and arrangement of neural progenitor cells that generate central

neurons (termed neuroblasts, NBs) is highly conserved despite a remarkable diversity of

insect body plans and behaviors (Wheeler, 1891; Bate, 1976; Hartenstein and Campos-

Ortega, 1984; Doe and Goodman, 1985; Tamarelle et al., 1985; Booker and Truman, 1987;

Truman and Bate, 1988; Shepherd and Bate, 1990; Doe, 1992; Truman, 1996; Truman and

Ball, 1998; Biffar and Stollewerk, 2014). In the ventral nerve cord (VNC – functionally

equivalent to the vertebrate spinal cord), the medial neuroblast (MNB) in particular stands

out as an unpaired NB located at the midline in the posterior end of each segment, while all

other NBs come in pairs arranged in a bilaterally symmetric array across the midline (**Figure

1A,B,C**). Its easy-to-locate position made the MNB identifiable in all developing insects

described from as early as 1891 by W.M. Wheeler (Wheeler, 1891), and spanning all insect

orders from wingless silverfish to locusts, beetles, moths, and flies (Bate, 1976; Hartenstein

and Campos-Ortega, 1984; Doe and Goodman, 1985; Tamarelle et al., 1985; Booker and

Truman, 1987; Truman and Bate, 1988; Shepherd and Bate, 1990; Doe, 1992; Truman and

Ball, 1998; Biffar and Stollewerk, 2014). The MNB gives rise to two populations of neurons,

one GABAergic and one octopaminergic, which are also homologous across insects (Rowell,

1976; Siegler et al., 1991, 2001; Thompson and Siegler, 1991; Campbell et al., 1995;

Stevenson and Spörhase-Eichmann, 1995; Siegler and Pankhaniya, 1997; Witten and

Truman, 1998; Jia and Siegler, 2002; Pflüger and Stevenson, 2005; Lacin et al., 2019). There

appears to be a relationship between cell number and function in these populations. Flying

insects have greater numbers of octopaminergic neurons within segments that control wings

(**Figure 1D**; Stevenson and Spörhase-Eichmann, 1995), while grasshoppers have more

GABAergic neurons in the fused metathoracic/abdominal ganglia, where they receive

auditory input from the abdomen (**Figure 1D**; Witten and Truman, 1998; Thompson and

Siegler 1991). Alongside differences in numbers of the same cell type between segments and

species, numbers of GABAergic and octopaminergic neurons found in one segment are never

equal (**Figure 1D,E**). This is especially intriguing as during development each GABAergic

neuron is a sister cell to an octopaminergic neuron, arising from one cell division and are

produced in equal number. (**Figure 1F**). The greater number of GABAergic cells in each

segment results from PCD targeting octopaminergic neurons in both grasshoppers (Jia and

Siegler, 2002) and fruit flies (Truman et al., 2010). The repeated elimination of one neuronal

sibling over the other is not restricted to the MNB lineage.

Figure 1. Median neuron numbers vary between insect species
(**A**) Cartoon of a young grasshopper embryo, here used as a depiction of a generic insect embryo (modified from Truman, 1996). Thoracic territory where one segments-worth of neuroectoderm generates an array of neuroblasts (grey box). (**B**) Schematic of neuroblast array showing bilaterally symmetric neuroblasts organised in 7 rows and 6 columns and a single median neuroblast (MNB) (modified from Truman, 1996). (**C**) Schematic of a lineage derived from a neuroblast (NB). Every NB buds off a ganglion mother cell (GMC) which undergoes a terminal division to generate 2 neurons with distinct cell fates, an A cell (black) and a B cell (grey). As the A and B cells result from a single division, one cannot be produced without the other. After several rounds of GMC divisions, a lineage produced by a single NB is composed of two half-lineages: 'hemilineage A' made up of all the A cells and 'hemilineage B' made up of B cells. Arrow indicates relative age, with newly-born cells located close to the NB. (**D**) Schematic showing summary of thoracic midline neuron population data from 5 orders of insects (colour coded in the phylogenetic tree and in the figure key bellow) including the primitive wingless firebrat *Thermobia domestica*, the cockroach *Periplaneta americana*, the cricket

Gryllus bimaculatus, the locust *Schistocerca gregaria*, the moth *Manduca sexta*, the fruit fly *Drosophila melanogaster*, the blowfly *Calliphora vicina*, the horse lousefly *Hippobosca equina* and the flightless bee louse fly *Braula coeca* and swift lousefly *Crataerina pallida*. The numbers of GABAergic neurons (black cells) and octopaminergic neurons (grey cells) produced by the MNB is given for each thoracic segment. Except for the moth, a higher number of octopaminergic neurons can be found in winged segments in flying insects (yellow boxes). Cell numbers in this homologous lineage vary both between segments and species. Data on firebrats, cockroaches, crickets, locusts, moths, fruit flies and blowflies are compiled from Monastirioti et al., 1995; Stevenson and Spörhase-Eichmann, 1995; Witten and Truman, 1998; Schlurmann and Hausen, 2003; Lacin et al., 2019; and unpublished data from Dacks, Pflüger and Hildebrand (AM Dacks, personal communication, May 2020), while data on horse, swift and bee louseflies are from our own work. **(E)** Cartoon of *Drosophila* mesothoracic midline lineage populations with 'hemilineage A' cells revealed by GABA immunoreactivity (black) and 'hemilineage B' revealed by octopamine immunoreactivity (grey). **(F)** Schematic of MNB during development with 'hemilineage A' cells (black) and 'hemilineage B' (grey). The first neurons in hemilineage B cells survive (they are the oldest). After this point, all hemilineage B cells are removed by PCD whereas 'hemilineage A' cells from the same GMC division are left intact. **(G)** Schematic representation of the pattern of hemilineage-specific cell death in one hemisegment in the mesothorax. Each circle represents one lineage produced by one NB. Numbers represent postembryonic lineage nomenclature established by Truman et al. (2004). Shade circle refers to doomed hemilineage. Lineage 0 is the postembryonic name given to the MNB lineage.

In the *Drosophila* embryo a first wave of neurogenesis generates the larval nervous system (Booker and Truman, 1987; Truman and Bate, 1988) after which the majority of NBs become quiescent. Following reactivation from quiescence NBs produce neurons throughout larval life until the early pupal stages. These postembryonic neurons - which make up most of the adult CNS – extend simple neuritic processes into the neuropil and stall until the pupal-adult transition when they grow complex arborisations, synapsing with their target cells (Truman, 1990). In the VNC, NBs bud off a ganglion mother cell (GMC) which undergoes a terminal division to generate 2 neurons with distinctly different cell fates (an A cell and a B cell). As the A and B cells result from a single division, one cannot be produced without the other. After several rounds of GMC divisions, a lineage produced by a single NB is composed of two half-lineages: 'hemilineage A' made up of all the A cells and 'hemilineage B' made up of B cells (**Figure 1C**). Hemilineages act as functional units in adult flies (Truman et al., 2004, 2010; Lin et al., 2010; Harris et al., 2015; Shepherd et al., 2016, 2019; Lacin et al., 2019). For example, in the MNB lineage, hemilineage A cells mature into GABAergic local interneurons while hemilineage B cells become efferent octopaminergic

neurons (**Figure 1F**). Our previous work showed that a common fate of postembryonic

neurons is PCD affecting approximately 40% of VNC hemilineages (**Figure 1G**) (Truman et

al., 2010), this is also seen in the brain (Kumar et al., 2009; Bertet et al., 2014). The pattern

of PCD is stereotypical and targets the same hemilineages across individuals. Taken together,

the breadth of PCD suggests its major role in shaping the final makeup of the adult nervous

system, while its stereotypy points towards a heritable genetic basis. We therefore propose

that changes in neural circuits may results from heritable alterations in the extent and pattern

of PCD in hemilineages.

Here we show an extensive role for PCD in patterning the insect CNS, from primitive

firebrats to the most derived true flies. We see evidence of hemilineage-specific PCD in

primitively wingless firebrats and hippoboscid louseflies, suggesting that it is deployed

widely. We search for changes in hemilineage programs of cell death during the evolution of

flightlessness in true flies *Braula coeca* (the bee louse) and *Crataerina pallida* (the swift

louse), and show that PCD is likely responsible for reductions in flight hemilineages within

thoracic networks. Mimicking an evolutionary role for reducing PCD, we show that blocking

PCD in the octopaminergic hemilineage produced by the MNB in *Drosophila melanogaster*

results in mature differentiated 'undead' neurons that survive into adulthood, elaborate

complex arborizations and are functional. Using new tools, we also show that this PCD takes

place early, very soon after neurons are born. Our work highlights the importance of viewing

hemilineages as functional units of neurodevelopment in all insects and shows that their

alteration through an early mode of PCD can lead to adaptive changes in central circuits

during evolution.

RESULTS

Early mode of neuronal cell death is found in many different insects.

Because PCD selectively eliminates hemilineages in the fruit fly, and has been reported to

kill off immature octopaminergic neurons produced by the MNB in the grasshopper (Jia and

Siegler, 2002), we wondered if PCD occurs in the development of the CNS in a 'primitive'

wingless insect. Using TUNEL labelling in the firebrat *Thermobia domestica* (**Figure 2A,B**),

we found dying cells close to many NBs in all thoracic neuromeres at 50-55% of embryonic

development (**Figure 2C,D,E**). We also looked for evidence of PCD during VNC

development in the flightless dipteran *Crataerina pallida*, the swift lousefly (**Figure 2F**).

Crataerina is a viviparous haematophagous ectoparasite of the swift *Apus apus* (Hagan,

1951; Bequaert, 1952; Hutson, 1984; Walker and Rotherham, 2010a; Walker and Rotherham,

2010b). We focused on the swift lousefly, firstly to ascertain if the loss of flight during

evolution resulted in increased PCD of flight hemilineages, and secondly, as it is a dipteran

fly, we anticipated that we could use our specialist knowledge and immunohistochemical

tools established in *Drosophila*.

164   The swift lousefly is viviparous, females carry a single larva inside their abdomen

where it feeds on 'milk' from fatty glands until pupariation, when the mother deposits the

prepupa just before hardening of its pupal case (**Figure 2F,G**). Similar to the tsetse fly

(Truman, 1990), neurodevelopment is significantly delayed compared to typical dipteran flies

- the nervous system only acquires dipteran larval features many days after pupariation

(**Figure 2H**). Using EdU to label proliferating cells and immunostaining for active Dcp-1

(**Figure 2H,I,J**), we found dying cells located close to NBs throughout the 24 days of pupal

neurogenesis (manuscript in preparation). In the MNB lineage, which is easily identified by

its medial position and projection pattern in the neuropil, we found cell death in thoracic

segments at all time points examined, from day 4 after pupariation to day 23 (**Figure 2I,J**).

Figure 2. Cell death during neurogenesis is found in many insects.

(**A**) Adult firebrat. Scale bar, 1mm. (**B**) Maximum intensity projection of DAPI staining in a wholemount firebrat embryo (*Thermobia domestica*) at 50-55% of embryonic development. Scale bar, 100 µm. (**C**) Dying cells in the thoracic VNC of a firebrat embryo labelled using TUNEL (magenta) and Syncrip (white) antibodies. Syncrip was used here as a proxy for Neuroglian staining to reveal lineages. Scale bar, 50 µm. (**D**) Dying cells (arrowheads) are located close to NBs (*). Scale bar, 10 µm. (**E**) Dying cells (magenta, TUNEL) are located in the cortex of the VNC, where neurogenesis takes place, and not in the neuropil (green, Phalloidin) in a firebrat embryo. Scale bar, 50 µm. (**F**) Dorsal view of an adult swift lousefly with vestigial wings (top). Ventral view of a female pregnant

with a prepupa (bottom). Scale bars, 1 mm. **(G)** Swift lousefly larva (top) and day 30 swift lousefly pupa removed from its puparial case (bottom). Scale bars, 1 mm **(H)** EdU labels proliferating cells and antibody-labelling for active Dcp-1 reveals dying cells in the CNS of a swift lousefly pupa 9 days after pupariation. Scale bar, 50 µm. **(I)** Dying cells in lineage 0 labelled with antibodies for active Dcp-1 are located close to proliferating cells (e.g., NB*) throughout neurogenesis at Day 5 (top left), Day 9 (top right), Day 16 (bottom left) and Day 23 (bottom right) after pupariation. Scale bar, 10 µm. $n = 1$ each. **(J)** The occurrence of active Dcp-1-positive cells in lineage 0 in T1, T2 and T3 from Day 4 to Day 23 after pupariation in swift lousefly pupae ($n = 1$ each). Each black box indicates one occurrence. **(K)** Wild-type octopaminergic neurons in hemilineage 0B in a *Drosophila melanogaster* VNC labelled with CD8::GFP driven by *TDC2-GAL4* (orange arrowheads). Scale bar, 50 µm. **(L)** Octopaminergic neurons in hemilineage 0B in a swift lousefly VNC labelled with antibodies for tyramine β-hydroxylase (TBH, orange arrowheads). Fluorescence in the neuropil is derived from secondary antibodies trapped in the tracheal system and does not mark the true presence of TBH protein. Scale bar, 50 µm. **(M)** Quantification of T2/T1 number of octopaminergic neurons in fruit flies and swift louseflies shows that swift louseflies have lost the T2-specific higher numbers typical of flying dipterans ($P* = 0.012$, Mann-Whitney. $n = 7$ fruit flies, $n = 3$ swift louseflies). Bars represent mean ± standard deviation. **(N)** Cluster of cell bodies belonging to hemilineage 0A (dashed outline) labelled with antibodies for GABA and **(O)** cell bodies belonging to hemilineage 0B labelled with TBH antibodies in the prothorax (T1) of a swift lousefly.

Similar to cockroaches, which are not efficient fliers (Eckert et al., 1992), the flightless swift louseflies have the same number of octopaminergic neurons in each thoracic segment (see **Figure 1D**). In flying insects these octopaminergic neurons are required for flight maintenance and initiation (Roeder, 2005) and display variability, with more neurons found in winged segments (Stevenson and Spörhase-Eichmann, 1995). We labelled lousefly octopaminergic neurons from hemilineage 0B using antibodies for tyramine β-hydroxylase (Monastirioti et al., 1996) and by comparing the ratio of octopaminergic cells in the mesothorax (winged segment) and the prothorax (lacks wings), we found that, unlike flying dipterans, the swift lousefly has lost segment-specific variability of cell numbers (fruit fly [2.2±0.2, n = 7] versus swift lousefly [1.1±0.2, n = 3], p = 0.012 , Mann-Whitney U = 0, Mann- Whitney t-test) (**Figure 2K,L,M,N,O**). In the sister hemilineage 0A we found a considerably larger number of GABAergic neurons (**Figure 2N**), suggesting that PCD is responsible for the selective elimination of the octopaminergic hemilineage.

Our observations in firebrats and louseflies suggest that PCD during neurogenesis may be a universal and ancestral feature that sculpts the nervous system of all insects. To

further explore if changes in PCD have been deployed during evolution to accommodate

adaptive modifications to behaviour, we next searched for evidence of increased PCD in

other flight hemilineages of flightless dipterans.

Changes in hemilineage programs of PCD in flightless dipterans suggest adaptive

modifications to neural circuits

To explore the possibility that changes in the pattern and/or extent of PCD are adaptive, we

looked for evidence of evolutionary modifications in the VNC of other species. We

specifically asked whether lineages known to function in flight circuitry might be modified in

flightless insects. Using antibodies for Neuroglian, we compared homologous hemilineages

involved in the flight circuits of two flightless dipterans, the swift lousefly *Crataerina pallida*

and the bee lousefly *Braula coeca* (**Figure   3A,B,C,D**). *Braula*, a close relative of

drosophilids, is wingless, lacks halteres and has an extremely reduced thorax (**Figure 3A**).

Bee louseflies spend their entire adult life as kleptoparasites on the honeybee *Apis mellifera*

(Imms, 1942; McAlister, 2018).

Figure 3. Neuronal cell death sculpts the wing circuitry of wingless dipterans.

(**A**) Dorsal (top) and side view (bottom) of an adult bee lousefly with a reduced thorax lacking wings and halteres. Scale bar, 1 mm. (**B-D**) Hemilineage VNC fiber tracts labelled with Neuroglian in a fruit fly (**B**), a swift lousefly (**C**) and a bee lousefly (**D**). Shown are hemilineages 6A and 7B which are reduced in the bee lousefly in all three thoracic segments (T1, T2, T3). Scale bars, 50 μm. (**E-G**) 3A and 3B hemilineage fiber tracts labelled with Neuroglian in a fruit fly (**top left**) and a bee lousefly (**top right**). Shown are maximum intensity projections, chosen to best display hemilineages, from cross-section T1 (**E**), T2 (**F**) and frontal perspectives T3 (**G**). Quantifications of 3A/3B hemilineage bundle diameter ratios in fruit flies and bee louseflies are given below ($P**** < 0.0001$ in T1, independent samples t-test, $P*** = 0.0001$ for T2, Welch's t-test, $P* = 0.0044$, Welch's t-test. $n = 7$ fruit flies each, $n = 8$ bee louseflies each). Bars represent mean ± standard deviation.

Comparing the neuroglian-labelled axon bundle width of homologous hemilineages

between flightless and flying species, we found that hemilineages 3B, 5B, 6A, 7B, 11B, 12A

and 19B, which innervate the wing neuropil (Shepherd et al., 2019), are reduced in bee

louseflies, but not in swift louseflies (**Figure 3B,C,D**). Among these, 3B, 6A, 11B, 12A and

19B belong to lineages in which both hemilineages survive in fruit flies (see schematic in

Figure 1G). A difference in axon bundle diameter between sister hemilineages could indicate

a difference in cell number, possibly established by PCD during development. We chose to

quantify the ratio of axon bundle diameters in lineage 3 because we expected a reduction in

hemilineage 3B, which innervates the wing neuropil, but no change in hemilineage 3A,

which projects into the leg neuropil (Shepherd et al., 2019). The fiber tracts of 3A and 3B

originate as a common bundle and split only in the intermediate neuropil. After they split, the

sister fiber tracts sit in the same plane before they defasciculate, making them easy to trace

and compare (**Figure 3E,F,G**). We found that the ratio between sister hemilineages A and B

is significantly higher in bee louseflies compared to fruit flies, indicating that hemilineage

3B, which controls flight-related behaviours, is severely reduced in these flightless flies (T1:

fruit fly [0.6 ± 0.1, n = 7] versus bee louse [1.8 ± 0.4, n = 8], $p < 0.0001$, t = -8.084,

independent samples t-test; T2: fruit fly [0.7 ± 0.1, n = 7] versus bee louse [1.7 ± 0.4, n = 8], p =

0.0001 , F = 42.22, Welch's t-test; T3: fruit fly [1.1 ± 0.4, n = 7] versus bee louse [1.7 ± 0.2, n =

8], p = 0.0044 , F = 14.84, Welch's t-test). Even though we cannot make a precise inference

of cell numbers in each hemilineage, as the fiber tract of hemilineage 3B appears frayed

while 3A is more compact in both species, our results clearly show there has been an adaptive

reduction of 3B, associated with the loss of flight machinery.

Blocking programmed cell death in *Drosophila* generates identifiable, differentiated

populations of undead neurons

After observing the outcome of increased PCD in hemilineages consistent with adaptations to

the loss of flight in bee louse and swift louse, we wondered if, conversely, by only reducing

PCD, we could generate novel functional expansions of a hemilineage. To explore this, we

made use of the powerful genetic tools available in *Drosophila* to block PCD in the MNB

lineage to determine if 'undead' cells survive into adulthood, elaborate their neurites and

acquire a distinctive neurotransmitter identity.

281   From our previous work (Truman et al., 2010), we know that during postembryonic

neurogenesis MNB hemilineage A survives, expresses Engrailed (Truman et al., 2004) and

differentiates into GABAergic interneurons (Lacin et al., 2019). Because during embryonic

development the MNB hemilineage B produces a small number of octopaminergic neurons,

we hypothesised that preventing PCD would generate additional octopaminergic neurons in

the later postembryonic phase of neurogenesis. In postembryonic nomenclature, neurons

generated by the MNB are collectively called lineage 0 and therefore we will refer to

octopaminergic neurons generated by the MNB as hemilineage 0B.

289   Using the octopaminergic neuron driver, *TDC2-GAL4,* we observed a 4 to 9-fold

increase in the number of octopaminergic neurons in the thoracic VNC of *H99/XR38* adult

flies deficient for proapoptotic genes ($hid^{+/-}$, $grim^{+/-}$, $rpr^{-/-}$ and $skl^{+/-}$) (White et al., 1994;

Peterson et al., 2002) compared with wild-type control animals (**Figure 4A,B,C,D**), (T1: 20.9

± 2.3, Mann-Whitney U = 0, p = 0.0002; T2: 26.3 ± 4.4, Mann-Whitney U = 0, p = 0.0004;

T3: 27.5 ± 3.5, Mann-Whitney U = 0, p = 0.0004; n = 11 each). These 'undead' neurons also

express the vesicular glutamate transporter VGlut (**Figure 4E,F**), just like wild-type

octopaminergic neurons (Greer et al., 2005).

297   Ideally, to label and manipulate dying neurons from hemilineage 0B, we require a

specific driver line expressed only in the newly-born doomed neurons. To test if we could use

TDC2-GAL4 to label and manipulate dying neurons from hemilineage 0B during their

development, we performed a timeline of expression in wild-type and *H99/XR38* flies

(**Figure S1**). Unfortunately, even though undead neurons are generated from L2 onwards in

H99/XR38 flies, the *TDC2-GAL4* is only active in the undead cells days later. Gradually, in

pupae, TDC2-GAL4 expression reveals the remaining undead B cells (**Figure S1C,D**). We

concluded that the TDC2 driver line cannot be used to visualize and manipulate newly-born

postembryonic 'doomed cells'. Instead, TDC2-GAL4 allowed us to accurately reveal

'undead' hemilineage 0B neurons *but* only in the adult (see cartoon **Figure 4G**).

Figure 4. Blocking death results in differentiated undead neurons in the medial neuroblast
lineage
(**A**) CD8::GFP expression driven by *TDC2-GAL4* in octopaminergic neurons from hemilineage 0B in
the VNC of wild-type (WT, left) and PCD-blocked adult flies (*H99/XR38* deficient for $hid^{+/-}$, $grim^{+/-}$,
$rpr^{-/-}$ and $skl^{+/-}$, right). Scale bar, 50 µm. (**B-D**) Quantifications of the number of *TDC2-GAL4*-positive
octopaminergic neurons in the VNC of WT and *H99/XR38* adult flies. Bars represent mean ± standard
deviation. (**B**) ****P* = 0.0002 in T1, (**C**) ****P* = 0.0004 in T2, (**D**) ****P* = 0.0004 in T3, Mann-
Whitney. Scale bar, 10 µm. *n* = 11 each. Mann-Whitney. Scale bar, 10 µm. *n* = 11 each. (**E**)
Antibodies for the vesicular glutamate transporter VGlut and (**F**) GFP expression driven by the
glutamatergic driver line *OK371-GAL4* label both WT and undead (*H99/XR38*) octopaminergic
neurons. Orange arrowheads indicate cell bodies. Scale bars, 10 µm (VGlut), 50 µm (OK371). (**G**)
Schematic of TDC2-GAL4 expression in postembryonic lineage 0 in wildtype and *H99/XR38* third
instar larvae and adults (left panels). Postembryonic hemilineage B populations only start expressing
TDC2-GAL4 in early pupal development and maintain it throughout adult life. MARCM mosaic
clones that are homozygous for a null *Dronc* allele lack GAL80. These show robust expression of
GAL4 in small numbers of surviving postembryonic hemilineage B cells (right panel). In adult WT
and *H99/XR38* flies, GFP is expressed in both embryonically-born and postembryonic TDC2-positive
neurons, while in MARCM flies GFP is *only* present in postembryonic cells. (**H-I**) CD8::GFP
expression driven by *TDC2-GAL4* in WT (**G**) and *H99/XR38* (**H**). WT and undead primary neurites

project dorsally and branch extensively in the dorsal neuropil. In WT neurons the primary neurite
bifurcates at the dorsal midline, while undead neurons are unable to bifurcate and turn to one side
(orange arrowheads). In *H99/XR38* flies which contain both WT and undead neurons, the primary
neurite to one side is thicker (orange arrowhead). Both WT and undead neurons join thoracic nerves
(white arrows). Scale bars, 50 µm. **(J)** Reconstructed arborizations of undead neurons expressing
CD8::GFP driven by *TDC2-GAL4* in flies bearing MARCM clones homozygous for the loss of
function allele *dronc$^{\Delta A8}$* (in which PCD is blocked). The 3D-rendered image is tilted at a 45° angle.
Undead neurons have somata that are located at the ventral midline (orange arrowhead), branch
extensively in the neuropil (green), have a turning primary neurite (orange), and project to the
periphery through a thoracic nerve (arrow). Scale bar, 50 µm.
338

To ensure sparse labelling and precise manipulation of doomed cells from

hemilineage 0B, we generated postembryonic *TDC2-GAL4*-expressing MARCM clones

homozygous for the loss-of-function allele *Dronc$^{\Delta A8}$* (in which PCD is inhibited) (Kondo et

al., 2006; Truman et al., 2010). This strategy guarantees that, even though cell death can be

rescued in other lineages, it is only *TDC2*-positive postembryonic neurons that express UAS-

based tools.

Analysis of the projection patterns of undead neurons revealed that they display both

common and distinct features compared to their wild-type embryonically-born counterparts.

Similar to wild-type octopaminergic cells (Monastirioti et al., 1995), the undead neurons

have cell bodies located ventrally at the midline, at the posterior border of the thoracic

segment (**Figure 4H,I,J**), project a primary neurite in the dorsal-most region of the neuropil,

the tectulum (Court et al., 2017), and join thoracic nerves (Pauls et al., 2018) (**Figures**

4H,I,J and **S2**). Unlike wild-type cells which bifurcate and branch extensively in the

tectulum, the primary neurite of undead neurons fails to bifurcate, branches in both dorsal

and ventral regions of the neuropil and sends projections to neighbouring segments (**Figures**

4J and **S2**). As we describe in **Figure S1**, a few wild-type octopaminergic neurons are

produced in all thoracic segments during postembryonic neurogenesis in lineage 0. We

propose that the very few bilateral projecting neurons we encounter in our clones are wild-

type cells. To avoid any uncertainty when performing our behavioural experiments (below),

we excluded flies which contained a bifurcating neuron in undead MARCM clones. Thus,

using MARCM clonal approaches we show that undead neurons in hemilineage 0B become

octopaminergic, elaborate complex neurites and join thoracic nerves.

Undead neurons are functional and integrate into motor networks

We next asked if these differentiated undead neurons are functional. To address this, we

tested if activating undead neurons with the warm temperature-gated ion channel TrpA1 in

headless adult *Drosophila* could elicit behaviours (**Figure 5 and Video S1**). In *TDC2-GAL4*

positive control flies (expressing TrpA1 in wild-type embryonic-born octopaminergic

neurons), thermogenetic stimulation induced long bouts of locomotion (**Figure 5C,D** and

Video S1). Importantly, we found *UAS-TrpA1* negative controls (in the absence of a GAL4)

and MARCM control flies (containing no GAL4-positive clones) did not walk in response to

temperature elevation (**Figure 5B,C,D** and **Video S1**). We found that the activation of

undead neurons expressing TrpA1 caused decapitated males to walk (**Figures 5B,C,D,E** and

Video S1). These data are consistent with the observation that octopamine applied to the

exposed anterior notum of decapitated flies causes walking (Yellman et al., 1997) and

suggests that undead neurons are functional and capable of releasing neurotransmitters in the

CNS. The extent of walking was greater in positive control flies than in the undead MARCM

condition (compare panels in **Figure 5C** and **Video S1**). This is likely because, alongside

activating octopaminergic neurons in the VNC, in positive controls we also stimulated the

severed axons of octopaminergic cells in the brain which send descending projections to the

VNC.

Figure 5. Undead neurons are functional: thermogenetic activation of undead neurons induces walking in decapitated *Drosophila*.

(**A**) Schematic of behavioural assay with TrpA1 activation. Decapitated flies are placed on a thermoelectric plate connected to a DC power supply, exposed to a temperature ramp (right panel), and filmed from above using a digital camera. (**B**) Examples of a stationary MARCM Control fly and a walking fly with undead neurons. Images represent maximum intensity projections of 13 frames at 0.3 fps tracing the centroid over time (orange line). Scale bar, 5 mm. (**C**) Fly body tracks generated by identifying the geometric centre of the fly body in each frame and storing the centre coordinates, plotted as a continuous line, one for each fly (walking or stationary) for negative controls (*UAS-TrpA1*), positive controls (*TDC2>TrpA1*), MARCM control flies and flies with MARCM clones of Undead neurons. Each trace represents one individual fly. Scale bar, 5 mm. (**D**) Quantification of the percentage of flies that walked per experimental group. $P^{***} = 0.0002$ for negative versus positive control, $P^{***} = 0.0002$ for positive controls versus MARCM Control, $P^* = 0.0242$ for MARCM control versus undead neurons, $P^* = 0.0135$ for negative control versus undead neurons, Pearson's chi-squared corrected for multiple comparisons using a Bonferroni correction. $n = 19$ for negative controls, $n = 18$ for positive controls, $n = 22$ for MARCM control, $n = 17$ for undead neurons. n numbers for each group are given below and the number of flies which walked is shown above each bar. (**E**) Quantification of the number of walking undead neuron flies split into 3 anatomical subgroups according to the location of MARCM clones in T1, T2, or T3. $P^{ns} = 0.2628$, Pearson's chi-squared. $n = 6$ for T1, $n = 4$ for T2, $n = 7$ for T3. Numbers at the base are the number of walking flies. The percentage is shown above each bar.

To determine if undead neurons are integrated into thoracic motor circuits, we

recorded the activity of mixed undead and wild-type octopaminergic neuron populations

expressing GCaMP6s (an activity reporter) and tdTomato (an anatomical fiduciary) in intact

H99/XR38 flies during tethered behaviour on a spherical treadmill (Chen et al., 2018) (**Figure**

6A,B). In these animals we observed conspicuous increases in neural activity during air-puff-

induced walking in both wild-type controls (**Figure S3**) and in *H99/XR38* flies (**Figure**

6C,D,E and **Videos S2-4**). Because undead neurons outnumber their wild-type counterparts

by a ratio of 6.5 to 1, these results imply that both neuronal types are active in *H99/XR38*

flies. Supporting this, we observed an increase in GCaMP6s fluorescence across all

subregions along the width of the primary neurite bundle (**Figure S4** and **Video S4**). Taken

together these data reveal that 'undead' neurons in the adult fly are functional and integrate

into motor networks.

Figure 6. Undead neurons integrate into VNC networks: undead neurons are active during
naturalistic behaviours in intact adult *Drosophila*.

(**A**) Schematic of the dorsal thoracic dissection and approach for ventral nerve cord functional
imaging in tethered, adult flies. (**B**) Location of the imaging region of interest (grey box) with respect
to a schematic of the adult CNS. (**C**) Raw 2-photon image of *TDC2-GAL4-positive* neurons co-
expressing tdTomato (red) and GCaMP6s (cyan) in *H99/XR38* flies (**left**). Region-of-interest used to

calculate %ΔR/R is outlined (yellow). Standard deviation z-projection of a dorsal-ventral image stack
of the functional imaging region-of-interest in **b** (**right**). Scale bar, 50 µm. **(D)** Representative
behavioural and functional imaging data in *H99/XR38* flies. Shown are: CO_2 stimulation (black),
%ΔR/R (ratio of GCaMP6s / tdTomato) signal (green), and ball rotations indicating forward walking
(red), sideways walking (blue), and turning (purple). The behaviour of the fly was classified as either
walking (light blue), or resting (gray) by applying a threshold on ball rotation speed. **(E) (left)**
Individual (green) and average (black) %ΔR/R traces within each behavioural epoch for walking (n =
82) and resting (n = 86) events processed from 750 s of imaging data . Solid green circles indicate the
end of a behavioural epoch. The average trace (black line) was calculated for only periods with 4 or
more traces. (**right**) Normalised average %ΔR/R traces for three different flies during walking and
resting. The average (black) trace is the same as in the left panel.

An early and rapid mode of developmental cell death eliminates significant numbers of

newly-born neurons throughout postembryonic development in *Drosophila*

After observing the potential for blocking PCD to incorporate new neurons, we wondered

which specific type of PCD is responsible, reasoning that our knowledge of how nervous

systems evolve can only be enriched by pinpointing the exact developmental processes

involved. The majority of studies on neuronal PCD in insects have focused on its role at

metamorphic transitions, where it eliminates fully differentiated neurons either at puparium

formation (Truman et al., 1994) or in adults post-eclosion (Kimura and Truman, 1990;

Draizen et al., 1999). Both of these remodelling events are gated by ecdysteroids. However,

finding dying cells close to NBs in firebrats and swift louseflies, combined with similar

observations in *Drosophila* (Truman et al., 2010), made us consider that hemilineage-specific

PCD takes place early, in newly-born neurons. This has been difficult to evaluate on a cell-

by-cell basis within a complex nervous system.

To interrogate the postembryonic neuronal death we see here, we built a novel

genetically encoded effector caspase probe called SR4VH (Figure 7A,B). SR4VH consists of

a membrane-bound red fluorescent protein (Src::RFP) and a yellow fluorescent protein with a

strong nuclear localisation signal from histone H2B (Venus::H2B) linked together by four

tandem repeats of the amino acid sequence DEVD. When effector caspases cleave the DEVD

site, Venus accumulates in the nucleus while RFP remains bound to the cell membrane

(Figure 5B). This reporter is similar in design to Apoliner (Bardet et al., 2008), but has

different subcellular localization signals as well as four tandem caspase cleavage sites instead

of one. We also found that tethering the probe to the membrane with the myristoylation

signal from Src means that there is no excess signal accumulation in the Golgi apparatus

(Mukherjee et al., in prep). The nuclear localisation signal from H2B allows for highly

efficient sequestration of cleaved Venus in the nucleus even in late stages of apoptosis, when

the nuclear membrane is likely compromised.

Using the GAL4/UAS system and the neuroblast driver *Worniu-GAL4*, we found we

could visualize postembryonic neurogenesis and label up to 20 of the most recently born

progeny from a single NB (this is due to GAL4 and reporter perdurance). The number of

progeny we can detect at any one time using *Worniu-GAL4* varies from 10 to 20, most likely

as a result of differential proliferation rates across lineages. We confirmed that SR4VH is

reliable as a reporter for cell death in larvae by analysing its expression pattern in all lineages

of postembryonic neurons in the thoracic VNC and comparing it to our previous work on

MARCM homozygous mutant clones of the initiator caspase *Dronc* (Truman et al., 2010).

(**Figures 7C,D,E,F and S5**). We found dying cells associated with lineages in the brain and

VNC throughout the whole of postembryonic neurogenesis (**Figures 5D,E,F**), which lasts for

3.5 days, from mid-2^{nd} instar (L2) to 12h after pupariation. As previously suggested (Truman

et al., 2010), the time course of PCD indicates that cells die early – very soon after they are

born – often before they have even extended a neuritic process. This death appears to be

unlike the 'trophic' PCD found in vertebrates, where a neuron extends a process, interacts

with its target cell and dies in the absence of appropriate survival signals. In support of an

early onset of PCD, we were able to see sequential stages of cell death, dependent on the

distance from the NB (**Figures 7G,H,I** and **S5B**). Older cells located further away from the

NB appear to be at a more advanced stage of PCD indicated by the complete translocation of

Venus from the membrane to the nucleus (Cell 3 in **Figure 7G,H**) and by the accumulation

of RFP-positive dead cell membranes close to the lineage bundle (arrowheads in **Figure**

7E,F).

Figure 7. Early neuronal cell death occurs throughout postembryonic development.

(**A**) Simple schematic of PCD in *Drosophila*. Elements disrupted in this study, or used as a PCD
readout are underlined. (**B**) Schematic of effector caspase reporter SR4VH (top) and idealised
fluorescence patterns in two cells with distinct caspase activity (bottom): RFP (magenta) and Venus
(green) present in the cell membrane (Line 1) versus RFP at the cell membrane and Venus
accumulation in the nucleus of a cell undergoing programmed cell death (PCD) (Line 2). (**C-E**)
SR4VH driven by *Worniu-GAL4* reveals dying cells in the central nervous system (CNS) throughout
postembryonic neurogenesis in a 2nd instar (**C**, n = 9), early 3rd instar (**D**, n = 8) and wandering 3rd
instar (**E**, n = 18) larva. Right panels show examples of lineages with dying cells located close to the
NB (*) in the brain (top) and ventral nerve cord (VNC, bottom). Arrows indicate dying cell,

arrowheads indicate RFP-positive dead cell membranes. Scale bars, 50 µm (left panels), 10 µm (right panels). **(F)** SR4VH driven by *Worniu-GAL4* reveals younger cells at earlier stages of cell death (Cell 2) are located closer to the MNB than older cells at later stages of cell death (Cell 3), which are closer to the lineage bundle (arrowhead). Image represents a single optical section. Scale bar, 10 µm. **(G)** Fluorescence intensity profiles (normalised to the maximum value along the line for each channel) plotted along the lines indicated in **(F)**. **(H)** Schematic representing successive stages of cell death correlated with distance from NB and cell age in doomed lineages as reported with SR4VH. **(I)** Revealing markers for lineage 0. Immature hemilineage A progeny revealed with anti-Engrailed (En, blue – arrowheads) and immature hemilineage B progeny labelled with anti-Vestigial (Vg, green – arrows) in a third instar *Worniu-GAL4; UAS- CD8::GFP* larva (white). Expression of markers is mutually exclusive. Scale bars, 10 µm. **(J)** SR4VH driven by *Worniu-GAL4* together with antibodies for Engrailed (En, blue) reveal that only Engrailed-negative cells from hemilineage 0B undergo PCD (white arrowheads) during postembryonic development in the thoracic VNC. Scale bar, 10 µm. *n* = 6.

The number of dying cells within a doomed lineage varied from 1 to 8, with most lineages containing 1-2 dying cells from a total of 10-20 cells labelled with *Worniu-GAL4* (dying cells/lineage: 1.3 ± 1.5 given as average ± standard deviation; n = 444 doomed lineages from 5 VNCs). From a total of 444 doomed lineages, 243 harboured more than one dying cell, of which 148 displayed a progression of cell death (**Figure 7G,H**). Truman and Bate approximated the cell cycle of an NB to 55 minutes and that of a GMC to 6.5 hours, with 7 GMCs present in a proliferating lineage at all times (Truman and Bate, 1988). Therefore, after subtracting the NB and GMCs from clusters of 10-20 *Worniu-GAL4*-labelled cells, 2-12 will be neurons which resulted from 1-6 divisions, each separated in time by 55 minutes. This means that PCD was initiated early, at some time between 0 and 5.5 hours after neurons were born.

To look at death during the development of the MNB lineage (lineage 0) we imaged SR4VH in wandering L3 larvae and used molecular markers to identify members of hemilineage 0A and 0B. The transcription factors Engrailed/Invected (En/Inv) are known to be expressed in immature and fully differentiated interneurons of hemilineage A (Truman et al., 2004; Lacin et al., 2014, 2019; Allen et al., 2020). As previously reported, the mature differentiated octopaminergic neurons found in hemilineage B express the transcription factor Vestigial (Vg) (Landgraf et al., 2003). Here we find that a small number of immature

postembryonic neurons (about 3-5) in close proximity to the MNB also express Vg (**Figure**

7J.). Within these immature neurons the expression of Vg and En are mutually exclusive.

Using *Worniu-GAL4* to drive SR4VH we found that only the engrailed-negative cells are

undergoing apoptosis (**Figure 7K**), i.e. the same small number of cells that express Vg. Their

proximity to the MNB suggests that Vg-positive B cells (i.e. immature octopaminergic

neurons) undergo an early death, very soon after they are born.

DISCUSSION

Midline neurons show hemilineage-based variations in different species

536    The midline neurons within the VNC of insects have long been a source of interest

because they are homologous across species, yet show a diversity in cell numbers correlated

with body form and function (Stevenson and Spörhase-Eichmann, 1995; Witten and Truman,

1998; Lacin et al., 2019). For example, flying insects have greater numbers of midline

octopaminergic neurons within segments that control wings (Konings et al., 1988; Siegler et

al., 1991, 2001; Eckert et al., 1992; Stevenson et al., 1992; Spörhase-Eichmann et al., 1992;

Thompson and Siegler, 1993; Campbell et al., 1995; Monastirioti et al., 1995; Siegler and

Pankhaniya, 1997; Jia and Siegler, 2002; Schlurmann and Hausen, 2003), while grasshoppers

have more GABAergic neurons in the metathoracic/abdominal ganglia, which receives

auditory input (Witten and Truman, 1998; Thompson and Siegler 1991). These two midline

neuronal populations are derived from the same neuroblast, the MNB, which buds off

multiple GMCs, each dividing once to generate a GABAergic (A cell) and an octopaminergic

(B cell) neuron. Both A and B neurons are generated in equal numbers but, in all cases, the

numbers of GABAergic ('hemilineage A') and octopaminergic ('hemilineage B') neurons

within one segment are never the same. The greater numbers of GABAergic cells within each

segment has been shown in grasshoppers (Jia and Siegler, 2002) and fruit flies (Truman et al.,

2010) to be the result of removal by PCD of large numbers of cells from hemilineage B.

As suggested by our previous work, the 'hemilineage' emerges as a discrete

developmental unit that shows common features of gene expression and function (Truman et

al., 2010; Harris et al., 2015; Lacin and Truman, 2016; Lacin et al., 2019; Shepherd et al.,

2019). Therefore, we refer here to the PCD found in the MNB lineage of grasshoppers as

'hemilineage-specific PCD'. Following our observations of PCD during development in other

insects, together with a vast body of knowledge on homology in insect nervous systems

(Thomas et al., 1984; Kutsch and Breidbach, 1994), we suggest that variations in neural

circuits between species is very likely set up by modifying hemilineages, with PCD playing a

major role.

Cell death during neurogenesis is widespread across insects

Our data show that PCD is extensive and widespread during neurogenesis in the CNS of

insects, from the primitive firebrats *Thermobia domestica*, to true flies *Drosophila*

melanogaster and the swift lousefly *Crataerina pallida.*

We wondered whether alterations in cell death may have contributed to adaptations in the

VNC of flightless dipterans and found that a greater extent of PCD in the MNB lineage is

responsible for abolishing the segment-specific difference in octopaminergic cell numbers in

the swift lousefly *Crataerina pallida* – which has lost flight and gained adaptations to a

parasitic lifestyle (Hagan, 1951; Bequaert, 1952; Hutson, 1984; Lehane, 2008; Walker and

Rotherham, 2010b; Petersen et al., 2018). As octopaminergic neurons are involved in flight-

related behaviours (Duch and Pflüger, 1999; Roeder, 2005; Brembs et al., 2007), we suggest

that PCD has been co-opted in this lineage during the evolution of flightlessness in the swift

louse. We believe that this PCD takes place early, in newly born neurons, as we have seen

dying cells close to the MNB in all of our 20 pupae, from day 4 to day 23. Therefore, the

decrease in octopaminergic cell number in the mesothoracic segment of swift louseflies is

likely the result of increased hemilineage-specific PCD during evolution.

Hemilineage-specific reductions in fiber diameter in the bee lousefly

In the wingless bee lousefly *Braula coeca,* we found clear reductions in the thickness of fibre

tracts in several hemilineage bundles which in *Drosophila* are associated with flight-related

behaviours (Harris et al., 2015). It remains for us to determine if early PCD takes place in

these specific bee lice lineages during development and causes the reduction in bundle

diameter that we see in flight hemilineages. As with most parasitic insects, bee and swift

louseflies are impossible to maintain in the laboratory in the absence of their hosts and

procuring them is a challenge (e.g. bee louseflies are now only found on two islands in the

UK, while collecting swift louseflies is restricted to the summer months due to *Apus apus*

migrations). Nonetheless, our observations that PCD is widespread across insects

complements our findings in bee louseflies, strongly suggesting that an extensive PCD in

flight hemilineages accompanied the loss of flight during evolution. Interestingly, the

reduction in fibre diameter we see in bee louseflies was not evident in swift louseflies. This

difference is likely due to the more significant changes to body plan in bee louseflies, i.e. a

complete loss of the flight apparatus during evolution. Swift louseflies however still maintain

vestigial wings and halteres (Walker and Rotherham, 2010), whereas bee louseflies have a

severely reduced thorax, completely lacking wings, halteres and flight muscles (McAlister,

2018).

Blocking death results in functional neurons that integrate into adult networks

Here we blocked PCD within the MNB lineage (called lineage 0 in the postembryonic

literature) and found that 'doomed' neurons become octopaminergic, generate arborizations,

and target the tectulum neuropil (Court et al., 2017). Previous work has shown that

octopamine can induce and maintain rhythmic behaviours such as stepping movements and

flight muscle contractions in locusts (Sombati and Hoyle, 1984) and walking, wing flicking

and hindleg grooming in decapitated fruit flies (Yellman et al., 1997). Consistently, we find

that our 'undead' hemilineage 0B cells can induce walking when activated thermogenetically

in headless flies and show calcium activity during naturalistic bouts of locomotion. Thus, by

blocking death, we have 'resurrected' functional neurons that are able to integrate into

thoracic motor networks. Although this 'dialling up' of cell numbers is artificial in our

system, it reveals how doomed neurons possess cryptic cellular phenotypes that can emerge

when death is blocked, advocating for the evolvability of such a hemilineage based system.

While these undead hemilineage 0B octopaminergic neurons share many conserved

features with wild-type cells, the variations we see in their morphology may act as a substrate

for evolutionary change. A recent study has linked structural changes in the ventral nerve

cord (VNC) with changes in behaviour between strains of *Drosophila melanogaster* (Mellert

et al., 2016). Mellert *et al.,* show that hemilineage 12A in the mesothorax has variable bundle

morphologies and that these correlate well with the time of flight initiation. Flight is used as

an escape response and can be instrumental for predator evasion, one of the major

evolutionary forces which have selected for flight in insects in the first place (Dudley, 2002).

With this in mind, it seems plausible that changes in either neuron number and/or innovations

in 'undead' neuron structure could affect adult behaviour and be ultimately adaptive.

Recently it has been shown that undead sensory neurons that are functional, integrate, and

appear to be tuned to specific odours (Prieto-Godino et al., 2020). Importantly, Prieto-Godino

et al. also show that there is cell number variation in this neuronal population across

drosophilids and that blocking PCD in *melanogaster* results in the survival of mosquito-like

CO_2-sensing neurons in the maxillary palps. This suggests that both the central and peripheral

nervous system may use similar modes of early PCD to sculpt circuits during the evolution of

true flies.

Hemilineage-specific cell death occurs in newly born neurons

To help us understand more about the patterning of PCD we designed and used a new

effector caspase probe SR4VH allowed us to interrogate the extent and dynamics of

hemilineage-specific cell death. It shows us that an early-onset PCD is responsible for the

elimination of postembryonic neurons in the fly VNC, that this happens throughout the entire

3.5 days of postembryonic neurogenesis, and is hemilineage-specific. Although PCD has

been reported as a fate within lineages in the embryo (Karcavich and Doe, 2005; Rogulja-

Ortmann et al., 2007), the impact of this 'early' and hemilineage-specific PCD on the

construction of the adult network has yet to be fully appreciated. This type of PCD is

responsible for removing almost half of all postembryonic neurons that are born in the fly

(Truman et al., 2010). Until now, the most frequently reported type of neuronal death

described in insects has been the hormonally-regulated PCD that removes mature neurons

during the narrow developmental windows at the beginning of metamorphosis and within the

first day after adult eclosion (Draizen et al., 1999; Lee et al., 2013). We now know from our

data that these make up only a small fraction, compared to the total number of neuronal

deaths in the fly.

Our SR4VH probe allows us to see that newly born neurons initiate cell death within

the first 5.5 hours after birth. We can capture different stages of cell death: with young cells

at very initial stages of PCD located closer to the neuroblast, while older cells at more

advanced stages of PCD with RFP labelled cell membranes found close to the lineage bundle.

Importantly, death happens before neurites have extended, strongly suggesting that this PCD

is not an analogue of neurotrophic death, found in vertebrates - where neuron-target

interactions play a major role in the decisions of cell survival (Dekkers and Barde, 2013).

This the dynamics of hemilineage-specific cell death we see with SR4VH fits well with the

patterns that we find in *Thermobia domestica* and *Crataerina pallida* suggesting that they too

undergo a death with rapid onset following division.

The critical question that this data brings into focus is how early-onset PCD is

orchestrated, especially that an early intrinsically-determined mode of cell death seems to be

widespread across animals, from *C. elegans* to mice (Southwell et al., 2012; Fricker et al.,

2018). Early PCD likely involves a combination of intrinsic patterning and cell-cell

interactions between sibling neurons which ultimately deploy the activity of proapoptotic

genes *rpr*, *hid*, *grim* and/or *skl*. Previously, patterning genes such as *Ubx* have been shown to

contribute to the survival of hemilineages in the thoracic VNC in a parasegment-specific

manner (Marin et al., 2012), while the transcription factor Unc-4 has been recently

demonstrated to provide neural identity to hemilineages involved in flight (Lacin et al.,

2020). Such spatial patterning is also required to establish NB identity which, in turn, can

determine which hemilineage is maintained and which dies. In the developing optic lobe

Bertet *et al*. have shown that the temporal sequence of transcription factors expressed in

neuroblasts (inherited by the GMC and newly-born neurons) produces a switch in the

selective survival of one hemilineage over the other (Bertet et al., 2014). In this manner, the

changes to NB identity which have been documented in other insects (Biffar and Stollewerk,

2014), despite a conserved NB array and progeny, could in turn influence the pattern of PCD.

Alongside, cell-cell interactions between newly-born neurons could influence fate choices.

The requirement of interactions between newly-born siblings in determining asymmetric

fates has been shown in the grasshopper VNC (Doe et al., 1985; Kuwada and Goodman,

1985), although this exact same mechanism has not been demonstrated in *Drosophila*.

The spatio-temporal pattern of death revealed with SR4VH shows us that

understanding the molecular control of hemilineage-based death is the key question going

forward and is likely to provide insight into how networks evolve.

Conclusions

Here we have shown that undead neurons elaborate complex arborizations, express distinct

transmitter identities and function. We find hemilineage-based, 'early' programmed cell

death is widespread during the development of the CNS of insects from the primitive

Firebrats, to most derived true flies. Early cell death appears to be a specific subtype of PCD

present across animals. Understanding how early PCD is specified across species should help

us elucidate how nervous systems are built and evolve. Our exploration of homologous

lineages in flightless dipterans shows that changes in the extent and pattern of PCD are

evident with changes in body plan. As the evolutionary changes seen in neural networks

ultimately result from heritable differences in developmental processes, our future

endeavours will be directed towards elucidating how genetic programs are deployed to

establish the pattern of PCD. The cellular leitmotif of hemilineage-based cell death, we

present here, provides us with something tangible that we can search for. Thus, we suggest

that viewing the evolution of insect nervous systems through the lens of the 'hemilineage'

will be critical for understanding how development brings about adaptive changes in neural

network motifs.

SUPPLEMENTARY FIGURES

Figure S1. Postembryonic development of lineage 0B in wild-type and *H99/XR38* flies.
(**A**) CD8::GFP expression driven by *TDC2-GAL4* in octopaminergic neurons from hemilineage 0B in the VNC of wild-type flies in T1 (1st row), T2 (2nd row) and T3 (3rd row) during pupal development at 2 hours after puparium formation (2h APF, 1st column), 16h APF (2nd column) and 25 hAPF (3rd column). One embryonically-born cell is replaced during postembryonic neurogenesis in T1 and T2, while roughly 4 postembryonic octopaminergic neurons are added in T2. White arrows indicate newly-born neurons with faint GFP and no RedStinger visible at 2hAPF (arrows). RedStinger fluorescence becomes visible at 16-25hAPF, as the protein matures (arrows). Scale bar, 10 µm. (**B**) Accumulations of GFP and RedStinger (orange arrow) in the proximity of the prothoracic cell cluster, alongside the presence of one cell with dim RedStinger fluorescence (white arrow) suggest one embryonic cell has died at 16hAPF. Scale bar, 10 µm. (**C**) CD8::GFP expression driven by *TDC2-GAL4* in octopaminergic neurons from hemilineage 0B in the VNC of *H99/XR38* flies in T1 (1st row), T2 (2nd row) and T3 (3rd row) during pupal development at 2h APF (1st column) and 25h APF (2nd column). Not all undead neurons present in the adult have matured by 25h APF. Scale bar, 10 µm. (**D**) Graphical representation of the number of *TDC2-GAL4*-expressing neurons throughout development

in wild-type (magenta) and *H99/XR38* mutant flies (green), where X represents time and Y represents
the number of cells. Individual lines in the graph depict cell numbers in different thoracic segments.
Error bars represent standard deviation. The inset represents a compressed idealised graphical
representation for wild-type flies; marks on the X axis are the same; numbers at branch points
represent cell numbers. In wild-type flies both T1 and T3 acquire an additional cell at 2h APF and
lose one by the end of metamorphosis (magenta). Roughly half of undead neurons are present at 25
hAPF, with the other half maturing by the end of metamorphosis (green).
722

Comment on S1 data: *The implication of additional postembryonic octopaminergic neurons*

born in the in the wild-type larva. It is well known that there are 3-4 extra octopaminergic

neurons generated postembryonically in the wing bearing second (T2) thoracic neuromere in

adult flies compared to the 3 cells found in each thoracic segment in larvae [29]. Until our

inquiry into the timeline of postembryonic neurogenesis, it was thought that all

octopaminergic neurons born in the embryo persist into adulthood, but to our surprise, that is

not the case. We noticed that one postembryonic octopaminergic cell is born in wild-type

flies in both the first (T1) and third (T3) thoracic segments, bringing the total number of cells

to 4 (**Figure S1A**,**D**). Instead, the total number of 3 cells per cluster in T1 and T3 in adult

flies is achieved via *hormonally gated metamorphic cell death of an embryonically born*

mature neuron at larval-pupal stages (**Figure S1A,B,D**). We have shown here for the first

time that wild-type octopaminergic neurons are produced in lineage 0 in all three thoracic

neuromeres during postembryonic neurogenesis. This fits well with our observations that

some *Dronc*-null singleton clones generated using MARCM showed bilateral symmetry. As

the majority of these were in the mesothorax (see also **Figure S2A,D**), where it is more likely

to label wild-type neurons purely by chance (with up to 4 postembryonic neurons produced

here and only 1 each in T1 and T3), we think these may be indeed wild-type. Furthermore,

we only ever encountered mixed MARCM clones containing one bilaterally-symmetric and

several turning neurites in the mesothorax (see also **Figure S2A,D**). For behavioural

experiments we excluded all MARCM *Dronc* clones with bifurcating neurons to ensure that

we only interpreted the activity of undead neurons.

Figure S2. Quantification of undead neuron morphology.
(**A**) Quantification of the primary neurite path. Most undead neurons arborize to the left. (**B**) Quantification of neuropil innervation. Most undead neurons branched extensively in the dorsal neuropil. Undead neurons in T3 never innervated the ventral neuropil. Few undead neurons from each thoracic segment projected neurites into adjacent segments. (**C**) Quantification of efferent neurites. Most undead neurons joined a thoracic nerve heading towards the periphery. Some undead neurons in T1 sent branches through the cervical connective. (**D**) Quantification of the number of undead neurons in MARCM clones for each thoracic segment color-coded by primary neurite path. The number of undead neurons varied across MARCM clones, with up to 13 in T1 and T2 and up to 3 in T3. Most single-celled MARCM clones contained undead neurons with a wild-type morphology (beige circles) and most undead neurons with a wild-type morphology were encountered in T2. Mixed clones (half-filled circles) made up of one bifurcating neuron and several neurons turning either left or right were encountered only in T2.

Figure S3. Wild-type octopaminergic neurons are active during walking in intact adult *Drosophila*.
(**A**) Raw 2-photon image of *TDC2-GAL4-positive* neurons co-expressing tdTomato (red) and GCaMP6s (cyan) in wild-type flies (**top**). Region-of-interest used to calculate %ΔR/R is outlined (yellow). Standard deviation z-projection of a dorsal-ventral image stack of the functional imaging region-of-interest in **b** (**bottom**). Scale bar, 50 µm. (**B**) Representative behavioural and functional imaging data in wild-type flies. Shown are: CO_2 stimulation (black, no stimulation), %ΔR/R (ratio of GCaMP6s / tdTomato) signal (green), and ball rotations indicating forward walking (red), sideways walking (blue), and turning (purple). The behaviour of the fly was classified as either walking (light blue), or resting (grey) by applying a threshold on ball rotation speed. (**C**) (**left**) Individual (green) and average (black) %ΔR/R traces within each behavioural epoch for walking (n = 77) and resting (n = 80) events processed from 720 s of imaging data . Solid green circles indicate the end of a behavioural epoch. The average trace (black line) was calculated for only periods with 4 or more traces. (**right**) Normalised average %ΔR/R traces for three different flies during walking and resting. The average (black) trace is the same as in the left panel.

Figure S4. Subregion analysis of calcium signals along the width of a primary neurite in a H99/XR38 animal.

(**A**) Time-averaged projection of the imaging plane for *TDC2-GAL4*-positive neurons co-expressing tdTomato (red) and GCaMP6s (cyan) in a *H99/XR38* animal. Regions-of-interest used to calculate %ΔR/R (color-coded) are overlaid on top of time-projected and optic flow registered 2-photon images. Scale bar, 50 µm. (**B**) Representative behavioural and functional imaging data for this animal. Shown are: CO_2 stimulation (black), %ΔR/R (ratio of GCaMP6s / tdTomato) signal (color-coded for each ROI as in panel **a**), and ball rotations indicating forward walking (red), sideways walking (blue), and turning (purple). The behaviour of the fly was classified as either walking (light blue), or resting (grey) by applying a threshold on ball rotation speed.

Figure S5. SR4VH reveals successive stages of cell death in lineages with doomed hemilineages.
(**A**) Quantification of lineages with dying cells as reported with SR4VH in each thoracic segment from 5 wandering 3^{rd} instar larvae. Back bars represent the number of samples in which dying cells were encountered in at least one hemisegment for that lineage. Grey bars represent lineages with no record of hemilineage-specific cell death as reported in Truman et al., 2010. (**B**) Antibodies for active Dcp-1 label cells located close to the lineage bundle in doomed lineages from larvae expressing SR4VH driven by *Worniu-GAL4*. Right panels show examples of doomed lineages 0, 18 and 23 containing cells at subsequent stages of cell death. White arrows indicate: nuclear Venus without cleaved Dcp-1 (early-stage), Venus and Dcp-1 colocalization (mid-stage), and pyknotic cells/dead cell membranes with RFP and cleaved Dcp-1 (late-stage). Scale bars, 10 µm. $n = 7$.

SUPPLEMENTARY VIDEOS

Video S1

Video recordings of control and 'undead' decapitated flies during thermogenetic activation. Examples from each behavioural category showing responses to heat-activation in negative control (top left), positive control (top right), MARCM control (bottom left) and undead neurons (bottom right). MARCM control and undead neuron animals were used for

extracting the centroid trace provided in **Fig. 3, B and C**. The increase in temperature is

displayed in the bottom right corner. Frames represent recordings from 30 to 70s.

Video S2.

Recording of 2-photon calcium imaging in undead neurons. Synchronized front and side

camera behaviour videography (**bottom-right**), and 2-photon imaging data (**top-right**) used

for the data analysis provided in **Fig. 3, H-J**.

Video S3.

Z-stack of the imaging area for GCaMP6s activity in undead neurons. A videography

showing the imaging plane at different depths of the prothoracic segment corresponding to

Figure 3H. The thicker left branch likely includes mostly undead neurons.

Video S4.

Subregion neuronal activity patterns during walking. Imaging data used for analysis in **Fig.**

S4. Shown are %ΔR/R traces for R

MATERIALS AND METHODS

<u>Animals</u>

We used the following *Drosophila melanogaster* stocks: *Worniu-GAL4; Dr/TM3,*

Ubx-LacZ, Sb (BL 56553), *Tdc2-GAL4* (BL 9313), *OK371-GAL4* (BL 26160), *UAS-SR4VH*

(described here), *UAS-CD8::GFP* (BL 5137), *UAS-tdTomato-p2A-GCaMP6s* (28) (kind gift

from M. Dickinson), *H99/TM3, Sb* (BL 1576), *XR38/TM3*, Sb, *If/CyO; dronc$^{\Delta A8}$,*

FRT2A/TM6β, Tb, Hu (*37*) and *hs-flp; ; TubP-GAL80, FRT2A/TM3, Sb* (*16*).

Firebrat adults of *Thermobia domestica* were obtained from Buzzard Reptile and

Aquatics (buzzardreptile.co.uk) and reared on a diet of fish flakes and wholemeal bran at

40°C in darkness inside a humid plastic container. A staging series was calculated by time to

hatching.

Crataerina pallida swift lousefly adults were collected from swift (*Apus apus*) nesting

boxes fitted behind the louvres of belfry windows from churches in Cambridgeshire and

Suffolk (UK) with the help of local conservationists Simon Evans, Richard Newell and Bill

Murrells. Swift louseflies were kept at 20°C on a 12h dark:12h light cycle until dissected.

Pregnant females, recognised by their enlarged and translucent abdomen through which

larvae or prepupae could be detected, were kept separately and checked daily for pupa

ejection. The day in which a pupa was laid was defined as 'Day 0' of external development

(outside the mother's abdomen).

Braula coeca bee lousefly adults were obtained from a black bee (*Apis mellifera*

mellifera) colony on the Isle of Colonsay, UK (kind gift from A. Abrahams). Bee louseflies

were shipped by post in small cages containing worker bees feeding on bee fondant. The

black bees and bee louseflies were anesthetized by placing the cage on a CO_2 pad and the bee

louseflies were removed for dissection.

Construction of *UAS-SR4VH*

SR4VH was constructed by standard molecular biology procedures. It comprises the

myristylation signal of *Drosophila* Src64B (amino acids 1-95), a monomeric red fluorescent

protein mRFP1 (*38*), a linker that contains four DEVD sites, a yellow florescent protein

Venus (*39*), and a nuclear localization signal of *Drosophila* histone H2B (amino acids 1-51).

While the design is similar to the previously reported caspase probe Apoliner (*40*), the

Src64B myristylation signal and the H2B NLS offers better membrane and nuclear

localization, respectively, and four DEVD sites are expected to provide higher sensitivity.

854 The probe was cloned in pUAST (*41*) and introduced into the *Drosophila* genome by P

855 element-mediated transformation.

857 <u>Immunohistochemistry and chemical staining</u>

858  *Drosophila* larvae were dissected in PBS without anaesthesia. Firebrat embryos were

859 removed from their chorion and dissected using minuten pins. *Drosophila*, swift lousefly and

860 bee lousefly adults were anesthetised on ice, briefly submerged in absolute ethanol and

861 dissected in PBS. Swift lousefly pupae were immobilised on double sided sticky tape,

862 removed from their pupal case using forceps and dissected in PBS without anaesthesia.

863 Samples were fixed in 3.6% paraformaldehyde in PBS for 30 min (larvae and pupae) or 1h (

864 adults), washed 3 times in 0.3% PBST (0.3% Triton-X100 in PBS, Sigma-Aldrich), blocked

865 in 5% goat serum (Sigma-Aldrich) in PBST for 1h and incubated with primary antibodies in

866 block for 1-3 days at 4°C (*Drosophila*, bee louseflies, swift lousefly pupae), room

867 temperature (firebrats) or 37°C to increase antibody penetration (swift lousefly adults; block

868 supplemented with 0.02% NaN$_3$ to prevent microbial growth). Samples were then washed 4

869 times throughout the day in PBST and incubated with secondary antibodies in block for a

870 further 1-3 days, followed by final washes in PBST and PBS. Brains and VNCs were

871 mounted on poly-L-lysine-coated coverslip, dehydrated in increasing serial concentrations of

872 ethanol (15%, 30%, 70%, 80%, 90% and twice in 100%) for 5min each, dipped once in

873 xylene, then incubated twice for 5min in fresh xylene. A droplet of DePeX (EMS) was added

874 on top of the mounted sample and the coverslip was placed face-down on a glass slide.

875  We used the following primary antibodies: chicken anti-GFP (1:500; ab13970,

876 Abcam), mouse anti-Neuroglian (1:50; BP 104, Developmental Studies Hybridoma Bank),

877 rabbit anti-cleaved *Drosophila* Dcp1 (1:100; Asp216, Cell Signaling), guinea pig anti-

878 Syncrip (1:100; kind gift from I. Davis; to label neuroblasts and early progeny in lineages -

Jim Truman, personal communication), mouse anti-Engrailed/Invected (1:2; 4D9,

Developmental Studies Hybridoma Bank), rabbit anti-DVGLUT C-terminus (*43*) (1:5000;

AB_2490071, kind gift from H. Aberle), rat anti-tyramine β-hydroxylase (*30*) (1:50; TBH,

kind gift from M. Monastirioti), rabbit anti-vsg (1:400; kind gift from Sean Carroll and

Kirsten Gruss) and rabbit anti-GABA (1:100; AB_572234, Imunostar).

Secondary antibodies were Alexa Fluor® 488-conjugated goat anti-chicken (1:500;

A11039, Invitrogen, Thermo Fisher Scientific), Alexa Fluor® 488-conjugated goat anti-

rabbit (1:500; A11070, Invitrogen, Thermo Fisher Scientific), Cy^{TM}3-conjugated donkey

anti-rabbit (1:500; 711-006-152, Jackson ImmunoResearch), Cy^{TM}5-conjugated donkey anti-

mouse (1:500; 715-006-151, Jackson ImmunoResearch), Alexa Fluor® 488-conjugated

donkey anti-rat cross-adsorbed against mouse (1:100; 712-545-153, Jackson

ImmunoResearch), Alexa Fluor® 488-conjugated donkey anti-guinea pig (1:500; A11073,

Invitrogen, Thermo Fisher Scientific).

In firebrat embryos we detected dying cells using the Click-iT Plus TUNEL assay kit

(C10618, *life* technologies). To stain cell nuclei and neuropil, firebrat samples were incubated

with DAPI (1:10000; D9542, Sigma-Aldrich) and Phalloidin-488 (1:100, *life* technologies) in

PBST for 30 min at room temperature. Incubations were carried out following secondary

antibody treatment. Samples were then washed in PBST and PBS, and mounted.

<u>EdU treatment</u>

To label proliferating cells and their progeny we used the Click-iT EdU imaging Kit

(C10337, *life* technologies). Freshly dissected nervous systems from swift lousefly pupae

were incubated in EdU 1:1000 in PBS at room temperature for 1-3h on a shaker, rinsed with

PBS and fixed in cold buffered formaldehyde 3.6% in PBS for 30min. Samples were then

stained using the immunohistochemistry protocol described above. The colour reaction for

EdU was carried out as instructed by the vendor after the secondary antibodies were washed

out.

<u>Generation of Undead neuron MARCM clones</u>

To induce mitotic clones of undead neurons, rescued from programmed cell death, we

used the mosaic analysis with a repressible cell marker technique (*43*). 0-4h first instar larvae

resulting from crossing females of the genotype *hs-flp; ; TubP-GAL80, FRT2A/TM3, Sb* with

; TDC2-GAL4, UAS-CD8::GFP, UAS-TrpA1; dronc$^{\Delta A8}$, FRT2A/TM6β, Tb, Hu males were

heat-shocked at 37°C in a plastic food vial placed in a water bath for either 1h or 45min,

followed by 45min at room temperature and a second incubation period at 37°C for 30min.

After heat-shock, larvae were immediately returned to 23 or 25°C. Cell death was blocked in

clones homozygous for the loss-of-function allele of the initiator caspase dronc. Because we

used the octopaminergic driver line *TDC2-GAL4* to induce the expression of CD8::GFP and

TrpA1, we were able to visualise and thermogenetically activate only postembryonic neurons

of hemilineage 0B. A small number of wild-type octopaminergic neurons are born during

postembryonic neurogenesis (1 in T1 and T3, 4-5 in T2, data now shown). To ensure the

characterisation of undead neurons only, MARCM clones including a bilaterally symmetrical

primary neurite were excluded from analysis.

<u>Thermogenetic activation and video recordings</u>

Prior to recordings, 2-6-day-old males of *Drosophila melanogaster* (*hs-flp/+; Tdc2-*

GAL4, UAS-CD8::GFP, UAS-dTRPA1/+; dronc$^{\Delta A8}$, FRT2A/TubP-GAL80, FRT2A) reared at

23 or 25°C in a 12h:12h light:dark cycle were anesthetised on ice and decapitated using a pair

of micro spring scissors in under 3min. We used males as we found they are more responsive

to octopamine release by thermogenetic activation than females (data not shown). The

headless flies were brushed back into a food vial placed on its side and left to recover for at

least 1 hour. To generate the heat ramp required to thermogenetically activate undead

neurons, we used a 12V thermoelectric Peltier plate (model: TEC1-12706, size: 40mm x

40mm x 3.6mm) connected to a DC power supply (HY3005D, Rapid Electronics) set at a

constant current of 0.46A, with a variable voltage, calibrated using an infrared laser

thermometer (N92FX, Maplin). These settings generated a temperature ramp which lasted

70s from 22°C to 34°C. Videos were recorded at 25 fps using a Sony NEX-5N digital camera

(kindly provided by Ian Wynne) mounted to a stereo microscope. A piece of graph paper was

used for spatial calibration. To match the presence of undead neurons with behaviour, each

decapitated fly used for thermogenetic activation was indexed and prepared for dissection

and immunostaining.

<u>Two-photon calcium imaging in behaving intact flies</u>

The method for *in vivo* two-photon imaging of the VNC in behaving adult *Drosophila*

is described in (*28*). Briefly, flies were anesthetized through cooling and then mounted onto

custom imaging stages. The dorsal thoracic cuticle was removed and indirect flight muscles

were left to degrade over the course of 1h. Subsequently, the proventriculus and salivary

glands were resected to gain optical access to the VNC.

Horizontal sections of the T1 leg ganglion were imaged using galvo-galvo scanning.

For control animals, the bifurcation point of TDC-positive neurites were imaged to

circumvent ROI disappearances caused by movement. For animals harbouring undead *TDC2-*

GAL4-positive neurons, the thickest branch of the axonal bifurcation was chosen because

they were most likely to contain undead neurites. Image dimensions ranged between

512×512 and 320×320, resulting in 1.6 to 3.4 fps data acquisition. Imaging areas ranged

between 92×92 µm and 149×149 µm. Laser power was held at ~8 mW.

<u>Data analysis for 2-photon imaging in behaving *Drosophila*</u>

Python scripts (modified from (*28*)) were used to extract ROI fluorescence traces and to compute spherical treadmill ball rotations. Walking epochs were determined by placing a threshold on ball rotations, which were first converted into anterior-posterior ($v_{forward}$) and medial-lateral (v_{side}) speeds (1 rot s^{-1} = 31.42 mm s^{-1}) and into degrees s^{-1} (1 rot s^{-1} $\Box$ = $\Box$ 360° s^{-1}) for yaw ($v_{rotation}$) movements. Thresholds were 0.12 mm, 0.12 mm, and 5 degrees, respectively. Periods below these thresholds were considered 'resting' while other periods were considered 'walking'. Fluorescence traces for epochs with the same behaviour were aligned by start point to compute average %ΔR/R traces for specific actions.

To calculate fluorescence traces for small subregions-of-interest across neuritic bundles containing both undead and wild-type neurites, images were registered using an optic flow method described in (*28*). This registration served to minimize motion artefacts. Analysis was limited to a period with no warping artefacts and no ROI disappearance. Subregions were manually selected as small circular ROIs across the neuritic bundle of the registered image. Fluorescence values were then computed from each sub-ROI.

<u>Confocal imaging and image processing</u>

Images were acquired using a Zeiss LSM 510 or a Zeiss LSM 800 confocal microscope at a magnification of 20x or 40x with optical sections taken at 1µm intervals. The resulting images were examined and processed using Fiji (https://imagej.net/Fiji). Some images were manually cropped using the Freehand Selection tool to remove debris or to cut out neuronal lineages in *; Worniu-GAL4, UAS-SR4VH;* samples.

<u>Fluorescence intensity plots</u>

To generate fluorescence intensity along Line plots, we used the Plot Profile tool in

Fiji to extract raw fluorescence intensity values for the RFP and Venus channels. The values

were imported into MATLAB (R2018a, MathWorks) and normalised by dividing all

fluorescence intensity values to the maximum value encountered along each Line. In this

manner, all fluorescence intensity along Line plots have a common scale from 0 to 1, with 1

being the highest value encountered along that Line.

<u>Analysis of thermogenetic activation</u>

Decapitated flies were considered to be walking if they covered a distance greater

than 1 body length and moved their legs in a coordinated sequence from T3 to T2 to T1 at

least once on each side (11). Forward, backward and sideway movements were all interpreted

as walking when both aforementioned conditions were respected. To generate fly body traces

video recordings were imported in MATLAB (R2018a, MathWorks) and the centroid of the

decapitated fly (located on the scutellum) was extracted from each frame using a custom-

written script. Each frame was converted into a grayscale image, its contrast enhanced using

contrast-limited adaptive histogram equalization, filtered using a gaussian smoothing kernel

with a standard deviation of 4, binarized using a custom threshold and the geometric centre of

the fly body automatically extracted and stored in an array. To confirm that the centroid

detection was accurate, a red dot with the centroid coordinates was superimposed onto each

frame of the original recording and the annotated movie was saved for manual inspection.

1001

<u>Quantification of 3A/3B bundle diameters</u>

For calculating 3A/3B hemilineage bundle diameter ratios in fruit flies and bee

louseflies, we generated transverse rendered maximum intensity projections of inverted

grayscale confocal stacks for the pro- and mesothorax (T1 and T2) and frontal projections for

the metathorax (T3). Optical sections were selected to include the common lineage bundle

and the individual hemilineage bundles after their split. Diameter measurements were taken

at the widest point within 5 µm of the bundle split using the Straight Line tool in Fiji and

ratios were calculated by dividing the diameter of hemilineage 3A to that of 3B.

1010

<u>Statistical analysis</u>

For comparing neuron numbers, 3A/3B bundle diameter and T2/T1 number of

neurons, data was tested for normal distribution using the Kolmogorov-Smirnov test and

visualisation of Normal Q-Q plots. Differences between groups were analysed using either

the independent samples t-test for normally distributed data, Welch's test if data failed to

meet the homogeneity of variances assumption or Mann-Whitney t-tests if data failed to meet

the normality and homogeneity of variances assumptions of the independent samples t-test.

For comparing the number of flies which walked in each experimental group, we

performed a Pearson chi-squared test and interpreted the resulting exact significance if the

minimum expected count was greater than 5, or the Fisher's Exact Test 2-sided significance

if the minimum expected count was lower than 5 in at least one cell of the contingency table.

To correct for multiple comparisons we performed a Bonferroni correction (i.e. p values were

multiplied by 6, the total number of pairwise tests).

All statistical tests were performed in SPSS Statistics 23 (IBM) with an α set at 0.05. In

all figures, bars represent means ± standard deviation; $P^* < 0.05$, $P^{***} < 0.001$, P^{ns} = not

significant.

1027

1028

1029

ACKNOWLEDGEMENTS

We would like to thank Richard Benton and Lucia Prieto-Godino for discussions and sharing

data. We thank Kristin White and Bloomington Drosophila Stock Center (NIH

P40OD018537) for sharing flies; Maria Monastirioti, Hermann Aberle, Ilan Davis, Kirsten

Gruss, Sean Carroll and Developmental Studies Hybridoma Bank (NICHD of the NIH,

University of Iowa) for antibodies. We are indebted to Simon Evans, Richard Newell and Bill

Murrells for their kind help in collecting swift lice and Andrew Abrams for sending us bee

lice. We are grateful to Andrew M. Dacks and Hans-Joachim Pflüger for providing

unpublished data on *Manduca* octopaminergic neurons. We would like to thank Ian Wynne

for his camera. We also thank Matthias Landgraf, David Shepherd, Jon Clarke and Sanjay

Sane for reading the manuscript. Funding: Williams: BBSRC BB/P025552/1 and

BB/L022672/1. Ramdya: SNSF Project Grant: 175667; Eccellenza Grant: 181239; R'Equip

Grant: 177102.

1043

AUTHOR CONTRIBUTIONS

D.W. and S.P. conceived the study and designed the experiments, apart from *in vivo* imaging;

collected *C.pallida* in the field, and wrote the first draft of the manuscript. S.P. performed the

experiments apart from *in vivo* imaging. P.R. and C.C. conceived and performed the *in*

vivo imaging experiment. S.K. designed and made the genetically encoded caspase reporter

SR4VH. C.S. generated the *T.domestica* data. P.R. and S.K. edited the final manuscript

drafts.

1051

1052   **COMPETING INTERESTS**

1053   Authors declare no competing interests.

1054

1055

REFERENCES

Allen, A.M., Neville, M.C., Birtles, S., Croset, V., Treiber, C.D., Waddell, S., Goodwin, S.F.,
2020. A single-cell transcriptomic atlas of the adult Drosophila ventral nerve cord.
Elife 9. doi:10.7554/eLife.54074
Bardet, P.-L., Kolahgar, G., Mynett, A., Miguel-Aliaga, I., Briscoe, J., Meier, P., Vincent, J.-
P., 2008. A fluorescent reporter of caspase activity for live imaging. Proc. Natl. Acad.
Sci. USA 105, 13901–13905. doi:10.1073/pnas.0806983105
Bate, M.C., 1976. Embryogenesis of an insect nervous system I. A map of the thoracic and
abdominal neuroblasts in Locusta migratoria. Journal of Embryology and
Experimental Morphology 35, 107–123.
Bequaert, J.C., 1952. The Hippoboscidae or Louse-Flies (Díptera) of Mammals and Birds.
Part I. Structure, Physiology and Natural History. Entomologica Americana 32, 1–
209.
Bertet, C., Li, X., Erclik, T., Cavey, M., Wells, B., Desplan, C., 2014. Temporal patterning of
neuroblasts controls Notch-mediated cell survival through regulation of Hid or
Reaper. Cell 158, 1173–1186. doi:10.1016/j.cell.2014.07.045
Biffar, L., Stollewerk, A., 2014. Conservation and evolutionary modifications of neuroblast
expression patterns in insects. Dev. Biol. 388, 103–116.
doi:10.1016/j.ydbio.2014.01.028
Booker, R., Truman, J.W., 1987. Postembryonic neurogenesis in the CNS of the tobacco
hornworm, Manduca sexta. I. Neuroblast arrays and the fate of their progeny during
metamorphosis. J. Comp. Neurol. 255, 548–559. doi:10.1002/cne.902550407
Brembs, B., Christiansen, F., Pflüger, H.J., Duch, C., 2007. Flight initiation and maintenance
deficits in flies with genetically altered biogenic amine levels. J. Neurosci. 27, 11122–
11131. doi:10.1523/JNEUROSCI.2704-07.2007
Campbell, H.R., Thompson, K.J., Siegler, M.V., 1995. Neurons of the median neuroblast
lineage of the grasshopper: a population study of the efferent DUM neurons. J. Comp.
Neurol. 358, 541–551. doi:10.1002/cne.903580407
Charvet, C.J., Striedter, G.F., Finlay, B.L., 2011. Evo-devo and brain scaling: candidate
developmental mechanisms for variation and constancy in vertebrate brain evolution.
Brain Behav. Evol. 78, 248–257. doi:10.1159/000329851
Chen, C.-L., Hermans, L., Viswanathan, M.C., Fortun, D., Aymanns, F., Unser, M.,
Cammarato, A., Dickinson, M.H., Ramdya, P., 2018. Imaging neural activity in the
ventral nerve cord of behaving adult Drosophila. Nat. Commun. 9, 4390.
doi:10.1038/s41467-018-06857-z
Court, R.C., Armstrong, J.D., Borner, J., Card, G., Costa, M., Dickinson, M., Duch, C.,
Korff, W., Mann, R., Merritt, D., Murphey, R., Namiki, S., Seeds, A., Shepherd, D.,
Shirangi, T., Simpson, J., Truman, J., Tuthill, J., Williams, D., 2017. A Systematic
Nomenclature for the Drosophila Ventral Nervous System. BioRxiv.
doi:10.1101/122952

Dekkers, M.P.J., Barde, Y.-A., 2013. Developmental biology. Programmed cell death in
neuronal development. Science 340, 39–41. doi:10.1126/science.1236152
Doe, C.Q., 1992. Molecular markers for identified neuroblasts and ganglion mother cells in
the Drosophila central nervous system. Development 116, 855–863.
Doe, C.Q., Goodman, C.S., 1985. Early events in insect neurogenesis: I. Development and
segmental differences in the pattern of neuronal precursor cells. Dev. Biol. 111, 193–
205. doi:10.1016/0012-1606(85)90445-2
Doe, C.Q., Kuwada, J.Y., Goodman, C.S., 1985. From epithelium to neuroblasts to neurons:
the role of cell interactions and cell lineage during insect neurogenesis. Philos. Trans.
R. Soc. Lond. B, Biol. Sci. 312, 67–81. doi:10.1098/rstb.1985.0178
Draizen, T.A., Ewer, J., Robinow, S., 1999. Genetic and hormonal regulation of the death of
peptidergic neurons in the Drosophila central nervous system. Dev. Neurobiol.
Duch, C., Pflüger, H.J., 1999. DUM neurons in locust flight: a model system for amine-
mediated peripheral adjustments to the requirements of a central motor program.
Journal of Comparative Physiology A: Sensory, Neural, and Behavioral Physiology
184, 489–499. doi:10.1007/s003590050349
Dudley, R., 2002. Chapter 6: Evolution of flight and flightlessness, in: The' ' biomechanics of
insect flight: Form, function, evolution. Princeton University Press, Princeton, N.J,
pp. 261–301.
Eckert, M., Rapus, J., Nürnberger, A., Penzlin, H., 1992. A new specific antibody reveals
octopamine-like immunoreactivity in cockroach ventral nerve cord. J. Comp. Neurol.
322, 1–15. doi:10.1002/cne.903220102
Fricker, M., Tolkovsky, A.M., Borutaite, V., Coleman, M., Brown, G.C., 2018. Neuronal
Cell Death. Physiol. Rev. 98, 813–880. doi:10.1152/physrev.00011.2017
Greer, C.L., Grygoruk, A., Patton, D.E., Ley, B., Romero-Calderon, R., Chang, H.-Y.,
Houshyar, R., Bainton, R.J., Diantonio, A., Krantz, D.E., 2005. A splice variant of the
Drosophila vesicular monoamine transporter contains a conserved trafficking domain
and functions in the storage of dopamine, serotonin, and octopamine. J. Neurobiol.
64, 239–258. doi:10.1002/neu.20146
Hagan, H.R., 1951. Chapter 8: Adenotrophic Viviparity - Diptera (Pupipara), in: Embryology
of the Viviparous Insects. The Ronald Press Company, New York, pp. 159–205.
Harris, R.M., Pfeiffer, B.D., Rubin, G.M., Truman, J.W., 2015. Neuron hemilineages provide
the functional ground plan for the Drosophila ventral nervous system. Elife 4.
doi:10.7554/eLife.04493
Hartenstein, V., Campos-Ortega, J.A., 1984. Early neurogenesis in wild-typeDrosophila
melanogaster. Wilhelm Roux'. Archiv. 193, 308–325. doi:10.1007/BF00848159
Herculano-Houzel, S., Manger, P.R., Kaas, J.H., 2014. Brain scaling in mammalian evolution
as a consequence of concerted and mosaic changes in numbers of neurons and
average neuronal cell size. Front. Neuroanat. 8, 77. doi:10.3389/fnana.2014.00077
Horder, T.J., 1989. Syllabus For An Embryological Synthesis, in: Wake, D.B., Roth, G.
(Eds.), Complex Organismal Functions: Integration and Evolution in Vertebrates.
Wiley.
Hutson, A.M., 1984. Keds, flat-flies and bat-flies, in: Handbooks for the identification of
British insects. Royal Entomological Society, London.
Imms, A.D., 1942. On Braula coeca Nitsch and its affinities. Parasitology 34, 88.
doi:10.1017/S0031182000016012
Jia, X.X., Siegler, M.V.S., 2002. Midline lineages in grasshopper produce neuronal siblings
with asymmetric expression of Engrailed. Development 129, 5181–5193.
Karcavich, R., Doe, C.Q., 2005. Drosophila neuroblast 7-3 cell lineage: a model system for
studying programmed cell death, Notch/Numb signaling, and sequential specification

of ganglion mother cell identity. J. Comp. Neurol. 481, 240–251. doi:10.1002/cne.20371

Kimura, K.I., Truman, J.W., 1990. Postmetamorphic cell death in the nervous and muscular systems of Drosophila melanogaster. J. Neurosci. 10, 403–401.

Kondo, S., Senoo-Matsuda, N., Hiromi, Y., Miura, M., 2006. DRONC coordinates cell death and compensatory proliferation. Mol. Cell. Biol. 26, 7258–7268. doi:10.1128/MCB.00183-06

Konings, P.N., Vullings, H.G., Geffard, M., Buijs, R.M., Diederen, J.H., Jansen, W.F., 1988. Immunocytochemical demonstration of octopamine-immunoreactive cells in the nervous system of Locusta migratoria and Schistocerca gregaria. Cell Tissue Res. 251, 371–379. doi:10.1007/BF00215846

Kumar, A., Bello, B., Reichert, H., 2009. Lineage-specific cell death in postembryonic brain development of Drosophila. Development 136, 3433–3442. doi:10.1242/dev.037226

Kutsch, W., Breidbach, O., 1994. Homologous Structures in the Nervous Systems of Arthropoda, in: Advances in Insect Physiology Volume 24, Advances in Insect Physiology. Elsevier, pp. 1–113. doi:10.1016/S0065-2806(08)60082-X

Kuwada, J.Y., Goodman, C.S., 1985. Neuronal determination during embryonic development of the grasshopper nervous system. Dev. Biol. 110, 114–126. doi:10.1016/0012-1606(85)90069-7

Lacin, H., Chen, H.-M., Long, X., Singer, R.H., Lee, T., Truman, J.W., 2019. Neurotransmitter identity is acquired in a lineage-restricted manner in the Drosophila CNS. Elife 8. doi:10.7554/eLife.43701

Lacin, H., Truman, J.W., 2016. Lineage mapping identifies molecular and architectural similarities between the larval and adult Drosophila central nervous system. Elife 5, e13399. doi:10.7554/eLife.13399

Lacin, H., Williamson, W.R., Card, G.M., Skeath, J.B., Truman, J.W., 2020. Unc-4 acts to promote neuronal identity and development of the take-off circuit in the Drosophila CNS. Elife 9. doi:10.7554/eLife.55007

Lacin, H., Zhu, Y., Wilson, B.A., Skeath, J.B., 2014. Transcription factor expression uniquely identifies most postembryonic neuronal lineages in the Drosophila thoracic central nervous system. Development 141, 1011–1021. doi:10.1242/dev.102178

Landgraf, M., Sánchez-Soriano, N., Technau, G.M., Urban, J., Prokop, A., 2003. Charting the Drosophila neuropile: a strategy for the standardised characterisation of genetically amenable neurites. Dev. Biol. 260, 207–225. doi:10.1016/S0012-1606(03)00215-X

Lee, G., Sehgal, R., Wang, Z., Nair, S., Kikuno, K., Chen, C.-H., Hay, B., Park, J.H., 2013. Essential role of grim-led programmed cell death for the establishment of corazonin-producing peptidergic nervous system during embryogenesis and metamorphosis in Drosophila melanogaster. Biol. Open 2, 283–294. doi:10.1242/bio.20133384

Lehane, M.J., 2008. Chapter 9: The blood-sucking insect groups, in: The Biology of Blood-Sucking in Insects. Cambridge University Press, Cambridge, pp. 202–258. doi:10.1017/CCOL0521836085.009

Lin, S., Lai, S.-L., Yu, H.-H., Chihara, T., Luo, L., Lee, T., 2010. Lineage-specific effects of Notch/Numb signaling in post-embryonic development of the Drosophila brain. Development 137, 43–51. doi:10.1242/dev.041699

Marin, E.C., Dry, K.E., Alaimo, D.R., Rudd, K.T., Cillo, A.R., Clenshaw, M.E., Negre, N., White, K.P., Truman, J.W., 2012. Ultrabithorax confers spatial identity in a context-specific manner in the Drosophila postembryonic ventral nervous system. Neural Dev. 7, 31. doi:10.1186/1749-8104-7-31

McAlister, E., 2018. Chapter 9: The parasites, in: The secret life of flies. Natural History Museum, London, pp. 185–211.

Mellert, D.J., Williamson, W.R., Shirangi, T.R., Card, G.M., Truman, J.W., 2016. Genetic and environmental control of neurodevelopmental robustness in drosophila. PLoS One 11, e0155957. doi:10.1371/journal.pone.0155957

Monastirioti, M., Gorczyca, M., Rapus, J., Eckert, M., White, K., Budnik, V., 1995. Octopamine immunoreactivity in the fruit fly Drosophila melanogaster. J. Comp. Neurol. 356, 275–287. doi:10.1002/cne.903560210

Monastirioti, M., Linn, C.E., White, K., 1996. Characterization of Drosophila tyramine beta-hydroxylase gene and isolation of mutant flies lacking octopamine. J. Neurosci. 16, 3900–3911.

Pauls, D., Blechschmidt, C., Frantzmann, F., El Jundi, B., Selcho, M., 2018. A comprehensive anatomical map of the peripheral octopaminergic/tyraminergic system of Drosophila melanogaster. Sci. Rep. 8, 15314. doi:10.1038/s41598-018-33686-3

Petersen, D.S., Kreuter, N., Heepe, L., Büsse, S., Wellbrock, A.H.J., Witte, K., Gorb, S.N., 2018. Holding tight to feathers - structural specializations and attachment properties of the avian ectoparasite Crataerina pallida (Diptera, Hippoboscidae). J. Exp. Biol. 221. doi:10.1242/jeb.179242

Peterson, C., Carney, G.E., Taylor, B.J., White, K., 2002. reaper is required for neuroblast apoptosis during Drosophila development. Development 129, 1467–1476.

Pflüger, H.J., Stevenson, P.A., 2005. Evolutionary aspects of octopaminergic systems with emphasis on arthropods. Arthropod Struct Dev 34, 379–396. doi:10.1016/j.asd.2005.04.004

Prieto-Godino, L.L., Silbering, A.F., Khallaf, M.A., Cruchet, S., Bojkowska, K., Pradervand, S., Hansson, B.S., Knaden, M., Benton, R., 2020. Functional integration of "undead" neurons in the olfactory system. Sci. Adv. 6, eaaz7238. doi:10.1126/sciadv.aaz7238

Rakic, P., 2009. Evolution of the neocortex: a perspective from developmental biology. Nat. Rev. Neurosci. 10, 724–735. doi:10.1038/nrn2719

Ramdya, P., Benton, R., 2010. Evolving olfactory systems on the fly. Trends Genet. 26, 307–316. doi:10.1016/j.tig.2010.04.004

Roeder, T., 2005. Tyramine and octopamine: ruling behavior and metabolism. Annu Rev Entomol 50, 447–477. doi:10.1146/annurev.ento.50.071803.130404

Rogulja-Ortmann, A., Lüer, K., Seibert, J., Rickert, C., Technau, G.M., 2007. Programmed cell death in the embryonic central nervous system of Drosophila melanogaster. Development 134, 105–116. doi:10.1242/dev.02707

Rowell, H.F., 1976. The cells of the insect neurosecretory system: constancy, variability, and the concept of the unique identifiable neuron, in: Advances in insect physiology. Elsevier, pp. 63–123. doi:10.1016/S0065-2806(08)60254-4

Schlurmann, M., Hausen, K., 2003. Mesothoracic ventral unpaired median (mesVUM) neurons in the blowfly Calliphora erythrocephala. J. Comp. Neurol. 467, 435–453. doi:10.1002/cne.10930

Shepherd, D., Bate, C.M., 1990. Spatial and temporal patterns of neurogenesis in the embryo of the locust (Schistocerca gregaria). Development.

Shepherd, D., Harris, R., Williams, D., Truman, J.W., 2016. Postembryonic Lineages of the Drosophila Ventral Nervous System: Neuroglian expression reveals the adult hemilineage associated fiber tracts in the adult thoracic neuromeres. J. Comp. Neurol. 524, 2677–2695. doi:10.1002/cne.23988

Shepherd, D., Sahota, V., Court, R., Williams, D.W., Truman, J.W., 2019. Developmental organization of central neurons in the adult Drosophila ventral nervous system. J. Comp. Neurol. 527, 2573–2598. doi:10.1002/cne.24690

Siegler, M.V., Manley Jr., P.E., Thompson, K.J., 1991. Sulphide silver staining for
endogenous heavy metals reveals subsets of dorsal unpaired median (DUM) neurons
in insects. J Exp Biol 157, 565–571.
Siegler, M.V., Pankhaniya, R.R., 1997. Engrailed protein is expressed in interneurons but not
motor neurons of the dorsal unpaired median group in the adult grasshopper. J. Comp.
Neurol. 388, 658–668.
Siegler, M.V., Pankhaniya, R.R., Jia, X.X., 2001. Pattern of expression of engrailed in
relation to gamma-aminobutyric acid immunoreactivity in the central nervous system
of the adult grasshopper. J. Comp. Neurol. 440, 85–96. doi:10.1002/cne.1371
Sombati, S., Hoyle, G., 1984. Generation of specific behaviors in a locust by local release
into neuropil of the natural neuromodulator octopamine. J. Neurobiol. 15, 481–506.
doi:10.1002/neu.480150607
Southwell, D.G., Paredes, M.F., Galvao, R.P., Jones, D.L., Froemke, R.C., Sebe, J.Y.,
Alfaro-Cervello, C., Tang, Y., Garcia-Verdugo, J.M., Rubenstein, J.L., Baraban, S.C.,
Alvarez-Buylla, A., 2012. Intrinsically determined cell death of developing cortical
interneurons. Nature 491, 109–113. doi:10.1038/nature11523
Spörhase-Eichmann, U., Vullings, H.G., Buijs, R.M., Hörner, M., Schürmann, F.W., 1992.
Octopamine-immunoreactive neurons in the central nervous system of the cricket,
Gryllus bimaculatus. Cell Tissue Res. 268, 287–304.
Stevenson, P.A., Pflüger, H.J., Eckert, M., Rapus, J., 1992. Octopamine immunoreactive cell
populations in the locust thoracic-abdominal nervous system. J. Comp. Neurol. 315,
382–397. doi:10.1002/cne.903150403
Stevenson, P.A., Spörhase-Eichmann, U., 1995. Localization of octopaminergic neurones in
insects. Comp. Biochem. Physiol. A, Physiol. 110, 203–215. doi:10.1016/0300-
9629(94)00152-J
Tamarelle, M., Haget, A., Ressouches, A., 1985. Segregation, division, and early patterning
of lateral thoracic neuroblasts in the embryos of Carausius morosus Br. (Phasmida :
Lonchodidae). International Journal of Insect Morphology and Embryology 14, 307–
317. doi:10.1016/0020-7322(85)90045-5
Thomas, J.B., Bastiani, M.J., Bate, M., Goodman, C.S., 1984. From grasshopper to
Drosophila: a common plan for neuronal development. Nature 310, 203–207.
doi:10.1038/310203a0
Thompson, K.J., Siegler, M.V., 1991. Anatomy and physiology of spiking local and
intersegmental interneurons in the median neuroblast lineage of the grasshopper. J.
Comp. Neurol. 305, 659–675. doi:10.1002/cne.903050409
Thompson, K.J., Siegler, M.V., 1993. Development of segment specificity in identified
lineages of the grasshopper CNS. J. Neurosci. 13, 3309–3318.
Truman, J.W., 1990. Metamorphosis of the central nervous system of Drosophila. J.
Neurobiol. 21, 1072–1084. doi:10.1002/neu.480210711
Truman, J.W., 1996. Metamorphosis of the Insect Nervous System, in: Gilbert, L., Tata, J.,
Atkinson, B. (Eds.), Metamorphosis: Postembryonic Reprogramming of Gene
Expression in Amphibian and Insect Cells. Academic Press.
Truman, J.W., Ball, E.E., 1998. Patterns of embryonic neurogenesis in a primitive wingless
insect, the silverfish, Ctenolepisma longicaudata: comparison with those seen in
flying insects. Dev. Genes Evol. 208, 357–368. doi:10.1007/s004270050192
Truman, J.W., Bate, M., 1988. Spatial and temporal patterns of neurogenesis in the central
nervous system of Drosophila melanogaster. Dev. Biol. 125, 145–157.
doi:10.1016/0012-1606(88)90067-X

Truman, J.W., Moats, W., Altman, J., Marin, E.C., Williams, D.W., 2010. Role of Notch
1293   signaling in establishing the hemilineages of secondary neurons in Drosophila
1294   melanogaster. Development 137, 53–61. doi:10.1242/dev.041749
1295   Truman, J.W., Schuppe, H., Shepherd, D., Williams, D.W., 2004. Developmental architecture
1296   of adult-specific lineages in the ventral CNS of Drosophila. Development 131, 5167–
1297   5184. doi:10.1242/dev.01371
1298   Truman, J.W., Talbot, W.S., Fahrbach, S.E., Hogness, D.S., 1994. Ecdysone receptor
1299   expression in the CNS correlates with stage-specific responses to ecdysteroids during
1300   Drosophila and Manduca development. Development 120, 219–234.
1301   Walker, M D, Rotherham, I.D., 2010a. Characteristics of Crataerina pallida (Diptera:
1302   Hippoboscidae) populations; a nest ectoparasite of the common swift, Apus apus
1303   (Aves: Apodidae). Exp Parasitol 126, 451–455. doi:10.1016/j.exppara.2010.05.019
1304   Walker, Mark D, Rotherham, I.D., 2010b. The common swift louse fly, Crataerina pallida: an
1305   ideal species for studying host-parasite interactions. J. Insect Sci. 10, 193.
1306   doi:10.1673/031.010.19301
1307   Wheeler, W.M., 1891. Neuroblasts in the arthropod embryo. J Morphol 4, 337–343.
1308   doi:10.1002/jmor.1050040305
1309   White, K., Grether, M.E., Abrams, J.M., Young, L., Farrell, K., Steller, H., 1994. Genetic
1310   control of programmed cell death in Drosophila. Science 264, 677–683.
1311   doi:10.1126/science.8171319
1312   Witten, J.L., Truman, J.W., 1998. Distribution of GABA-like immunoreactive neurons in
1313   insects suggests lineage homology. J. Comp. Neurol. 398, 515–528.
1314   doi:10.1002/(SICI)1096-9861(19980907)398:4<515::AID-CNE4>3.0.CO;2-5
1315   Yellman, C., Tao, H., He, B., Hirsh, J., 1997. Conserved and sexually dimorphic behavioral
1316   responses to biogenic amines in decapitated Drosophila. Proc. Natl. Acad. Sci. USA
1317   94, 4131–4136. doi:10.1073/pnas.94.8.4131