Bridging the TB data gap: *in silico* extraction of rifampicin-resistant tuberculosis diagnostic test results from whole genome sequence data

Kamela Charmaine S. Ng[1,2], Jean Claude S. Ngabonziza[1,3], Pauline Lempens[1], Bouke Catherine de Jong[1], Frank van Leth[2,4], Conor Joseph Meehan[1,5]

[1] Mycobacteriology Unit, Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium
[2] Amsterdam Institute for Global Health and Development, University of Amsterdam, Amsterdam, The Netherlands
[3] Department of Biomedical Services, Rwanda Biomedical Center, Kigali, Rwanda
[4] Department of Global Health, Amsterdam UMC, Location Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands
[5] School of Chemistry and Biosciences, University of Bradford, Bradford, United Kingdom

Corresponding Author:
Kamela Ng
Nationalestraat 155, Antwerp, 2000, Belgium
Email address: kng@itg.be

Abstract

Background. *Mycobacterium tuberculosis* rapid diagnostic tests (RDTs) are widely employed in routine laboratories and national surveys for detection of rifampicin-resistant (RR)-TB. However, as next generation sequencing technologies have become more commonplace in research and surveillance programs, RDTs are being increasingly complemented by whole genome sequencing (WGS). While comparison between RDTs is difficult, all RDT results can be derived from WGS data. This can facilitate longitudinal analysis of RR-TB burden regardless of the data generation technology employed. By converting WGS to RDT results, we enable comparison of data with different formats and sources particularly for countries that employ different diagnostic algorithms for drug resistance surveys. This allows national TB control programs (NTPs) and epidemiologists to exploit all available data for improved RR-TB surveillance.

Methods. We developed the Python-based TB Genome to Test (TBGT) tool that transforms WGS-derived data into laboratory-validated results of the primary RDTs – Xpert MTB/RIF, XpertMTB/RIF Ultra, Genotype MDRTB*plus* v2.0, and Genoscholar NTM + MDRTB II. The tool was validated through RDT results of RR-TB strains with diverse resistance patterns and geographic origins and applied on routine-derived WGS data.

Results. The TBGT tool correctly transformed the SNP data into the RDT results and generated tabulated frequencies of the RDT probes as well as rifampicin susceptible cases. The tool supplemented the RDT probe reactions output with the RR-conferring mutation based on identified SNPs.

Conclusion. Overall, the TBGT tool allows the NTP to assess whether currently implemented RDTs adequately detect RR-TB in their setting. With its feature to transform WGS to RDT results and enable longitudinal RR-TB data analysis, the TBGT tool may bridge the gap between and among data from periodic surveys, continuous surveillance, research, and routine tests, and may be integrated within the existing national connectivity platform for use by the NTP and epidemiologists to improve setting-specific RR-TB control. The TBGT source code and accompanying documentation is available at https://github.com/KamelaNg/TBGT.

Introduction

Resistance to rifampicin (RIF), the most potent anti-tuberculosis (TB) drug, hampers TB control. Rifampicin-resistant (RR)-TB persists as an urgent public health crisis as only 29% of estimated RR-TB patients world-wide were detected and notified in 2017. Further, 18% of previously treated TB patients were found to have RR-TB (WorldHealthOrganization 2018). The WHO endorsed the following rapid molecular RR-TB diagnostic test (RDTs) to address this concern: Xpert MTB/RIF (hereinafter called Xpert Classic) and the new version Xpert MTB/RIF Ultra (hereinafter called Ultra) [Cepheid, Sunnyvale, USA] that employ real-time polymerase chain reaction and molecular beacon technology (Blakemore et al. 2010); and the line probe assays - Genotype MDRTB*plus* v2.0 (hereinafter called LPA-Hain) [Hain Lifescience GmbH, Nehren, Germany]; and Genoscholar NTM + MDRTB II (hereinafter called LPA-Nipro) [NIPRO

Corporation, Osaka, Japan] (Dheda et al. 2017). These tests detect RR-conferring mutations
within the rifampicin resistance determining region (RRDR) or hotspot of the *rpoB* gene (Andre
et al. 2017; Blakemore et al. 2010; Ng et al. 2018a; Ng et al. 2018c), and revolutionized
detection of RR-TB, allowing for prompt identification of patients who need to undergo adapted
treatment. Xpert Classic is the most widely deployed RDT globally, implemented as the initial
diagnostic tool for all presumptive pulmonary TB patients by 32 out of 48 countries in at least
one of the high burden country lists (WorldHealthOrganization 2018). The wide utilization of the
tests in both low and high burden TB countries resulted in the production of large volumes of
RDT data, although comparisons within and between countries can be difficult due to use of
differing technologies.
88
The utility of next generation sequencing technologies has been widely studied for improved
detection of drug-resistant TB in diverse laboratory settings worldwide (Gardy & Loman 2018).
Next generation sequencing-based whole genome sequencing (WGS) of *Mycobacterium*
tuberculosis (*Mtb*) has been shown to accurately detect RR-TB by calling relevant single
nucleotide polymorphisms (SNPs) in the *rpoB* gene (Coll et al. 2018; Miotto et al. 2017). WGS
is already being widely implemented in the United Kingdom, the Netherlands, and New York,
aimed towards completely replacing phenotypic drug susceptibility testing in the clinic
(CRyPTICConsortium et al. 2018; de Viedma 2019). WGS implemented in high burden TB
countries was shown to accurately estimate the prevalence of DR-TB (Zignol et al. 2018). The
conventional periodic TB drug resistance surveys have been gathering data representative of the
Mtb population in poor resource settings, while high income countries typically apply continuous
surveillance (WorldHealthOrganization 2018; Zignol et al. 2018), to help improve the choice of
standard TB treatment before full drug sensitivity profile is known (WorldHealthOrganization
2018).
103
The increasing use of WGS in research and public health initiatives can lead to a disconnect from
the RDT-based data being generated routinely in the clinic, and the majority of surveys,
widening the *Mtb* data gap. We aim to bridge this *Mtb* data gap by transforming WGS data into
each of the related RDT data outputs, allowing longitudinal analysis of RR-TB prevalence and
underlying mutations regardless of the switch between data generation technologies. This will
allow end-users to compare data of varying formats from different technologies, and analyze
previous historical strains with current isolates representative of the entire TB patient population
in the country. Information obtained from this analysis may unearth years' worth of data
revealing RR-TB burden, circulating RR-conferring mutations, and associated RDT probe
patterns in a specific setting, critical for improved RR-TB control.
114
115   We present the TB Genome to Test (TBGT) tool, a Python 3-based set of scripts that rapidly
116   transform WGS data type to RDT and mutation reports. The modules automatically generate
117   frequencies of rifampicin-susceptible (RS) and RR-TB samples detected and supplement the

118 RDT probes output with the detected RR-conferring mutation in the format of 'wild-type amino
119 acid-codon number-mutant amino acid'.
120

Materials & Methods

122 We developed the TBGT tool, a Python 3-executable that converts *Mtb* WGS data in the form of
123 variant call format (VCF) or MTBseq (Kohl et al. 2018) tab files into the most likely output that
124 would be observed from Xpert Classic, Xpert Ultra, LPA-Hain, and LPA-Nipro based on
125 previously validated work (Ng et al. 2018a; Ng et al. 2018c). The TBGT tool can be accessed
126 through https://github.com/KamelaNg/TBGT.
127

The TBGT tool

The TBGT tool can be run on any python-enabled operating system with no additional
prerequisites. The TBGT workflow is shown in Figure 1. The input is a folder of files derived
from a SNP calling pipeline, either in standard VCF format or tab format as output from
MTBseq. The TBGT tool assumes the standard H37Rv NC000962.3 genome was used for
calling these SNPs. If not, the user may remap the genome positions to the specific RR-TB-
related codons using a tab delimited mapping file. An example of this tab-separated file is
bundled with the tool. By default, the module will run all the RDTs on the input files.
The generated output file includes the Sample name, RIF resistance or susceptibility, the
associated mutant codon position and mutation pattern, and a series of 0's or 1's indicating
absence or presence of the capturing probe for the RDTs, and the RR-conferring mutation (An
example output is given in Table 2). A summary table with counts and proportions of detected
RS-TB cases, RDT probes, and RR-conferring mutations is also generated (Supplemental File
Table S1).
142

Validation and sample application of the TBGT modules

We simulated VCF files for internal validation of the TBGT modules A randomly chosen VCF
file was edited in Notepad ++ 7.6.3 (https://notepad-plus-plus.org/) to contain all previously
tested and validated RR-conferring mutations (Andre et al. 2017; Miotto et al. 2017) mapped to
known RDT results (Ng et al. 2018a; Ng et al. 2018c) (Table 1). We generated files with single
and multiple RR mutations to ensure the tool is robust for all scenarios. These simulated VCFs
are provided with the tool.
150

To show real world applicability, the TBGT tool was run on WGS of RS and RR -TB WHO
Tropical Disease Research (TDR) strains with diverse resistance patterns and geographic origins
stored in the Belgian Coordinated Collections of Microorganisms in the Institute of Tropical
Medicine including the 47 TDR-TB strains tested in the previous validation of the RDTs against
the available *rpoB* Sanger sequences of the strains (Ng et al. 2018a; Ng et al. 2018c; Vincent et
al. 2012). The fastQ files (ENA accession PRJEB31023) for these samples were run through the
MTBseq pipeline with default settings (Kohl et al. 2018) to generate the tab files for input to

TBGT. Additionally, WGS from 324 phenotypically RR-TB isolates of retreatment TB patients
in Kinshasa, the Democratic Republic of Congo (DRC) collected in 2005-2010 (Meehan et al.
2018), and 233 WGS isolated 1991- 2010 from Rwanda were subjected to the MTBseq pipeline
and run through the TBGT tool.
162
163   We were then able to extract the associated Xpert results from the 1991 to 2010 Rwandan dataset
164   and compare it with actual Xpert results from 2012 to 2017 (Ng et al. 2018b).
165

Results

167   We tested the TBGT tool on a 64-bit Windows 10 Enterprise computer with a 2.50 GHz
168   processor and a 8.00 GB of RAM. The running time was 39 milliseconds for 1 MTBseq tab file
169   and 530 milliseconds for a VCF file.
170
The TBGT tool correctly transformed the WGS-derived SNPs in the VCF and MTBseq tab files
into the laboratory-validated RDT probe reactions (Table 2), and accurately detected all all
previously validated RR-conferring mutations in the simulated VCF and MTBseq clinical WGS
data from the TDR, DRC (Figures 2A and 2B), and Rwanda (Figures 3A and 3B) strains,
including double nucleotide changes – two SNPs covering two loci or genome positions – such
as the CAC → AGC and CAC → TCC His445Ser mutations and combinations of any two
unlinked mutations. The generated frequency and proportion tables (Supplemental File Table S1)
showed the distribution of the RS and RR-TB strains, the RDT probe reactions, and the RR-
conferring mutations detected.
180
The *in silico* extracted Xpert results from the Rwandan WGS collected in 1991-2010 revealed
that the majority of RR-TB cases detected were linked with Xpert Classic probe E (Figure 3A).
This observation was consistent with the actual Xpert results gathered in 2012 to 2017. Xpert
Classic probe E was also predominantly seen in samples from Kinshasa, DRC (Figure 2A,
Supplemental File Table S1). Mutation S450L, linked with Xpert Classic probe E, was the major
RR-conferring mutation observed in both settings (Figures 2B and 3B).
187

Discussion

In this work, we developed and validated TBGT, a Python-implemented tool that extracts RR-
TB RDT results from WGS-derived SNP data. If a country performed prior drug resistance
surveys with a different diagnostic algorithm, the TBGT tool allows for comparing data of
different formats and sources, making it possible for longitudinal analysis of previous historical
data and current test results, such as periodic DRS results conducted as cross-sectional surveys,
often with different technology from the previous one.
195
The TBGT tool allowed us to analyze almost 3 decades' worth of Xpert data from Rwanda. The
longitudinal analysis of extracted Xpert results from the 1991-2010 Rwandan dataset and actual

Xpert results from 2012-2017 is a proof-of-concept that the TBGT tool provides an exceptional
opportunity to compare and analyze old and new data produced by different technologies. The
predominant Xpert Classic probe E observed in Rwanda is supported by the associated RR
mutation Ser450Leu (Tables 1 and 2) being present in the primary circulating multidrug-resistant
TB clone in this setting (Ngabonziza et al. 2018), and also most frequently associated with global
RR-TB (Cohen et al. 2015; Coll et al. 2018; Georghiou et al. 2016).

The genome-based approach of the TBGT tool also allows for reporting of disputed mutations
that confer occult RR and are frequently missed by the Mycobacterium Growth Indicator Tube
phenotypic DST (Ng et al. 2018a; Van Deun et al. 2015). For instance, disputed mutation
Leu452Pro, epidemiologically linked with an extensively drug-resistant TB outbreak in
KwaZulu-Natal, South Africa in 2005 (Cohen et al. 2015; Ioerger et al. 2009), was detected in
some strains from Kinshasa, DRC and Rwanda (Figures 2B and 3B, Supplemental File).
Mutation Leu452Pro is captured by delayed Xpert Classic probe E, denoting partially inhibited
probe E fluorescence (Lawn & Nicol 2011; Ng et al. 2018a), and specifically identified by the
unique combination of Xpert Ultra probe rpoB4B and corresponding melting temperature shift
(Table 1), provided sufficient *Mtb* DNA is detected. Leu452Pro was reported to be missed in
clinical samples by LPA-Hain due its end-probe location (Al-Mutairi et al. 2011; Rigouts et al.
2013), thus contributing to the RR-TB detection gap. The ability of the TBGT tool to rapidly and
accurately detect disputed mutations is therefore important.

As a supplementary feature, the TBGT tool picks up any RR-conferring mutation present in the
VCF or MTBseq tab file, and may help assess whether RDTs sufficiently detect RR-TB cases in
specific settings.

The TBGT tool may combine clinical data stored in large aggregated files and collected through
the routine laboratory and periodic survey RDT results, with WGS-derived data from national
surveillance programs to create a larger picture of the RS-TB and RR-TB burden in a country.
Through longitudinal analysis of previous historical and current RR-TB data, the TBGT tool may
then allow the NTPs and epidemiologists to identify unusual frequencies of RR-TB cases, RDT
probe reactions, and RR-conferring mutations that define a potential transmission hotspot. The
setting-specific RS and RR-TB counts and proportions generated by the TBGT tool may bring
about an extensive global database with years' worth of data for continuous statistical modeling
analyses and surveillance investigations.

Potentially, the TBGT tool may be integrated in the national TB diagnostic algorithm through the
existing connectivity platform or the newly implemented WHO cloud-based software (Dean
2019). The prospective application of the TBGT tool may bridge and transform the *Mtb* data gap
into action points for RR-TB clinicians to provide appropriate care for the individual TB/RR-TB
patient, and the NTP, public health officials, and policy makers to intervene at the population-

level for improved and sustained RR-TB control (Gardy & Loman 2018). The TBGT modules
could also be expanded to report non-RRDR RR-causing mutations and include other TB drugs –
e.g. isoniazid, pyrazinamide, and fluoroquinolones.
241

Conclusions

Our TBGT tool leverages on the improved access to next generation sequencing technologies in
this genomic epidemiology era of TB, and complement *Mtb* WGS by rapidly transforming WGS
files that store genomic sequence variations to validated outputs of the RR-TB RDTs. The
prospective application of the TBGT tool within a nationwide genomic epidemiology program
will bridge the *Mtb* data gap among routine RDT data, research setting WGS, and periodic
survey and continuous surveillance *rpoB* sequencing and WGS data. This may result in efficient
longitudinal analysis of the circulating RDT probes and underlying distribution of RR-conferring
mutations. The TBGT tool may also help assess whether currently implemented RDT(s)
adequately detect the RR-TB cases in the country. Linking the TBGT tool to the RR-TB
algorithm we previously developed (Ng et al. 2018b) will help identify unusual frequencies of
RR-TB cases as well as rare RDT probes and RR mutations that define a potential transmission
hotspot, critical for active surveillance of transmitted strains in a population by the NTP.
255

Acknowledgements

257 We thank Button Ricarte for his insightful feedback on the manuscript.
258

ADDITIONAL INFORMATION AND DECLARATIONS
Funding
This work was supported by the Erasmus Mundus Joint Doctorate Fellowship grant
2016-1346 to K.C.S.N.

Competing Interests
The authors declare that they have no competing interests.
266

Author Contributions
K. C. S. Ng conceived and designed the study, wrote, implemented, troubleshooted, and finalized
the scripts, generated simulated VCF files, performed internal validation and real-world
application of the TBGT tool, analyzed and interpreted the data, wrote first draft of the
manuscript, revised and approved final paper.
J. C. S. Ngabonziza generated MTBseq tab files of Rwanda WGS for validation of the modules,
reviewed drafts of the manuscript, and approved final paper.
P. L. curated the TDR WGS, reviewed drafts of the manuscript, and approved final paper.
B. C. de Jong reviewed drafts of the manuscript and approved final paper.
F. van Leth reviewed drafts of the manuscript and approved final paper.

277 C. J. Meehan conceived and designed the study, refined, troubleshooted, and finalized the
278 scripts, generated MTBseq tab files of TDR and Kinshasa, DRC WGS for validation of the
279 modules, authored and reviewed drafts of the manuscript, and approved final paper.
280

281 **Data Availability**
282 All scripts and data for validation may be accessed through:
283 https://github.com/KamelaNg/TBGT.
284

285 **Supplemental Information**
286 Supplemental information for this article can be found online at:
287

References

Al-Mutairi NM, Ahmad S, and Mokaddas E. 2011. Performance comparison of four methods for
detecting multidrug-resistant Mycobacterium tuberculosis strains. Int J Tuberc Lung Dis 15:110-
115.
Andre E, Goeminne L, Cabibbe A, Beckert P, Kabamba Mukadi B, Mathys V, Gagneux S,
Niemann S, Van Ingen J, and Cambau E. 2017. Consensus numbering system for the rifampicin
resistance-associated *rpoB* gene mutations in pathogenic mycobacteria. Clin Microbiol Infect
23:167-172. 10.1016/j.cmi.2016.09.006
Blakemore R, Story E, Helb D, Kop J, Banada P, Owens MR, Chakravorty S, Jones M, and
Alland D. 2010. Evaluation of the analytical performance of the Xpert MTB/RIF assay. J Clin
Microbiol 48:2495-2501. 10.1128/JCM.00128-10
Cohen KA, Abeel T, Manson McGuire A, Desjardins CA, Munsamy V, Shea TP, Walker BJ,
Bantubani N, Almeida DV, Alvarado L, Chapman SB, Mvelase NR, Duffy EY, Fitzgerald MG,
Govender P, Gujja S, Hamilton S, Howarth C, Larimer JD, Maharaj K, Pearson MD, Priest ME,
Zeng Q, Padayatchi N, Grosset J, Young SK, Wortman J, Mlisana KP, O'Donnell MR, Birren
BW, Bishai WR, Pym AS, and Earl AM. 2015. Evolution of Extensively Drug-Resistant
Tuberculosis over Four Decades: Whole Genome Sequencing and Dating Analysis of
Mycobacterium tuberculosis Isolates from KwaZulu-Natal. PLoS Med 12:e1001880.
10.1371/journal.pmed.1001880
Coll F, Phelan J, Hill-Cawthorne GA, Nair MB, Mallard K, Ali S, Abdallah AM, Alghamdi S,
Alsomali M, Ahmed AO, Portelli S, Oppong Y, Alves A, Bessa TB, Campino S, Caws M,
Chatterjee A, Crampin AC, Dheda K, Furnham N, Glynn JR, Grandjean L, Minh Ha D, Hasan R,
Hasan Z, Hibberd ML, Joloba M, Jones-Lopez EC, Matsumoto T, Miranda A, Moore DJ,
Mocillo N, Panaiotov S, Parkhill J, Penha C, Perdigao J, Portugal I, Rchiad Z, Robledo J, Sheen
P, Shesha NT, Sirgel FA, Sola C, Oliveira Sousa E, Streicher EM, Helden PV, Viveiros M,
Warren RM, McNerney R, Pain A, and Clark TG. 2018. Genome-wide analysis of multi- and
extensively drug-resistant Mycobacterium tuberculosis. Nat Genet 50:307-316. 10.1038/s41588-
017-0029-0

CRyPTICConsortium, the GP, Allix-Beguec C, Arandjelovic I, Bi L, Beckert P, Bonnet M, Bradley P, Cabibbe AM, Cancino-Munoz I, Caulfield MJ, Chaiprasert A, Cirillo DM, Clifton DA, Comas I, Crook DW, De Filippo MR, de Neeling H, Diel R, Drobniewski FA, Faksri K, Farhat MR, Fleming J, Fowler P, Fowler TA, Gao Q, Gardy J, Gascoyne-Binzi D, Gibertoni-Cruz AL, Gil-Brusola A, Golubchik T, Gonzalo X, Grandjean L, He G, Guthrie JL, Hoosdally S, Hunt M, Iqbal Z, Ismail N, Johnston J, Khanzada FM, Khor CC, Kohl TA, Kong C, Lipworth S, Liu Q, Maphalala G, Martinez E, Mathys V, Merker M, Miotto P, Mistry N, Moore DAJ, Murray M, Niemann S, Omar SV, Ong RT, Peto TEA, Posey JE, Prammananan T, Pym A, Rodrigues C, Rodrigues M, Rodwell T, Rossolini GM, Sanchez Padilla E, Schito M, Shen X, Shendure J, Sintchenko V, Sloutsky A, Smith EG, Snyder M, Soetaert K, Starks AM, Supply P, Suriyapol P, Tahseen S, Tang P, Teo YY, Thuong TNT, Thwaites G, Tortoli E, van Soolingen D, Walker AS, Walker TM, Wilcox M, Wilson DJ, Wyllie D, Yang Y, Zhang H, Zhao Y, and Zhu B. 2018. Prediction of Susceptibility to First-Line Tuberculosis Drugs by DNA Sequencing. N Engl J Med 379:1403-1415. 10.1056/NEJMoa1800474

de Viedma DG. 2019. Pathways and strategies followed in the genomic epidemiology of Mycobacterium tuberculosis. Infect Genet Evol. 10.1016/j.meegid.2019.01.027

Dean A. 2019. Sequencing for the surveillance of drug-resistant TB. WHO Global TB Programme.

Dheda K, Gumbo T, Maartens G, Dooley KE, McNerney R, Murray M, Furin J, Nardell EA, London L, Lessem E, Theron G, van Helden P, Niemann S, Merker M, Dowdy D, Van Rie A, Siu GK, Pasipanodya JG, Rodrigues C, Clark TG, Sirgel FA, Esmail A, Lin HH, Atre SR, Schaaf HS, Chang KC, Lange C, Nahid P, Udwadia ZF, Horsburgh CR, Jr., Churchyard GJ, Menzies D, Hesseling AC, Nuermberger E, McIlleron H, Fennelly KP, Goemaere E, Jaramillo E, Low M, Jara CM, Padayatchi N, and Warren RM. 2017. The epidemiology, pathogenesis, transmission, diagnosis, and management of multidrug-resistant, extensively drug-resistant, and incurable tuberculosis. Lancet Respir Med. 10.1016/S2213-2600(17)30079-6

Gardy JL, and Loman NJ. 2018. Towards a genomics-informed, real-time, global pathogen surveillance system. Nat Rev Genet 19:9-20. 10.1038/nrg.2017.88

Georghiou SB, Seifert M, Catanzaro D, Garfein RS, Valafar F, Crudu V, Rodrigues C, Victor TC, Catanzaro A, and Rodwell TC. 2016. Frequency and Distribution of Tuberculosis Resistance-Associated Mutations between Mumbai, Moldova, and Eastern Cape. Antimicrob Agents Chemother 60:3994-4004. 10.1128/AAC.00222-16

Ioerger TR, Koo S, No EG, Chen X, Larsen MH, Jacobs WR, Jr., Pillay M, Sturm AW, and Sacchettini JC. 2009. Genome analysis of multi- and extensively-drug-resistant tuberculosis from KwaZulu-Natal, South Africa. PLoS One 4:e7778. 10.1371/journal.pone.0007778

Kohl TA, Utpatel C, Schleusener V, De Filippo MR, Beckert P, Cirillo DM, and Niemann S. 2018. MTBseq: a comprehensive pipeline for whole genome sequence analysis of Mycobacterium tuberculosis complex isolates. PeerJ 6:e5895. 10.7717/peerj.5895

Lawn SD, and Nicol MP. 2011. Xpert(R) MTB/RIF assay: development, evaluation and
implementation of a new rapid molecular diagnostic for tuberculosis and rifampicin resistance.
Future Microbiol 6:1067-1082. 10.2217/fmb.11.84
Miotto P, Tessema B, Tagliani E, Chindelevitch L, Starks AM, Emerson C, Hanna D, Kim PS,
Liwski R, Zignol M, Gilpin C, Niemann S, Denkinger CM, Fleming J, Warren RM, Crook D,
Posey J, Gagneux S, Hoffner S, Rodrigues C, Comas I, Engelthaler DM, Murray M, Alland D,
Rigouts L, Lange C, Dheda K, Hasan R, Ranganathan UDK, McNerney R, Ezewudo M, Cirillo
DM, Schito M, Koser CU, and Rodwell TC. 2017. A standardised method for interpreting the
association between mutations and phenotypic drug resistance in Mycobacterium tuberculosis.
Eur Respir J 50. 10.1183/13993003.01354-2017
Ng KC, Meehan CJ, Torrea G, Goeminne L, Diels M, Rigouts L, de Jong BC, and Andre E.
2018a. Potential Application of Digitally Linked Tuberculosis Diagnostics for Real-Time
Surveillance of Drug-Resistant Tuberculosis Transmission: Validation and Analysis of Test
Results. JMIR Med Inform 6:e12. 10.2196/medinform.9309
Ng KCS, Ngabonziza JCS, Meehan CJ, Migambi P, de Jong BC, Cobelens F, and van Leth F.
2018b. Automated algorithm for early identification of rifampicin-resistant tuberculosis
transmission hotspots in Rwanda. The International Journal of Tuberculosis and Lung Disease
22:S1 – S652.
Ng KCS, van Deun A, Meehan CJ, Torrea G, Driesen M, Gabriels S, Rigouts L, Andre E, and de
Jong BC. 2018c. Xpert Ultra Can Unambiguously Identify Specific Rifampin Resistance-
Conferring Mutations. J Clin Microbiol 56. 10.1128/JCM.00686-18
Ngabonziza JCS, Rigouts L, Mazarati JB, Uwizeye C, Nyaruhirira AU, Torrea G, de Jong BC,
and Meehan CJ. 2018. Transmission drives the increase of multidrug-resistant tuberculosis in
Rwanda. The International Journal of Tuberculosis and Lung Disease 22:S1 – S652.
Rigouts L, Gumusboga M, de Rijk WB, Nduwamahoro E, Uwizeye C, de Jong B, and Van Deun
A. 2013. Rifampin resistance missed in automated liquid culture system for Mycobacterium
tuberculosis isolates with specific *rpoB* mutations. J Clin Microbiol 51:2641-2645.
10.1128/JCM.02741-12
Van Deun A, Aung KJ, Hossain A, de Rijk P, Gumusboga M, Rigouts L, and de Jong BC. 2015.
Disputed *rpoB* mutations can frequently cause important rifampicin resistance among new
tuberculosis patients. Int J Tuberc Lung Dis 19:185-190. 10.5588/ijtld.14.0651
Vincent V, Rigouts L, Nduwamahoro E, Holmes B, Cunningham J, Guillerm M, Nathanson CM,
Moussy F, De Jong B, Portaels F, and Ramsay A. 2012. The TDR Tuberculosis Strain Bank: a
resource for basic science, tool development and diagnostic services. Int J Tuberc Lung Dis
16:24-31. 10.5588/ijtld.11.0223
WorldHealthOrganization. 2018. Global tuberculosis report.
Zignol M, Cabibbe AM, Dean AS, Glaziou P, Alikhanova N, Ama C, Andres S, Barbova A,
Borbe-Reyes A, Chin DP, Cirillo DM, Colvin C, Dadu A, Dreyer A, Driesen M, Gilpin C, Hasan
R, Hasan Z, Hoffner S, Hussain A, Ismail N, Kamal SMM, Khanzada FM, Kimerling M, Kohl
TA, Mansjo M, Miotto P, Mukadi YD, Mvusi L, Niemann S, Omar SV, Rigouts L, Schito M,

Sela I, Seyfaddinova M, Skenders G, Skrahina A, Tahseen S, Wells WA, Zhurilo A, Weyer K,
Floyd K, and Raviglione MC. 2018. Genetic sequencing for surveillance of drug resistance in
tuberculosis in highly endemic countries: a multi-country population-based surveillance study.
Lancet Infect Dis 18:675-683. 10.1016/S1473-3099(18)30073-2.

Figure 1. The TBGT tool workflow.

Figure 2A. Distribution of rifampicin-sensitive (RS) samples and Xpert Classic probes among rifampicin-resistant tuberculosis (RR-TB) isolates in Kinshasa, DRC from 2005 to 2010, detected by the TBGT tool.

Figure 2B. Distribution of RR-conferring mutations in Kinshasa, DRC from 2005 to 2010, detected by the TBGT tool.

Figure 3A. Distribution of rifampicin-sensitive (RS) samples and Xpert Classic probes among rifampicin-resistant tuberculosis (RR-TB) isolates in Rwanda from 1991 to 2010, determined by the TBGT tool.

Figure 3B. Distribution of RR-conferring mutations in Rwanda from 1991 to 2010, detected

by the TBGT tool.

Table 1. Rifampicin resistance (RR)-conferring mutations and associated rapid diagnostic test (RDT) results, previously tested and validated, and used as basis for developing the TBGT tool.

RR-TB RDT							
Xpert Classic	Xpert Ultra		LPA-Hain		LPA-Nipro		RR mutation
Capturing probe	Capturing probe	Melting temperature shift	Absent (WT) probe	Developing (MUT) probe	Absent (WT) probe	Developing (MUT) probe	
ND	ND	ND	ND	ND	ND	ND	Val170Phe
Probe A	rpoB1	3.5	WT1		S1		Ser428Arg
Probe A	rpoB1	5.9-6.3	WT2		S1		Leu430Pro
Probe A, Probe B	rpoB1	2.9	WT2		S1		Ser431Gly
Probe B	rpoB1	3.4	WT2, WT3		S1		Gln432Glu
Probe B	rpoB2	3.2	WT3		S2		Met434Ile
Probe B	rpoB2	3.3	WT3		S2		Met434Thr
Probe B	rpoB1	6.3	WT3		S2		Met434Val
Probe B	rpoB2	2.8	WT3, WT4		S2		Asp435Glu
Probe B delayed	rpoB2	5.3	WT3, WT4		S2		Asp435Phe
Probe B	rpoB2	3.3	WT3, WT4		S2		Asp435Gly
Probe B	rpoB2	3.5-3.7	WT3, WT4	MUT1	S2	R2	Asp435Val
Probe B delayed	rpoB2	4.0-4.4	WT3, WT4		S2		Asp435Tyr
Probe C	rpoB2	6.4	WT4		S2		Asn437Asp
Probe C	rpoB2; rpoB3	3.0; 2.3	WT5, WT6		S3		Ser441Leu
Probe C	rpoB2; rpoB3	4.7; 2.3	WT5, WT6		S3		Ser441Gln
Probe D	rpoB3	3.7-3.9	WT7	MUT2B	S4	R4b	His445Asp
Probe D	rpoB3	4.9	WT7		S4		His445Gly
Probe D	rpoB3	3.5-3.6	WT7		S4		His445Leu
Probe D	rpoB3	3.4-3.5	WT7		S4		His445Asn

| RR-TB RDT | | | | | | | RR mutation |
| Xpert Classic | Xpert Ultra | | LPA-Hain | | LPA-Nipro | | |
Capturing probe	Capturing probe	Melting temperature shift	Absent (WT) probe	Developing (MUT) probe	Absent (WT) probe	Developing (MUT) probe	
Probe D	rpoB3	3.6	WT7		S4		His445Gln (CAG)
Probe D	rpoB3	4.1	WT7		S4		His445Gln (CAA)
Probe D delayed	rpoB3	1.9	WT7		S4		His445Arg
Probe D	rpoB3	4.7	WT7		S4		His445Ser
Probe D	rpoB3	4.9	WT7		S4		His445Thr
Probe D	rpoB3	3.2-3.3	WT7	MUT2A	S4	R4a	His445Tyr
Probe D	rpoB4B	5.0	WT7		S4		Lys446Gln
Probe E	rpoB3	4.0	WT8		S5		Ser450Phe
Probe E	rpoB3; rpoB4A	2.5-2.9; 6.0-6.5	WT8	MUT3	S5	R5	Ser450Leu
Probe E	rpoB3; rpoB4A	2.3-2.7; 3.3-3.7	WT8		S5		Ser450Trp
Probe E delayed	rpoB4B	5.7-6.1	WT8		S5		Leu452Pro
ND	ND	ND	ND	ND	ND	ND	Ile491Phe

Capturing probe, RDT probe associated with the RR-conferring mutation; Melting temperature shift, difference between mutant (MUT) and wild-type (WT) melting temperatures; ND, 'not detected', refers to mutations outside the RR determining region, not detected by the RR-TB RDTs.

Table 2. Example of combined results from the TBGT tool rapid diagnostic test (RDT) modules supplemented by the rifampicin resistance (RR)-conferring mutations detected

| Filename | RIF Resistance | Codon number | Codon | Xpert Classic | | Xpert Ultra | | LPA-Hain | | LPA-Nipro | | RR mutation |
				Capturing probe	Probe pattern	Capturing probe, Melting temperature shift	Probe pattern	Capturing probe	Probe pattern	Capturing probe	Probe pattern	
DRC-052577	DETECTED	450	TTG	Probe E	1 1 1 1 0 1	rpoB3,rpoB4A; 2.5-2.9,6.0-6.5	1 1 0 0 1	WT8, MUT3	1 1 1 1 1 1 1 0 0 0 0 1	S5, R5	1 1 1 1 0 0 0 0 1	Ser450Leu
DRC-091003	DETECTED	452	CCG	Probe E delayed	1 1 1 1 1 0	rpoB4B; 5.7-6.1	1 1 1 1 0	WT8	1 1 1 1 1 1 1 0 0 0 0 0	S5	1 1 1 1 0 0 0 0 0	Leu452Pro
1993-09004	DETECTED (only by *rpoB* Sanger sequencing module)	170	TTC	NOT DETECTED		NOT DETECTED		NOT DETECTED		NOT DETECTED		Val170Phe
DRC-101308	DETECTED (only by *rpoB* Sanger sequencing module)	491	TTC	NOT DETECTED		NOT DETECTED		NOT DETECTED		NOT DETECTED		Ile491Phe