Increased risk of infections in Addison's disease and congenital adrenal hyperplasia: a

primary care database cohort study

Alberto S. Tresoldi[1,2,3], Dana Sumilo[4], Mary Perrins[4], Konstantinos A. Toulis[4], Alessandro

Prete[1,2], Narendra Reddy[5], John A.H. Wass[6], Wiebke Arlt[1,2,7*#], Krishnarajah Nirantharakumar[4*]

*joint senior authors   #corresponding author

[1]Institute of Metabolism and Systems Research, College of Medical and Dental Sciences,

University of Birmingham, Birmingham, United Kingdom

[2]Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham,

United Kingdom

[3]Endocrinology, Diabetology and Medical Andrology Unit, Humanitas Research Hospital,

Rozzano (Milan), Italy

[4]Institute of Applied Health Research, College of Medical and Dental Sciences, University of

Birmingham, Birmingham, United Kingdom

[5]Department of Diabetes & Endocrinology, University Hospitals of Leicester NHS Trust,

Leicester Royal Infirmary, Leicester, UK

[6]Oxford Centre for Diabetes, Endocrinology and Metabolism, Churchill Hospital, Oxford, United

Kingdom

[7]NIHR Birmingham Biomedical Research Centre, University Hospitals Birmingham NHS

Foundation Trust and University of Birmingham, Birmingham, UK

Short title: Infection risk in primary adrenal insufficiency

Keywords: primary adrenal insufficiency · Addison's disease · congenital adrenal hyperplasia ·

infections · antimicrobials · primary care

Please address all correspondence to:

Professor Wiebke Arlt

Institute of Metabolism and Systems Research

College of Medical and Dental Sciences

University of Birmingham

Medical School IBR Tower, Rm 236

Birmingham, B15 2TT, United Kingdom

Email w.arlt@bham.ac.uk

Funding: This work was supported by the Medical Research Council UK (Program Grant

0900567, to W A). K.N. is a UK Research and Innovation (UKRI)/Health Data Research (HDR)

UK Innovation Clinical Fellow. W.A. receives support from the National Institute of Health

Research (NIHR) Birmingham Biomedical Research Centre. The views expressed in this

publication are those of the authors and not necessarily those of the NIHR or the Department of

Health and Social Care UK.

Disclosure Statement: W.A. serves as scientific consultant to Diurnal Ltd. and Spruce

Biosciences Inc.. All other authors have no conflict of interest to declare.

ABSTRACT

Context: Mortality and infection-related hospital admissions are increased in patients with primary adrenal insufficiency (PAI). However, the risk of primary care-managed infections in patients with PAI is unknown.

Objective: To estimate infection risk in PAI due to autoimmune Addison's disease (AD) and congenital adrenal hyperplasia (CAH) in a primary care setting.

Design: Retrospective cohort study using UK data collected from 1995 to 2018.

Main outcome measures: Incidence of lower respiratory tract infections (LRTIs), urinary tract infections (UTIs), gastrointestinal infections (GIIs), and prescription counts of antimicrobials in adult PAI patients compared to unexposed controls.

Results: A diagnosis of PAI was established in 1580 AD patients (mean age 51.7 years) and 602 CAH patients (mean age 35.4 years). All AD patients and 42% of CAH patients were prescribed glucocorticoids, most frequently hydrocortisone in AD (82%) and prednisolone in CAH (50%). AD and CAH patients exposed to glucocorticoids, but not CAH patients without glucocorticoid treatment, had a significantly increased risk of LRTIs (adjusted incidence rate ratio AD 2.11 [95% confidence interval 1.64-2.69], CAH 3.23 [1.21-8.61]), UTIs (AD 1.51 [1.29-1.77], CAH 2.20 [1.43-3.34]), and GIIs (AD 3.80 [2.99-4.84], CAH 1.93 [1.06-3.52]). This was mirrored by increased prescription of antibiotics (AD 1.73 [1.69-1.77], CAH 1.77 [1.66-1.89]) and antifungals (AD 1.89 [1.74-2.05], CAH 1.91 [1.50-2.43]).

Conclusions: There is an increased risk of infections and antimicrobial use in PAI in the primary care setting at least partially linked to glucocorticoid treatment. Future studies will need to address whether more physiological glucocorticoid replacement modes could reduce this risk.

Précis

Using data from 1580 AD patients and 602 CAH patients collected in a UK primary care

database from 1995 to 2018, we identified increased risk of infections and antimicrobial

prescription counts.

INTRODUCTION

Primary adrenal insufficiency (PAI) is a severe and potentially life-threatening condition caused

by the failure of the adrenal cortex to produce glucocorticoids and, in most cases,

mineralocorticoids, which occurs in the setting of adrenal disease (1). The two most frequent

causes of PAI are autoimmune adrenalitis (Addison's disease, AD) and congenital adrenal

hyperplasia (CAH).

The prognosis of patients with PAI has improved considerably after life-saving glucocorticoid

replacement therapy became available in the 1950s; however, an increased risk of death has

been described in both AD and CAH patients even in recent years (2,3). In patients with AD, this

has been attributed to adrenal crisis- and infection-related mortality (4). Another study has

reported an increased use of antimicrobial agents and infection-related hospital admissions in

patients with PAI (5). Recent evidence suggests that the increased risk of infections in these

patients could be explained by an impairment of natural killer cell function (6), which may be

caused by the non-physiological delivery of glucocorticoids by currently available preparations

and an associated change in clock gene expression patterns in immune cells (6, 23).

No studies have estimated yet the overall risk of common infections in people with PAI, i.e.

infections that are primarily managed in the primary care setting and usually do not require

hospital admission. However, such infections potentially expose PAI patients to significant risk

of adrenal crisis. Therefore, this study aimed to assess the risk of common types of primary

care-managed infections, namely infections of the lower respiratory tract, the urinary tract, and

gastrointestinal infections, and the use of antimicrobials in the primary care setting in patients

with PAI, comparing AD to CAH and CAH patients with and without glucocorticoid therapy.

MATERIALS AND METHODS

Study design and setting

We conducted a population-based, retrospective, open cohort study to determine the infection risk of patients with AD and CAH in the primary care setting. We assessed the risk of lower respiratory tract infections (LRTIs), urinary tract infections (UTIs), gastrointestinal infections (GIIs), and the counts of antimicrobial prescriptions. We used data from The Health Improvement Network (THIN) database, comprising anonymized electronic medical records from UK general practitioner (GP) practices covering over 5% of the UK population. THIN holds data on demographic characteristics, clinical diagnoses, physical measurements, laboratory results and drug prescriptions recorded using clinical Read code system. Patients registered in THIN have similar age and sex distributions to the general UK population and, therefore, THIN data are well suited for epidemiological studies (7,8).

Study population and period

Our study population consisted of two "exposed" cohorts, comprising adult patients (≥18 years old) diagnosed with AD or CAH according to selected Read codes (see Appendix for the codes used) (9,10). We excluded patients who were at any time point coded with a code consistent with other causes of adrenal insufficiency. In the AD cohort, we included only patients who had at least one prescription of both glucocorticoids (accepting glucocorticoids commonly used in AD) and mineralocorticoids. We also subdivided the CAH cohort in two sub-cohorts: (a) patients who had at least one glucocorticoid prescription at any point (using the same glucocorticoid codes used for AD patients) and (b) patients who were never prescribed with any glucocorticoid therapy, since patients with CAH do not always require glucocorticoid therapy. For every exposed patient, we randomly selected two individuals from a pool of patients matched for age, sex and GP practice who did not have a Read code consistent with PAI at any point before or during the observation period.

The study period extended from 1 January 1995 to 1 January 2018. Patients were eligible for

inclusion one year from the latest of the following dates: study start date, patient registration

date with the GP practice, and practice eligibility date (the date when practices have

implemented an electronic medical record and have passed the assessment for acceptable data

quality). The one-year lag period was applied to ensure there was enough time to document all

information accurately after registration with the practice or after a practice was deemed eligible

to take part. To ensure acceptable data quality, practices were required to have used the

electronical health record system for at least one year and have acceptable mortality reporting

(11).

The index date, i.e. the date when follow-up commences, was defined as the date of diagnosis

for newly diagnosed patients or, if they were already diagnosed with PAI, the date when they

registered with an eligible GP practice. Patients were followed from index date up until the

earliest of the following dates: outcome of interest (only for estimating the incidence of

infections), patient transfer date from practice, patient death, practice's last data collection date,

and study end date.

Outcomes

For the first outcome, the incidence of infections, we used the Read codes that identify cases of

LRTIs, UTIs and GIIs. These infections were chosen because they are the most common type

of infections evidenced in general population and they are frequently diagnosed in primary care

(12). We then calculated the occurrence of this outcome in the different cohorts.

For the second outcome, antimicrobial use, we used the codes for antibiotics and antifungals as

classified in the British National Formulary. We then calculated the total number of prescriptions

for every antimicrobial in each cohort.

For each of the study groups we analyzed age, sex, body mass index (BMI), smoking status,

Townsend Deprivation Index (a measure of deprivation within a population) (13), Charlson

Comorbidity Index (a method of classifying comorbidities to predict mortality in primary care)

(14,15), and type of glucocorticoids prescribed at baseline. For the AD cohort, given the

frequent association with autoimmune conditions, we also evaluated the prevalence of

associated autoimmune comorbidities.

Statistical analysis

Descriptive statistics were used to summarize the baseline characteristics for the exposed and

unexposed groups of patients. Categorical variables were investigated using Chi-square test

and continuous variables were analyzed using t-test.

Adjusted incidence rate ratios (aIRRs) for specific infections and antimicrobial prescriptions

were calculated after adjustment for age, sex, smoking status, BMI, Townsend Deprivation

Index, and Charlson Comorbidity Index, using multivariate Poisson regression analysis.

Statistical analyses were conducted using Stata version 14.2 (Stata Corp, College Station

Texas, USA) and GraphPad Prism 7.04 (GraphPad Software Inc, San Diego, CA).

Ethical approval

The THIN database obtained ethical approval from the South East Multicentre Research Ethics

Committee in 2003. The present study was reviewed and approved (study reference:

18THIN063) by the THIN Scientific Review Committee in July 2018.

RESULTS

Baseline characteristics of the AD cohort

In total, 1580 patients fulfilled the AD criteria; these were matched with 3158 unexposed

individuals (Table 1). The mean age of AD patients was 51.7 years and the majority were

women (57.8%). Compared to unexposed individuals, AD patients had a lower median BMI,

while the Townsend Deprivation Index did not differ significantly between the two groups. The

Charlson Comorbidity Index showed that AD patients had an increased burden of comorbidities

compared to the matched population; this included a higher prevalence of autoimmune

comorbidities, including autoimmune thyroid diseases, type 1 diabetes mellitus, ulcerative

colitis, celiac disease and pernicious anemia (Table 1).

*** Baseline characteristics of the CAH cohort ***

In total, 602 patients fulfilled the CAH criteria and were subdivided into 254 glucocorticoid-

treated patients (42.2%) and 348 patients not on glucocorticoids (57.8%). These were matched

with 508 and 696 unexposed controls, respectively (Table 2).

The majority of CAH patients were female (72.3%), with a lower mean age in glucocorticoid-

treated patients at cohort entry (33.4 vs. 36.9 years). CAH patients had a higher median BMI

compared to controls, and this was evident for both glucocorticoid-treated sub-cohort and the

CAH sub-cohort never treated with glucocorticoids. CAH patients were more frequently

overweight or obese (60.3% vs. 44.2% in matched controls, p $\square$0.001), and this was observed

both in glucocorticoid-treated and untreated CAH patients (59.1 and 61.1%, respectively). The

Townsend Deprivation Index and the Charlson Comorbidity Index did not differ between CAH

patients and controls.

*** Glucocorticoid prescription ***

The most commonly prescribed type of glucocorticoid in the AD cohort was hydrocortisone

(1296 patients, 82%), followed by prednisolone (187 patients, 11.8%). Only a minority of

patients were prescribed cortisone acetate (91 patients, 5.8%, no longer available in the UK)

and dexamethasone (6 patients, 0.4%).

In the glucocorticoid-treated CAH cohort, prednisolone was most commonly prescribed (127

patients, 50.0%), followed by hydrocortisone (96 patients, 37.8%), with a small minority

receiving dexamethasone (15 patients, 5.9%) or cortisone acetate (11 patients, 4.3%). Only five

CAH patients (2%) were prescribed a combination of short- and long-acting glucocorticoids.

Risk of Infections

The risk of LRTIs, UTIs and GIIs was significantly increased in the AD cohort compared to unexposed patients, with the highest relative risk observed for GIIs (adjusted incidence rate ratio (aIRR) 3.80 [95% CI 2.99-4.84]) followed by LRTIs (aIRR 2.11 [95% CI 1.64-2.69]) and UTIs (aIRR 1.51 [95% CI 1.29-1.77]) (Table 3 and Figure 1).

In the overall CAH cohort, there was a significantly increased risk of UTIs and LRTIs (aIRR 1.40 [95% CI 1.06-1.85] and 2.36 [95% CI 1.25-4.42], respectively), with no difference in GI infections (Table 4 and Figure 1). However, when analyzing the population accordingly to glucocorticoid use, only patients exposed to glucocorticoids had an increased risk of infections, with the highest risk observed for LRTIs (aIRR 3.23 [95% CI 1.21-8.61]) followed by UTIs (aIRR 2.20 [95% CI 1.43-3.4]) and GIIs (aIRR 1.93 [95% CI 1.06-3.52]) (Table 4 and Figure 1). In contrast, infection risk in CAH patients not treated with glucocorticoids did not differ from that observed in the matched background population.

Antimicrobial prescriptions

Prescription rates of antibiotics and antifungals were increased in patients with AD (aIRR 1.73 [95% CI 1.69-1.77] and 1.89 [95% CI 1.74-2.05], respectively) (Table 5 and Figure 1).

Similarly, we observed increased antimicrobial prescription rates in CAH patients, with a higher prescription rate glucocorticoid-treated patients (antibiotics: aIRR 1.77 [95% CI 1.66-1.89]; antifungals: aIRR 1.91 [95% CI 1.50-2.43]) than in CAH patients not exposed to glucocorticoids (antibiotics: aIRR 1.15 [95% CI 1.08-1.23]; antifungals: aIRR 1.44 [95% CI 1.18-1.83]) (Table 6 and Figure 1).

DISCUSSION

In this population-based study we found that the risk of three common infections (lower respiratory tract, urinary tract, and gastrointestinal infections) was increased in the primary care setting in patients with PAI, as compared to population-based matched controls. This was also supported by our finding of increased prescription rates of antimicrobials in in patients with PAI. Moreover, we found that CAH patients not receiving glucocorticoids did not have an increased risk of infections, indicating that glucocorticoid therapy might at least partly drive the increased infection risk observed in PAI. To our knowledge, our study is the first to analyze the risk of infection in PAI according to different etiologies and also the first to evaluate these outcomes in a primary care setting.

Previous studies have described an increased infection-related mortality in patients with AD (2,4), but not in CAH patients (3). This was attributed to infections representing a possible trigger for a fatal adrenal crisis. Smans and colleagues reported an increase of the use of antimicrobials and of infection-related hospital admissions in PAI (5); however, the authors focused on hospital-treated infections only, possibly overestimating the actual incidence of this complication due to a lower threshold for admission in PAI patients. In addition, information on the actual etiology of PAI was not available in this study, as PAI was diagnosed on the basis of concomitant glucocorticoid and mineralocorticoid prescriptions, which did not allow to differentiate between AD, CAH, and other causes of PAI.

Until recently, it was unclear whether the observed increase in infection episodes in patients with PAI is related to the underlying disease itself or to the non-physiological delivery of glucocorticoid replacement by currently available glucocorticoid preparations. AD patients frequently also suffer from other autoimmune comorbidities (16), and this was confirmed in our study, with more prevalent autoimmune disease in AD patients than in matched controls. However, in CAH patients, there is only marginal evidence of an imbalance of immune function

(17), and as we found similar increases in infection risk in the CAH cohort, potential etiology-
related immune function is unlikely to explain the increased susceptibility to infections we
observed.

Supraphysiological glucocorticoid doses, as usually administered in the context of chronic
inflammatory disease, is well known to cause changes in the immune system, with consequently
increased risk of bacterial and fungal infections (18). However, this has not been demonstrated
for the physiological replacement doses generally used in patients with PAI. Still, currently
available glucocorticoid replacement therapy does not provide a physiological substitution, with
significant peaks and troughs of cortisol availability during the day following oral intake of
immediate release glucocorticoid preparations. In addition, significant heterogeneity exists in the
management of glucocorticoid replacement in clinical practice; a recent paper recorded 25
different regimens with which glucocorticoid therapy is administered in AD patients receiving a
daily hydrocortisone of 20 mg (19). Therefore, it would come as no surprise that also
physiological dose glucocorticoid therapy is not free of side effects, if administered in a non-
physiological delivery pattern. An improvement in metabolic outcomes after switching from
standard cortisol replacement to more physiological cortisol replacement via continuous
subcutaneous hydrocortisone was previously demonstrated in both AD and CAH patients
(20,21).

Some recent papers have indeed suggested that adverse changes in immune function might
occur with glucocorticoid replacement in PAI. A recent paper documented significantly
decreased natural killer cytotoxicity in patients with PAI (6), which was present in both patients
with autoimmune adrenalitis and those with PAI following bilateral adrenalectomy, indicating that
the underlying etiology did not play a role in these changes in immune function. A recent
randomized control trial including patients with primary and secondary adrenal insufficiency
reported a reduction in respiratory tract infections with modified-release hydrocortisone (22).
However, this was a secondary outcome, based on self-reported questionnaires on infections

and not verified against medical records, thereby providing only limited evidence. A study on

immune function in the same cohort reported dysregulation of circadian gene expression in

peripheral blood mononuclear cells in the PAI patients at baseline, which attenuated after the

switch to modified-release hydrocortisone therapy (23). The findings of our study, including both

patients with AD and CAH, suggest that exogenous glucocorticoid is at least a contributory

factor to the increased infection risk we observed, given that no significant increase in infection

risk was observed in the CAH patients not receiving glucocorticoid therapy.

Both our AD and CAH populations had increased prescription rates for antibiotics and

antifungals. Interestingly, increased prescription rates were also noted in the CAH patients not

receiving glucocorticoid treatment, albeit to a much lower extent. This could possibly be

explained by a lower threshold for prescribing antimicrobials due to the perceived risk of adrenal

crisis in CAH patients; in fact, up to 60% of non-classic CAH patients, who usually do not

receive chronic glucocorticoid replacement, have been reported to have at least partial

glucocorticoid deficiency as assessed by cosyntropin testing (24).

The highest increase in risk of infection in our AD cohort was seen in GIIs, while for the CAH

cohort on glucocorticoids the most significant increase in risk was seen in LRTIs and UTIs;

however, the differences between the three infection groups was not statistically significant. This

may be explained by the age difference between AD and CAH patients, with mean ages of 51.7

and 35.4 years, respectively. Indeed, LRTIs and UTIs are more frequently diagnosed in older

patients (25,26), and this was noted in our matched populations as well (population matched for

AD: LRTIs 4.3%, UTIs 12.5%; population matched for CAH patients on glucocorticoids: LRTIs

1.4%, UTIs 8.5%). Therefore, the higher aIRR of LRTIs and UTIs in CAH patients is probably

related to a difference in age-related background risk.

Our AD cohort had an age and sex distribution similar to the one reported in other papers (2,5),

and the types of prescribed glucocorticoid preparations at baseline in this cohort were not

different from the ones reported in a recent worldwide survey (27). Our CAH cohort was

younger that the AD cohort, consistent with the different etiology of these two diseases, and the

types of glucocorticoids prescribed was similar to those reported in the cross-sectional UK

CaHASE study (28), with the possible exception of lower numbers of dexamethasone users in

our study. Taking this into account, our results can be assumed to be representative of the UK

AD and CAH populations.

Our study has several strengths. We used a large population-based sample of patients of both

sexes, across all adult age groups, with very strict inclusion and exclusion criteria, allowing us to

include only patients with a true diagnosis of AD and CAH. Using the cohort study design,

allowed us to look at longitudinal occurrence of infections and antimicrobial use. There are also

some limitations to our study. Firstly, all data relies on the accurate recording of diagnoses by

GPs and this could have resulted in some degree of misclassification of the exposed cohorts

and of the different episodes of infection. Secondly, we could not evaluate the influence of

different doses or types of glucocorticoid on the outcomes of interest due to the methodology

used and due to the small number of infection events when further subdividing our populations

according to type of glucocorticoid. Thirdly, even though we tried to assess the impact of

associated comorbidities by adjusting for Charlson Comorbidity Index, this does not exclude the

possibility that some other confounders not accounted for in our analyses might have influenced

our results. Lastly, though there is some evidence of an immune-modulatory effect of androgens

(29), we could not take this into account in our population as we had no data on DHEA

replacement therapy in AD patients, since this is a hospital-prescribed drug in the UK; similarly,

we did not have data on biochemical control of androgen excess in the CAH patients.

Our findings have several practical implications. Firstly, given the confirmation of a higher risk of

infections in patients with PAI due to AD and CAH, all healthcare professionals involved in the

care of PAI patients should have a heightened alertness for the possibility of infections in these

patients. This may also provide a case for recommending a vaccination strategy in PAI, e.g.

against *Streptococcus pneumoniae*, the leading cause of LRTIs in adults (30), in order to reduce

the risk of these infections and related morbidity and mortality. Secondly, our paper provides

additional evidence that non-physiological delivery of glucocorticoid replacement by currently

available preparations represents a risk factor for the development of infections. This supports

the case for a therapeutic shift towards more physiological replacement therapy options in these

patients (31). Future studies will have to clarify whether achieving a more physiological delivery

of glucocorticoid replacement will decrease the risk of infections in PAI, with the potential to

result in reduced morbidity and mortality in these patients.

ACKNOWLEDGMENTS

This work was supported by the Medical Research Council UK (Program Grant 0900567, to W

A). K.N. is a UK Research and Innovation (UKRI)/Health Data Research (HDR) UK Innovation

Clinical Fellow. W.A. receives support from the National Institute of Health Research (NIHR)

Birmingham Biomedical Research Centre. The views expressed in this publication are those of

the authors and not necessarily those of the NIHR or the Department of Health and Social Care

UK.

REFERENCES

1. Bancos I, Hahner S, Tomlinson J, Arlt W. Diagnosis and management of adrenal insufficiency. *The lancet Diabetes & endocrinology*. 2015;3(3):216-226.

2. Erichsen MM, Lovas K, Fougner KJ, Svartberg J, Hauge ER, Bollerslev J, Berg JP, Mella B, Husebye ES. Normal overall mortality rate in Addison's disease, but young patients are at risk of premature death. *European journal of endocrinology*. 2009;160(2):233-237.

3. Falhammar H, Frisén L, Norrby C, Hirschberg AL, Almqvist C, Nordenskjöld A, Nordenström A. Increased Mortality in Patients With Congenital Adrenal Hyperplasia Due to 21-Hydroxylase Deficiency. *The Journal of Clinical Endocrinology & Metabolism*. 2014;99(12):E2715-E2721.

4. Bergthorsdottir R, Leonsson-Zachrisson M, Oden A, Johannsson G. Premature mortality in patients with Addison's disease: a population-based study. *The Journal of clinical endocrinology and metabolism*. 2006;91(12):4849-4853.

5. Smans LC, Souverein PC, Leufkens HG, Hoepelman AI, Zelissen PM. Increased use of antimicrobial agents and hospital admission for infections in patients with primary adrenal insufficiency: a cohort study. *European journal of endocrinology*. 2013;168(4):609-614.

6. Bancos I, Hazeldine J, Chortis V, Hampson P, Taylor AE, Lord JM, Arlt W. Primary adrenal insufficiency is associated with impaired natural killer cell function: a potential link to increased mortality. *European journal of endocrinology*. 2017;176(4):471-480.

7. Lewis JD, Schinnar R, Bilker WB, Wang X, Strom BL. Validation studies of the health improvement network (THIN) database for pharmacoepidemiology research. *Pharmacoepidemiology and drug safety*. 2007;16(4):393-401.

8. Blak BT, Thompson M, Dattani H, Bourke A. Generalisability of The Health Improvement Network (THIN) database: demographics, chronic disease prevalence and mortality rates. *Informatics in primary care*. 2011;19(4):251-255.

9.   Iqbal K, Halsby K, Murray RD, Carroll PV, Petermann R. Glucocorticoid
management of adrenal insufficiency in the United Kingdom: assessment using
real-world data. *Endocrine connections.* 2018;8(1):20-31.

10.   Jenkins-Jones S, Parviainen L, Porter J, Withe M, Whitaker MJ, Holden SE,
Morgan CL, Currie CJ, Ross RJM. Poor compliance and increased mortality,
depression and healthcare costs in patients with congenital adrenal hyperplasia.
European journal of endocrinology. 2018;178(4):309-320.

11.   Maguire A, Blak BT, Thompson M. The importance of defining periods of
complete mortality reporting for research using automated data from primary
care. *Pharmacoepidemiology and drug safety.* 2009;18(1):76-83.

12.   Fleming DM, Cross KW, Barley MA. Recent changes in the prevalence of
diseases presenting for health care. *The British journal of general practice : the*
journal of the Royal College of General Practitioners. 2005;55(517):589-595.

13.   UK Data Service Census from https://census.ukdataservice.ac.uk/get-
data/related/deprivation.

14.   Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying
prognostic comorbidity in longitudinal studies: development and validation.
Journal of chronic diseases. 1987;40(5):373-383.

15.   Crooks CJ, West J, Card TR. A comparison of the recording of comorbidity in
primary and secondary care by using the Charlson Index to predict short-term
and long-term survival in a routine linked data cohort. *BMJ open.*
2015;5(6):e007974.

16.   Betterle C, Dal Pra C, Mantero F, Zanchetta R. Autoimmune adrenal insufficiency
and autoimmune polyendocrine syndromes: autoantibodies, autoantigens, and
their applicability in diagnosis and disease prediction. *Endocrine reviews.*
2002;23(3):327-364.

17.   Parlato F, Pisano G, Brillante M, Ferrone R, Cavalcanti MR, Cosentini E, Misiano
G, Brai M, Bellastella A. Immunological pattern in patients with 21-hydroxylase
deficiency. *Journal of endocrinological investigation.* 1994;17(8):635-639.

18.   Fardet L, Petersen I, Nazareth I. Common Infections in Patients Prescribed
381   Systemic Glucocorticoids in Primary Care: A Population-Based Cohort Study.
382   *PLoS medicine*. 2016;13(5):e1002024.
383   19.   Murray RD, Ekman B, Uddin S, Marelli C, Quinkler M, Zelissen PM, the EUAIRI.
384   Management of glucocorticoid replacement in adrenal insufficiency shows
385   notable heterogeneity - data from the EU-AIR. *Clinical endocrinology*.
386   2017;86(3):340-346.
387   20.   Gagliardi L, Nenke MA, Thynne TR, von der Borch J, Rankin WA, Henley DE,
388   Sorbello J, Inder WJ, Torpy DJ. Continuous subcutaneous hydrocortisone
389   infusion therapy in Addison's disease: a randomized, placebo-controlled clinical
390   trial. *The Journal of clinical endocrinology and metabolism*. 2014;99(11):4149-
391   4157.
392   21.   Mallappa A, Nella AA, Sinaii N, Rao H, Gounden V, Perritt AF, Kumar P, Ling A,
393   Liu CY, Soldin SJ, Merke DP. Long-term use of continuous subcutaneous
394   hydrocortisone infusion therapy in patients with congenital adrenal hyperplasia.
395   *Clinical endocrinology*. 2018.
396   22.   Isidori AM, Venneri MA, Graziadio C, Simeoli C, Fiore D, Hasenmajer V,
397   Sbardella E, Gianfrilli D, Pozza C, Pasqualetti P, Morrone S, Santoni A, Naro F,
398   Colao A, Pivonello R, Lenzi A. Effect of once-daily, modified-release
399   hydrocortisone versus standard glucocorticoid therapy on metabolism and innate
400   immunity in patients with adrenal insufficiency (DREAM): a single-blind,
401   randomised controlled trial. *The lancet Diabetes & endocrinology*. 2018;6(3):173-
402   185.
403   23.   Muller L, Quinkler M. Adrenal disease: Imitating the cortisol profile improves the
404   immune system. *Nature reviews Endocrinology*. 2018;14(3):137-139.
405   24.   Stoupa A, Gonzalez-Briceno L, Pinto G, Samara-Boustani D, Thalassinos C,
406   Flechtner I, Beltrand J, Bidet M, Simon A, Piketty M, Laborde K, Morel Y,
407   Bellanne-Chantelot C, Touraine P, Polak M. Inadequate cortisol response to the
408   tetracosactide (Synacthen(R)) test in non-classic congenital adrenal hyperplasia:
409   an exception to the rule? *Hormone research in paediatrics*. 2015;83(4):262-267.

25. Ahmed H, Farewell D, Jones HM, Francis NA, Paranjothy S, Butler CC.
Incidence and antibiotic prescribing for clinically diagnosed urinary tract infection
in older adults in UK primary care, 2004-2014. *PloS one.* 2018;13(1):e0190521.
26. Hak E, Rovers MM, Kuyvenhoven MM, Schellevis FG, Verheij TJ. Incidence of
GP-diagnosed respiratory tract infections according to age, gender and high-risk
co-morbidity: the Second Dutch National Survey of General Practice. *Family*
practice. 2006;23(3):291-294.
27. Forss M, Batcheller G, Skrtic S, Johannsson G. Current practice of glucocorticoid
replacement therapy and patient-perceived health outcomes in adrenal
insufficiency - a worldwide patient survey. *BMC endocrine disorders.* 2012;12:8.
28. Arlt W, Willis DS, Wild SH, Krone N, Doherty EJ, Hahner S, Han TS, Carroll PV,
Conway GS, Rees DA, Stimson RH, Walker BR, Connell JM, Ross RJ. Health
status of adults with congenital adrenal hyperplasia: a cohort study of 203
patients. *The Journal of clinical endocrinology and metabolism.*
2010;95(11):5110-5121.
29. Trigunaite A, Dimo J, Jorgensen TN. Suppressive effects of androgens on the
immune system. *Cellular immunology.* 2015;294(2):87-94.
30. Use of 13-valent pneumococcal conjugate vaccine and 23-valent pneumococcal
polysaccharide vaccine for adults with immunocompromising conditions:
recommendations of the Advisory Committee on Immunization Practices (ACIP).
MMWR Morbidity and mortality weekly report. 2012;61(40):816-819.
31. Whitaker M, Debono M, Huatan H, Merke D, Arlt W, Ross RJ. An oral
multiparticulate, modified-release, hydrocortisone replacement therapy that
provides physiological cortisol exposure. *Clinical endocrinology.* 2014;80(4):554-
561.

Table 1: Baseline characteristics of patients with Addison's disease (AD) and matched unexposed patients.

	AD patients (n = 1580)	Matched unexposed patients (n = 3158)
Age years, mean ± SD	51.7 ± 18.5	51.7 ± 18.5
Male sex, n. (%)	666 (42.2%)	1.330 (42.1%)
BMI	Total n = 1227	Total n = 2264
median (IQR)	24.3 (21.5-27.6)*	25.5 (22.6-28.9)
□18.5 kg/m2, n. (%)	87 (7.1%)	51 (2.3%)
18.5-25 kg/m^2, n. (%)	606 (49.4%)	973 (43.0%)
25-30 kg/m^2, n. (%)	362 (29.5%)	791 (34.9%)
≥30 kg/m^2, n. (%)	172 (14.0%)	449 (19.8%)
Missing, n.	353	894
Smoking status	Total = 1384	Total = 2679
Non-smoker, n. (%)	882 (63.7%)*	1548 (57.8%)
Ex-smoker, n. (%)	250 (18.1%)	490 (18.3%)
Smoker, n. (%)	252 (18.2%)*	641 (23.9%)
Missing, n.	196	479
Townsend Deprivation Index	Total = 1373	Total = 2780
1 (least deprived), n. (%)	350 (25.5%)	743 (26.7%)
2, n. (%)	306 (22.3%)	590 (21.2%)
3, n. (%)	290 (21.1%)	607 (21.8%)
4, n. (%)	255 (18.6%)	471 (16.9%)
5 (most deprived), n. (%)	172 (12.5%)	369 (13.3%)
Missing, n.	207	378
Charlson Comorbidity Index		
0 (no comorbidities), n. (%)	863 (54.6%)*	2263 (71.7%)
1, n. (%)	377 (23.9%)*	536 (17.0%)
≥2 (more comorbidities), n. (%)	340 (21.5%)*	359 (11.4%)
Associated autoimmune comorbidities		
Hyperthyroidism, n. (%)	40 (2.5%)*	16 (0.5%)
Hypothyroidism, n. (%)	457 (28.9%)*	122 (3.9%)
Rheumatoid arthritis, n. (%)	25 (1.6%)	37 (1.2%)
Type 1 diabetes mellitus, n. (%)	127 (8.0%)*	15 (0.5%)
Inflammatory bowel disease, n. (%)	29 (1.8%)*	22 (0.7%)
Crohn's disease, n. (%)	9 (0.6%)	9 (0.3%)
Ulcerative colitis, n. (%)	20 (1.3%)*	11 (0.4%)
Coeliac disease, n. (%)	25 (1.6%)*	10 (0.3%)
Multiple sclerosis, n. (%)	n < 5	8 (0.3%)
Pernicious anaemia, n. (%)	41 (2.6%)*	15 (0.5%)
Systemic lupus erythematosus, n. (%)	6 (0.4%)	n < 5

* p <0.05 for AD patients vs. unexposed cohort.

Table 2: Baseline characteristics of the CAH patients (overall cohort and sub-cohorts divided by glucocorticoid therapy and no glucocorticoid therapy) and respective matched unexposed controls.

	CAH cohort (n = 602)	Matched unexposed cohort (n = 1204)	CAH cohort on glucocorticoids (n = 254)	Matched unexposed cohort (n = 508)	CAH cohort not on glucocorticoids (n = 348)	Matched unexposed cohort (n = 696)
Age, years, mean ± SD	35.4 ± 16.3	35.5 ± 16.2	33.4 ± 15.1	33.5 ± 15.0	36.9 ± 17.0[†]	37.0 ± 16.9
Male sex, n. (%)	167 (27.7%)	334 (27.7%)	80 (31.5%)	160 (31.5%)	87 (25.0%)	174 (25.0%)
BMI	Total = 438	Total = 835	Total = 186	Total = 340	Total = 252	Total = 495
median (IQR)	26.9 (23.2-31.2)*	24.0 (21.0-28.0)	27.0 (23.2-32.0)*	24.0 (21.3-27.9)	26.9 (23.2-30.9)*	24.4 (21.8-28.3)
☐18.5, n. (%)	12 (2.7%)	35 (4.2%)	8 (4.3%)	9 (2.6%)	4 (1.6%)	26 (5.2%)
18.5-25, n. (%)	162 (37.0%)	431 (51.6%)	68 (36.6%)	187 (55.0%)	94 (37.3%)	244 (49.3%)
25-30, n. (%)	133 (30.4%)	224 (26.8%)	52 (28.0%)	90 (26.5%)	81 (32.1%)	134 (27.1%)
≥30, n. (%)	131 (29.9%)	145 (17.4%)	58 (31.2%)	54 (15.9%)	73 (29.0%)	91 (18.4%)
Missing, n.	164	369	68	168	96	201
Smoking status	Total = 524	Total = 1048	Total = 212	Total = 423	Total = 312	Total = 625
Non-smoker, n. (%)	349 (66.6%)	670 (63.9%)	138 (65.1%)	277 (65.5%)	211 (67.6%)	393 (62.9%)
Ex-smoker, n. (%)	71 (13.6%)	142 (13.6%)	28 (13.2%)	50 (11.8%)	43 (13.8%)	92 (14.7%)
Smoker, n. (%)	104 (19.9%)	236 (22.5%)	46 (21.7%)	96 (22.7%)	58 (18.6%)	140 (22.4%)
Missing, n.	78	156	42	85	36	71
Townsend Deprivation Index	Total = 516	Total = 1057	Total = 223	Total = 459	Total = 293	Total = 598
1 (least deprived), n. (%)	123 (23.8%)	235 (22.2%)	52 (23.3%)	107 (23.3%)	71 (24.2%)	128 (21.4%)
2, n. (%)	107 (20.7%)	222 (21.0%)	46 (20.6%)	92 (20.0%)	61 (20.8%)	130 (21.7%)
3, n. (%)	114 (22.1%)	232 (22.0%)	51 (22.9%)	99 (21.6%)	63 (21.5%)	133 (22.2%)
4, n. (%)	104 (20.2%)	225 (21.3%)	44 (19.7%)	98 (21.4%)	60 (20.5%)	127 (21.2%)
5 (most deprived), n. (%)	68 (13.2%)	143 (13.5%)	30 (13.5%)	63 (13.7%)	38 (13.0%)	80 (13.4%)
Missing, n.	86	147	31	49	55	98
Charlson Comorbidity Index						
0 (no comorbidities), n. (%)	457 (75.9%)	907 (75.3%)	187 (73.6%)	394 (77.6%)	270 (77.6%)	513 (73.7%)
1, n. (%)	103 (17.1%)	242 (20.1%)	50 (19.7%)	96 (18.9%)	53 (15.2%)	146 (21.0%)
≥2 (more comorbidities), n. (%)	42 (7.0%)	55 (4.6%)	17 (6.7%)	18 (3.5%)	25 (7.2%)	37 (5.3%)

* p <0.05 for CAH patients vs. unexposed cohort.

[†] p <0.05 for CAH patients on glucocorticoids vs. CAH patients not on glucocorticoids.

Table 3: Absolute and relative risk of infections in AD patients and matched cohort.

	AD cohort (n = 1580)	Matched unexposed cohort (n = 3158)
Lower respiratory tract infections		
Outcome events, n. (%)	130 (8.2)	137 (4.3)
Person-years	10,337	22,836
Crude incidence rate/1000-person years	12.58	6.00
Follow-up years, median (IQR)	4.87 (1.78-10.20)	5.78 (2.37-11.12)
Unadjusted incidence rate ratio (95% CI) p-value	2.10 (1.65-2.66) p <0.001	
Adjusted incidence rate ratio (95% CI)† p-value	2.11 (1.64-2.69) p <0.001	
Urinary tract infections		
Outcome events, n. (%)	282 (17.9)	396 (12.5)
Person-years	9,248	21,003
Crude incidence rate/1000-person years	30.49	18.85
Follow-up years, median (IQR)	4.09 (1.44-9.14)	5.08 (2.11-10.26)
Unadjusted incidence rate ratio (95% CI) p-value	1.62 (1.39-1.88) p <0.001	
Adjusted incidence rate ratio (95% CI)† p-value	1.51 (1.29-1.77) p <0.001	
Gastro-intestinal infections		
Outcome events, n. (%)	194 (12.3)	110 (3.5)
Person-years	9,598	22,662
Crude incidence rate/1000-person years	20.21	4.85
Follow-up years, median (IQR)	4.49 (1.67-9.31)	5.63 (2.37-11.03)
Unadjusted incidence rate ratio (95% CI) p-value	4.16 (3.30-5.26) p <0.001	
Adjusted incidence rate ratio (95% CI)† p-value	3.80 (2.99-4.84) p <0.001	

†Adjusted for age, gender, smoking status, BMI, Townsend Deprivation Index and Charlson Comorbidity Index.

Table 4: Absolute and relative risk of infections in CAH patients and matched control cohort.

	CAH cohort (n = 602)	Matched unexposed cohort (n = 1204)	CAH cohort on glucocorticoids (n = 254)	Matched unexposed cohort (n = 508)	CAH cohort not on glucocorticoids (n = 348)	Matched unexposed cohort (n = 696)
Lower respiratory tract infections						
Outcome events, n. (%)	22 (3.7)	19 (1.6)	12 (4.7)	7 (1.4)	10 (2.9)	12 (1.7)
Person-years	3843	7842	1924	3567	1919	4275
Crude incidence rate/1000-person years	5.72	2.42	6.24	1.96	5.21	2.81
Follow-up years, median (IQR)	4.83 (1.92-9.54)	5.10 (2.04-9.80)	6.00 (2.60-12.06)	5.17 (2.45-10.77)	4.07 (1.54-8.00)	4.93 (1.86-9.37)
Unadjusted incidence rate ratio (95% CI) p-value	2.36 (1.28-4.36) p = 0.01		3.18 (1.25-8.07) p = 0.02		1.86 (0.80-4.30) p = 0.15	
Adjusted incidence rate ratio (95% CI)[†] p-value	2.36 (1.25-4.43) p = 0.01		3.23 (1.21-8.61) p = 0.02		1.82 (0.77-4.30) p = 0.17	
Urinary tract infections						
Outcome events, n. (%)	83 (13.8)	130 (10.8)	45 (17.7)	43 (8.5)	38 (10.9)	87 (12.5)
Person-years	3478	7217	1709	3374	1769	3843
Crude incidence rate/1000-person years	23.87	18.01	26.33	12.75	21.48	22.64
Follow-up years, median (IQR)	4.15 (1.59-8.80)	4.64 (1.83-8.80)	4.97 (2.09-9.77)	4.84 (2.22-9.77)	3.42 (1.24-7.53)	4.18 (1.69-7.90)
Unadjusted incidence rate ratio (95% CI) p-value	1.32 (1.01-1.74) p = 0.05		2.07 (1.36-3.14) p <0.001		0.95 (0.65-1.39) p = 0.79	
Adjusted incidence rate ratio (95% CI)[†] p-value	1.40 (1.06-1.85) p = 0.02		2.20 (1.43-3.34) p <0.001		1.00 (0.68-1.48) p = 0.99	
Gastro-intestinal infections						
Outcome events, n. (%)	29 (4.8)	52 (4.3)	23 (9.1)	23 (4.5)	6 (1.7)	29 (4.2)
Person-years.	3773	7678	1825	3486	1948	4192
Crude incidence rate/1000-person years	7.69	6.77	12.60	6.60	3.08	6.92
Follow-up years, median (IQR)	4.70 (1.83-9.63)	4.94 (2.04-9.69)	5.82 (2.12-11.38)	5.14 (2.47-10.38)	4.09 (1.50-8.11)	4.79 (1.86-9.13)
Unadjusted incidence rate ratio (95% CI) p-value	1.13 (0.72-1.79) p = 0.59		1.91 (1.07-3.40) p = 0.03		0.45 (0.18-1.07) p = 0.07	
Adjusted incidence rate ratio (95% CI)[†] p-value	1.10 (0.69-1.75) p = 0.70		1.93 (1.06-3.52) p = 0.03		0.45 (0.18-1.08) p = 0.07	

[†]Adjusted for age, gender, smoking status, BMI, Townsend Deprivation Index and Charlson Comorbidity Index.

Table 5: Antimicrobial prescriptions counts in AD patients compared to the matched control cohort.

	AD cohort (n = 1580)	Matched unexposed cohort (n = 3158)
Antibiotic prescriptions		
Count of prescriptions, n.	13286	15884
Person-years	10767	23308
Count rates (per 1000 years)	1234	681
Follow-up years, median (IQR)	5.12 (1.95-10.77)	5.90 (2.54-11.33)
Unadjusted incidence rate ratio (95% CI) p-value	1.81 (1.77-1.85) p <0.001	
Adjusted incidence rate ratio (95% CI)[†] p-value	1.73 (1.69-1.77) p <0.001	
Antifungal prescriptions		
Count of prescriptions, n.	1191	1213
Person-years	10767	23308
Count rates (per 1000 years)	111	52
Follow-up years, median (IQR)	5.12 (1.95-10.77)	5.90 (2.54-11.33)
Unadjusted incidence rate ratio (95% CI) p-value	2.13 (1.96-2.30) p <0.001	
Adjusted incidence rate ratio (95% CI)[†] p-value	1.89 (1.74-2.05) p <0.001	

[†]Adjusted for age, gender, smoking status, BMI, Townsend Deprivation Index and Charlson Comorbidity Index.

Table 6: Antimicrobial prescription counts in CAH patients compared to the matched control cohort.

	CAH cohort (n = 602)	Matched unexposed cohort (n = 1204)	CAH cohort on glucocorticoids (n = 254)	Matched unexposed cohort (n = 508)	CAH cohort not on glucocorticoids (n = 348)	Matched unexposed cohort (n = 696)
Antibiotics prescriptions						
Count of prescriptions, n.	3543	4930	2134	2088	1409	2842
Person-years	3941	7957	1974	3611	1967	4346
Count rates (per 1000 years)	899	619	1081	578	716	654
Follow-up years, median (IQR)	4.85 (1.95-10.32)	5.23 (2.09-9.90)	6.12 (2.67-12.48)	5.32 (2.54-10.87)	4.12 (1.54-8.30)	5.20 (1.88-9.62)
Unadjusted incidence rate ratio (95% CI) p-value	1.45 (1.39-1.51) p <0.001		1.87 (1.76-1.99) p <0.001		1.10 (1.03-1.17) p = 0.01	
Adjusted incidence rate ratio (95% CI)[†] p-value	1.46 (1.39-1.52) p <0.001		1.77 (1.66-1.89) p <0.001		1.15 (1.08-1.23) p <0.001	
Antifungals prescriptions						
Count of prescriptions, n.	302	330	159	130	143	200
Person-years	3941	7957	1974	3611	1967	4346
Count rates (per 1000 years)	77	41	81	36	73	46
Follow-up years, median (IQR)	4.85 (1.95-10.32)	5.23 (2.09-9.90)	6.12 (2.67-12.48)	5.32 (2.54-10.87)	4.12 (1.54-8.30)	5.20 (1.88-9.62)
Unadjusted incidence rate ratio (95% CI) p-value	1.85 (1.58-2.16) p <0.001		2.24 (1.77-2.82) p <0.001		1.58 (1.27-1.96) p <0.001	
Adjusted incidence rate ratio (95% CI)[†] p-value	1.69 (1.44-1.99) p <0.001		1.91 (1.50-2.43) p <0.001		1.47 (1.18-1.83) p <0.001	

[†]Adjusted for age, gender, smoking status, BMI, Townsend Deprivation Index and Charlson Comorbidity Index.

Figure 1 - Forest plot of outcomes. Panel A: adjusted Incidence Rate Ratio (aIRR) for infections and antimicrobial prescriptions in AD and CAH cohorts. Panel B: aIRR for infections and antimicrobial prescriptions in CAH patients separately for patients with and without chronic glucocorticoid treatment.