Extensive intraspecies cryptic variation in an ancient embryonic gene regulatory network

Yamila N. Torres Cleuren[1,2]*, Chee Kiang Ewe[2], Kyle C. Chipman[2], Emily Mears[1], Cricket G. Wood[2], Coco A.E. Al-Alami[1], Melissa R. Alcorn[2], Thomas L. Turner[3], Pradeep M. Joshi[2], Russell G. Snell[1], and Joel H. Rothman[1,2,3]

1. School of Biological Sciences, University of Auckland, Auckland, New Zealand

2. Department of MCD Biology and Neuroscience Research Institute, University of California Santa Barbara, CA, USA

3. Department of Ecology, Evolution, and Marine Biology, University of California Santa Barbara, CA, USA

*Current address: Computational Biology Unit, Department of Informatics, University of Bergen, Bergen, 5020 Norway

Author for correspondence: joel.rothman@lifesci.ucsb.edu

Keywords: SKN-1, *C. elegans*, endoderm, GWAS, development, gene regulatory networks, Genotype-By-Sequencing, cryptic variation

ABSTRACT

Innovations in metazoan development arise from evolutionary modification of gene regulatory networks (GRNs). We report widespread cryptic variation in the requirement for two key regulatory inputs, SKN-1/Nrf2 and MOM-2/Wnt, into the *C. elegans* endoderm GRN. While some natural variants show a nearly absolute requirement for these two regulators, in others, most embryos differentiate endoderm in their absence. GWAS and analysis of recombinant inbred lines reveal multiple genetic regions underlying this broad phenotypic variation. We observe a reciprocal trend, in which genomic variants, or knockdown of endoderm regulatory genes, that result in a high SKN-1 requirement often show low MOM-2/Wnt requirement and *vice-versa,* suggesting that cryptic variation in the endoderm GRN may be tuned by opposing requirements for these two key regulatory inputs. These findings reveal that while the downstream components in the endoderm GRN are common across metazoan phylogeny, initiating regulatory inputs are remarkably plastic even within a single species.

INTRODUCTION

While the core regulatory machinery that specifies embryonic germ layers and major organ identity in the ancestor of modern animals is largely conserved in all extant animals, GRN architecture must be able to accommodate substantial plasticity to allow for evolutionary innovation in developmental strategies, changes in selective pressures, and genetic drift [1,2]. Genetic variation, often with neutral effects on fitness, provides for plasticity in GRN structure and implementation [2]. Although studies of laboratory strains of model organisms with a defined genetic background have been highly informative in identifying the key regulatory nodes in GRNs that specify developmental processes [3–5], these approaches do not reveal the evolutionary basis for plasticity in these networks. The variation and incipient changes in GRN function and architecture can be discovered by analyzing phenotypic differences resulting from natural genetic variation present in distinct isolates of a single species [6–8].

The endoderm has been proposed to be the most ancient of the three embryonic germ layers in metazoans [9,10], having appeared prior to the advent of the Bilateria about 600 Mya [11]. It follows, therefore, that the GRN for endoderm in extant animals has undergone substantial modifications over the long evolutionary time span since its emergence. However, the core transcriptional machinery for endoderm specification and differentiation appears to share common mechanisms across metazoan phylogeny. For example, cascades of GATA-type transcription factors function to promote endoderm development not only in triploblastic animals but also in the most basal metazoans that contain a digestive tract [12–16]. Among the many observations supporting a common regulatory mechanism for establishing the endoderm, it has been found that the endoderm-determining GATA factor, END-1, in the nematode *C. elegans*, is sufficient to activate

endoderm development in cells that would otherwise become ectoderm in Xenopus [17]. This

indicates that the role of GATA factors in endoderm development has been preserved since

the nematodes and vertebrates diverged from a common ancestor that lived perhaps 600

Mya.

To assess the genetic basis for evolutionary plasticity and cryptic variation underlying

early embryonic germ layer specification, we have analyzed the well-described GRN for

endoderm specification in *C. elegans*. The E cell, which is produced in the very early *C. elegans*

embryo, is the progenitor of the entire endoderm, which subsequently gives rise exclusively

to the intestine. The EMS blastomere at the four-cell stage divides to produce the E founder

cell and its anterior sister, the MS founder cell, which is the progenitor for much of the

mesoderm [18]. Both E and MS fates are determined by maternally provided SKN-1, an

orthologue of the vertebrate Nrf2 bZIP transcription factor [19–21]. In the laboratory N2

strain, elimination of maternal SKN-1 function (through either knockdown or knockout)

results in fully penetrant embryonic lethality as a result of misspecification of EMS cell

descendants. In these embryos, the fate of MS is transformed to that of its cousin, the

mesectodermal progenitor C cell. E cells similarly adopt a C cell-like fate in a majority, but not

all, of these embryos [19]. SKN-1 initiates mesendoderm development via the GRN in E and

MS cells in part by activating zygotic expression of the MED-1/2 divergent GATA transcription

factors [22,23]. This event mobilizes a cascade of GATA factors in the E cell lineage that

ultimately direct intestinal differentiation [21,24–26].

This differential requirement for SKN-1 in endoderm (E) and mesoderm (MS)

development is determined by its combinatorial action with triply redundant Wnt, MAPK, and

Src signaling systems, which act together to polarize EMS [27–30]. MOM-2/Wnt acts through

the MOM-5/Frizzled receptor, mobilizing WRM-1/β-catenin, resulting in its cytoplasmic

accumulation in the posterior side of EMS. WRM-1, together with LIT-1/NLK kinase, alters

both the nucleocytoplasmic distribution and activity of the Wnt effector POP-1/Tcf [30–32],

converting it from a repressor of endoderm in the MS cell lineage to an activator in the E cell

lineage [33–38]. Loss of MOM-2 expression in the laboratory N2 strain results in a partial

gutless phenotype, while removal of both MOM-2 and SKN-1, through either knockdown or

knockout, leads to a completely penetrant loss of gut [30], revealing their genetically

redundant roles.

The regulatory relationship between SKN-1 and POP-1, the effector of Wnt signaling,

shows substantial variation even in species that diverged 20-40 million years ago, suggesting

significant evolutionary plasticity in this key node in the endoderm GRN. *C. elegans* embryos

lacking maternal POP-1 always make gut, both in the normal E cell lineage and in the MS cell

lineage. However, in embryos lacking both SKN-1 and POP-1, endoderm is virtually never

made, implying that these two factors constitute a Boolean "OR" logic gate. In contrast,

removal of either SKN-1 or POP-1 alone in *C. briggsae* causes >90% of embryos to lack gut,

indicative of an "AND" logic gate (Fig. 1A, B) [39].

In this study, we sought to determine whether the plasticity in regulatory logic of the

two major inputs into endoderm development is evident within a single species. The

availability of many naturally inbred variants (isotypes) of *C. elegans* that show widespread

genomic variation [40–42], provides a genetically rich resource for investigating potential

quantitative variation in developmental GRNs. We report here that the requirement for

activation of the endoderm GRN by SKN-1 or MOM-2, but not POP-1, is highly variable

between natural *C. elegans* isolates, and even between closely related isotypes. Genome-

wide association studies (GWAS) in isolates from the natural populations and targeted

analysis of recombinant inbred lines (RILs), revealed that a multiplicity of loci and their

interactions are responsible for the variation in the developmental requirement for SKN-1

and MOM-2. We found a complex, but frequently reciprocal requirement for SKN-1 and

MOM-2 among variants underlying these phenotypes: loci associated with a high

requirement for SKN-1 in endoderm development tend to show a more relaxed requirement

for MOM-2 and *vice-versa*. Consistent with this finding, three other endoderm regulatory

factors, RICT-1, PLP-1, and MIG-5, show similar inverse relationships between these two GRN

inputs. These findings reveal that the activation of the GRN network for a germ layer, one of

the most critical early developmental switches in embryos, is subject to remarkable genetic

plasticity within a species and that the dynamic and rapid change in network architecture

reflects influences distributed across many genetic components that affect both SKN-1 and

Wnt pathways.

Fig 1. Endoderm regulatory pathway and scoring of gut differentiation.

A) Under normal conditions, signaling from the posterior P_2 cell (Wnt, MAPK and Src) results in asymmetric cortical localization of Wnt signaling pathway components in EMS leading to POP-1 asymmetry in the descendants of EMS, with high levels of nuclear POP-1 in anterior MS and low levels of nuclear POP-1 in the posterior, E, daughter cell. In the anterior MS cell, high nuclear POP-1 represses the END genes, allowing SKN-1 to activate MS fate. In the posterior E cell, which remains in contact with P2, POP-1 is converted to an activator and, along with SKN-1, activates the END genes, resulting in endoderm fate. Loss of *skn-1,* either by RNAi or in loss-of-function mutants, causes 100% of the embryos to arrest; in 70% of the arrested embryos, EMS gives rise to two C-like cells, while in the remaining 30% only MS is converted to a C fate; the posterior daughter retains its E fate. Loss of *mom-2* leads to 100% embryonic arrest with a partially penetrant E→MS cell fate transformation, resulting in two MS-like daughter cells in ~72% of the embryos. (B) Regulatory logic of SKN-1 and POP-1 in E specification in *C. elegans, C. briggsae* and a hypothetical intermediate state. POP-1* denotes the activated state. (C-H) Gut visualization in embryos affected by *skn-1* RNAi. (C-E) arrested embryos without endoderm, (F-H) arrested embryos with endoderm. (C, F) DIC images of arrested embryos ~12 hours after egg laying. (D, G) the same embryos expressing the gut-specific *elt-2::GFP* reporter, and (E,H) birefringent gut granules under polarized light. All embryos showing gut birefringence also show *elt-2::GFP* expression. (I, J) Fields of arrested *skn-1(RNAi)* embryos in wild isolate strains JU1491 (I) and JU440 (J), which reflect the extremes in the spectrum of requirement of SKN-1 in gut development at 0.9% and 60%, respectively.

143 **MATERIALS AND METHODS**

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (C. elegans)	Wild isolate refer to supplementary file 1	CGC		
strain, strain background (C. elegans)	JJ185	CGC		dpy-13(e184) skn-1(zu67) IV; mDp1 (IV; f).
strain, strain background (C. elegans)	JR3666	This study		(elt-2::GFP) X ; (ifb-2::CFP) IV
strain, strain background (C. elegans)	EU384	CGC		dpy-11(e1180) mom-2(or42) V/nT1 [let-?(m435)] (IV;V).
strain, strain background (C. elegans)	JJ1057	CGC		pop-1(zu189) dpy-5(e61)/hT1 I; him-5(e1490)/hT1 V.
strain, strain background (C. elegans)	KQ1366	CGC		rict-1(ft7) II.
strain, strain background (C. elegans)	SU351	CGC		mig-5(rh94)/mIn1 [dpy-10(e128) mIs 14] II.
strain, strain background (C. elegans)	RB1711	CGC		plp-1(ok2155) IV.
strain, strain background (C. elegans)	JR3493-JR3590	This study		N2xMY16 RILs

strain, strain background (C. elegans)	MT3414	CGC		dpy-20(e1282) unc-31(e169) unc-26(e205) IV.
strain, strain background (C. elegans)	DA491	CGC		dpy-20(e1282) unc-30(e191) IV.
strain, strain background (C. elegans)	JR2750	[43]		bli-6(Sc16)unc-22(e66)/unc-24(e138)fus-1(w13) dpy-20(e2017)IV
strain, strain background (C. elegans)	JR3812 (NIL 1)	This study		NIL N2XMY16
strain, strain background (C. elegans)	JR3813 (NIL 2)	This study		NIL N2XMY16
strain, strain background (C. elegans)	JR3814 (NIL 3)	This study		NIL N2XMY16
strain, strain background (C. elegans)	JR3815 (NIL 4)	This study		NIL N2XMY16
strain, strain background (C. elegans)	JR3816 (NIL 5)	This study		NIL N2XMY16
strain, strain background (C. elegans)	JR3817 (NIL 6)	This study		NIL N2XMY16
antibody	MH33	DSHB		
antibody	AHP418	Serotec Bio-Rad		
antibody	ab150116	Abcam		Goat Anti-Mouse IgG H&L (Alexa Fluor® 594)
antibody	ab150077	Abcam		Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488)
software, algorithm	R v 3.2.3	The R Foundation		
software, algorithm	PLINK	http://pngu.mgh.harvard.edu/purcell/plink/		

C. elegans strains and maintenance

All wild isolates, each with a unique haplotype [42], were obtained from the Caenorhabditis Genetics Center (CGC) (see Supplementary File 1). Worm strains were maintained as described [44] and all experiments were performed at 20°C unless noted otherwise.

RNAi

Feeding-based RNAi experiments were performed as described [45]. RNAi clones were obtained from either the Vidal [46] or Ahringer libraries [47]. RNAi bacterial strains were grown at 37°C in LB containing 50 μg/ml ampicillin. The overnight culture was then diluted 1:10. After 4 hours of incubation at 37°C, 1 mM of IPTG was added and 60μl was seeded onto 35mm agar plates containing 1 mM IPTG and 25 μg/ml carbenicillin. Seeded plates were allowed to dry and used within five days. Five to 10 L4 animals were placed on RNAi plate. 24 hours later, they were transferred to another RNAi plate and allowed to lay eggs for four or 12 hours (12 hours for *skn-1* RNAi and four hours for the other RNAi). The adults were then removed, leaving the embryos to develop for an extra 7-9 hours. Embryos were quantified and imaged on an agar pad using a Nikon Ti-E inverted microscope. We chose to perform RNAi on agar plates to maximize sensitivity, robustness, and reproducibility of the assay, as liquid culture RNAi can introduce variability owing to aggregation and settling of bacteria, which affects RNAi efficacy [48]. In addition, performing RNAi on agar plates allowed us to collect large numbers of embryos with which to quantify gut formation (as described below).

Antibody staining

The embryonic gut cells and nuclei of all cells were stained with MH33 (mouse anti-IFB-2, deposited to the DSHB by Waterston, R.H.) and AHP418 (rabbit anti-acetylated histone H4,

Serotec Bio-Rad) respectively. Fixation and permeabilization were carried out as described

previously [49]. Goat anti-mouse Alexa Fluor® 594 and goat anti-rabbit Alexa Fluor® 488

secondary antibodies were used at 1:1000 dilution.

Quantification of endoderm specification

Gut was scored by the presence of birefringent gut granule in arrested embryos [50,51]. For

skn-1(RNAi), the laboratory strain N2, which shows invariable ~30% of embryos with

endoderm, was used as a control for all experiments.

Introgression of *skn-1(zu67)*, *pop-1(zu189)*, and *mom-2(or42)* alleles into wild isolate

backgrounds

To introgress *skn-1(zu67)* into wild isolates (WI), males from the wild isolate strains were

crossed to JJ186 *dpy-13(e184) skn-1(zu67) IV; mDp1 (IV;f)* hermaphrodites. mDp1 is a free

duplication maintained extrachromosomally that rescues the Dpy and lethal phenotypes of

dpy-13(e184) and *skn-1(zu67)* respectively. mDp1 segregates in a non-Mendelian fashion and

animals that have lost the free duplication are Dpy and produce dead offspring. Wild type F1

hermaphrodites that have lost the free duplication, as determined by the presence of 1/4 Dpy

progeny in the F2 generation, were selected. 10 single non-Dpy F2 hermaphrodite

descendants from F1 animals heterozygous for *skn-1(zu67)* (2/3 of which are expected to be

of the genotype *WI dpy-13(+) skn-1(+) / dpy-13(e184) skn-1(zu67)* were backcrossed to their

respective parental wild strain. 10 F3 hermaphrodites were picked to individual plates. Half

of the F3 cross progeny are expected to be heterozygous for *dpy-13(e184) skn-1(zu67)*, as

evidenced by presence of F4 Dpy progeny that produced dead embryos. Non-Dpy siblings

were used to continue the introgression as described. This strategy was repeated for at least

five rounds of introgression. The embryonic gutless phenotype in the progeny of the Dpy

animals was quantified.

Similarly, to introgress *pop-1(zu189)* or *mom-2(or42)* alleles into wild isolates, JJ1057

pop-1(zu189) dpy-5(e61)/hT1 I; him-5(e1490)/hT1V or EU384 *dpy-11(e1180) mom-2(or42)*

V/nT1 [let-?(m435)] (IV;V) were used, respectively. The mutant strain was crossed to the wild

isolates. Non-Dpy F2 animals heterozygous for the chromosomal mutation were selected and

backcrossed to their respective parental wild strain for at least four rounds of introgression

for *pop-1* and seven rounds for *mom-2*. The embryonic gutless phenotype in the progeny of

the Dpy animals was quantified, as above.

Statistical Analyses: GWAS

All data were analyzed and plotted using R software v 3.2.3 (https://www.r-

project.org/). GWAS for both phenotypes was performed using *C. elegans* wild isolates and

a previously published SNP map containing 4,690 SNPs [42] with the EMMA R package. P-

values were calculated using mixed model analysis [52] (emma.REML.t() function) and

identity-by-state (IBS) kinship matrix to account for population structure. For *skn-1* and *mom-*

2 RNAi phenotypic data, a genome-wide permutation-based FDR was also calculated for the

EMMA results from 10,000 permuted values [53,54].

Phylogenetic and geographical analyses

Phylogenetic trees were constructed from 4,690 polymorphisms using R package "ape"

[55]. Neighbor-joining algorithm based on pairwise distances was used. Phylogenetic signal

(Pagel's λ statistics) was measured using "phylosig()" function in phytools R package [56,57].

Statistical significance of λ was obtained by comparing the likelihood a model accounting for

the observed λ with the likelihood of a model that assumes complete phylogenetic

independence.

Geographic information for strains were obtained from Andersen *et al.* [42], available in

Supplementary File 1, together with the corresponding *skn-1(RNAi)* and *mom-2(RNAi)*

phenotypes.

Correlation Analysis

To test for the relationship between *mom-2 (RNAi)* and *skn-1 (RNAi)* phenotypic data, the

differences between median phenotypic values for each SNP were calculated independently

on a genome-wide level for the wild isolates. In order to correct for LD, SNPs were pruned

with PLINK (http://pngu.mgh.harvard.edu/purcell/plink/) [58] and only a subset of SNPs was

used for the correlation analyses. Outliers were removed from the calculations by using z-

score with a cutoff of 1.96 (i.e., 95% of values fall within ±1.96 in a normal distribution).

RIL construction and Genotype-By-Sequencing (GBS)

Recombinant inbred lines (RILs) were created by crossing an N2 hermaphrodite and an MY16

male. 120 F2 progeny were cloned to individual plates and allowed to self-fertilize for 10

generations. A single worm was isolated from each generation to create inbred lines. A total

of 95 lines were successfully created and frozen stocks were immediately created and kept at

-80°C (Supplementary File 2), prior to DNA sequencing.

DNA was extracted using Blood and Tissue QIAGEN kit from worms from each of the RILs

grown on four large NGM plates (90x15mm) with OP50 *E. coli* until starved (no more than a

day). Samples were submitted in 96-well plate format at 10 ng/µl < n < 30 ng/µl. GBS libraries

were constructed using digest products from ApeKI (GWCGC), using a protocol modified from

[59]. After digestion, the barcoded adapters were ligated and fragments < 100bp were

sequenced as single-end reads using an Illumina HiSeq 2000 lane (100 bp, single-end reads).

SNP calling was performed using the GBSversion3 pipeline in Trait Analysis by aSSociation,

Evolution and Linkage (TASSEL) [60]. Briefly, fastq files were aligned to reference genome

WS252 using BWA v. 0.7.8-r455 and SNPs were filtered using vcftools [61]. Samples with

greater than 90% missing data and SNPs with minor allele frequencies (mAF) of <1% were

excluded from analysis, identifying 27,396 variants.

QTL mapping using R/qtl

Variants identified by GBS pipeline were filtered to match the SNPs present in the parental

MY16 strain (using vcftools –recode command), and variants were converted to a 012 file

(vcftools –012 command). Single-QTL analysis was performed in R/QTL [62] using 1770

variants and 95 RILs. Significant QTL were determined using Standard Interval Mapping

(scanone() "em") and genome-wide significance thresholds were calculated by permuting the

phenotype (N =1,000). Change in log-likelihood ratio score of 1.5 was used to calculate 95%

confidence intervals and define QTL regions [63]. SNP data for the RILs and their

corresponding phenotypes used in analysis are shown in Supplementary Files 2 and 3.

Creation of Near-Isogenic Lines (NILs)

Three N2-derived mutant strains were used to introgress regions from chromosome IV from

N2 into the MY16 strain background and vice-versa. For both types of crosses, N2 was always

used as the maternal line. The following strains were used:

● DA491: *dpy-20(e1282) unc-30(e191) IV.*

● JR2750: *bli-6(sc16) unc-22(e66)/unc-24(e138)fus-1(w13) dpy-20(e2017) IV*. Worms

segregating Bli Unc were selected for crosses.

● MT3414: *dpy-20(e1282) unc-31(e169) unc-26(e205) IV*.

To introgress the N2 region into the MY16 genetic background, hermaphrodites from the N2-

derived strains containing genetic markers flanking the genomic region of interest were

crossed with MY16 males (Supplementary Fig. 7). After successful mating, 10 F1

heterozygotes were isolated and allowed to self. After 24 hours, the F1 adults were removed

from the plate, and the F2 hermaphrodites left to develop to young adults. F2 animals

homozygous for the region being introgressed were selected as young adults and crossed

with MY16 males. This process was repeated until ten rounds of introgression were

completed. These new lines were preserved at -80C.

Introgression of MY16 region into an N2 background began with the same initial cross as

above. F1 heterozygous males were crossed with N2 hermaphrodites containing phenotypic

markers near the region being introgressed (Supplementary Fig. 7). After successful mating,

the F1 parents were removed and the F2 generation was left to develop until heterozygous

males were visible. F2 heterozygous males were crossed with hermaphrodites from the N2-

derived strain. This process was repeated until ten successful introgressions were completed.

To homozygose the introgressed MY16 regions, worms were singled and allowed to self until

a stable wildtype population was obtained. These new lines preserved at -80C.

NILs were genotyped to test for correct introgression of the desired regions by Sanger

sequencing of 10 markers spaced along chromosome IV (carried out by the Centre for

Genomics and Proteomics, University of Auckland). Upon confirmation of their genetic

identity, one NIL was used to further dissect the QTL region by segregating the visual markers

(Dpy and Unc).

RESULTS

Extensive natural cryptic variation in the requirement for SKN-1 in endoderm specification

within the *C. elegans* species

The relationship between SKN-1 and Wnt signaling through POP-1 in the endoderm

GRN has undergone substantial divergence in the *Caenorhabditis* genus [39]. While neither

input alone is absolutely required for endoderm specification in *C. elegans*, each is essential

in *C. briggsae*, which has been estimated to have diverged from *C. elegans* ~20-40 Mya

[64,65]. In contrast to the *C. elegans* N2 laboratory strain, removal of either SKN-1 or POP-1

alone results in fully penetrant conversion of the E founder cell fate into that of the

mesectodermal C blastomere and of E to MS fate, respectively, in *C. briggsae* [39]. These

findings revealed that the earliest inputs into the endoderm GRN are subject to substantial

evolutionary differences between these two species (Fig. 1B). We sought to determine

whether incipient evolutionary plasticity in this critical node at the earliest stages of

endoderm development might be evident even within a single species of the *Caenorhabditis*

genus by assessing their requirement in *C. elegans* wild isolates and testing whether the

quantitative requirements of each input were correlated.

Elimination of detectable maternal SKN-1 from the laboratory N2 strain by either a

strong (early nonsense) chromosomal mutation (*skn-1(zu67)*), or by RNAi knockdown, results

in a partially penetrant phenotype: while the E cell adopts the fate of the C cell in the majority

of embryos, and gut is not made, ~30% of arrested embryos undergo strong gut

differentiation, as evidenced by the appearance of birefringent, gut-specific rhabditin

granules, or expression of *elt-2::GFP*, a marker of the developing and differentiated intestine

(Fig. 1C-H). In our experimental conditions, we found that RNAi of *skn-1* in different N2-

derived mutant strains gave highly reproducible results: 100% of the embryos derived from

skn-1(RNAi)-treated mothers arrest (n>100,000) and 32.0 ± 1.9% of the arrested embryos

exhibited birefringent gut granules (Fig. 2A; Supplementary Fig. 1) over many trials by

separate investigators. We found that the LSJ1 laboratory strain, which is derived from the

same original source as N2, but experienced very different selective pressures in the

laboratory owing to its constant propagation in liquid culture over 40 years [66], gave virtually

identical results to that of N2 (31.0% ± s.d 1.2%), implying that SKN-1-independent endoderm

formation is a quantitatively stable trait. The low variability in this assay, and high number of

embryos that can be readily examined (≥500 embryos per experiment), provides a sensitive

and highly reliable system with which to analyze genetic variation in the endoderm GRN

between independent *C. elegans* isolates.

To assess variation in SKN-1 requirement within the *C. elegans* species, we analyzed

the outcome of knocking down SKN-1 by RNAi in 96 unique *C. elegans* wild isolates [42].

Owing to their propagation by self-fertilization, each of the isolates (isotypes) is a naturally

inbred clonal population that is virtually homozygous and defines a unique haplotype. The

reported estimated population nucleotide diversity averages 8.3×10^{-4} per bp [42], and we

found that a substantial fraction (29/97) of isotypes were quantitatively indistinguishable in

phenotype between the N2 and LSJ1 laboratory strains (Fig. 2A, Supplementary File 1). We

found that all strains, with the exception of the RNAi-resistant Hawaiian CB4856 strain,

invariably gave 100% embryonic lethality *with skn-1(RNAi)*. However, we observed dramatic

variation in the fraction of embryos with differentiated gut across the complete set of strains,

ranging from 0.9% to 60% (Fig. 2A). Repeated measurements with >500 embryos per replicate

per strain revealed very high reproducibility (Supplementary Fig. 1), indicating that even small

differences in the fraction of embryos generating endoderm could be reproducibly measured.

Further, we found that some wild isolates that were subsequently found to have identical

genome sequences also gave identical results.

Although birefringent and autofluorescent rhabditin granules have been used as a

marker of gut specification and differentiation in many studies [50,51], it is conceivable that

the variation in fraction of embryos containing this marker that we observed might reflect

variations in gut granule formation rather than in gut differentiation *per se*. We note that

embryos from all strains showed a decisive "all-or-none" phenotype: i.e., they were either

strongly positive for gut differentiation or completely lacked gut granules, with virtually no

intermediate or ambiguous phenotypes. A threshold of gene activity in the GRN has been

shown to account for such an all-or-none switch in gut specification [22,67,68]. This

observation is inconsistent with possible variation in gut granule production: if SKN-1-

depleted embryos were defective in formation of the many granules present in each gut cell,

one might expect to observe gradations in numbers or signal intensity of these granules

between gut cells or across a set of embryos. Nonetheless, we extended our findings by

analyzing the expression of the gut-specific intermediate filament IFB-2, a marker of late gut

differentiation, in selected strains representing the spectrum of phenotypes observed (Fig.

2B). As with gut granules, we found that embryos showed all-or-none expression of IFB-2. In

all cases, we found that the fraction of embryos containing immunoreactive IFB-2 was not

significantly different (Fisher's exact test, p-values > 0.05) from the fraction containing gut

granules, strongly suggesting that the strains vary in endoderm specification *per se* and

consistent with earlier studies of SKN-1 function [19,22].

Although we found that *skn-1(RNAi)* was 100% effective at inducing embryonic

lethality in all strains (with the exception of the RNAi-defective Hawaiian strain, CB4856), it is

conceivable that, at least for the strains that showed a weaker phenotype than for N2 (i.e.,

higher number of embryos specifying endoderm), the variation observed between strains was

attributable to differences in RNAi efficacy rather than in the endoderm GRN. Indeed, studies

with N2 and CB4856 showed that germline RNAi sensitivity is a quantitative trait, involving

the Argonaute-encoding *ppw-1* gene and additional interacting loci present in some wild

isolates [69,70]. To address this possibility, we introgressed the strong loss-of-function *skn-*

1(zu67) chromosomal mutation into five wild isolates whose phenotypes spanned the

spectrum observed (ranging from 2% of embryos with differentiated gut for MY16 to 50% for

MY1) (Fig. 2C). In all cases, we found that introgression of the allele through five rounds of

backcrosses resulted in a quantitative phenotype that was similar or indistinguishable from

that observed with *skn-1(RNAi)*. The phenotypes of the introgressed allele were significantly

different (p-values <0.01) from that of the parental N2 *skn-1(zu67)* strain, except for DL238,

whose *skn-1(RNAi)* phenotype was indistinguishable from that of N2. The results obtained by

introgression from four of the isotypes (CX11262, DL238, EG4724 and MY1), were not

statistically different (Student t-test, p-values >0.05) from the corresponding RNAi

knockdown results (Fig. 2C) (i.e., the phenotype was suppressed or enhanced relative to N2

in these genetic backgrounds to the same extent as with *skn-1(RNAi)*). However, while the

MY16 *skn-1(zu67)* strain shifted in the predicted direction (i.e., became stronger) as

compared to the N2 strain, it showed a weaker phenotype than was evident by RNAi

knockdown, even following eight rounds of introgression. Regardless, diminished RNAi

efficacy in MY16 cannot explain the large difference between the *skn-1(RNAi)* phenotype of

N2 and MY16, as the latter phenotype is, in fact, much stronger, not weaker, than the former.

As described below, we identified a modifier locus in the MY16 strain that is closely linked to

the *skn-1* gene; it therefore seems likely that the N2 chromosomal segment containing this

modifier was carried with the *skn-1(zu67)* mutation through the introgression crosses,

thereby explaining the somewhat weaker phenotype of the introgressed allele in MY16. We

conclude that the extreme variation in *skn-1(RNAi)* phenotype between the wild isolates

tested results from *bona fide* cryptic variation in the endoderm GRN, rather than differences

in RNAi efficacy.

We note that the strength of *skn-1(RNAi)* phenotype does not correlate with

phylogenetic relatedness between the strains (Pagel's λ = 0.42, p-value = 0.14). For example,

while some closely related strains (e.g., MY16 and MY23) showed a similar phenotype, other

very closely related strains (e.g., JU1491 and JU778) showed phenotypes on the opposite ends

of the phenotypic spectrum (Fig. 3A). We also did not observe any clear association between

geographical distribution and *skn-1 (RNAi)* phenotype (Fig. 3B). These findings suggest that

the initiating inputs into the endoderm GRN is subject to rapid intraspecies evolutionary

divergence.

Fig. 2: Quantitative effects of loss of *skn-1* on endoderm formation.

(A) Spectrum of *skn-1(RNAi)* effects across the *C. elegans* isolates. The effects of *skn-1(RNAi)* are quantified as the average percentage of arrested embryos with endoderm (y-axis). All wild isolates treated with *skn-1(RNAi)* resulted in 100% embryonic arrest (n >500 embryos per replicate per isotype and at least two replicates per isotype). (B) Comparison of *skn-1(RNAi)* phenotype using two different gut markers (birefringent gut granules and MH33 staining of IFB-2) in five different genetic backgrounds. In all cases, no significant statistical difference was found between the two quantitative methods. Fisher's exact test (NS p-value>0.05). (C) Comparison of *skn-1(RNAi)* and *skn-1(zu67)* effects on endoderm development in six different genetic backgrounds. For each color-coded strain, the first value is of the *skn-1(RNAi)* results (five replicates), while the second is the result for the *skn-1(zu67)* allele introgression (10 replicates). For all strains (with the exception of MY16), no significant statistical difference was found between the RNAi knockdown and corresponding *skn-1(zu67)* allele effects on endoderm development. Student t-test (NS p-value > 0.05, * p-value < 0.05).

Fig. 3: SKN-1 requirement does not correlate with genotypic relatedness or geographical location.

(A) *skn-1(RNAi)* phenotype of 97 isolates arranged with respect to the neighbor-joining tree constructed using 4,690 SNPs and pseudo-rooted to QX1211. Red asterisk indicates an example of closely related strains (MY23 and MY16) with similar phenotype, while black asterisks indicate example sister strains (JU778 and JU1491; JU561 and JU1652) with distinct phenotype. Phylogenetic relatedness and phenotype are not significantly correlated (Pagel's λ = 0.42, p-value = 0.14). (B) Worldwide distribution of *skn-1(RNAi)* phenotype across 97 wild isolates. Each circle represents a single isotype.

Cryptic variation in the quantitative requirement for MOM-2/Wnt, but not POP-1, in

endoderm development

The switch in the relationship of the SKN-1 and Wnt inputs between *C. elegans* ("OR"

operator) and *C. briggsae* ("AND" operator) [39], and the extensive variation in the

requirement for SKN-1 seen across *C. elegans* isolates, raised the possibility that the

quantitative requirement for Wnt components might vary between unique isolates of *C.*

elegans. It has been shown that signaling from Ras pathway varies in different *C. elegans* wild

isolates and hyperactive Wnt signaling can compensate for reduced Ras activity in the vulva

signaling network [6,71]. Given that removal of the maternal Wnt input also results in a

partially penetrant gut defect (through either knock-out or knockdown of Wnt signaling

components), it is conceivable that a compensatory relationship may exist between the SKN-

1 and Wnt inputs. We investigated this possibility by examining the requirement for the

MOM-2/Wnt ligand in the same wild isolates that were tested for the SKN-1 gut

developmental requirement. Indeed, we observed broad variation in the requirement for

MOM-2/Wnt in activation of the endoderm GRN between isotypes. *mom-2(RNAi)* of 94

isotypes resulted in embryonic arrest, indicating that, as with *skn-1(RNAi), mom-2(RNAi*) was

effective at least by the criterion of lethality. Two isotypes, CB4853 and EG4349, did not

exhibit *mom-2(RNAi)*-induced lethality and were omitted from further analyses. In the

affected strains, the fraction of *mom-2(RNAi)* embryos with differentiated gut varied from

~40% to ~99% (Fig. 4A, Supplementary File 1). As with *skn-1(RNAi)*, the *mom-2(RNAi)*

phenotype of isotypes N2, JU440, and JU1213 was further confirmed by immunostaining with

IFB-2 (Fig 4B), again demonstrating that birefringence of gut granules is a reliable proxy for

endoderm formation for this analysis.

To assess whether the observed variation in the *mom-2(RNAi)* phenotype reflected

differences in the GRN or RNAi efficacy, the *mom-2(or42)* allele was introgressed into three

different genetic backgrounds chosen from the extreme ends of the phenotypic spectrum.

mom-2(RNAi) of the laboratory N2 strain resulted in the developmental arrest of embryos. Of

those, ~90% contained differentiated endoderm, a result that was highly reproducible. In

contrast, the introgression of an apparent loss-of-function allele, *mom-2(or42)*, into the N2

strain results in a more extreme phenotype: only ~28% of embryos show endoderm

differentiation (Fig. 4C) [30]. While this discrepancy can partly be explained by incomplete

RNAi efficacy, it is notable that the penetrance of *mom-2* alleles vary widely [30]. We

observed strain-specific variation in embryonic lethality response to RNAi of *mom-2* between

the different isotypes. However, we found that the *mom-2(or42)* introgressed strains show

qualitatively similar effects to those observed with *mom-2* RNAi. For example, the *mom-*

2(or42) allele introgressed into the isotype JU1213 background resulted in a low fraction of

arrested embryos with gut (5.7% ± s.d 2.4%; n=2292), a more extreme effect than was seen

with RNAi (34.0% ± s.d 1.5%; n=1876). This is the strongest phenotype that has been reported

for any known *mom-2* allele. On the other hand, introgression of the *mom-2* mutation gave

rise to a significantly higher fraction of embryos with endoderm in isotypes DL226 (55.2% ±

s.d 1.2%, n=1377) and PB303 (65.5% ± s.d 4.9%, n=2726), relative to the laboratory strain N2

(29.1% ± s.d 3.1%; n=1693), consistent with the RNAi phenotypes (Fig. 4C). These findings

indicate that the differential requirement for MOM-2 is, at least in part, attributable to

genetic modifiers in these strains. As with *skn-1(RNAi),* we found no correlation between the

mom-2(RNAi) phenotype and phylogenetic relatedness or geographical distribution

(Supplementary Fig. 5), suggesting rapid intraspecies developmental system drift.

As the MOM-2/Wnt signal is mediated through the POP-1 transcription factor, we

sought to determine whether the requirement for POP-1 might also vary between isolates.

We found that, while *pop-1(RNAi)* resulted in 100% embryonic lethality across all 96 RNAi-

sensitive isolates, 100% of the arrested embryos contained a differentiated gut (n>500 for

each isolate scored) (results not shown). Thus, all isolates behave similarly to the N2 strain

with respect to the requirement for POP-1. These results were confirmed by introgressing a

strong loss-of-function *pop-1(zu189)* allele into four wild isolates (N2, MY16, JU440, and

KR314) (Supplementary Fig. 2). The lack of variation in endoderm specification after loss of

POP-1 is not entirely unexpected. As has been observed in a *pop-1(-)* mutant strain,

elimination of the endoderm-repressive role of POP-1 in the MS lineage (which is not

influenced by the P_2 signal) supersedes its endoderm activating role in the presence of SKN-

1. Indeed, the original observation that all *pop-1(-)* embryos in an N2 background contain gut

masked the activating function for POP-1, which is apparently only in the absence of SKN-1

[33,35,37]. It is likely that, as with the N2 strain, gut arises from both E and MS cells in all of

these strains; however, as we have scored only for presence or absence of gut, it is

conceivable that the E lineage is not properly specified in some strains, a possibility that

cannot be ruled out without higher resolution analysis.

Our results contrast with those of Paaby *et al.* [72], who reported that RNAi of 29

maternal-effect genes across a set of 55 wild isolated in liquid culture resulted in generally

weaker effects on lethality than we observed. This difference is likely attributable to

diminished and variable RNAi efficacy in the latter study owing to the different culture

methods used (see MATERIALS AND METHODS) [48,73]. To assess this possibility further, we

compared our results with those of Paaby *et al.* (2015) and found no correlation between the

variation in fraction of embryos with gut and the lethality observed in that report with both

mom-2(RNAi) and *skn-1(RNAi)* (Pearson's R = 0.19, p = 0.23; Pearson's R = 0.22, p = 0.17,

respectively). In addition, Paaby *et al.* (2015) found that the genetically divergent strain

QX1211 consistently showed weak penetrance across all targeted genes, while under our

experimental conditions, QX1211 exhibited a slightly stronger skn-1(RNAi)phenotype (25.2%

vs. 32.0%, Fisher's exact test p-value = 0.03) and a similar mom-2(RNAi) phenotype (90% *vs*.

90%, Fisher's exact test p-value = 0.9) compared to the N2 strains with fully penetrant

lethality in all cases.

Fig. 4: Widespread variation in the *mom-2(RNAi)* phenotype.

(A) Spectrum of *mom-2(RNAi)* effects across the *C. elegans* isolates. The effects of *mom-2(RNAi)* are quantified as the average percentage of arrested embryos with endoderm (y-axis). Each column represents the mean for each wild isolate (n >500 embryos were scored for each experiment with at least two replicates per isotype). (B) Comparison of *mom-2(RNAi)* phenotype using two different gut markers (birefringent gut granules and MH33 immunostaining of IFB-2) in three different genetic backgrounds. In all cases, no significant statistical difference was found between the two quantitative methods. Fisher's exact test (NS p-value > 0.05). (C) Comparison of the effect of *mom-2(or42)* on endoderm development after introgression into four different genetic backgrounds. At least three independent introgressed lines were studied for each wild isotype. The results were compared *to N2; mom-2(or42)*. Student t-test (*** p-value < 0.001).

Genome-wide association studies (GWAS) and analysis of RILs identify multiple genomic

regions underlying variation in the two major endoderm GRN inputs

We sought to examine the genetic basis for the wide variation in SKN-1 and Wnt

requirements across *C. elegans* isolates and to evaluate possible relationships in the variation

seen with the SKN-1 and Wnt inputs by performing GWAS using the available SNP markers

and map [42], adjusting for population structure by using Efficient Mixed-Model Analysis

(EMMA) (Fig. 5A, B) [52,74]. This approach identified two significant closely-located positions

on chromosome IV that underlie the variation in SKN-1 requirement (Fig. 5A, Supplementary

Table 1).

GWAS of the *mom-2(RNAi)* variation proved more challenging because this phenotype

showed a highly skewed distribution (Shapiro-Wilk' test W =0.8682, p-value = $1.207\mathrm{X}10^{-7}$)

(Supplementary Fig. 3). While GWAS did not reveal any genomic regions for the *mom-2(RNAi)*

variation that exceeded an FDR of 5%, we found that the most strongly associated loci for the

mom-2(RNAi) phenotype also showed large effects for *skn-1(RNAi)* (Fig. 5C). In particular, we

observed substantial overlap in the p-values for individual SNPs from *skn-1(RNAi)* and *mom-*

2(RNAi) in the central region of chromosome IV (Supplementary Fig. 4), raising the possibility

that common genetic factors might underlie these phenotypes.

Fig. 5. Genome-Wide Association Studies of *skn-1(RNAi)* and *mom-2(RNAi)* phenotypes.

(A) Manhattan plot of *skn-1(RNAi)* GWAS. The red line indicates a genome-wide 1.5% FDR (permutation-based FDR, from 10,000 permutated results). Black line represents 3.0% FDR. The y axis is the −$\log_{10}$ of p-value. (B) Manhattan plot of *mom-2 (RNAi)* EMMA. The y axis is the −log10 of p-value. Genomic regions are shown on the x-axis. (C) Effect plots of the most strongly-linked SNPs from *mom-2(RNAi)* GWAS at position 3,362,389 bp on chromosome I and position 17,924,783 bp on chromosome V. Horizontal lines within each box represent the median, and the boxes represent 25th–75th percentile.

In an effort to narrow in on causal loci underlying the *skn-1(-)* and *mom-2(-)* phenotypic variation, and to assess possible relationships between these two GRN inputs, we prepared and analyzed 95 recombinant inbred lines (RILs) between two *C. elegans* isotypes, N2 and MY16. These strains were chosen for their widely varying differences in requirement for both inputs (see MATERIALS AND METHODS). In contrast to the very low variation seen between multiple trials of each parental strain, analysis of the RNAi-treated RIL strains (>500 embryos/RIL) revealed a very broad distribution of phenotypes. We found that, while some RILs showed phenotypes similar to that of the two parents, many showed intermediate phenotypes and some were reproducibly more extreme than either parent, indicative of transgressive segregation [75]. For *skn-1(RNAi)*, the phenotype varied widely across the RILs,

with 1 to 47% of embryos containing gut (Fig. 6A, Supplementary File 2). This effect was even

stronger with *mom-2(RNAi)*, for which virtually the entire possible phenotypic spectrum was

observed across a selection of 31 RILs representing the span of *skn-1(RNAi)* phenotypes. The

mom-2(RNAi) phenotypes ranged from RILs showing 3% of embryos with gut to those

showing 92% (Fig. 6A). In all RILs, *skn-1(RNAi)* and *mom-2(RNAi)* resulted in 100% lethality. It

is noteworthy that one RIL (JR3572, Supplementary File 2) showed a nearly completely

penetrant gutless phenotype, an effect that is much stronger than has been previously

observed for *mom-2(-)* [30]*.* These results indicate that a combination of natural variants can

nearly eliminate a requirement for MOM-2 altogether, while others make it virtually essential

for endoderm development. Collectively, these analyses reveal that multiple quantitative

trait loci (QTL) underlie SKN-1- and MOM-2-dependent endoderm specification.

To identify QTLs from the recombinant population, we performed linkage mapping for

both phenotypes using interval mapping (see MATERIALS AND METHODS). For *skn-1(RNAi)*,

two major peaks were revealed on chromosomes II and IV (above 1% FDR estimated from

1,000 permutations). Two minor loci were found on chromosomes I and X (suggestive linkage,

above 20% FDR) (Fig 6B). For *mom-2(RNAi)*, two major independent QTL peaks were found

on chromosomes I and II (above the 5% FDR estimated from 1,000 permutations). Although

the candidate peaks observed on chromosome IV for *skn-1(RNAi)* (Fig. 6B) did not appear to

overlap with those for *mom-2(RNAi)*, overlap was observed between the chromosomes I and

II candidate regions for these two phenotypes. These QTLs show large individual effects on

both phenotypes (Fig. 6C),

Fig 6. Quantitative genetic analysis of *mom-2(RNAi)* and *skn-1(RNAi)* phenotype in Recombinant Inbred Lines (RILs) between N2 and MY16.

(A) *mom-2(RNAi)* (left) and *skn-1(RNAi)* (right) phenotype of RILs. The phenotype of the parental strains, MY16 and N2 are shown by red and blue lines, respectively. (B) QTL analyses (interval mapping) of *skn-1(RNAi)* (blue line) and *mom-2(RNAi)* (red line) phenotype shown in (A). Genomic regions are shown on the x-axis and LOD score is shown on the y-axis. Significance thresholds for *mom-2(RNAi)* and *skn-1(RNAi)* at 5% and 20% linkage represented in red and blue dashed lines, respectively. (C) Effect plots of significant SNPs from *mom-2(RNAi)* and *skn-1(RNAi)*, indicated by chromosome number and color, showing the direction of the allelic effects. Confidence intervals for the average phenotype in each genotype group are shown.

Potential cryptic relationships between SKN-1 and MOM-2 inputs

The preceding findings unveiled wide cryptic variation in the requirements for both

SKN-1 and MOM-2/Wnt in the endoderm GRN, raising the possibility that the variation

affecting the two inputs might be related. Indeed, comparisons of the GWAS and QTL

mapping results for *skn-1* and *mom-2* showed an overlap in candidate QTL regions on

chromosome I, II and IV (Fig. 5, Fig 6, Supplementary Fig. 4), suggesting a possible connection

between the genetic basis underlying these two traits. It is conceivable that some genetic

backgrounds are generally more sensitive to loss of either input (e.g., the threshold for

activating the GRN is higher) and others more robust to single-input loss. Alternatively, a

higher requirement for one input might be associated with a relaxed requirement for the

other, i.e., a reciprocal relationship.

As an initial assessment of these alternatives, we examined whether the requirements

for SKN-1 and MOM-2 across the strains were significantly correlated. This analysis revealed

no strong relationship between the cryptic variation in the requirement for these inputs seen

across all the strains (Spearman correlation R=0.18, p-value=0.07) (Fig. 7A). This apparent lack

of correlation at the level of strains is not unexpected, as many factors likely contribute to the

cryptic variation and the comparison reflects the collective effect of all causal loci in the

genome of each strain (Fig. 5, 6).

We next sought to examine possible relationships between the two GRN inputs at

higher resolution by comparing association of specific genetic regions with the quantitative

requirement for each input. We used the available sequencing data for all isotypes tested [42]

and examined the impact of each allele on the *skn-1(RNAi)* and *mom-2(RNAi)* phenotypes,

correcting for outliers and using a pruned SNP map (see MATERIALS AND METHODS). We

found a weak positive correlation (Pearson's R = 0.21, p= p-value < 2.2e-16, Supplementary

Fig. 6) between the allelic effects. One possible explanation for this observation might be that

variants across the set of wild isolates may generally influence the threshold for activating the

positive feedback loops that lock down gut development [49,67], thereby altering the

sensitivity for regulatory inputs into the endoderm pathway. Alternatively, although evidence

for variation in germline RNAi sensitivity among *C. elegans* wild isolates is lacking (except for

CB4856, which has been omitted from our study) [69,76], and we have shown above that

variation in SKN-1 and MOM-2 requirement reflects in large part cryptic genetic differences

in the endoderm GRN, it remains possible that a fraction of the variation found in the two

phenotypes tested is attributable to varying RNAi penetrance, which may underlie the minor

positive correlation between *skn-1(RNAi)* and *mom-2(RNAi)* phenotypes.

In contrast, analysis of the N2/MY16 RILs uncovered a potential reciprocal relationship

between the requirements for SKN-1 and MOM-2: we observed a negative correlation

between the *skn-1(RNAi)* and *mom-2(RNAi)* phenotypes across the genome (Fig 7B, genome-

wide Pearson's R = -0.35, p = 0.001, correcting for LD and outliers as above; correlation

without chromosome IV R = -0.59, p < 0.001). This finding suggested that at least some

quantitative variants result in opposing effects on the requirement for SKN-1 and MOM-2.

While a reciprocal relationship was observed generally across the genome spanning

five of the chromosomes, we observed the opposite correlation on chromosome IV (Pearson's

R = 0.83, p-value = 1.695×10^{-5}). No correlation was observed for chromosome IV with the wild

isolates (Pearson's R = 0.08, NS, Supplementary Fig. 6). As there is a major QTL on

chromosome IV for the SKN-1 requirement and there is substantial overlap in the same region

with the GWAS analysis of the *skn-1(RNAi)* and *mom-2(RNAi)* phenotypes, we sought to

dissect further the relationship between the requirement for MOM-2 and SKN-1 in this

region. We created six near-isogenic lines (NILs) in which the QTL region for the *skn-1(RNAi)*

phenotype on chromosome IV from N2 was introgressed into the MY16 background, and *vice-*

versa (Supplementary Fig. 7). Control lines were created from the same crosses at the same

generation by selecting the original parental region (e.g., selecting for the N2 region in an N2

background and MY16 in MY16 background). We found that the region affects the *skn-*

1(RNAi) phenotype as expected: the N2 region increased the fraction of gut in an MY16

background, and the MY16 regions decreased this fraction in an N2 background. However,

for *mom-2(RNAi),* while introgressing the N2 region in MY16 dramatically changed the

phenotype (Fig. 7C), we found that the MY16 region was not sufficient to alter the phenotype

in an N2 background. We created segregant NILs in which one of the genetic markers was lost

(see MATERIALS AND METHODS) and found that replacing the N2 region with the

corresponding MY16 region in all cases results in a stronger *mom-2(RNAi)* phenotype.

However, for the *skn-1(RNAi)* phenotype six of nine segregants showed the opposite effect:

i.e., a weaker phenotype (Fig. 7D), revealing that when contributing variants were separated

by recombination, a reciprocal effect was frequently seen. These observations suggest that

complex genetic interactions between variants on chromosome IV might mask the potential

reciprocal effects that were observed on the other chromosomes.

Fig. 7: Correlation of *skn-1(RNAi)* and *mom-2(RNAi)* allelic differences.

(A) Comparison of *skn-1(RNAi)* and *mom-2(RNAi)* phenotype in 94 natural isolates tested. No correlation was found (Spearman correlation R=0.1844, p-value=0.07). Each dot corresponds to a wild isolate. Y-axis, *skn-1(RNAi)* phenotype, x-axis, *mom-2(RNAi)* phenotype. (B) Genome-wide correlation of *skn-1(RNAi)* and *mom-2(RNAi)* allelic differences in the N2xMY16 RILs. Each dot represents a SNP. Chromosomes are color-coded with their Pearson's R values represented (NS = Not Significant, *** = p-value < 0.001). Regression line in grey. (C) Six different NILs were created for chromosome IV, each of which was compared with a control NIL from the same cross (e.g., MY16 in MY16 as control for N2 in MY16). A schematic of the introgressed regions is represented below the plots. Percentage of *skn-1(RNAi)* or *mom-2(RNAi)* embryos with gut is represented. (D) Changes in phenotype for both *skn-1(RNAi)* and *mom-2(RNAi)* following recombination in segregant NILs, which a schematic representation of the segregant NILs below the plot.

Multiple factors reciprocally regulate the requirement for SKN-1 and MOM-2/Wnt

While the above findings revealed that the relationship between the requirement for

SKN-1 and MOM-2 may be complicated by genetic interactions, our results raised the

possibility of compensatory relationships between them. To further assess this possibility, we

tested other candidate genes that reside in the QTL regions and that have been implicated in

endoderm development [77–79]. We found that loss of RICT-1, the *C. elegans* orthologue of

the human RICTOR (Rapamycin-insensitive companion of mTOR; [80]), a component of the

TORC2 complex, which has been shown to antagonize SKN-1 function [77], results in opposite

effects on *skn-1(-)* and *mom-2(-)* mutants (Fig. 8A). Specifically, while *rict-1(RNAi)* suppresses

the absence of gut in *skn-1(zu67)* embryos (*skn-1(zu67)*: 34.3% ± s.d 4.1% with gut *vs. skn-*

1(zu67); *rict-1(RNAi)*: 48.3% ± s.d 4.9%; p=<0.001), we found that it *enhances* this phenotype

in *mom-2(or42)* mutants (*mom-2(or42)*: 23.8% ± s.d 2.0%; vs. *mom-2(or42)*; *rict-1(RNAi)*:

11.2% ± s.d 3.2%; p<0.001). Confirming this effect, a similar outcome was observed when

SKN-1 was depleted by RNAi in *rict-1(ft7)* chromosomal mutants (*skn-1(RNAi)*: 31.6% ± s.d

4.3% with gut vs. *rict-1(ft7); skn-1(RNAi)*: 45.9% ± s.d 6.3%; p<0.05) (Fig. 8A). Similarly, RNAi

depletion of PLP-1, the *C. elegans* homologue of the Pur alpha transcription factor that has

been shown to bind to and regulate the *end-1* promoter [78], reciprocally affects the outcome

of removing these two inputs in the same direction: loss of PLP-1 function suppresses the *skn-*

1(-) phenotype (to 48.0% ± s.d 6.6%), and strongly enhances the *mom-2* phenotype (to 6.9%

± s.d 1.6%). Again, this result was confirmed by RNAi of *skn-1* in a *plp-1(ok2156)* chromosomal

mutant (Fig. 8B). Thus, as observed with the effect across the genome with natural variants,

we observed a substantial reciprocal effect of both of these genes on loss of SKN-1 and MOM-

2.

We also observed a reciprocal effect on the SKN-1 and Wnt inputs with MIG-

5/*dishevelled*, a component of the Wnt pathway that acts downstream of the Wnt receptor

[79]; however, in this case the effect was in the opposite direction as seen for RICT-1 and PLP-

1. Loss of MIG-5 as a result of chromosomal mutation or RNAi leads to *enhancement* of the

skn-1(-) phenotype (*mig-5(rh94); skn-1(RNAi)*: 6.6% ± s.d 2.3%; *skn-1(zu67); mig-5(RNAi)*:

9.4% ± s.d 1.4%) and *suppression* of the *mom-2(-)* phenotype (88.6% ± s.d 4.0%) (Fig 8C).

Together, these findings reveal that, as observed with many of the N2/MY16 RILs

variants across most of the genome, RICT-1, PLP-1, and MIG-5 show opposite effects on the

phenotype of removing SKN-1 and MOM-2, suggesting a trend toward genetic influences that

reciprocally influence the outcome in the absence of these two inputs.

677
Fig 8. Reciprocal effects of RICT-1, PLP-1, and MIG-5 on *skn-1(-)* and *mom-2(-)* phenotypes

(A, B) Loss of RICT-1 or PLP-1 enhances the *mom-2(or42)* loss-of-endoderm phenotype and suppresses
skn-1(zu67) and *skn-1(RNAi)* phenotype. (C) Loss of MIG-5 enhances the *skn-1(zu67)* and *skn-1(RNAi)*
phenotype and suppresses *mom-2(or42)* phenotype. At least three replicates were performed per

experiment and >200 embryos per experiment. Student t-test (*** p-value<0.001). Data represented
with Standard Deviations.

DISCUSSION

The remarkable variety of forms associated with the ~36 animal phyla [81] that emerged from a common metazoan ancestor >600 Mya is the product of numerous incremental changes in GRNs underlying the formation of the body plan and cell types [1,82]. Here, we describe an unexpectedly broad divergence in the deployment of SKN-1/Nrf and MOM-2/Wnt signaling in generating the most ancient germ layer, the endoderm, within wild isolates of a single animal species, *C. elegans*. In this study, we report five major findings: 1) while the quantitative requirement for two distinct regulatory inputs that initiate expression of the endoderm GRN (SKN-1 and MOM-2) are highly reproducible in individual *C. elegans* isolates, there is wide cryptic variation between isolates. 2) Cryptic variation in the requirement for these regulatory factors shows substantial differences even between closely related strains, suggesting that these traits are subject to rapid evolutionary change in this species. 3) Quantitative genetic analyses of natural and recombinant populations revealed multiple loci underlying the variation in the requirement for SKN-1 and MOM-2 in endoderm specification. 4) The requirements for SKN-1 and MOM-2 in endoderm specification is frequently reciprocal in their relation to other genetic factors. 5) *rict-1*, *plp-1*, *and mig-5* reciprocally influence the outcome of *skn-1(-)* and *mom-2(-)*, substantiating the reciprocal influences on the two GRN inputs. These findings reveal prevalent plasticity and complexity underlying SKN-1 and MOM-2/Wnt regulatory inputs in mobilizing a conserved system for endoderm specification. Thus, while the core genetic toolkit for the development of the endoderm, the most ancient of the three germ layers, appears to have been preserved for well over half a billion years, the molecular regulatory inputs that initiate its expression in *C. elegans* vary extremely rapidly over short evolutionary time scales within the species.

Multigenic variation in the requirement for SKN-1 and MOM-2

Quantitative analyses of the wild isolates and RILs revealed that multigenic factors are responsible for the difference in requirement for SKN-1 and MOM-2 between isotypes. Notably, we observed substantial overlap on chromosome IV in the GWAS analyses of the *skn-1* and *mom-2* requirements in wild isotypes (Fig. 5, Supplementary Fig. 4) and on chromosome II from analyses using RILs (Fig. 6B). This finding raises the possibility that some QTLs may influence requirement for both inputs into the endoderm specification pathway: as SKN-1 and Wnt converge to regulate expression of the *end-1/3* genes, it is conceivable that common genetic variants might modulate the relative strength or outcome of both maternal inputs. However, our findings do not resolve whether these genetic variants act independently to influence the maternal regulatory inputs. Genetic interactions are often neglected in large-scale genetic association studies [83] owing in part to the difficulty in confirming them [83]. Many studies [84–87], including ours here, showed that epistasis can strongly influence the behavior of certain variants upon genetic perturbation. In addition, selection on pleiotropically acting loci facilitates rapid developmental system drift. Together, epistasis and selection on pleiotropic loci play important roles in the evolution of natural populations [88–91].

Potential compensatory relationships between SKN-1 and MOM-2/Wnt

Although we did not observe a direct correlation between the *skn-1(-)* and *mom-2(-)* phenotypes across the isotypes studied here, we found a negative correlation across much of the genome for the N2 X MY16 RILs (Fig. 6, 7). Further, while GWAS and QTL analysis of natural and inbred lines, respectively, did not reveal a causal region in chromosome IV for *mom-*

2(RNAi) variation, analysis of NILs results did uncover at least one QTL affecting this

phenotype. Moreover, while broad regions of the chromosome showed a positive correlation

between the SKN-1 and MOM-2 requirements, isolation of variants in NILs revealed an inverse

requirement for these inputs for at least some regions on this chromosome. These results

reflect the limitations of genome-wide studies of complex genetic traits: in the case of

chromosome IV, several closely linked loci appear to influence both the SKN-1 and MOM-2

requirements.

Our findings raise the possibility that the SKN-1 and MOM-2/Wnt inputs might

compensate for each other and that genetic variants that enhance the requirement for one

of the inputs may often relax the requirement for the other. Such reciprocality could reflect

cross-regulatory interactions between these two maternal inputs or could be the result of

evolutionary constraints imposed by selection on these genes, which act pleiotropically in a

variety of processes. Further supporting this possibility, we identified two genes, *rict-1* and

plp-1, that show similar inverse effects on the requirements from *skn-1* and *mom*-2:

debilitation of either gene enhances the phenotype of *mom-2(-)* and suppresses that of *skn-*

1(-). RICT-1 function extends lifespan in *C. elegans* through the action of SKN-1[77], and loss

of RICT-1 rescues the misspecification of the MS and E blastomeres and lethality of *skn-1(-)*

embryos [77], consistent with our finding. We previously reported that PLP-1, a homologue

of the vertebrate transcription factor pur alpha, binds to the *end-1* promoter and acts in

parallel to the Wnt pathway and downstream of the MAPK signal [78], thereby promoting gut

formation. PLP-1 shows a similar reciprocal relationship with SKN-1 and MOM-2 as with RICT-

1 (Fig. 8). Given that PLP-1 binds at a *cis* regulatory site in *end-1* near a putative POP-1 binding

site [78], and that SKN-1 also binds to the *end-1* regulatory region [68], it is conceivable that

this reciprocality might reflect integration of information at the level of transcription factor

binding sites. As the architecture of the GRN is shaped by changes in *cis*-regulatory sequences

[1,3], analyzing alterations in SKN-1 and Wnt/POP-1 targets among *C. elegans* wild isolates

may provide insights into how genetic changes are accommodated without compromising the

developmental output at microevolutionary time scale.

MIG-5, a *dishevelled* orthologue, functions in the Wnt pathway in parallel to Src

signaling to regulate asymmetric cell division and endoderm induction [29,79]. We found that

the loss of *mig-5* function enhances the gut defect of *skn-1(-)* and suppresses that of the *mom-*

2(-), the opposite reciprocal relationship to that of *rict-1* and *plp-1,* and consistent with a

previous report (Fig. 8) [29]. These effects were not observed in embryos lacking function of

dsh-2, the orthologue of *mig-5* (data not shown), supporting a previous study that showed

overlapping but non-redundant roles of MIG-5 and DSH-2 in EMS spindle orientation and gut

specification [79]. Recent studies showed that Dishevelled can play both positive and negative

roles during axon guidance [92,93]. Dishevelled, upon Wnt-activation, promotes

hyperphosphorylation and inactivation of Frizzled receptor to fine-tune Wnt activity. It is

tempting to speculate that MIG-5 may perform similar function in EMS by downregulating

activating signals (Src or MAPK), in the absence of MOM-2.

We hypothesize that compensatory mechanisms might evolve to fine-tune the level

of gut-activating regulatory inputs. Successful developmental events depend on tight spatial

and temporal regulation of gene expression. For example, anterior-posterior patterning in the

Drosophila embryo is determined by the local concentrations of the Bicoid, Hunchback, and

Caudal transcription factors [94]. We postulate that SKN-1 and Wnt signaling is modulated so

that the downstream genes, *end-1/3*, which control specification and later differentiation of

endoderm progenitors, are expressed at optimal levels that ensure normal gut development.

Suboptimal END activity leads to poorly differentiated gut and both hypo- and hyperplasia in

the gut lineage [95–97]. Hyper- or hypo-activation of Wnt signaling has been implicated in

cancer development [98], bone diseases [99,100], and metabolic diseases [101,102],

demonstrating the importance of regulating the timing and dynamics of such developmental

signals within a quantitatively restricted window.

Cryptic variation and evolvability of GRNs

This study revealed substantial cryptic genetic modifications that alter the relative

importance of two partially redundant inputs into the *C. elegans* endoderm GRN, leading to

rapid change in the developmental network architecture (Fig. 9). Such modifications may

occur through transitional states that are apparent even within this single species. For

example, the finding that POP-1 is not required for gut development even in a wild isolate

(e.g., MY16) that, like *C. briggsae*, shows a near-absolute requirement for SKN-1 may reflect

a transitional state between the two species: *i.e.*, a nearly essential requirement for SKN-1

but non-essential requirement for POP-1, an effect not previously seen in either species. In

addition, duplicated GATA factors (the MEDs, ENDs, and ELTs) and partially redundant

activating inputs (SKN-1, Wnt, Src, and MAPK) in endoderm GRN, provide an opportunity for

genetic variation to accumulate and "experimentation" of new regulatory relationships

without diminishing fitness [2,102,103].

Redundancy in the regulatory inputs may act to 'rescue' an initial mutation and allow

for secondary mutations that might eventually lead to rewiring of the network. For example,

loss of either MyoD or Myf5, two key regulators of muscle differentiation in metazoans,

produces minimal defects in myogenesis as a result of compensatory relationship between

the myogenic factors [104]. In vertebrates, gene duplication events have resulted in an

expansion of Hox genes to a total of >200, resulting in prevalent redundancy [105–107]. This

proliferation of redundant genes provides opportunities for evolutionary experimentation,

subsequent specialization of new functions, and developmental system drift [107,108]. In *C.*

elegans, loss of GAP-1 (a Ras inhibitor) or SLI-1 (a negative regulator of EGFR signaling) alone

does not produce obvious defects, while double mutations lead to a multivulva phenotype

[109]. Similar redundant relationships between redundant partners exist in many other

contexts in the animal. Notably, the relative importance of Ras, Notch, and Wnt signals in

vulva induction differ in various genetic backgrounds [6,71] and physiological conditions

[110,111], resulting in flexibility in the system. While vulval development in *C. elegans,* when

grown under standard laboratory conditions, predominantly favors utilization of the EGF/Ras

signaling pathway [110], Wnt is the predominant signaling pathway in the related *Pristionchus*

pacificus, which is ~250 MY divergent [112,113]. In addition, while *Cel-lin-17* functions

positively to transduce the Wnt signal, *Ppa-lin-17/Fz* antagonizes Wnt signaling and instead

the Wnt signal is transmitted by *Ppa-lin-18/Ryk*, which has acquired a novel SH3 domain not

present in the *C. elegans* ortholog [114]. Thus, extensive rewiring of signaling networks and

modularity of signaling motifs contribute to developmental systems drift [108,115].

The broad cryptic variation may drive developmental system drift, giving rise to GRN

architectures that differ in the relative strength of the network components. Our finding that

the key regulatory inputs that initiate the endoderm GRN show dramatic plasticity is

consistent with comparative transcriptomic studies that demonstrate high gene expression

variability and divergence during early embryonic stages in fly [116–118], worm [117–120],

Xenopus [118,121,122], zebrafish [118,122], and mouse [118,122]. Therefore, early

developmental events may be highly evolvable, in part due to weak purifying selection on

maternal-effect genes [123,124]. This is in accordance with the "hourglass" concept of

embryonic development [125–127], in which divergent developmental mechanisms during

early embryogenesis converge on a more constant state (i.e., a "phylotypic stage" at the

molecular regulatory level). Indeed, unlike the terminal differentiation factor ELT-2, upstream

MEDs and ENDs genes are present only in closely related Caenorhabditis species

[15,97,128,129]. This is likely attributable either to positive selection during early embryonic

and later larval stages or to developmental constraints. Analysis of developmental gene

expression in mutation accumulation lines, which have evolved in the absence of any positive

selection, showed similarity to the developmental hourglass model of evolvability, consistent

with strong developmental constraints on the phylotypic stage [120]. However, they do not

rule out the possibility that early and late stages of development might be more adaptive and

therefore subject to positive selection. It will be of interest to learn the degree to which the

divergence in network architecture might arise as a result of differences in the environment

and selective pressures on different *C. elegans* isotypes.

Fig. 9. Simplified models for potential cryptic compensatory relationship between the SKN-1 and MOM-2/Wnt regulatory inputs in the endoderm GRN.

Accumulation of cryptic genetic modifications drives rapid rewiring of the GRN, causing broad variation of SKN-1 and MOM-2/Wnt dependence in endoderm (E) specification among *C. elegans* isotypes. Wnt-signaled POP-1 (indicated by *) acts as an E activator, while unmodified POP-1 in the MS blastomere acts as a repressor of E fate in all *C. elegans* isotypes. The relative strength of the inputs is indicated by the thickness of the arrow. RICT-1, PLP-1 and MIG-5 reciprocally influence the outcome in the absence of the two inputs

ACKNOWLEDGEMENTS

We thank members of the Rothman lab, especially Sagen Flowers and Kristoffer C. Mellingen

for experimental assistance, and Snell labs, particularly Dr. Kien Ly, for helpful advice and

feedback. We thank Dr. Kathy Ruggiero (University of Auckland, New Zealand) for helpful

advice on GWAS methodology and Dr. James McGhee (University of Calgary, Canada) for

providing the MH33 antibody. Nematode strains used in this work were provided by the

Caenorhabditis Genetics Center, which is funded by the National Institutes of Health - Office

of Research Infrastructure Programs (P40 OD010440). Y.N.T.C was supported during part of

this work by a University of Auckland Doctoral Scholarship. This work was supported by grants

from the NIH (#1R01HD082347 and # 1R01HD081266) to J.H.R.

COMPETING INTERESTS

The authors declare no competing or financial interests.

REFERENCES

1. Peter IS, Davidson EH. Evolution of gene regulatory networks controlling body plan development. Cell. Elsevier; 2011;144: 970–85. doi:10.1016/j.cell.2011.02.017

2. Félix M-A, Wagner A. Robustness and evolution: concepts, insights and challenges from a developmental model system. Heredity (Edinb). Nature Publishing Group; 2008;100: 132–140. doi:10.1038/sj.hdy.6800915

3. Davidson EH, Levine MS. Properties of developmental gene regulatory networks. Proc Natl Acad Sci U S A. National Academy of Sciences; 2008;105: 20063–6. doi:10.1073/pnas.0806007105

4. Oliveri P, Tu Q, Davidson EH. Global regulatory logic for specification of an embryonic cell lineage. Proc Natl Acad Sci U S A. National Academy of Sciences; 2008;105: 5955–62. doi:10.1073/pnas.0711220105

5. Peter IS, Davidson EH. Assessing regulatory information in developmental gene regulatory networks. Proc Natl Acad Sci U S A. National Academy of Sciences; 2017;114: 5862–5869. doi:10.1073/pnas.1610616114

6. Milloz J, Duveau F, Nuez I, Felix M-A. Intraspecific evolution of the intercellular signaling network underlying a robust developmental system. Genes Dev. 2008;22: 3064–3075. doi:10.1101/gad.495308

7. Nunes MDS, Arif S, Schlötterer C, McGregor AP. A perspective on micro-evo-devo: progress and potential. Genetics. Genetics; 2013;195: 625–34. doi:10.1534/genetics.113.156463

8. Phinchongsakuldit J, MacArthur S, Brookfield JFY. Evolution of Developmental Genes: Molecular Microevolution of Enhancer Sequences at the Ubx Locus in Drosophila and Its Impact on Developmental Phenotypes. Mol Biol Evol. 2003;21: 348–363. doi:10.1093/molbev/msh025

9. Hashimshony T, Feder M, Levin M, Hall BK, Yanai I. Spatiotemporal transcriptomics reveals the evolutionary history of the endoderm germ layer. Nature. Nature Publishing Group; 2015;519: 219–222. doi:10.1038/nature13996

10. Rodaway A, Patient R. Mesendoderm: An Ancient Germ Layer? Cell. Cell Press; 2001;105: 169–172. doi:10.1016/S0092-8674(01)00307-5

11. Peterson KJ, Lyons JB, Nowak KS, Takacs CM, Wargo MJ, McPeek MA. Estimating metazoan divergence times with a molecular clock. Proc Natl Acad Sci U S A. National Academy of Sciences; 2004;101: 6536–41. doi:10.1073/pnas.0401670101

12. Martindale MQ, Pang K, Finnerty JR. Investigating the origins of triploblasty: "mesodermal" gene expression in a diploblastic animal, the sea anemone Nematostella vectensis (phylum, Cnidaria; class, Anthozoa). Development. The Company of Biologists Ltd; 2004;131: 2463–74. doi:10.1242/dev.01119

13. Boyle MJ, Seaver EC. Developmental expression of foxA and gata genes during gut formation in the polychaete annelid, Capitella sp. I. Evol Dev. 2008;10: 89–105. doi:10.1111/j.1525-142X.2007.00216.x

14. Boyle MJ, Seaver EC. Expression of FoxA and GATA transcription factors correlates with regionalized gut development in two lophotrochozoan marine worms: Chaetopterus (Annelida) and Themiste lageniformis (Sipuncula). Evodevo. 2010;1: 2. doi:10.1186/2041-

9139-1-2

15. Gillis WJ, Bowerman B, Schneider SQ. Ectoderm- and endomesoderm-specific GATA
transcription factors in the marine annelid Platynereis dumerilli. Evol Dev. 2007;9: 39–50.
doi:10.1111/j.1525-142X.2006.00136.x

16. Davidson EH, Rast JP, Oliveri P, Ransick A, Calestani C, Yuh C-H, et al. A Provisional Regulatory
Gene Network for Specification of Endomesoderm in the Sea Urchin Embryo. Dev Biol.
2002;246: 162–190. doi:10.1006/dbio.2002.0635

17. Shoichet SA, Malik TH, Rothman JH, Shivdasani RA. Action of the Caenorhabditis elegans
GATA factor END-1 in Xenopus suggests that similar mechanisms initiate endoderm
development in ecdysozoa and vertebrates. Proc Natl Acad Sci U S A. National Academy of
Sciences; 2000;97: 4076–81. Available: http://www.ncbi.nlm.nih.gov/pubmed/10760276

18. Sulston JE, Schierenberg E, White JG, Thomson JN. The embryonic cell lineage of the
nematode Caenorhabditis elegans. Dev Biol. 1983;100: 64–119. Available:
http://www.ncbi.nlm.nih.gov/pubmed/6684600

19. Bowerman B, Eaton BA, Priess JR. skn-1, a maternally expressed gene required to specify the
fate of ventral blastomeres in the early C. elegans embryo. Cell. 1992;68: 1061–75. Available:
http://www.ncbi.nlm.nih.gov/pubmed/1547503

20. Bowerman B, Draper BW, Mello CC, Priess JR. The maternal gene skn-1 encodes a protein
that is distributed unequally in early C. elegans embryos. Cell. Elsevier; 1993;74: 443–52.
doi:10.1016/0092-8674(93)80046-H

21. Maduro MF, Rothman JH. Making Worm Guts: The Gene Regulatory Network of the
Caenorhabditis elegans Endoderm. Dev Biol. Academic Press; 2002;246: 68–85.
doi:10.1006/DBIO.2002.0655

22. Maduro MF, Broitman-Maduro G, Mengarelli I, Rothman JH. Maternal deployment of the
embryonic SKN-1 → MED-1,2 cell specification pathway in C. elegans. Dev Biol. Academic
Press; 2007;301: 590–601. doi:10.1016/J.YDBIO.2006.08.029

23. Maduro MF, Meneghini MD, Bowerman B, Broitman-Maduro G, Rothman JH. Restriction of
mesendoderm to a single blastomere by the combined action of SKN-1 and a GSK-3beta
homolog is mediated by MED-1 and -2 in C. elegans. Mol Cell. 2001;7: 475–85. Available:
http://www.ncbi.nlm.nih.gov/pubmed/11463373

24. Maduro MF. Gut development in C. elegans. Semin Cell Dev Biol. 2017;66: 3–11.
doi:10.1016/j.semcdb.2017.01.001

25. Wiesenfahrt T, Osborne Nishimura E, Berg JY, McGhee JD. Probing and rearranging the
transcription factor network controlling the *C. elegans* endoderm. Worm. 2016;5: e1198869.
doi:10.1080/21624054.2016.1198869

26. McGhee J. The C. elegans intestine. WormBook. 2007; 1–36. doi:10.1895/wormbook.1.133.1

27. Meneghini MD, Ishitani T, Carter JC, Hisamoto N, Ninomiya-Tsuji J, Thorpe CJ, et al. MAP
kinase and Wnt pathways converge to downregulate an HMG-domain repressor in
Caenorhabditis elegans. Nature. Nature Publishing Group; 1999;399: 793–797.
doi:10.1038/21666

28. Shin TH, Yasuda J, Rocheleau CE, Lin R, Soto M, Bei Y, et al. MOM-4, a MAP kinase kinase
kinase-related protein, activates WRM-1/LIT-1 kinase to transduce anterior/posterior polarity
signals in C. elegans. Mol Cell. 1999;4: 275–80. Available:

http://www.ncbi.nlm.nih.gov/pubmed/10488343

29.   Bei Y, Hogan J, Berkowitz LA, Soto M, Rocheleau CE, Pang KM, et al. SRC-1 and Wnt signaling
act together to specify endoderm and to control cleavage orientation in early C. elegans
embryos. Dev Cell. 2002;3: 113–25. Available:
http://www.ncbi.nlm.nih.gov/pubmed/12110172

30.   Thorpe CJ, Schlesinger A, Carter JC, Bowerman B. Wnt signaling polarizes an early C. elegans
blastomere to distinguish endoderm from mesoderm. Cell. 1997;90: 695–705. Available:
http://www.ncbi.nlm.nih.gov/pubmed/9288749

31.   Nakamura K, Kim S, Ishidate T, Bei Y, Pang K, Shirayama M, et al. Wnt signaling drives WRM-
1/beta-catenin asymmetries in early C. elegans embryos. Genes Dev. Cold Spring Harbor
Laboratory Press; 2005;19: 1749–54. doi:10.1101/gad.1323705

32.   Rocheleau CE, Yasuda J, Shin TH, Lin R, Sawa H, Okano H, et al. WRM-1 Activates the LIT-1
Protein Kinase to Transduce Anterior/Posterior Polarity Signals in C. elegans. Cell. Cell Press;
1999;97: 717–726. doi:10.1016/S0092-8674(00)80784-9

33.   Owraghi M, Broitman-Maduro G, Luu T, Roberson H, Maduro MF. Roles of the Wnt effector
POP-1/TCF in the C. elegans endomesoderm specification gene network. Dev Biol. NIH Public
Access; 2010;340: 209–21. doi:10.1016/j.ydbio.2009.09.042

34.   Huang S, Shetty P, Robertson SM, Lin R. Binary cell fate specification during C. elegans
embryogenesis driven by reiterated reciprocal asymmetry of TCF POP-1 and its coactivator -
catenin SYS-1. Development. 2007;134: 2685–2695. doi:10.1242/dev.008268

35.   Maduro MF, Lin R, Rothman JH. Dynamics of a Developmental Switch: Recursive Intracellular
and Intranuclear Redistribution of Caenorhabditis elegans POP-1 Parallels Wnt-Inhibited
Transcriptional Repression. Dev Biol. Academic Press; 2002;248: 128–142.
doi:10.1006/DBIO.2002.0721

36.   Phillips BT, Kidd AR, King R, Hardin J, Kimble J. Reciprocal asymmetry of SYS-1/beta-catenin
and POP-1/TCF controls asymmetric divisions in Caenorhabditis elegans. Proc Natl Acad Sci.
2007;104: 3231–3236. doi:10.1073/pnas.0611507104

37.   Maduro MF, Kasmir JJ, Zhu J, Rothman JH. The Wnt effector POP-1 and the PAL-1/Caudal
homeoprotein collaborate with SKN-1 to activate C. elegans endoderm development. Dev
Biol. 2005;285: 510–523. doi:10.1016/j.ydbio.2005.06.022

38.   Shetty P, Lo M-C, Robertson SM, Lin R. C. elegans TCF protein, POP-1, converts from
repressor to activator as a result of Wnt-induced lowering of nuclear levels. Dev Biol.
2005;285: 584–592. doi:10.1016/j.ydbio.2005.07.008

39.   Lin KT-H, Broitman-Maduro G, Hung WWK, Cervantes S, Maduro MF. Knockdown of SKN-1
and the Wnt effector TCF/POP-1 reveals differences in endomesoderm specification in C.
briggsae as compared with C. elegans. Dev Biol. 2009;325: 296–306.
doi:10.1016/j.ydbio.2008.10.001

40.   Félix M-A, Braendle C. The natural history of Caenorhabditis elegans. Curr Biol. Elsevier;
2010;20: R965-9. doi:10.1016/j.cub.2010.09.050

41.   Cook DE, Zdraljevic S, Roberts JP, Andersen EC. CeNDR, the *Caenorhabditis elegans* natural
diversity resource. Nucleic Acids Res. 2017;45: D650–D657. doi:10.1093/nar/gkw893

42.   Andersen EC, Gerke JP, Shapiro JA, Crissman JR, Ghosh R, Bloom JS, et al. Chromosome-scale
selective sweeps shape Caenorhabditis elegans genomic diversity. Nat Genet. NIH Public

Access; 2012;44: 285–90. doi:10.1038/ng.1050

43.   Kontani K, Moskowitz IPG, Rothman JH. Repression of Cell-Cell Fusion by Components of the
C. elegans Vacuolar ATPase Complex. Dev Cell. 2005;8: 787–794.
doi:10.1016/j.devcel.2005.02.018

44.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77: 71–94. Available:
http://www.ncbi.nlm.nih.gov/pubmed/4366476

45.   Kamath RS, Ahringer J. Genome-wide RNAi screening in Caenorhabditis elegans. Methods.
Academic Press; 2003;30: 313–321. doi:10.1016/S1046-2023(03)00050-1

46.   Rual J-F, Ceron J, Koreth J, Hao T, Nicot A-S, Hirozane-Kishikawa T, et al. Toward Improving
Caenorhabditis elegans Phenome Mapping With an ORFeome-Based RNAi Library. Genome
Res. 2004;14: 2162–2168. doi:10.1101/gr.2505604

47.   Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, et al. Systematic functional
analysis of the Caenorhabditis elegans genome using RNAi. Nature. Nature Publishing Group;
2003;421: 231–237. doi:10.1038/nature01278

48.   Gomez-Amaro RL, Valentine ER, Carretero M, LeBoeuf SE, Rangaraju S, Broaddus CD, et al.
Measuring Food Intake and Nutrient Absorption in Caenorhabditis elegans. Genetics.
Genetics Society of America; 2015;200: 443–54. doi:10.1534/genetics.115.175851

49.   Sommermann EM, Strohmaier KR, Maduro MF, Rothman JH. Endoderm development in
Caenorhabditis elegans: The synergistic action of ELT-2 and -7 mediates the
specification→differentiation transition. Dev Biol. 2010;347: 154–166.
doi:10.1016/j.ydbio.2010.08.020

50.   Clokey G V, Jacobson LA. The autofluorescent "lipofuscin granules" in the intestinal cells of
Caenorhabditis elegans are secondary lysosomes. Mech Ageing Dev. 1986;35: 79–94.
Available: http://www.ncbi.nlm.nih.gov/pubmed/3736133

51.   Hermann GJ, Schroeder LK, Hieb CA, Kershner AM, Rabbitts BM, Fonarev P, et al. Genetic
Analysis of Lysosomal Trafficking in Caenorhabditis elegans. Mol Biol Cell. 2005;16: 3273–
3288. doi:10.1091/mbc.E05

52.   Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, et al. Efficient Control of
Population Structure in Model Organism Association Mapping. Genetics. 2008;178: 1709–
1723. doi:10.1534/genetics.107.080101

53.   Millstein J, Volfson D. Computationally efficient permutation-based confidence interval
estimation for tail-area FDR. Front Genet. Frontiers Media SA; 2013;4: 179.
doi:10.3389/fgene.2013.00179

54.   Hansen E, Kerr KF. A Comparison of Two Classes of Methods for Estimating False Discovery
Rates in Microarray Studies. Scientifica (Cairo). 2012;2012: 1–9. doi:10.6064/2012/519394

55.   Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R language.
Bioinformatics. Narnia; 2004;20: 289–290. doi:10.1093/bioinformatics/btg412

56.   Revell LJ. phytools: an R package for phylogenetic comparative biology (and other things).
Methods Ecol Evol. John Wiley & Sons, Ltd (10.1111); 2012;3: 217–223. doi:10.1111/j.2041-
210X.2011.00169.x

57.   Ives AR, Midford PE, Garland T. Within-Species Variation and Measurement Error in
Phylogenetic Comparative Methods. Oakley T, editor. Syst Biol. 2007;56: 252–270.
doi:10.1080/10635150701313830

58.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: A Tool Set
1032   for Whole-Genome Association and Population-Based Linkage Analyses. Am J Hum Genet.
1033   2007;81: 559–575. doi:10.1086/519795

1034   59.   Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. A Robust, Simple
1035   Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. Orban L, editor. PLoS
1036   One. Public Library of Science; 2011;6: e19379. doi:10.1371/journal.pone.0019379

1037   60.   Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. TASSEL: software for
1038   association mapping of complex traits in diverse samples. Bioinformatics. 2007;23: 2633–
1039   2635. doi:10.1093/bioinformatics/btm308

1040   61.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call
1041   format and VCFtools. Bioinformatics. Oxford University Press; 2011;27: 2156–8.
1042   doi:10.1093/bioinformatics/btr330

1043   62.   Broman KW, Sen S. A Guide to QTL Mapping with R/qtl. New York, NY: Springer New York;
1044   2009; 1–20. doi:10.1007/978-0-387-92125-9

1045   63.   Broman KW, Wu H, Sen S, Churchill GA. R/qtl: QTL mapping in experimental crosses.
1046   Bioinformatics. 2003;19: 889–90. Available: http://www.ncbi.nlm.nih.gov/pubmed/12724300

1047   64.   Zhao Z, Boyle TJ, Bao Z, Murray JI, Mericle B, Waterston RH. Comparative analysis of
1048   embryonic cell lineage between Caenorhabditis briggsae and Caenorhabditis elegans. Dev
1049   Biol. NIH Public Access; 2008;314: 93–9. doi:10.1016/j.ydbio.2007.11.015

1050   65.   Cutter AD. Divergence Times in Caenorhabditis and Drosophila Inferred from Direct Estimates
1051   of the Neutral Mutation Rate. Mol Biol Evol. Oxford University Press; 2008;25: 778–786.
1052   doi:10.1093/molbev/msn024

1053   66.   Sterken MG, Snoek LB, Kammenga JE, Andersen EC. The laboratory domestication of
1054   Caenorhabditis elegans. Trends Genet. NIH Public Access; 2015;31: 224–31.
1055   doi:10.1016/j.tig.2015.02.009

1056   67.   Raj A, Rifkin SA, Andersen E, van Oudenaarden A. Variability in gene expression underlies
1057   incomplete penetrance. Nature. Nature Publishing Group; 2010;463: 913–918.
1058   doi:10.1038/nature08781

1059   68.   Zhu J, Hill RJ, Heid PJ, Fukuyama M, Sugimoto A, Priess JR, et al. end-1 encodes an apparent
1060   GATA factor that specifies the endoderm precursor in Caenorhabditis elegans embryos.
1061   Genes Dev. Cold Spring Harbor Laboratory Press; 1997;11: 2883–96. Available:
1062   http://www.ncbi.nlm.nih.gov/pubmed/9353257

1063   69.   Tijsterman M, Okihara KL, Thijssen K, Plasterk RH. PPW-1, a PAZ/PIWI Protein Required for
1064   Efficient Germline RNAi, Is Defective in a Natural Isolate of C. elegans. Curr Biol. Cell Press;
1065   2002;12: 1535–1540. doi:10.1016/S0960-9822(02)01110-7

1066   70.   Pollard DA, Rockman M V. Resistance to Germline RNA Interference in a Caenorhabditis
1067   elegans Wild Isolate Exhibits Complexity and Nonadditivity. G3 Genes, Genomes, Genet. G3:
1068   Genes, Genomes, Genetics; 2013;3: 941–947. doi:10.1534/G3.113.005785

1069   71.   Gleason JE, Korswagen HC, Eisenmann DM. Activation of Wnt signaling bypasses the
1070   requirement for RTK/Ras signaling during C. elegans vulval induction. Genes Dev. Cold Spring
1071   Harbor Laboratory Press; 2002;16: 1281–90. doi:10.1101/gad.981602

1072   72.   Paaby AB, White AG, Riccardi DD, Gunsalus KC, Piano F, Rockman M V. Wild worm
1073   embryogenesis harbors ubiquitous polygenic modifier variation. Elife. eLife Sciences

Publications Limited; 2015;4: e09178. doi:10.7554/eLife.09178

73.   Çelen İ, Doh JH, Sabanayagam CR. Effects of liquid cultivation on gene expression and
phenotype of C. elegans. BMC Genomics. BioMed Central; 2018;19: 562. doi:10.1186/s12864-
018-4948-7

74.   Wang J, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, et al. An estimator for
pairwise relatedness using molecular markers. Genetics. Genetics; 2002;160: 1203–15.
doi:10.1534/genetics.167.1.531

75.   Rieseberg LH, Widmer A, Arntz AM, Burke B. The genetic architecture necessary for
transgressive segregation is common in both natural and domesticated populations. Philos
Trans R Soc B Biol Sci. 2003;358: 1141–1147. doi:10.1098/rstb.2003.1283

76.   Félix M-A, Ashe A, Piffaretti J, Wu G, Nuez I, Bélicard T, et al. Natural and Experimental
Infection of Caenorhabditis Nematodes by Novel Viruses Related to Nodaviruses. Hodgkin J,
editor. PLoS Biol. Public Library of Science; 2011;9: e1000586.
doi:10.1371/journal.pbio.1000586

77.   Ruf V, Holzem C, Peyman T, Walz G, Blackwell TK, Neumann-Haefelin E. TORC2 signaling
antagonizes SKN-1 to induce C. elegans mesendodermal embryonic development. Dev Biol.
Academic Press; 2013;384: 214–227. doi:10.1016/J.YDBIO.2013.08.011

78.   Witze ES, Field ED, Hunt DF, Rothman JH. C. elegans pur alpha, an activator of end-1,
synergizes with the Wnt pathway to specify endoderm. Dev Biol. 2009;327: 12–23.
doi:10.1016/j.ydbio.2008.11.015

79.   Walston T, Tuskey C, Edgar L, Hawkins N, Ellis G, Bowerman B, et al. Multiple Wnt Signaling
Pathways Converge to Orient the Mitotic Spindle in Early C. elegans Embryos. Dev Cell. Cell
Press; 2004;7: 831–841. doi:10.1016/J.DEVCEL.2004.10.008

80.   Tatebe H, Shiozaki K. Evolutionary Conservation of the Components in the TOR Signaling
Pathways. Biomolecules. 2017;7: 77. doi:10.3390/biom7040077

81.   Adoutte A, Philippe H. The major lines of metazoan evolution: summary of traditional
evidence and lessons from ribosomal RNA sequence analysis. EXS. 1993;63: 1–30. Available:
http://www.ncbi.nlm.nih.gov/pubmed/8422536

82.   Carroll SB. Evo-Devo and an Expanding Evolutionary Synthesis: A Genetic Theory of
Morphological Evolution. Cell. 2008;134: 25–36. doi:10.1016/j.cell.2008.06.030

83.   Page GP, George V, Go RC, Page PZ, Allison DB. "Are We There Yet?": Deciding When One Has
Demonstrated Specific Genetic Causation in Complex Diseases and Quantitative Traits. Am J
Hum Genet. 2003;73: 711–719. doi:10.1086/378900

84.   Mackay TFC. Epistasis and quantitative traits: using model organisms to study gene-gene
interactions. Nat Rev Genet. NIH Public Access; 2014;15: 22–33. doi:10.1038/nrg3627

85.   Volis S, Shulgina I, Zaretsky M, Koren O. Epistasis in natural populations of a predominantly
selfing plant. Heredity (Edinb). Nature Publishing Group; 2011;106: 300–9.
doi:10.1038/hdy.2010.79

86.   Félix M-A. Cryptic Quantitative Evolution of the Vulva Intercellular Signaling Network in
Caenorhabditis. Curr Biol. 2007;17: 103–114. doi:10.1016/j.cub.2006.12.024

87.   Barkoulas M, van Zon JS, Milloz J, van Oudenaarden A, Félix M-A. Robustness and Epistasis in
the C. elegans Vulval Signaling Network Revealed by Pathway Dosage Modulation. Dev Cell.
Cell Press; 2013;24: 64–75. doi:10.1016/J.DEVCEL.2012.12.001

88.   Duveau F, Félix M-A. Role of Pleiotropy in the Evolution of a Cryptic Developmental Variation
1118   in Caenorhabditis elegans. Noor MAF, editor. PLoS Biol. Public Library of Science; 2012;10:
1119   e1001230. doi:10.1371/journal.pbio.1001230

1120   89.   Johnson NA, Porter AH. Evolution of branched regulatory genetic pathways: directional
1121   selection on pleiotropic loci accelerates developmental system drift. Genetica. Springer
1122   Netherlands; 2006;129: 57–70. doi:10.1007/s10709-006-0033-2

1123   90.   Phillips PC. Epistasis--the essential role of gene interactions in the structure and evolution of
1124   genetic systems. Nat Rev Genet. NIH Public Access; 2008;9: 855–67. doi:10.1038/nrg2452

1125   91.   Wagner GP, Zhang J. The pleiotropic structure of the genotype–phenotype map: the
1126   evolvability of complex organisms. Nat Rev Genet. Nature Publishing Group; 2011;12: 204–
1127   213. doi:10.1038/nrg2949

1128   92.   Shafer B, Onishi K, Lo C, Colakoglu G, Zou Y. Vangl2 Promotes Wnt/Planar Cell Polarity-like
1129   Signaling by Antagonizing Dvl1-Mediated Feedback Inhibition in Growth Cone Guidance. Dev
1130   Cell. Cell Press; 2011;20: 177–191. doi:10.1016/J.DEVCEL.2011.01.002

1131   93.   Zheng C, Diaz-Cuadros M, Chalfie M. Dishevelled attenuates the repelling activity of Wnt
1132   signaling during neurite outgrowth in Caenorhabditis elegans. Proc Natl Acad Sci U S A.
1133   National Academy of Sciences; 2015;112: 13243–8. doi:10.1073/pnas.1518686112

1134   94.   Rivera-Pomar R, Jäckle H. From gradients to stripes in Drosophila embryogenesis: filling in the
1135   gaps. Trends Genet. Elsevier Current Trends; 1996;12: 478–483. doi:10.1016/0168-
1136   9525(96)10044-5

1137   95.   Maduro MF, Broitman-Maduro G, Choi H, Carranza F, Wu AC-Y, Rifkin SA. MED GATA factors
1138   promote robust development of the C. elegans endoderm. Dev Biol. Academic Press;
1139   2015;404: 66–79. doi:10.1016/J.YDBIO.2015.04.025

1140   96.   Choi H, Broitman-Maduro G, Maduro MF. Partially compromised specification causes
1141   stochastic effects on gut development in C. elegans. Dev Biol. Academic Press; 2017;427: 49–
1142   60. doi:10.1016/J.YDBIO.2017.05.007

1143   97.   Maduro MF. Developmental robustness in the *Caenorhabditis elegans* embryo. Mol Reprod
1144   Dev. 2015;82: 918–931. doi:10.1002/mrd.22582

1145   98.   Zhan T, Rindtorff N, Boutros M. Wnt signaling in cancer. Oncogene. Nature Publishing Group;
1146   2017;36: 1461–1473. doi:10.1038/onc.2016.304

1147   99.   Jenkins ZA, van Kogelenberg M, Morgan T, Jeffs A, Fukuzawa R, Pearl E, et al. Germline
1148   mutations in WTX cause a sclerosing skeletal dysplasia but do not predispose to
1149   tumorigenesis. Nat Genet. 2009;41: 95–100. doi:10.1038/ng.270

1150   100.   Baron R, Gori F. Targeting WNT signaling in the treatment of osteoporosis. Curr Opin
1151   Pharmacol. Elsevier; 2018;40: 134–141. doi:10.1016/J.COPH.2018.04.011

1152   101.   Chen N, Wang J. Wnt/β-Catenin Signaling and Obesity. Front Physiol. 2018;9: 792.
1153   doi:10.3389/fphys.2018.00792

1154   102.   Schinner S. Wnt-signalling and the Metabolic Syndrome. Horm Metab Res. 2009;41: 159–163.
1155   doi:10.1055/s-0028-1119408

1156   103.   Frankel N, Davis GK, Vargas D, Wang S, Payre F, Stern DL. Phenotypic robustness conferred by
1157   apparently redundant transcriptional enhancers. Nature. 2010;466: 490–493.
1158   doi:10.1038/nature09158

1159   104.   Mohun T. Muscle differentiation. Curr Opin Cell Biol. 1992;4: 923–8. Available:
1160   http://www.ncbi.nlm.nih.gov/pubmed/1485959

1161   105.   Imai Y, Gates MA, Melby AE, Kimelman D, Schier AF, Talbot WS. The homeobox genes vox
1162   and vent are redundant repressors of dorsal fates in zebrafish. Development. 2001;128:
1163   2407–20. Available: http://www.ncbi.nlm.nih.gov/pubmed/11493559

1164   106.   Manley NR, Capecchi MR. Hox Group 3 Paralogous Genes Act Synergistically in the Formation
1165   of Somitic and Neural Crest-Derived Structures. Dev Biol. Academic Press; 1997;192: 274–
1166   288. doi:10.1006/DBIO.1997.8765

1167   107.   Nam J, Nei M. Evolutionary change of the numbers of homeobox genes in bilateral animals.
1168   Mol Biol Evol. NIH Public Access; 2005;22: 2386–94. doi:10.1093/molbev/msi229

1169   108.   True JR, Haag ES. Developmental system drift and flexibility in evolutionary trajectories. Evol
1170   Dev. John Wiley & Sons, Ltd (10.1111); 2001;3: 109–119. doi:10.1046/j.1525-
1171   142x.2001.003002109.x

1172   109.   Yoon CH, Chang C, Hopper NA, Lesa GM, Sternberg PW. Requirements of multiple domains of
1173   SLI-1, a Caenorhabditis elegans homologue of c-Cbl, and an inhibitory tyrosine in LET-23 in
1174   regulating vulval differentiation. Mol Biol Cell. American Society for Cell Biology; 2000;11:
1175   4019–31. Available: http://www.ncbi.nlm.nih.gov/pubmed/11071924

1176   110.   Braendle C, Félix M-A. Plasticity and Errors of a Robust Developmental System in Different
1177   Environments. Dev Cell. 2008;15: 714–724. doi:10.1016/j.devcel.2008.09.011

1178   111.   Grimbert S, Vargas Velazquez AM, Braendle C. Physiological Starvation Promotes
1179   Caenorhabditis elegans Vulval Induction. G3 (Bethesda). Genetics Society of America; 2018;8:
1180   3069–3081. doi:10.1534/g3.118.200449

1181   112.   Zheng M, Messerschmidt D, Jungblut B, Sommer RJ. Conservation and diversification of Wnt
1182   signaling function during the evolution of nematode vulva development. Nat Genet. 2005;37:
1183   300–304. doi:10.1038/ng1512

1184   113.   Tian H, Schlager B, Xiao H, Sommer RJ. Wnt Signaling Induces Vulva Development in the
1185   Nematode Pristionchus pacificus. Curr Biol. 2008;18: 142–146. doi:10.1016/j.cub.2007.12.048

1186   114.   Wang X, Sommer RJ. Antagonism of LIN-17/Frizzled and LIN-18/Ryk in Nematode Vulva
1187   Induction Reveals Evolutionary Alterations in Core Developmental Pathways. Sternberg PW,
1188   editor. PLoS Biol. 2011;9: e1001110. doi:10.1371/journal.pbio.1001110

1189   115.   Haag ES, Fitch DHA, Delattre M. From "the Worm" to "the Worms"
1190   and Back Again: The Evolutionary Developmental Biology of Nematodes. Genetics. Genetics
1191   Society of America; 2018;210: 397–433. doi:10.1534/genetics.118.300243

1192   116.   Kalinka AT, Varga KM, Gerrard DT, Preibisch S, Corcoran DL, Jarrells J, et al. Gene expression
1193   divergence recapitulates the developmental hourglass model. Nature. Nature Publishing
1194   Group; 2010;468: 811–814. doi:10.1038/nature09634

1195   117.   Gerstein MB, Rozowsky J, Yan K-K, Wang D, Cheng C, Brown JB, et al. Comparative analysis of
1196   the transcriptome across distant species. Nature. Nature Publishing Group; 2014;512: 445–
1197   448. doi:10.1038/nature13424

1198   118.   Levin M, Anavy L, Cole AG, Winter E, Mostov N, Khair S, et al. The mid-developmental
1199   transition and the evolution of animal body plans. Nature. Nature Publishing Group;
1200   2016;531: 637–641. doi:10.1038/nature16994

1201   119.   Levin M, Hashimshony T, Wagner F, Yanai I. Developmental Milestones Punctuate Gene

1202   Expression in the Caenorhabditis Embryo. Dev Cell. 2012;22: 1101–1108.
1203   doi:10.1016/j.devcel.2012.04.004

1204   120.   Zalts H, Yanai I. Developmental constraints shape the evolution of the nematode mid-
1205   developmental transition. Nat Ecol Evol. Nature Publishing Group; 2017;1: 0113.
1206   doi:10.1038/s41559-017-0113

1207   121.   Yanai I, Peshkin L, Jorgensen P, Kirschner MW. Mapping Gene Expression in Two Xenopus
1208   Species: Evolutionary Constraints and Developmental Flexibility. Dev Cell. 2011;20: 483–496.
1209   doi:10.1016/j.devcel.2011.03.015

1210   122.   Irie N, Kuratani S. Comparative transcriptome analysis reveals vertebrate phylotypic period
1211   during organogenesis. Nat Commun. Nature Publishing Group; 2011;2: 248.
1212   doi:10.1038/ncomms1248

1213   123.   Cruickshank T, Wade MJ. Microevolutionary support for a developmental hourglass: gene
1214   expression patterns shape sequence variation and divergence in *Drosophila*. Evol Dev. John
1215   Wiley & Sons, Ltd (10.1111); 2008;10: 583–590. doi:10.1111/j.1525-142X.2008.00273.x

1216   124.   Cutter AD, Garrett RH, Mark S, Wang W, Sun L. Molecular evolution across developmental
1217   time reveals rapid divergence in early embryogenesis. Evol Lett. John Wiley & Sons, Ltd; 2019;
1218   doi:10.1002/evl3.122

1219   125.   Kalinka AT, Varga KM, Gerrard DT, Preibisch S, Corcoran DL, Jarrells J, et al. Gene expression
1220   divergence recapitulates the developmental hourglass model. Nature. Nature Publishing
1221   Group; 2010;468: 811–814. doi:10.1038/nature09634

1222   126.   Raff RA. The shape of life : genes, development, and the evolution of animal form. University
1223   of Chicago Press; 1996.

1224   127.   Domazet-Lošo T, Tautz D. A phylogenetically based transcriptome age index mirrors
1225   ontogenetic divergence patterns. Nature. Nature Publishing Group; 2010;468: 815–818.
1226   doi:10.1038/nature09632

1227   128.   Coroian C, Broitman-Maduro G, Maduro MF. Med-type GATA factors and the evolution of
1228   mesendoderm specification in nematodes. Dev Biol. 2006;289: 444–455.
1229   doi:10.1016/j.ydbio.2005.10.024

1230   129.   Maduro MF, Hill RJ, Heid PJ, Newman-Smith ED, Zhu J, Priess JR, et al. Genetic redundancy in
1231   endoderm specification within the genus Caenorhabditis. Dev Biol. 2005;284: 509–522.
1232   doi:10.1016/j.ydbio.2005.05.016

1233

Supplementary figures

Supplementary Fig. 1: High reproducibility of skn-1(RNAi) **phenotypes in various** C. elegans **isotypes.**

A minimum of two replicates were obtained, with >500 embryos per replicate (N = 3). Box-plot represents median with range bars showing upper and lower quartiles.

Supplementary Fig. 2: The requirement for POP-1 in endoderm formation does not vary in three introgressed strains.

Strains are shown on the x-axis and fraction of arrested embryos with endoderm are shown on the y-axis. The orange bars represent the results from mutant lines: MY16;pop-1(zu189), JU440;pop-1(zu189), and KR314;pop-1(zu189). Four introgressed lines were created for each new mutant strain. Black bars represent pop-1(RNAi) results on the wild isolates indicated. >200 embryos were scored per experiment.

1249

1250 ***Supplementary Fig. 3: Histogram of mom-2(RNAi) phenotype among the 94 wild isolates.***

1251 *A beta-distribution is observed (skewed to the right). Shapiro-Wilk normality test (W=0.8682, p-*
1252 *value=1.207X10^{-7}).*

1253

1254

Supplementary Fig. 4: Comparison of EMMA p-values for both mom-2 and skn-1 RNAi phenotypes.

Heatmap of p-values for mom-2(RNAi) *(left) and* skn-1(RNAi) *(right) as calculated in the GWAS analyses (see Fig. 5A, B). Strength of association between genotype and endoderm formation phenotypes is represented as –log$_{10}$(p-value), here depicted as a heatmap (lighter colors – weaker association, darker colors – stronger association). An overlap (indicated by arrow head) is found in a small region of chromosome IV, but no further correlations are observed. Significant SNPs for* skn-1(RNAi) *are shown in Table S1.*

1262

Supplementary Fig. 5: MOM-2 requirement does not correlate with genotypic relatedness or geographical location.

(A) mom-2(RNAi) phenotype of 94 isolates arranged with respect to the neighbor-joining tree constructed using 4,690 SNPs and pseudo-rooted to QX1211. Red asterisk indicates an example of closely related strains (JU394 and CB4851) with similar phenotypes, while black asterisks indicate examples sister strains (JU792 and JU1242; JU1440 and JT11398) with distinct phenotypes (λ = 6.94X10^-05, p-value = 1). (B) Worldwide distribution of mom-2(RNAi) phenotype across 94 isolates. Each circle represents a single isolate.

1270

1271

1272

Supplementary Fig. 6: Correlation between the skn-1(RNAi) and mom-2(RNAi) allelic differences.

Each dot represents a SNP, colour-coded by chromosome. A small genome-wide correlation is observed, but significant for only three of the chromosomes individually. Represented here is a set of pruned SNPs (N = 321) to cover the whole genome, and corrected for LD (also used for calculations). Z-score is used to calculate median of each allelic group, correcting for outliers. Strength of correlation (Pearson's R) represented. Significance levels: non-significant, p-value < 0.05 (*), p-value < 0.01 (**), p-value < 0.001 (***).

1280

Supplementary Fig. 7: Introgression of N2 and MY16 regions in chromosome IV.

Near-Isogenic Lines (NILs) were built between strains N2 (purple) and MY16 (green), using visual mutations as markers, with their locations in cM indicated in each graph. (A) Introgression of chromosome IV region flanked by dpy-20 and unc-30. (B) Introgression of chromosome IV region flanked by dpy-20 and unc-26. (C) Introgression of chromosome IV region flanked by bli-6 and unc-22.

Supplementary Table S1: Significantly linked SNPs for skn-1(RNAi) GWAS.

SNP	EMMA -log(p)	N2 allele	Variant allele
IV: 5,079,371 bp	*4.957651645*	*A*	*G*
IV: 5,725,367 bp	*5.211140897*	*C*	*T*
IV: 5,761,153 bp	*5.211140897*	*G*	*A*
IV: 5,891,378 bp	*4.252324884*	*G*	*A*
IV: 5,920,597 bp	*4.720037892*	*T*	*A*
IV: 5,921,302 bp	*4.720037892*	*T*	*G*
IV: 5,921,510 bp	*4.252324884*	*C*	*T*
IV: 6,453,892 bp	*5.142174312*	*T*	*A*

IV: 6,511,989 bp	5.142174312	C	A
IV: 6,563,740 bp	4.678423021	C	T
IV: 7,453,945 bp	4.652004517	G	A
IV: 7,453,143 bp	4.181579989	A	G

1288