A new role for Notch in control of polarity and asymmetric cell division of developing T cells

Running Title: Notch controls cell polarity and is asymmetrically inherited during ACD

Mirren Charnley[1,2,3]*, Mandy Ludford-Menting[3], Kim Pham[1,3,4,5] and Sarah M. Russell[1,3,4,5]*

[1]Centre for Micro-Photonics, Faculty of Science, Engineering and Technology, Swinburne University of Technology, Hawthorn, Victoria 3122, Australia

[2]Biointerface Engineering, Faculty of Science, Engineering and Technology, Swinburne University of Technology, Hawthorn, Victoria 3122, Australia

[3]Immune Signalling Laboratory, Peter MacCallum Cancer Centre, Parkville, Victoria 3000, Australia

[4]Department of Pathology, The University of Melbourne, Parkville, Victoria 3010, Australia

[5]Sir Peter MacCallum Department of Oncology, The University of Melbourne, Parkville, Victoria 3010, Australia

*Corresponding authors:

Mirren Charnley, Centre for Micro-Photonics, Faculty of Science, Engineering and Technology, Swinburne University of Technology, Mail No H74 | PO Box 218, Hawthorn, Victoria, 3122, Australia, E-mail: mcharnley@swin.edu.au

Sarah M. Russell, Centre for Micro-Photonics, Faculty of Science, Engineering and Technology, Swinburne University of Technology, Mail No H74 | PO Box 218, Hawthorn, Victoria, 3122, Australia, E-mail: sarah.russell@petermac.org

Current address

M. Ludford-Menting: ACRF Translation Research Laboratory, Victorian Comprehensive Cancer Centre, Parkville, Victoria 3000, Australia

Kim Pham: Division of Immunology, The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria, 3052, Australia.
Department of Medical Biology, The University of Melbourne, Parkville, Victoria 3052 Australia.

Abstract

A fundamental question in biology is how single cells can reliably produce progeny of different cell types. Notch signalling frequently facilitates fate determination. Asymmetric cell division (ACD) often controls segregation of Notch signalling by imposing unequal inheritance of regulators of Notch. Here, we assessed the functional relationship between Notch and ACD in mouse T cell development. To attain immunological specificity, developing T cells must pass through a pivotal stage termed β-selection, which involves Notch signalling and ACD. We assessed functional interactions between Notch and ACD in T cell development using direct presentation of Notch ligands, DL1 and DL4, and pharmacological inhibition of Notch signalling.Contrary to prevailing models, we find Notch controls distribution of Notch1 itself and cell fate determinants, α-Adaptin and Numb. Notch and CXCR4 signalling cooperated to drive polarity during division. Thus, Notch signalling directly orchestrates ACD, and Notch1 is differentially inherited by sibling cells.

Introduction

Developing multicellular organisms must produce multiple cell types from one progenitor cell. This is achieved by sequential bifurcations in the fate of daughter cells following cell division. Fate bifurcation is driven by various means, including random differences in the daughters, different positional cues, or controlled segregation of fate determinants differentially into the two daughter cells, a process termed Asymmetric Cell Division (ACD) (Dewey et al., 2015; Schweisguth, 2015; Venkei et al., 2018). A common mediator of fate bifurcation, which uses all of these processes, is the Notch signalling pathway (Artavanis-Tsakonas et al., 1999; Fortini, 2009; Liu et al., 2017; Sjoqvist et al., 2017). Notch is a transmembrane receptor that responds to ligands belonging to the Delta and Jagged family, which are present on adjacent cells (Bray, 2016). Notch signalling directly regulates transcription to influence cell fate decisions such as proliferation, death, multi-potency, differentiation and self-renewal (Fortini, 2009). Understanding how Notch signals differently in different tissue contexts is a subject of intense investigation (Bray, 2016). Differential Notch signalling in two daughter cells often occurs indirectly through the differential inheritance of regulators of Notch (Bray, 2016). Indeed, some of the most frequently observed fate determinants controlled by ACD are the Notch regulators, Numb and α-Adaptin (Daeden et al., 2018; Dewey et al., 2015; Rhyu et al., 1994; Schweisguth, 2015). Consequently, the prevailing view is that, although an important executor of cell fate, Notch does not directly determine the direction of bifurcation in fate (Schweisguth, 2015; Venkei et al., 2018).

We considered the possibility that Notch might play a more active role in dictating the direction of fate bifurcation. One observation suggestive of a controlling role for Notch showed that Notch signalling was required for asymmetric distribution of Numb during ACD of *Drosophila* neural progenitors (Bhat, 2014; Pinto-Teixeira et al., 2014). We have previously observed that Notch was polarised during interphase in developing T cells (Pham et al., 2015), leading to two non-mutually exclusive hypotheses: (i) that Notch might remain asymmetric during division, and thus be asymmetrically partitioned in the daughters, and (ii) that Notch might control partitioning of cell fate determinants during division. We test those two hypotheses in this study using an *in vitro* model of T cell development.

T cell development is a highly tractable model for the study of cell fate, with well-established *in vitro* models and an extensive knowledge of the molecular pathways involved (Janas et al., 2010; Mohtashami et al., 2010; Schmitt et al., 2002; Van de Walle et al., 2013; Yui et al., 2014). Many aspects of T cell biology share mechanisms of fate control with other developmental systems, and both Notch signalling and ACD operate in developing and mature T cells (Chang et al., 2007; Oliaro et al., 2010; Tajbakhsh et al., 2009). ACD occurs specifically during the β-selection phase of T cell development (Pham et al., 2015). In this critical phase, the developing T cells (termed DN3 cells) must balance self-

renewal, genomic DNA recombination to produce a functional T cell receptor chain, and differentiation. Notch signalling is required for progression through β-selection and subsequent fate choices (Ciofani et al., 2004; Ciofani et al., 2005). We exploited two model systems to explore the role of Notch1 during division of developing T cells; the OP9-DL1 coculture, and surfaces functionalised with Fc-DL1 (Janas et al., 2010; Mohtashami et al., 2010; Schmitt et al., 2002; Shukla et al., 2017; Van de Walle et al., 2013). We show that the interaction of Notch1 with its ligands, DL1 and DL4, drives the polarisation of Notch1 itself, and consequently the polarisation of cell fate determinants, α-Adaptin and Numb. Together, these data show that Notch signalling directly acts as a polarity cue, and that Notch1 is differentially inherited by the daughters of an ACD.

Results

Notch signalling drives polarisation during interphase and cell division

To discern whether Notch is a passive participant or driver of polarity in developing T cells, we first assessed the role of Notch signalling on the polarisation of developing T cells in interphase. Using established methods (Janas et al., 2010; Mohtashami et al., 2010; Schmitt et al., 2002; Van de Walle et al., 2013), we drove the development of mouse T cells from hematopoietic stem cells using the OP9-DL1 stromal cell line that expresses the Notch ligand, DL1. To ensure that the cells were poised to, but had not yet entered the b-selection checkpoint, we sorted for the DN3a stage by flow cytometry and reseeded onto OP9-DL-1 cells. Notch signalling was disrupted using the γ-secretase inhibitor (dibenzazepine, DBZ) (De Kloe et al., 2014). As previously published (Ciofani et al., 2004), Notch inhibition in DN3 cells reduced differentiation to the DN4 and DP stages, reduced cell proliferation, and increased cell death (**Fig. S1**). To determine the role of Notch signalling in polarity, we performed immunofluorescence microscopy to assess the localisation of α-tubulin and markers of cell polarity on fixed DN3a cells in interphase (**Fig. S2**). We chose an early time point (15 hours) where inhibiting Notch signalling had little or no impact on the number of attached DN3a cells (**Fig. 1A**), but slightly reduced microtubule organising centre (MTOC) polarisation to the interface between the DN3a cell and stromal cell (**Fig. 1B**). As previously seen (Pham et al., 2015), Notch1 was polarised to the interface (**Fig 1C**). Remarkably, the polarisation of Notch1 was dependent upon its signalling capacity, as DBZ treatment substantially reduced Notch1 polarisation. CXCR4, the receptor for the chemokine CXCL12 which is required for progression through β-selection and regulates polarity in DN3 cells (Pham et al., 2015; Trampont et al., 2010), was also less polarised after DBZ treatment (**Fig. 1D**). The non-polarised control protein, CD25, was not affected by DBZ (**Fig. 1E**). Two cell fate regulators that are polarised during ACD of developing T cells, α-Adaptin and Numb (Pham et al., 2015), also failed to polarise upon DBZ treatment (**Fig. 1F, G**). Thus, inhibiting Notch signalling reduced polarity of Notch1, CXCR4, α-Adaptin and Numb from above 70% to between 24 and 42%, similar to the level of polarisation observed with the CD25 control (Fig. 1E). These data suggest that Notch signalling was required for the polarisation of DN3a cells.

We next asked whether Notch1 was polarised during cell division, and if Notch signalling was required. Dividing cells were scored as undergoing symmetric cell division (SCD) if the protein of interest was symmetrically distributed between the two daughter cells, and as undergoing ACD if the protein of interest was more concentrated in one of the daughter cells (**Fig. S3**). As seen previously (Pham et al., 2015), Numb and α-Adaptin were asymmetric in 54.7 to 65.7% of dividing cells, and we further show that Notch1 and CXCR4 were also asymmetric in 55.4 and 51.0% of dividing cells, while CD25 was asymmetric in only 29.2% (**Fig. 2A-E**). Inhibiting Notch signalling reduced the asymmetry of the

markers in dividing cells to the same level as the CD25 control. CXCR4 acts as a polarity cue for ACD (Pham et al., 2015), so the effect of Notch signalling on CXCR4 polarisation indicates that these cues could cooperate to trigger ACD. These data together indicate that Notch1 was polarised during division of DN3a cells, and suggest that Notch signalling is required for the polarisation of Notch1, CXCR4, α-Adaptin and Numb.

Surfaces functionalised with only the Notch ligand, DL1, were sufficient to induce polarisation during cell division

The pharmacological inhibition experiments above showed that γ-secretase was necessary for polarisation of DN3a cells during interphase and division. To confirm that these effects were through Notch, and to assess whether Notch signalling was sufficient for polarity, we used surfaces functionalised with the Notch ligand, DL1. DL1 was coupled to protein A (PA) coated surfaces via an Fc linker to control the orientation of DL1 relative to the cell (Makaraviciute et al., 2013; Song et al., 2012; Toda et al., 2011). Growth factors essential to DN3a survival and differentiation were included in the cultures (Janas et al., 2010), but these were dispersed throughout the culture media and only the Notch ligand was spatially restricted to the surface and consequently presented to the cells with a defined orientation. As others have shown (Janas et al., 2010), surfaces functionalised with Fc-DL1 supported DN3 differentiation, albeit not as efficiently as OP9-DL1 cells (**Fig. S4**). The frequency and duration of divisions on the Fc-DL1 surfaces was comparable to that on the OP9-DL1 cells (**Fig. S5**). After two hours on the Fc-DL1 surfaces, the MTOC, α-Adaptin and Numb were recruited to the interface and this was maintained at 15 hours (**Fig. S6**). These data demonstrate that the immobilised DL1 ligand alone was sufficient to trigger the recruitment of α-Adaptin and Numb to the functionalised surface.

We then asked whether the surfaces coated with Fc-DL1 were capable of triggering asymmetry during division. First, DN3a cells were cultured on surfaces functionalised with Fc-DL1 or PA for 15 hours, fixed and labelled with α-tubulin and either α-Adaptin, Numb or CD25, imaged and blind scored using fluorescence microscopy. α-Adaptin and Numb had equal or greater polarity as compared to OP9-DL1 cells, which was specific to surfaces functionalised with DL1 (**Fig. 3A-C**). Given that both Notch1 and its negative regulator, Numb, were polarised, we scored whether they were on the same or opposing sides of the dividing cell. Interestingly, for dividing cells with polarised Notch1 and Numb, 83% showed corecruitment of both proteins into the same daughter cell (**Fig. 3D**). This indicates that Notch ligation by DL1 drove polarisation during division.

To complement the analysis of the fixed cells and quantify the extent of asymmetry, DN3a cells were transduced with fluorescently tagged α-Adaptin and Numb, cultured in cell paddocks onto which OP9-DL1 stromal cells had been allowed to attach overnight or were coated with PA or Fc-DL1 and imaged using time lapse microscopy (Day et al., 2009). The fluorescence intensity was quantified in dividing cells and expressed as polarisation ratios (PR; calculated as the difference in fluorescence between the two halves divided by the sum of the fluorescence. A low PR value indicates that the fluorescence is symmetric in the dividing cells, a high PR value indicates that it is asymmetric). A diffuse marker was also transduced into the DN3a cells (either GFP or cherry) to determine the background level of asymmetry and remove any artefacts (e.g. out of focus cells) (Charnley et al., 2017; Pham et al., 2015; Shimoni et al., 2014). The over-expression of the protein of interest had little or no effect on proliferation, differentiation or the level of polarity (**Fig. S7**) (Pham et al., 2015; Pham et al., 2013). To ensure against possible artefacts related to the nature of the tag (Couturier et al., 2014; Short, 2014), we compared the polarity of α-Adaptin and Numb coupled to both GFP and cherry and saw no effect (**Fig. S8A, B**). Consequently, the data for the two fluorophores was combined (indicated as FP-α-Adaptin and FP-Numb). Dividing DN3a cells showed strong polarisation of both α-Adaptin (**Fig 3E, F**, **S8C**) and Numb (**Fig 3G, H**, **S8D**) on OP9-DL1 stromal cells. Dividing DN3a cells also showed strong polarisation of α-Adaptin and Numb when cultured on surfaces functionalised with DL1, which was comparable to that observed with OP9-DL1. Conversely, DN3a cells on the PA functionalised surfaces predominately divided symmetrically. As we previously observed with OP9-DL1 cells (Pham et al., 2015), asymmetric divisions occurred in DN3a cells but not in DN3b or DN2b cells (**Fig. S9**). Thus, it is possible to mimic the stromal cell interaction and induce cell polarity by the orientated presentation of a single protein, namely Fc-DL1. This indicates that the binding of Notch1 to its ligand was both necessary and sufficient to dictate the polarisation of α-Adaptin and Numb during division.

Notch ligands presented on functionalised beads coordinate polarity and mitotic spindle orientation during ACD

ACD requires that polarity is coupled with orientation of the mitotic spindle (Venkei et al., 2018). We next asked whether the protein asymmetry triggered by Notch ligation was coordinated with the mitotic spindle. ACD of developing T cells requires a mitotic spindle orthogonal to the interface with the stromal cell (Pham et al., 2015). To determine the orientation of the cue relative to the axis of division, we used beads to pinpoint the localisation of the polarity cue (Habib et al., 2013), functionalised using the same strategy as the flat surfaces. The proportion of polarised divisions observed for the PA and Fc-DL1 functionalised beads was similar to the proportion observed for the flat surfaces (**Fig. S10**). When the dividing cell was in contact with OP9-DL1 cells or Fc-DL1 functionalised beads, higher PRs correlated with greater division angles (**Fig. 4A, B**). Binning the data to separate cells based on division

angle revealed that divisions perpendicular to the cue (division angles greater than 45°) showed higher PR values, indicating polarised divisions, compared with divisions that were more parallel to the cue (**Fig. 4C, D**). This demonstrates that the protein asymmetry observed was coordinated with the orientation of the mitotic spindle, indicative of ACD. Thus, Notch ligands were necessary and sufficient to coordinate polarity and the spindle to orchestrate ACD.

Notch functionally interacts with other cues to orchestrate ACD

The only directional cue presented to DN3a cells that promoted polarisation in the above experiments was DL1. However, the CXCR4 ligand, CXCL12, is required for T cell differentiation (Janas et al., 2010; Tussiwand et al., 2011) and was included in our cultures. We have previously found that ACD induced by OP9-DL1 stromal cells depends upon CXCR4 signalling in DN3a cells (Pham et al., 2015). This, combined with our observations above that CXCR4 polarisation depends on Notch signalling and that surfaces functionalised with the Notch ligand, DL1, induce asymmetry of α-Adaptin and Numb in dividing DN3a cells, raises the question of how these two receptors cooperate. We therefore assessed whether cells cultured on Fc-DL1 functionalised surfaces in the presence of CXCL12 could still polarise after pharmacological inhibition of one or both of the CXCR4 and Notch signalling pathways, using AMD3100 or DBZ, respectively (**Fig. 5A**, **S11A-C**). Indeed, both Notch and CXCR4 signalling were required for optimal polarisation of α-Adaptin in dividing DN3a cells, with the least polarisation occurring when both inhibitors were present. The effect of disrupting CXCR4 signalling is intriguing given that CXCL12 is diffuse in the cultures, so we looked specifically at the localisation of the two receptors during division. As expected, Notch1 was polarised in response to surface-bound DL1, to a level that was comparable to OP9-DL1 cells (**Fig. 5B**). Notch1 polarisation was strongly inhibited by the Notch inhibitor, but only minimally affected by inhibition of CXCR4 signalling. Surprisingly, CXCR4 was also polarised in response to presentation of the Notch ligand, and this polarisation was dependent on signalling through both Notch1 and CXCR4 (**Fig. 5C**). These data indicate that Notch ligation can drive the asymmetric polarisation of CXCR4 and that CXCR4 and Notch cooperate as non-redundant cues for ACD, albeit with Notch playing a more dominant role. Collectively these data show that multiple cues can interact to enhance or reduce the level of ACD.

Given the complexity of interactions in the thymus (Petrie et al., 2007), we evaluated two other ligands for the regulation of ACD in DN3a cells. DL4 is an alternative ligand for Notch1 (Hozumi et al., 2008; Hozumi et al., 2004; Koch et al., 2008), and we assessed whether it behaved similarly to DL1. Indeed, both ligands induced equivalent polarisation of α-Adaptin in dividing DN3a cells (**Fig. 5D, S11D**). VCAM-1 contributes to stromal-DN3 interactions and its presence is associated with improved *in vitro* generation of T cells (Abe et al., 2010; Shukla et al., 2017). We therefore assessed the impact of Fc-

VCAM-1 on DL4-mediated polarisation during division by functionalising the surfaces with both ligands. Interestingly, the inclusion of Fc-VCAM-1 slightly reduced the level of polarisation (**Fig. 5D, E, S11E**). One possible explanation for this decrease is that the presence of VCAM-1 altered the interaction of the DL1 and CXCL12 ligands with their receptors. Indeed, on the Fc-DL4 + Fc-VCAM-1 surfaces the polarisation of both Notch1 and CXCR4 was also slightly reduced (**Fig. 5F, G**). This effect correlated with a decrease in polarisation at interphase of the MTOC and α-Adaptin (**Fig. S12A-C**), but was not associated with an effect on differentiation (**Fig. S12D-F**). These data show that the slight reduction in ACD observed when VCAM-1 was included in the functionalised surfaces was associated with reduced polarisation of Notch1 and CXCR4.

Discussion

It is well established that Notch controls diversification in cell fate in a number of developmental systems. However, most of the proposed mechanisms assume that Notch executes decisions that have been made stochastically or are imposed by unequal expression of regulators of Notch (Dewey et al., 2015; Schweisguth, 2015; Venkei et al., 2018). Here, we demonstrate that Notch1 can directly trigger asymmetry in daughter cells, by steering ACD and by partitioning itself asymmetrically in the two daughter cells (**Fig. 6**).

Two model systems were used to explore the role of Notch1 during division of developing T cells; the OP9-DL1 coculture, and surfaces functionalised with Fc-DL1. The OP9-DL1 coculture system has been extensively used to study T cell development *in vitro*, and mimics physiological early T cell development with high fidelity (Janas et al., 2010; Mohtashami et al., 2010; Schmitt et al., 2002; Van de Walle et al., 2013). Surfaces functionalised with Notch ligands have also provided insights into the mechanisms controlling thymocyte development (Dallas et al., 2005; Janas et al., 2010; Tussiwand et al., 2011). Further, there is a growing interest in using surfaces functionalised with Notch ligands for the *in vitro* generation of T cells for therapeutic treatments, such as cancer immunology ((Gehre et al., 2015; Shukla et al., 2017; Suzuki et al., 2006) and as reviewed in (Singh et al., 2018)). Recently it was demonstrated that it is possible to direct human hematopoietic stem cells along the T cell lineage utilising functionalised surfaces in serum-free conditions (Shukla et al., 2017), which would circumvent issues associated with xenogeneic culture and enable the generation of clinical-grade pro-T cells.

Functionalised surfaces for the controlled presentation of proteins have been widely used to study fundamental biology, including adhesion and migration (Charnley et al., 2012; Gavard et al., 2004; Kovacs et al., 2002; Silvestre et al., 2009), division (Charnley et al., 2013; den Elzen et al., 2009; Toyoshima et al., 2007), immune responses (Charnley et al., 2009; Grakoui et al., 1999; Manz et al., 2010; Mossman et al., 2005) and T cell development (Dallas et al., 2005; Janas et al., 2010; Tussiwand et al., 2011). Here, we demonstrated that these surfaces successfully mimicked another aspect of T cell development; developing T cells cultured on the surfaces underwent ACD and this process was coordinated by external cues and restricted to the DN3a stage of development. This work also highlights that functionalised surfaces can be used to manipulate the cells' microenvironment and systemically test the influence of extrinsic factors on ACD.

The interaction of Notch1 with the DL1 ligand drove the polarisation of fate determinants that are asymmetrically distributed in several models of ACD, namely α-Adaptin and Numb. Notch signalling

also triggered the polarisation of CXCR4, which is a polarity cue for ACD of DN3a cells (Pham et al., 2015), and the polarisation of Notch1 itself. Thus, the orientated presentation of the Notch1 ligand was sufficient to trigger polarity during division in developing T cells. The role of Notch in controlling localisation of Numb is analogous to that observed in *Drosophila* ganglion mother cells (Bhat, 2014), indicating that its role as a polarity cue could be more broadly conserved.

In addition to its role in controlling polarity, we demonstrate that, in dividing DN3a cells, Notch1 was asymmetrically distributed between the two daughter cells. Asymmetric partitioning of Notch1 itself during an ACD provides direct mechanism to establish variation in Notch signalling levels. Interestingly, Notch1 localisation is also controlled at cytokinesis in dividing sensory organ precursor cells (Coumailleau et al., 2009; Couturier et al., 2012; Trylinski et al., 2017). However, here Notch was localised at the interface between two daughter cells during division, rather than polarised to one side of the mother cell. Whether preferential recruitment of both Notch and Numb into the same daughter cell leads to a net loss or gain of Notch signalling is not known. Using developing T cells as a model system of ACD we demonstrated that Notch1 is asymmetrically inherited providing a direct and novel mechanism to establish differential Notch signalling in the two daughters (Fig 6).

Interestingly, Notch1 signalling is required for T cell development past the β-selection stage, but peaks at the DN3a stage (Mingueneau et al., 2013) and is dramatically downregulated following β-selection (Yashiro-Ohtani et al., 2009). The requirement for Notch1 in controlling ACD identified here is compatible with our finding that ACD is restricted to DN3a cells. A role for Notch1 in dictating ACD also provides a possible explanation for the enduring effects of Notch1 following its downregulation: perhaps one consequence of ACD is to enable one daughter to differentiate and expand (Swamy et al., 2016).

Notch1, pre-TCR and CXCR4 signalling cooperate to drive the developing T cell through β-selection, and the presence of both CXCR4 and Notch1 signalling is required for survival on the functionalised surfaces (Ciofani et al., 2004; Ciofani et al., 2005; Janas et al., 2010; Trampont et al., 2010). These results indicate that they also cooperate to dictate polarity and ACD in developing T cells. A functional interaction of Notch1 and CXCR4 has been demonstrated in a number of cell systems (Chiaramonte et al., 2015; Cong et al., 2017; Ho et al., 2017; Wang et al., 2009; Xiao et al., 2017), but the molecular basis for this interaction is not clear. The colocalisation of Notch1 and CXCR4 could indicate that they physically interact, either via a direct interaction of the receptors or their ligands. The striking loss of CXCR4 polarisation when Notch signalling is inhibited suggests that Notch is upstream of the

chemokine receptor. Taken together, these findings indicate that Notch1 and CXCR4 have complementary and non-redundant roles in dictating polarity to control ACD.

These experiments reveal a new paradigm by which Notch controls cell fate. In developing T cells, Notch plays a key role in dictating its own asymmetry during cell division and influencing asymmetry of other fate determinants. This work focused on developing T cells, but given the influence of Notch on Numb asymmetry in neural precursor cells, this might be a more general paradigm.

Acknowledgments

We thank Juan Carlos Zúñiga-Pflücker (University of Toronto, Sunnybrook Health Sciences Centre, Toronto, Canada), for OP9 stromal cell lines. This work was performed in part at the Biointerface Engineering Hub @ Swinburne, part of the Victorian node of the Australian National Fabrication Facility (ANFF), a company established under the National Collaborative Research Infrastructure Strategy to provide nano-and micro-fabrication facilities for Australia's researchers. We thank Louse Cheng (Peter MacCallum Cancer Centre) and Helena Richardson (La Trobe University) for comments on the manuscript.

Funding

We acknowledge support from the Swiss National Science Foundation (SNSF) (grants PA00P3_142120 and P300P3_154664 to MC), the Australian Research Council (FT0990405 to SMR), the National Health and Medical Research Council (APP1099140 to SMR).

Methods

Primary developing T cell co-culture

Mouse hematopoietic stem cells (isolated from C57BL/6 fetal liver at E14.5) were seeded onto OP9-DL1 stromal cells (received from Juan-Carlos Zúñiga-Pflücker, University of Toronto, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada) at a 1:1 ratio in 6 well plates (2×10^5) in Minimal Essential Medium Alpha Modification supplemented with foetal calf serum (10% v/v), glutamine (1 mM), β-mercaptoethanol (50 µM), sodium pyruvate (1 nM), HEPES (10 mM), penicillin/streptomycin (100 ng/mL), mouse interleukin 7 (IL-7, 1 ng/mL) and mouse FMS-like tyrosine kinase 3 (Flt-3, 5 ng/mL). Hematopoietic cells were harvested via forceful pipetting and co-cultured on fresh OP9-DL1 stromal cells every 3–8 days. All mice were maintained in a specific pathogen-free environment with food and water freely available. All experiments on mice were performed in accordance with the Animal Experimentation Ethics Committee of the Peter MacCallum Cancer Centre.

Retroviral transduction

Phoenix E cells (provided by Garry Nolan, Stanford University, Stanford, CA) were maintained at 37°C and at 10% CO_2 in Dulbecco's Minimal Essential Medium supplemented with foetal calf serum (10% v/v), L-glutamine (1 mM) and penicillin/streptomycin (100 ng/mL). Calcium phosphate transfection was performed on Phoenix E cells with 10-20 µg of the following pMSCV retroviral constructs: Cherry, GFP, GFP-Numb, Cherry-Numb, Cherry-α-Adaptin, and GFP-α-Adaptin in 10 cm dishes (Corning). Viral supernatant was harvested 48 h after transfection and added to 6 well plates that had been precoated with 15 µg/ml RetroNectin (Takara Bio Inc.) and blocked with 2% BSA. After addition of the viral supernatant, plates were spun at 2,000 *g* for 1 h and incubated for 1 h at 37°C. 5×10^5 hematopoietic cells (day 4 co-culture) were added, and plates were spun for 1 h at 1,200 *g*.

Flow cytometry

Developing T cell subsets were purified by staining for the cell surface markers CD44 and CD25 to discriminate between DN 1-4, CD28 to discriminate between DN3a and DN3b, and CD4 and CD8 to discriminate between the DN and DP populations. Hematopoietic cells were harvested from OP9 cocultures by forceful pipetting and stained on ice for one hour with the following antibodies: biotin-CD4 (1:2000; BD Pharmingen, 553045), biotin CD8a (1:2000; BD Pharmingen, 553029), PerCP/Cy5.5 CD44 (1:300; Biolegend 103032), APC-eFluor 780 CD25 (1:300; eBioscience, 47-0251-82) and Pe/Cy7 CD28 (1:200; Biolegend, 122014). DN3a cells were isolated on a FACS (FACS Aria III; BD Biosciences) based on surface expression (CD28^{lo}/CD25$^+$/CD44^{lo}/CD4$^-$/CD8$^-$). Retrovirally transduced cells were sorted 72 hours after transduction and also sorted on the basis of GFP and Cherry

fluorescence. For the analysis of T cell differentiation, cells were stained with Pacific Blue CD4 (1:600; BD Pharmingen, 558107), APC-eFluor 780 CD8a (1:300; eBioscience, 47-0081-82), PerCP/Cy5.5 CD44 (1:300; Biolegend 103032), APC CD25 (1:300; eBioscience, 17-0251-82), Pe/Cy7 CD28 (1:200; Biolegend, 122014) and Sytox green viability stain (30 nM; Molecular Probes, S34860).

Substrate functionalisation

Functionalised surfaces were used to individually present ligands to the developing T cells. Protein A (PA) was used to couple the Fc linked ligands to the surface to ensure optimum orientation of the ligand (Makaraviciute et al., 2013; Song et al., 2012; Toda et al., 2011). Glass coverslips, silicon dioxide beads (5 µm) or 24 well plates were coated with protein A (235 nM, 1 hour), rinsed with PBS and further functionalised, if required, with Fc-DL1 (86 nM; R&D Systems, 5026-DL), Fc-DL4 (86 nM; BioVision, P1163), Fc-VCAM-1 (19 nM; R&D Systems, 643VM) or Fc-DL4 and Fc-VCAM-1 (86 nM + 19 nM) for 30 minutes. The substrates were then washed in PBS and media before cell deposition. For live cell imaging PDMS cell paddocks (120 µm x 120 µm) (Day et al., 2009) were rendered hydrophilic by exposure to air plasma at 1.5×10^{-2} mbar for 10 minutes and placed into an 8 well chamber slide (Ibidi) prior to protein functionalisation.

Proliferation and development of DN3 cells on the functionalised surfaces

For the analysis of the ability of functionalised surfaces to support developing T cell proliferation and differentiation, 24 well plates were coated with protein A, protein A plus Fc-linked proteins or OP9-DL1 stromal cells. 1×10^5 DN3 cells were seeded into the wells and assessed at 1 and 5 days by flow cytometry.

Immunofluorescence and fixed image acquisition by confocal microscopy

2×10^4 DN3a cells were added to 8 well chamber slide (Thermo Fisher Scientific) precoated with protein A, protein A plus Fc-linked proteins or OP9-DL1 stromal cells and cultured for 2 or 15 hours. For stromal cell-free culture cells were cultured in the presence of Flt-3 (5 ng/mL), CXCL12 (10 nM) and IL-7 (10 ng/ml) (Janas et al., 2010). Cells were then fixed, permeabilized, and labelled with primary antibodies for α-tubulin (1:800; Rocklands, 600-401-880 or 200-301-880) and test proteins, Notch1 (1:200; Abcam, ab27526), α-Adaptin (1:200; Abcam, Ab2730), Numb (1:200; Abcam, Ab4147), CXCR4 (1:200; BD Pharmingen, 551968), CD25 (1:1000; BD Pharmingen, 553070), and the appropriate secondary antibodies and then mounted with Prolong gold antifade (Molecular Probes). The slides were examined using an Olympus FV3000 microscope (Olympus) and 40x objective lens (0.95 NA Oil Plan Apochromat). 3D images of the cells were acquired with a z distance of 0.5 µm and

maximum intensity projections of z sections created using ImageJ. For the analysis of polarisation during interphase α-tubulin staining was used to identify cells with the MTOC at the cell-cell / cell-substrate interface and the cell was blind scored depending on whether the protein of interest was diffuse or enriched at the MTOC. To determine polarisation during division mitotic cells were identified by the presence of a mitotic spindle and the cell division was assigned as symmetric if the fluorescence of the test protein was evenly distributed between the two daughter cells and asymmetric if the fluorescence was greater in one of the daughter cells than the other daughter cell.

Live cell image acquisition and analysis

For live cell imaging, 2×10^4 DN3a cells were added to 8 well chamber slide (Ibidi) with the PDMS cell paddocks precoated with protein A, Fc-linked proteins or OP9-DL1 stromal cells. For stromal cell-free culture cells were cultured in the presence of Flt-3 (5 ng/mL), CXCL12 (10 nM) and IL-7 (10 ng/mL) (Janas et al., 2010). Images were captured on a spinning disc confocal system fitted with an inverted microscope and temperature controlled chamber maintained at 37°C and 5% CO_2. Images are acquired using a 40x air objective (0.95 NA) and multiple stage positions are recorded every 3 min for 20 hours, with 8-slice z-stacks of 1 µm thickness. As described in (Charnley et al., 2017; Oliaro et al., 2010; Pham et al., 2015; Shimoni et al., 2014), the polarisation of fluorescent markers in the nascent daughter cells was quantified by measuring the total fluorescence intensity of each daughter cell and applying a "Polarization Ratio" (PR) equation: $PR = (\Sigma H1 - \Sigma H2)/ (\Sigma H1 + \Sigma H2)$; where the difference in total intensity between Daughter 1 ($\Sigma H1$) and Daughter 2 ($\Sigma H2$) is divided by the sum of intensities in both Daughter 1 (H1) and Daughter 2 (H2). PR of 0.17 was used as a cutoff to designate cells as undergoing ACD (PR > 0.17) or SCD (PR < 0.17). On the functionalised surfaces, only cells that had not interacted with another cell for at least 30 minutes prior to division were selected for analysis. This enabled the isolation of the effect of the immobilised protein on ACD. To compare the distribution of PR α-Adaptin values between the different cell culture platforms, outliers (which were identified by their high PR diffuse values and consisted of less than 10% of dividing cells) were removed (Charnley et al., 2017) and the data was re-plotted as violin plots using the kernel density function.

To assess the role of the orientation of the polarity cue, 5 µm silicon dioxide beads were coated with PA or PA + Fc-DL1 using the same protocol outlined for the 2D substrates. The orientation of division relative to the polarity cue was determined for dividing DN3a cells in contact with functionalised beads and OP9-DL1 stromal cells. For DN3a cells in contact with OP9-DL1 stromal cells, the division angle was determined by measuring the angle between the interface between the DN3a cell and OP9-DL1 stromal cell and the long axis of the dividing cell. For the functionalised beads, a line was drawn

between the centre of mass of the bead and the centre of mass of the dividing cell. The division angle was defined as the angle orthogonal to the angle between this line and the long axis of the dividing cell.

Notch and CXCR4 inhibition protocol

To disrupt Notch signalling we used the γ-secretase inhibitor, dibenzazepine (DBZ; 0.05 µM) (van Es et al., 2005). DBZ prevents the cleavage of γ-secretase substrates Notch, and so prevents translocation of the cytoplasmic portion of Notch to the nucleus and prevents signalling. To analyse the effect of Notch signalling on proliferation and differentiation, purified DN3 cells were co-cultured on OP9-DL1 stromal cells in the presence or absence of DBZ and assessed by flow cytometry. To determine the effect of Notch and CXCR4 inhibition on ACD, purified DN3a cells were cultured on OP9-DL1 stromal cells or Fc-DL1 functionalised surfaces in the presence of AMD3100 (2 µg/ml, Sigma-Aldrich) or DBZ (0.05 µM, Calbiochem) for approximately 30 mins prior to the start of the time lapse imaging or for 15 hours prior to fixing for immunofluorescence analysis.

Statistical Analysis

All data was collected from 2 to 8 independent experiments and shown as data are mean ± SEM. The following tests were used for the statistical analysis: Student's unpaired two-way t-test was used to determine the difference between mean values for the scatter plots, for the PR scatter plots from the time lapse microscopy experiments paired *t* tests were performed between control (diffuse protein) and test protein PR values and the violin plots were analysed for differences in the distribution of the data using Two-sample Kolmogorov-Smirnov (KS) test (Young, 1977). Level of statistical significance is indicated as: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure Legends

Figure 1. Intact Notch signalling is required for DN3a cell polarisation during interphase. DN3a cells were seeded on OP9-DL1 cells in the absence ('no treat') or presence of DBZ (0.05 µM) for 15 hours, fixed and stained for α-tubulin and Notch1, Numb, α-Adaptin, CXCR4 or CD25. Inhibiting Notch signalling **(A)** did not affect thymocyte attachment to the stromal cell and **(B)** only slightly reduced MTOC recruitment to the interface between the stromal cell and the DN3a cell. Representative projected z stack images are shown on the left, and quantification on the right. **(C)** Notch1 and **(D)** CXCR4 were recruited to the cell-cell interface in over half the cells and this was severely disrupted in the presence of DBZ. **(E)** CD25 was not polarised and unperturbed by inhibiting Notch signalling. The cell fate regulators **(F)** α-Adaptin and **(G)** Numb were also polarised and this was reliant on intact Notch signalling. Total number of cells analysed: MTOC = 809 and 589, Notch1 = 130 and 110, CXCR4 = 158 and 113, CD25 = 68, 67, α-Adaptin = 131 and 105 and Numb = 158 and 110, for no treat and DBZ, respectively. Bar = 10 µm, * indicates the interface between the DN3a and OP9-DL1 stromal cells. n = 3 independent experiments. All data are represented as mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 2. Intact Notch signalling is required for DN3a cell polarisation during cell division. DN3a cells were seeded on OP9-DL1 cells in the absence ('no treat') or presence of DBZ (0.05 µM) for 15 hours, fixed and stained for α-tubulin and either **(A)** Notch1, **(B)** CXCR4, **(C)** CD25, **(D)** α-Adaptin or **(E)** Numb. Representative projected z stack images are shown on the left, and quantification on the right. Inhibiting Notch signalling significantly reduced the percentage of cells undergoing ACD. Total number of cells analysed: Notch1 = 57 and 44, CXCR4 = 43 and 41, CD25 = 30 and 30, α-Adaptin = 46 and 56 and Numb = 40 and 45, for no treat and DBZ, respectively. Bar = 10 µm. n = 3-5 independent experiments. All data are represented as mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 3. Dividing DN3a cells polarise cell fate determinants, α-Adaptin and Numb on functionalised surfaces. DN3a cells were cultured on surfaces pre-coated with PA or PA + Fc-DL1 (DL1) or OP9-DL1 stromal cells for 15 hours before fixing and staining with α-tubulin and either **(A)** α-Adaptin, **(B)** Numb or **(C)** CD25. Representative projected z stack images are shown on the left, and quantification on the right. α-Adaptin and Numb were asymmetrically distributed in a high proportion of cells when cultured on OP9-DL1 stromal cells or on Fc-DL1 functionalised surfaces, but not on surfaces only functionalised with PA. CD25 was symmetrically distributed regardless of the surface. Total number of divisions analysed: α-Adaptin = 40, 57 and 42; Numb = 46, 42 and 45; CD25 = 86, 73 and 67 for OP9-DL1, PA and Fc-DL1, respectively. **(D)** To determine if Notch and Numb inherited into the same or opposing daughters cells, DN3a cells cultured on Fc-DL1 and stained with both Notch1

and Numb. Dividing cells which were asymmetric for both Notch and Numb were selected for analysis and scored as colocalised if both proteins were concentrated within the same nascent daughter cell. Total number of divisions analysed: Notch = 83; Colocalisation (coloc) = 44. **(E)** Projected z stack images from time lapse movies of dividing DN3a cells expressing GFP and cherry-α-Adaptin. **(F)** Absolute PR values were calculated as the absolute difference in fluorescence between the two halves divided by the sum of fluorescence. Cells with high diffuse PR (see Fig S8) values were removed and the data plotted in the form of violin plots using the kernel density function. Culturing DN3a cells on OP9-DL1 stromal cells and Fc-DL1 increased the proportion of dividing cells with high PRs. Number of divisions analysed: OP9-DL1 = 208, PA = 102 and Fc-DL1 = 109. **(G, H)** A similar trend was observed for DN3a cells transduced with Numb. Number of divisions analysed = OP9-DL1 = 148, PA = 48 and Fc-DL1 = 45. Bar = 10 µm. n = 3-8 independent experiments. All data are represented as mean ± SEM; *p < 0.05, **p < 0.01, ***p < 0.001. See Fig. S6 for PR scatter plots.

Figure 4. Polarity during division correlates with spindle orientation. DN3a cells expressing a diffuse protein (GFP or cherry) and test protein (α-Adaptin or Numb) were cultured in contact with 5 µm beads functionalised with Fc-DL1 (DL1) or OP9-DL1 stromal cells. There was a correlation between increased division angles and polarity α-Adaptin on **(A)** OP9-DL1 stromal cells **(B)** Fc-DL1 functionalised beads. The blue shaded region indicates divisions that were assigned as asymmetric when the 0.17 cut-off was applied. **(C, D)** The same trend was observed when we compared the spread in PR values for cells which divided with a division angle < 45° *versus* > 45°. Number of divisions analysed: OP9-DL1 = 200; Fc-DL1 beads = 51. n = 3-8 independent experiments; ***p < 0.001. See Fig. S10 for PR scatter and violin plots.

Figure 5. Different polarity cues cooperate to regulate ACD. DN3a cells expressing a diffuse protein (GFP or cherry) and α-Adaptin were cultured onto surfaces pre-coated with PA, Fc-DL1 (DL1), Fc-DL4 (DL4) or Fc-DL4 + Fc-VCAM-1 (D+V) and imaged using time-lapse microscopy for 20 hours. The addition of **(A)** either DBZ or AMD3100 (AMD) or a combination of both inhibitors caused a reduction in the level of ACD. Number of divisions: DBZ = 68, AMD3100 = 65, DBZ + AMD3100 = 51. **(B)** The polarisation of Notch1 in dividing cells was reduced by DBZ, but only slightly effected by AMD3100. **(C)** Conversely, CXCR4 polarisation was reduced by the inhibition of either Notch or CXCR4 signalling. Number of divisions analysed = 55, 48 and 42 for no treat, DBZ and AMD3100, respectively. **(D)** Fc-DL4 coated surfaces triggered ACD and the inclusion of Fc-VCAM-1 lead to a reduction in the asymmetric segregation of α-Adaptin during division. Number of divisions analysed: DL4 = 67; DL4 + VCAM = 58. The presence of VCAM-1 in the functionalised surfaces also resulted

in reduced polarisation of **(E)** α-Adaptin, **(F)** Notch1 and **(G)** CXCR4 in dividing cells. Number of divisions analysed: DL4 = 56; DL4 + VCAM = 44 and DL4 = 64 (α-Adaptin); DL4 = 67; DL4 + VCAM = 71 (Notch1 and CXCR4). n = 3 – 6 independent experiments; All data are represented as mean ± SEM; * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$. See Fig. S11 for PR scatter plots.

Figure 6. A model of the role of Notch in controlling cell polarity and ACD. Notch ligation of DL1, presented either by the OP9-DL1 stromal cell or DL1 functionalised surfaces, orchestrates polarisation during the division of DN3a cells. CXCR4 signalling cooperates with Notch to drive this asymmetry during division. This results in the differential inheritance of the cell fate determinants, α-Adaptin and Numb, CXCR4 and Notch1 itself.

Figure S1. Notch signalling is important for downstream DN3 proliferation, survival and differentiation. DN3 cells were seeded on OP9-DL1 cells in the absence (no treat) or presence of γ-secretase inhibitor (DBZ; 0.05 µM) for up to 8 days and assessed using flow cytometry. **(A)** The inclusion of DBZ resulted in a reduction in the number of cells after 5 and 8 days of culture and **(B)** viability. **(C)** Flow cytometry plots from one representative experiment. **(D)** Treatment with DBZ reduced differentiation, evidenced by a decrease in DN4s and DPs. *n* = 3 independent experiments, with flow cytometry plots from one representative experiment shown. All data are represented as mean ±SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure S2. Analysis of polarisation during interphase in DN3a cells. DN3a cells were seeded on OP9-DL1 cells in the absence ('no treat') or presence of DBZ (0.05 µM) for 15 hours, fixed and stained for α-tubulin and Notch1, Numb, α-Adaptin, CXCR4 or CD25. To determine polarity during interphase, cells in which the MTOC was recruited to the stromal interface were selected for analysis and the protein of interest was assigned as diffuse or polarised, where polarisation was defined as a clear enrichment of fluorescence at the interface. Bar = 10 µm, * indicates the interface between the DN3a and OP9-DL1 stromal cells.

Figure S3. Analysis of polarisation during cell division in DN3a cells. DN3a cells were seeded on OP9-DL1 cells in the absence ('no treat') or presence of DBZ (0.05 µM) for 15 hours, fixed and stained for α-tubulin and Notch1, Numb, α-Adaptin, CXCR4 or CD25. Dividing cells were identified from α-tubulin staining and scored as undergoing SCD if the protein of interest was symmetrically distributed between the two daughter cells or ACD if the protein of interest was more concentrated in one of the daughter cells. Bar = 10 µm, * indicates the interface between the DN3a and OP9-DL1 stromal cells.

Figure S4. DL1 functionalised surfaces support DN3 proliferation and differentiation. DN3 cells were cultured onto surfaces pre-coated with PA, Fc-DL1 (DL1) or OP9-DL1 stromal cells and analysed with flow cytometry. **(A)** Flow cytometry plots from one representative experiment. Cells were cultured on the three surfaces for 1 day **(B, C, D)** or 5 days **(E, F, G)** and assessed for expansion, viability, and differentiation. Surfaces functionalised with Fc-DL1 supported DN3 development, although the expansion and development was reduced relative to culture on OP9-DL1 cells. Conversely, PA alone was not able to support thymocyte culture, as indicated by a substantial reduction in proliferation, increased cell death and a failure to develop past the DN stage. $n = 5$ independent experiments, with flow cytometry plots from one representative experiment shown. All data are represented as mean $\pm$SEM; *p < 0.05, **p < 0.01, ***p < 0.001.

Figure S5. Functionalised surfaces support the division of DN3a cells. (A) DN3a cells were cultured on OP9-DL1, PA and Fc-DL1 (DL1) for up to 40 hours. The time of divisions (defined as time that cytokinesis occurred) were binned into 10 hours intervals and expressed relative to the total number of divisions on that surface. Cells divided throughout the movies, although there was a decrease in the number of divisions on the PA surfaces at later timepoints. **(B)** The time required for the DN3a cells to divide (defined as nuclear envelope breakdown to cytokinesis) was not affected by the cell culture platform. **(C)** CTV data for DN3a cells cultured on the functionalised surfaces and OP9-DL1 stromal cells revealed that thymocytes divided on all of the cell culture platforms. **(D)** However, after 48 hours division was reduced on the PA surfaces and to a lesser extent on the Fc-DL1 functionalised surfaces. $n = 2$-7 independent experiments, with flow cytometry plots from one representative experiment shown. All data are represented as mean $\pm$SEM.

Figure S6. DN3a cells cultured on surfaces coated with the Notch ligand, DL1, polarise α-Adaptin and Numb during interphase. DN3a cells were cultured onto surfaces pre-coated with protein A (PA) or protein A and Fc-DL1 (DL1) for 2 or 15 hours before fixing. **(A)** After 2 hours thymocyte attachment was unaffected by the substrate coating, but **(B)** MTOC recruitment to the cell-substrate interface was increased in the presence of Fc-DL1 *versus* PA. A similar trend was seen after 15 hours **(C, D)** Total number of cells analysed for 2 h = 115 (PA) and 122 (Fc-DL1) and 15 h = 77 (PA) and 157 (Fc-DL1). xy slice of the cell-substrate interface of DN3a cells on functionalised surfaces (PA or Fc-DL1) and stained with α-tubulin and either **(E)** α-Adaptin, **(F)** Numb or **(G)** CD25. α-Adaptin and Numb were recruited to the cell-cell interface on the Fc-DL1 functionalised substrates but not on the PA only substrates. Conversely, CD25 was not polarised regardless of the substrate. A similar trend in polarisation was observed at 2 and 15 h. Total number of cells analysed at 2 h: PA = 65, 74 and 39;

DL1 = 54, 84 and 37 for α-Adaptin, Numb or CD25, respectively. Total number of cells analysed at 15 h: PA = 54, 69 and 35 and DL1 = 79, 124 and 38 for α-Adaptin, Numb or CD25, respectively. Bar = 10 µm. n = 3 - 5 independent experiments. All data are represented as mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure S7. Overexpression of α-Adaptin or Numb does not affect proliferation, differentiation or polarisation during division. DN3a cells expressing a diffuse protein (GFP or cherry) and α-Adaptin or Numb were cultured on OP9-DL1 stromal cells for 5 days and assessed using flow cytometry. **(A)** Transduction with either α-Adaptin or Numb had no effect on proliferation. The overexpression of **(B)** α-Adaptin or **(C)** Numb also had little effect on differentiation. **(D)** DN3a cells expressing a diffuse protein (GFP or cherry) and α-Adaptin were cultured onto surfaces pre-coated with PA or Fc-DL1 (DL1) for 15 hours. Dividing cells were identified from α-tubulin staining and scored as polarised or symmetric. The proportion of cells that were polarised was comparable to the proportion observed when assessed by antibody labelling of endogenous protein (compare with Fig. 3). Total number of divisions analysed: OP9-DL1 = 97, 52 and 45; PA = 99, 59 and 40; Fc-DL1 = 107, 55 and 52 for GFP, α-Adaptin and Numb, respectively. n = 2-9 independent experiments, all data are represented as mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure S8. DN3a cells undergo polarised divisions on functionalised surfaces. (A) PR scatter plots for dividing DN3a cells transduced with either GFP and cherry-α-Adaptin or GFP-α-Adaptin and cherry and cultured in contact with OP9-DL1 stromal cells. **(B)** PR scatter plots for dividing DN3a cells transduced with either GFP and cherry-Numb or GFP-Numb and cherry and cultured in contact with OP9-DL1 stromal cells. PR was calculated as the difference in fluorescence between the two halves divided by the sum of fluorescence. The distribution of PR values was similar regardless of whether the test protein was coupled to GFP or cherry. PR scatter plots for dividing DN3a cells expressing a diffuse protein (GFP or cherry) and **(C)** α-Adaptin or **(D)** Numb cultured in contact with OP9-DL1 stromal cells, PA or Fc-DL1. Culturing DN3a cells on OP9-DL1 stromal cells and Fc-DL1 increased the proportion of dividing cells with high PR values for α-Adaptin and Numb. The blue shaded region indicates divisions that were assigned as asymmetric when the 0.17 cut-off was applied.

Figure S9. Polarised divisions triggered by the Fc-DL1 functionalised surfaces are restricted to the β-selection stage of T cell development. Polarised divisions were stage specific as indicated by a lack of polarised divisions in **(A)** DN3b cells and **(B)** DN2b cells. The blue shaded region indicates divisions that were classified as asymmetric. **(C)** Violin plot of DN3b and DN2b *versus* DN3a. n = 3 - 4 independent experiments; ***$p < 0.001$

Figure S10. Functionalised beads were used to explore the correlation between the spatial localisation of the polarity cue and spindle orientation. The orientation of mitotic spindle relative to the polarity cue was determined for dividing DN3a cells in contact with OP9-DL1 stromal cells and DL1 functionalised beads. **(A)** For DN3a cells cultured in contact with OP9-DL1 stromal cell the division angle was defined as the angle between the interface between the DN3a cell and OP9-DL1 stromal cell and the long axis of the dividing cell. **(B)** For the functionalised beads, a line was drawn between the centre of mass of the bead and the centre of mass of the dividing cell. The division angle was defined as the angle orthogonal to the angle between this line and the long axis of the dividing cell. PR scatter plots for dividing DN3a cells expressing a diffuse protein (GFP or cherry) and α-Adaptin and cultured in contact with 5 µm beads functionalised with **(C)** PA or **(D)** Fc-DL1 (DL1). Beads functionalised with Fc-DL1, but not PA, triggered ACD, similar to functionalised flat surfaces (compare with Fig. 3). The blue shaded region indicates divisions that were classified as asymmetric. **(D)** Data was plotted in the form of violin plots using the kernel density function. Total number of divisions analysed: DL1 = 109, DL1 bead = 94, PA = 102 and PA bead = 86.

Figure S11. Different polarity cues can promote or inhibit ACD. DN3a cells expressing a diffuse protein (GFP or cherry) and α-Adaptin were cultured onto surfaces pre-coated with PA, Fc-DL1 (DL1) or Fc-DL4 (DL4) and imaged using time-lapse microscopy for 20 hours. The addition of **(A)** DBZ or **(B)** AMD3100, to inhibit Notch or CXCR4 signalling, respectively, partially abolished ACD. **(C)** The inclusion of both inhibitors further reduced the level of ACD. **(D)** Fc-DL4 coated surfaces triggered ACD. **(E)** The inclusion of VCAM-1 lead to a reduction in ACD. Blue shaded regions in PR scatter plots indicates divisions that were assigned as asymmetric using the 0.17 cut-off. n = 2-4 independent experiments.

Figure S12. The addition of Fc-VCAM-1 to Fc-DL4 coated surfaces reduces polarisation during interphase. Sorted DN3a cells were cultured onto surfaces pre-coated with Fc-DL4, Fc-VCAM-1 or Fc-DL4 + Fc-VCAM-1 for 15 hours before fixing and staining with α-tubulin and α-Adaptin. **(A)** Thymocyte attachment was unaffected by the substrate coating, but **(B)** MTOC recruitment and **(C)** α-Adaptin polarisation to the cell-substrate interface was reduced in the presence of Fc-VCAM-1. Total number of divisions analysed: DL4 = 113, VCAM-1 = 59, DL4 + VCAM-1 = 61. DN3a cells were cultured onto surfaces pre-coated with Fc-DL4 or Fc-DL4 + Fc-VCAM-1 for 5 days before harvesting and analysis with flow cytometry. The inclusion of Fc-VCAM-1 had no effect on **(D)** cell number, **(E)** viability or **(F)** differentiation. All data are represented as mean ± SEM; $n = 3$ independent experiments, *$p < 0.05$, **$p < 0.01$.

References

Abe, N., Hozumi, K., Hirano, K.-i., Yagita, H., & Habu, S. (2010). Notch ligands transduce different magnitudes of signaling critical for determination of T-cell fate. *European Journal of Immunology, 40*(9), 2608-2617. doi:10.1002/eji.200940006

Artavanis-Tsakonas, S., Rand, M. D., & Lake, R. J. (1999). Notch signaling: cell fate control and signal integration in development. *Science, 284*(5415), 770-776.

Bhat, K. M. (2014). Notch signaling acts before cell division to promote asymmetric cleavage and cell fate of neural precursor cells. *Science signaling, 7*(348), ra101. doi:10.1126/scisignal.2005317

Bray, S. J. (2016). Notch signalling in context. *Nature Reviews Molecular Cell Biology, 17*(11), 722-735. doi:10.1038/nrm.2016.94

Chang, J. T., Palanivel, V. R., Kinjyo, I., Schambach, F., Intlekofer, A. M., Banerjee, A., Longworth, S. A., Vinup, K. E., Mrass, P., Oliaro, J., Killeen, N., Orange, J. S., Russell, S. M., Weninger, W., & Reiner, S. L. (2007). Asymmetric T lymphocyte division in the initiation of adaptive immune responses. *Science, 315*(5819), 1687-1691. doi:10.1126/science.1139393

Charnley, M., Anderegg, F., Holtackers, R., Textor, M., & Meraldi, P. (2013). Effect of Cell Shape and Dimensionality on Spindle Orientation and Mitotic Timing. *PLoS ONE, 8*(6), e66918. doi:10.1371/journal.pone.0066918

Charnley, M., Fairfull-Smith, K., Haldar, S., Elliott, R., McArthur, S. L., Williams, N. H., & Haycock, J. W. (2009). Generation of bioactive materials with rapid self-assembling resorcinarene-peptides. *Advanced Materials, 21*(28), 2909-2915. doi:10.1002/adma.200802731

Charnley, M., Kroschewski, R., & Textor, M. (2012). The study of polarisation in single cells using model cell membranes. *Integrative Biology, 4*(9), 1059-1071.

Charnley, M., & Russell, S. M. (2017). Imaging Asymmetric T Cell Division. In Cosima T. Baldari & Michael L. Dustin (Eds.), *The Immune Synapse: Methods and Protocols* (pp. 383-398). New York, NY: Springer New York.

Chiaramonte, R., Colombo, M., Bulfamante, G., Falleni, M., Tosi, D., Garavelli, S., De Simone, D., Vigolo, E., Todoerti, K., Neri, A., & Platonova, N. (2015). Notch pathway promotes ovarian cancer growth and migration via CXCR4/SDF1alpha chemokine system. *The international journal of biochemistry & cell biology, 66*, 134-140. doi:10.1016/j.biocel.2015.07.015

Ciofani, M., Schmitt, T. M., Ciofani, A., Michie, A. M., Cuburu, N., Aublin, A., Maryanski, J. L., & Zuniga-Pflucker, J. C. (2004). Obligatory role for cooperative signaling by pre-TCR and Notch during thymocyte differentiation. *Journal of Immunology, 172*(9), 5230-5239.

Ciofani, M., & Zuniga-Pflucker, J. C. (2005). Notch promotes survival of pre-T cells at the beta-selection checkpoint by regulating cellular metabolism. *Nature Immunology, 6*(9), 881-888. doi:10.1038/ni1234

Cong, Z., Wu, H., Guo, Z., Qin, T., Xu, Y., Jing, H., Wang, Y., & Shen, Y. (2017). High expression of C-X-C chemokine receptor 4 and Notch1 is predictive of lymphovascular invasion and poor prognosis in lung adenocarcinoma. *Tumor Biology, 39*(6), 1-10. doi:10.1177/1010428317708698

Coumailleau, F., Furthauer, M., Knoblich, J. A., & Gonzalez-Gaitan, M. (2009). Directional Delta and Notch trafficking in Sara endosomes during asymmetric cell division. *Nature, 458*(7241), 1051-1055. doi:10.1038/nature07854

Couturier, L., Trylinski, M., Mazouni, K., Darnet, L., & Schweisguth, F. (2014). A fluorescent tagging approach in Drosophila reveals late endosomal trafficking of Notch and Sanpodo. *The Journal of Cell Biology, 207*(3), 351-363. doi:10.1083/jcb.201407071

Couturier, L., Vodovar, N., & Schweisguth, F. (2012). Endocytosis by Numb breaks Notch symmetry at cytokinesis. *Nature cell biology, 14*(2), 131-139. doi:10.1038/ncb2419

Daeden, A., & Gonzalez-Gaitan, M. (2018). Endosomal Trafficking During Mitosis and Notch-Dependent Asymmetric Division. *Prog Mol Subcell Biol, 57*, 301-329. doi:10.1007/978-3-319-96704-2_11

Dallas, M. H., Varnum- Finney, B., Delaney, C., Kato, K., & Bernstein, I. D. (2005). Density of the Notch ligand Delta1 determines generation of B and T cell precursors from hematopoietic stem cells. *Journal of Experimental Medicine, 201*(9), 1361-1366. doi:10.1084/jem.20042450

Day, D., Pham, K., Ludford-Menting, M. J., Oliaro, J., Izon, D., Russell, S. M., & Gu, M. (2009). A method for prolonged imaging of motile lymphocytes. *Immunology and Cell Biology, 87*(2), 154-158. doi:10.1038/icb.2008.79

De Kloe, G. E., & De Strooper, B. (2014). Small Molecules That Inhibit Notch Signaling. In Hugo J. Bellen & Shinya Yamamoto (Eds.), *Notch Signaling: Methods and Protocols* (pp. 311-322). New York, NY: Springer New York.

den Elzen, N., Buttery, C. V., Maddugoda, M. P., Ren, G., & Yap, A. S. (2009). Cadherin adhesion receptors orient the mitotic spindle during symmetric cell division in mammalian epithelia. *Molecular Biology of the Cell, 20*(16), 3740-3750. doi:10.1091/mbc.E09-01-0023

Dewey, E. B., Taylor, D. T., & Johnston, C. A. (2015). Cell Fate Decision Making through Oriented Cell Division. *Journal of developmental biology, 3*(4), 129-157. doi:10.3390/jdb3040129

Fortini, M. E. (2009). Notch signaling: the core pathway and its posttranslational regulation. *Developmental Cell, 16*(5), 633-647. doi:10.1016/j.devcel.2009.03.010

Gavard, J., Lambert, M., Grosheva, I., Marthiens, V., Irinopoulou, T., Riou, J. F., Bershadsky, A., & Mege, R. M. (2004). Lamellipodium extension and cadherin adhesion: two cell responses to cadherin activation relying on distinct signalling pathways. *Journal of Cell Science, 117*(2), 257-270. doi:10.1242/jcs.00857

Gehre, N., Nusser, A., von Muenchow, L., Tussiwand, R., Engdahl, C., Capoferri, G., Bosco, N., Ceredig, R., & Rolink, A. G. (2015). A stromal cell free culture system generates mouse pro-T cells that can reconstitute T-cell compartments in vivo. *European Journal of Immunology, 45*(3), 932-942. doi:10.1002/eji.201444681

Grakoui, A., Bromley, S. K., Sumen, C., Davis, M. M., Shaw, A. S., Allen, P. M., & Dustin, M. L. (1999). The immunological synapse: A molecular machine controlling T cell activation. *Science, 285*(5425), 221-227. doi:10.1126/science.285.5425.221

Habib, S. J., Chen, B. C., Tsai, F. C., Anastassiadis, K., Meyer, T., Betzig, E., & Nusse, R. (2013). A localized Wnt signal orients asymmetric stem cell division in vitro. *Science, 339*(6126), 1445-1448. doi:10.1126/science.1231077

Ho, S.-Y., Ling, T.-Y., Lin, H.-Y., Liou, J. T.-J., Liu, F.-C., Chen, I. C., Lee, S.-W., Hsu, Y., Lai, D.-M., & Liou, H.-H. (2017). SDF-1/CXCR4 Signaling Maintains Stemness Signature in Mouse Neural Stem/Progenitor Cells. *Stem cells international, 2017*, 2493752-2493752. doi:10.1155/2017/2493752

Hozumi, K., Mailhos, C., Negishi, N., Hirano, K., Yahata, T., Ando, K., Zuklys, S., Hollander, G. A., Shima, D. T., & Habu, S. (2008). Delta-like 4 is indispensable in thymic environment specific for T cell development. *J Exp Med, 205*(11), 2507-2513. doi:10.1084/jem.20080134

Hozumi, K., Negishi, N., Suzuki, D., Abe, N., Sotomaru, Y., Tamaoki, N., Mailhos, C., Ish-Horowicz, D., Habu, S., & Owen, M. J. (2004). Delta-like 1 is necessary for the generation of marginal zone B cells but not T cells in vivo. *Nature Immunology, 5*, 638. doi:10.1038/ni1075

Janas, M. L., Varano, G., Gudmundsson, K., Noda, M., Nagasawa, T., & Turner, M. (2010). Thymic development beyond beta-selection requires phosphatidylinositol 3-kinase activation by CXCR4. *Journal of Experimental Medicine, 207*(1), 247-261. doi:10.1084/jem.20091430

Koch, U., Fiorini, E., Benedito, R., Besseyrias, V., Schuster-Gossler, K., Pierres, M., Manley, N. R., Duarte, A., Macdonald, H. R., & Radtke, F. (2008). Delta-like 4 is the essential, nonredundant ligand for Notch1 during thymic T cell lineage commitment. *J Exp Med, 205*(11), 2515-2523. doi:10.1084/jem.20080829

Kovacs, E. M., Ali, R. G., McCormack, A. J., & Yap, A. S. (2002). E-cadherin homophilic ligation directly signals through Rac and phosphatidylinositol 3-kinase to regulate adhesive contacts. *Journal of Biological Chemistry, 277*(8), 6708-6718. doi:10.1074/jbc.M109640200

Liu, K., Shen, D., Shen, J., Gao, S. M., Li, B., Wong, C., Feng, W., & Song, Y. (2017). The Super Elongation Complex Drives Neural Stem Cell Fate Commitment. *Developmental Cell, 40*(6), 537-551 e536. doi:10.1016/j.devcel.2017.02.022

Makaraviciute, A., & Ramanaviciene, A. (2013). Site-directed antibody immobilization techniques for immunosensors. *Biosensors and Bioelectronics, 50*, 460-471. doi:10.1016/j.bios.2013.06.060

Manz, B. N., & Groves, J. T. (2010). Spatial organization and signal transduction at intercellular junctions. *Nature Reviews Molecular Cell Biology, 11*(5), 342-352. doi:10.1038/nrm2883

Mingueneau, M., Kreslavsky, T., Gray, D., Heng, T., Cruse, R., Ericson, J., Bendall, S., Spitzer, M. H., Nolan, G. P., Kobayashi, K., von Boehmer, H., Mathis, D., Benoist, C., Immunological Genome, C., Best, A. J., Knell, J., Goldrath, A., Jojic, V., Koller, D., Shay, T., Regev, A., Cohen, N., Brennan, P., Brenner, M., Kim, F., Rao, T. N., Wagers, A., Heng, T., Ericson, J., Rothamel, K., Ortiz-Lopez, A., Mathis, D., Benoist, C., Bezman, N. A., Sun, J. C., Min-Oo, G., Kim, C. C., Lanier, L. L., Miller, J., Brown, B., Merad, M., Gautier, E. L., Jakubzick, C., Randolph, G. J., Monach, P., Blair, D. A., Dustin, M. L., Shinton, S. A., Hardy, R. R., Laidlaw, D., Collins, J., Gazit, R., Rossi, D. J., Malhotra, N., Sylvia, K., Kang, J., Kreslavsky, T., Fletcher, A., Elpek, K., Bellemare-Pelletier, A., Malhotra, D., & Turley, S. (2013). The transcriptional landscape of alphabeta T cell differentiation. *Nat Immunol, 14*(6), 619-632. doi:10.1038/ni.2590

Mohtashami, M., Shah, D. K., Nakase, H., Kianizad, K., Petrie, H. T., & Zuniga-Pflucker, J. C. (2010). Direct Comparison of Dll1-and Dll4-Mediated Notch Activation Levels Shows Differential Lymphomyeloid Lineage Commitment Outcomes. *Journal of Immunology, 185*(2), 867-876. doi:10.4049/jimmunol.1000782

Mossman, K. D., Campi, G., Groves, J. T., & Dustin, M. L. (2005). Altered TCR signaling from geometrically repatterned immunological synapses. *Science, 310*(5751), 1191-1193. doi:10.1126/science.1119238

Oliaro, J., Van Ham, V., Sacirbegovic, F., Pasam, A., Bomzon, Z. e., Pham, K., Ludford-Menting, M. J., Waterhouse, N. J., Bots, M., Hawkins, E. D., Watt, S. V., Cluse, L. A., Clarke, C. J. P., Izon, D. J., Chang, J. T., Thompson, N., Gu, M., Johnstone, R. W., Smyth, M. J., Humbert, P. O., Reiner, S. L., & Russell, S. M. (2010). Asymmetric Cell Division of T Cells upon Antigen Presentation Uses Multiple Conserved Mechanisms. *Journal of Immunology, 185*(1), 367-375. doi:10.4049/jimmunol.0903627

Petrie, H. T., & Zúñiga-Pflücker, J. C. (2007). Zoned Out: Functional Mapping of Stromal Signaling Microenvironments in the Thymus. *Annual Review of Immunology, 25*(1), 649-679. doi:10.1146/annurev.immunol.23.021704.115715

Pham, K., Shimoni, R., Charnley, M., Ludford-Menting, M. J., Hawkins, E. D., Ramsbottom, K., Oliaro, J., Izon, D., Ting, S. B., Reynolds, J., Lythe, G., Molina-Paris, C., Melichar, H., Robey, E., Humbert, P. O., Gu, M., & Russell, S. M. (2015). Asymmetric cell division during T cell development controls downstream fate. *Journal of Cell Biology, 210*(6), 933-950. doi:10.1083/jcb.201502053

Pham, K., Shimoni, R., Ludford-Menting, M. J., Nowell, C. J., Lobachevsky, P., Bomzon, Z., Gu, M., Speed, T. P., McGlade, C. J., & Russell, S. M. (2013). Divergent lymphocyte signalling revealed by a powerful new tool for analysis of time-lapse microscopy. *Immunology and Cell Biology, 91*(1), 70-81. doi:10.1038/icb.2012.49

Pinto-Teixeira, F., & Desplan, C. (2014). Notch activity in neural progenitors coordinates cytokinesis and asymmetric differentiation. *Sci Signal, 7*(348), pe26. doi:10.1126/scisignal.2005980

Rhyu, M. S., Jan, L. Y., & Jan, Y. N. (1994). Asymmetric distribution of numb protein during division of the sensory organ precursor cell confers distinct fates to daughter cells. *Cell, 76*(3), 477-491.

Schmitt, T. M., & Zuniga-Pflucker, J. C. (2002). Induction of T cell development from hematopoietic progenitor cells by delta-like-1 in vitro. *Immunity, 17*(6), 749-756.

Schweisguth, F. (2015). Asymmetric cell division in the Drosophila bristle lineage: from the polarization of sensory organ precursor cells to Notch-mediated binary fate decision. *Wiley Interdisciplinary Reviews. Developmental Biology, 4*(3), 299-309. doi:10.1002/wdev.175

Shimoni, R., Pham, K., Yassin, M., Ludford-Menting, M. J., Gu, M., & Russell, S. M. (2014). Normalized polarization ratios for the analysis of cell polarity. *PLoS ONE, 9*(6), e99885. doi:10.1371/journal.pone.0099885

Short, B. (2014). Red and green traffic signals. *The Journal of Cell Biology, 207*(3), 319. doi:10.1083/jcb.2073if

Shukla, S., Langley, M. A., Singh, J., Edgar, J. M., Mohtashami, M., Zuniga-Pflucker, J. C., & Zandstra, P. W. (2017). Progenitor T-cell differentiation from hematopoietic stem cells using Delta-like-4 and VCAM-1. *Nature Methods, 14*(5), 531-538. doi:10.1038/nmeth.4258

Silvestre, J., Kenis, P. J. A., & Leckband, D. E. (2009). Cadherin and integrin regulation of epithelial cell migration. *Langmuir, 25*(17), 10092-10099. doi:10.1021/la901109e

Singh, J., & Zuniga-Pflucker, J. C. (2018). Producing proT cells to promote immunotherapies. *Int Immunol, 30*(12), 541-550. doi:10.1093/intimm/dxy051

Sjoqvist, M., & Andersson, E. R. (2017). Do as I say, Not(ch) as I do: Lateral control of cell fate. *Developmental Biology*. doi:10.1016/j.ydbio.2017.09.032

Song, H. Y., Zhou, X., Hobley, J., & Su, X. (2012). Comparative study of random and oriented antibody immobilization as measured by dual polarization interferometry and surface plasmon resonance spectroscopy. *Langmuir, 28*(1), 997-1004. doi:10.1021/la202734f

Suzuki, T., Yokoyama, Y., Kumano, K., Takanashi, M., Kozuma, S., Takato, T., Nakahata, T., Nishikawa, M., Sakano, S., Kurokawa, M., Ogawa, S., & Chiba, S. (2006). Highly efficient ex vivo expansion of human hematopoietic stem cells using Delta1-Fc chimeric protein. *Stem Cells, 24*(11), 2456-2465. doi:10.1634/stemcells.2006-0258

Swamy, M., Pathak, S., Grzes, K. M., Damerow, S., Sinclair, L. V., van Aalten, D. M., & Cantrell, D. A. (2016). Glucose and glutamine fuel protein O-GlcNAcylation to control T cell self-renewal and malignancy. *Nat Immunol*. doi:10.1038/ni.3439

Tajbakhsh, S., Rocheteau, P., & Le Roux, I. (2009). Asymmetric cell divisions and asymmetric cell fates. *Annual review of cell and developmental biology, 25*, 671-699. doi:10.1146/annurev.cellbio.24.110707.175415

Toda, H., Yamamoto, M., Kohara, H., & Tabata, Y. (2011). Orientation-regulated immobilization of Jagged1 on glass substrates for ex vivo proliferation of a bone marrow cell population containing hematopoietic stem cells. *Biomaterials, 32*(29), 6920-6928. doi:10.1016/j.biomaterials.2011.05.093

Toyoshima, F., & Nishida, E. (2007). Integrin-mediated adhesion orients the spindle parallel to the substratum in an EB1-and myosin X-dependent manner. *EMBO Journal, 26*(6), 1487-1498. doi:10.1038/sj.emboj.7601599

Trampont, P. C., Tosello-Trampont, A.-C., Shen, Y., Duley, A. K., Sutherland, A. E., Bender, T. P., Littman, D. R., & Ravichandran, K. S. (2010). CXCR4 acts as a costimulator during thymic beta-selection. *Nature Immunology, 11*(2), 162-U186. doi:10.1038/ni.1830

Trylinski, M., Mazouni, K., & Schweisguth, F. (2017). Intra-lineage Fate Decisions Involve Activation of Notch Receptors Basal to the Midbody in Drosophila Sensory Organ Precursor Cells. *Current Biology, 27*(15), 2239-2247 e2233. doi:10.1016/j.cub.2017.06.030

Tussiwand, R., Engdahl, C., Gehre, N., Bosco, N., Ceredig, R., & Rolink, A. G. (2011). The preTCR-dependent DN3 to DP transition requires Notch signaling, is improved by CXCL12 signaling and is inhibited by IL-7 signaling. *European Journal of Immunology, 41*(11), 3371-3380. doi:10.1002/eji.201141824

Van de Walle, I., Waegemans, E., De Medts, J., De Smet, G., De Smedt, M., Snauwaert, S., Vandekerckhove, B., Kerre, T., Leclercq, G., Plum, J., Gridley, T., Wang, T., Koch, U., Radtke, F., & Taghon, T. (2013). Specific Notch receptor–ligand interactions control human TCR-αβ/γδ development by inducing differential Notch signal strength. *The Journal of Experimental Medicine, 210*(4), 683-697. doi:10.1084/jem.20121798

van Es, J. H., van Gijn, M. E., Riccio, O., van den Born, M., Vooijs, M., Begthel, H., Cozijnsen, M., Robine, S., Winton, D. J., Radtke, F., & Clevers, H. (2005). Notch/gamma-secretase inhibition turns

proliferative cells in intestinal crypts and adenomas into goblet cells. *Nature, 435*(7044), 959-963. doi:10.1038/nature03659

Venkei, Z. G., & Yamashita, Y. M. (2018). Emerging mechanisms of asymmetric stem cell division. *Journal of Cell Biology, 217*(11), 3785-3795. doi:10.1083/jcb.201807037

Wang, Y. C., Hu, X. B., He, F., Feng, F., Wang, L., Li, W., Zhang, P., Li, D., Jia, Z. S., Liang, Y. M., & Han, H. (2009). Lipopolysaccharide-induced maturation of bone marrow-derived dendritic cells is regulated by notch signaling through the up-regulation of CXCR4. *Journal of Biological Chemistry, 284*(23), 15993-16003. doi:10.1074/jbc.M901144200

Xiao, W., Gao, Z., Duan, Y., Yuan, W., & Ke, Y. (2017). Notch signaling plays a crucial role in cancer stem-like cells maintaining stemness and mediating chemotaxis in renal cell carcinoma. *Journal of Experimental & Clinical Cancer Research, 36*(1), 41. doi:10.1186/s13046-017-0507-3

Yashiro-Ohtani, Y., He, Y., Ohtani, T., Jones, M. E., Shestova, O., Xu, L., Fang, T. C., Chiang, M. Y., Intlekofer, A. M., Blacklow, S. C., Zhuang, Y., & Pear, W. S. (2009). Pre-TCR signaling inactivates Notch1 transcription by antagonizing E2A. *Genes Dev, 23*(14), 1665-1676. doi:10.1101/gad.1793709

Young, I. T. (1977). Proof without prejudice: use of the Kolmogorov-Smirnov test for the analysis of histograms from flow systems and other sources. *Journal of Histochemistry & Cytochemistry, 25*(7), 935-941. doi:10.1177/25.7.894009

Yui, M. A., & Rothenberg, E. V. (2014). Developmental gene networks: a triathlon on the course to T cell identity. *Nature Reviews Immunology, 14*, 529. doi:10.1038/nri3702

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Notch controls polarity
during cell division

Notch, Notch regulators,
and CXCR4 are
asymmetrically inherited

Notch ligand
CXCL12
Notch1
CXCR4
Numb
α-Adaptin

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

DN3a

OP9-DL1

Not polarised

Polarised

Supplementary Figure 4

A)

B)

D)

C)

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9

Supplementary Figure 10

A)

Division angle

DN3a

OP9-DL1

B)

X°

Division angle = 90 - x

C) PA

PR FP-α-Adaptin

p = 0.03
n = 94

PR FP-diffuse

D) DL1

PR FP-α-Adaptin

p < 0.001
n = 100

PR FP-diffuse

E)

PR FP-α-Adaptin

***

***

DL1   DL1 Bead   PA   PA Bead

Supplementary Figure 11

Supplementary Figure 12