Longevity of growth-arrested bacteria requires protein synthesis and a stringent response

Liang Yin[a], Hongyu Ma[b], Ernesto S. Nakayasu[c], Samuel H. Payne[d], David R. Morris[e] and Caroline S. Harwood[a†]

[a] Department of Microbiology, University of Washington, Seattle, USA.

[b] Department of Environmental Science and Engineering, Xi'an Jiaotong University, Xi'an, P.R. China

[c] Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA, USA

[d] Department of Biology, Brigham Young University, Provo, UT, USA

[e] Department of Biochemistry, University of Washington, Seattle, USA.

Keywords: *Rhodopseudomonas palustris*, ribosome, growth-arrest, translation, longevity, stringent response

Address correspondence to Caroline S. Harwood, csh5@uw.edu

Abstract

Gram-negative bacteria in infections, biofilms and industrial settings often stop growing due to nutrient depletion, immune responses or environmental stresses. Bacteria in this state tend to be tolerant to antibiotics and are often referred to as dormant. Here we describe the physiology of *Rhodopseudomonas palustris*, a phototrophic alpha-proteobacterium, in its growth arrested state. *R. palustris* can maintain almost full viability for over four months and we found that ATP synthesis, proteins synthesis and a stringent response were required for this remarkable longevity. *R. palustris* is an extreme example of a bacterial species that will stay alive for long periods of time as a relatively homogeneous populations of cells. There is evidence that other gram-negative species also continue to synthesis proteins during growth arrest and that a stringent response is required for their longevity as well. Our observations challenge that notion that growth-arrested cells are dormant and expands our mechanistic understanding of a crucial but understudied phase of the bacterial life cycle.

Introduction

Bacteria in nature often exist in viable but non-growing states without forming differentiated structures like spores (*1, 2*). Yet this crucial phase of the bacterial life cycle is underexplored. Factors that limit cell replication include nutrient limitation, competition within microbial communities and stresses like oxygen-depletion and extreme temperatures (*3-5*). Non-growing pathogenic bacteria present in human infections present challenges for treatment because antibiotics tend to target processes such as cell-wall biosynthesis or DNA replication that are active in growing, but not non-growing cells. One example of a bacterium that is recalcitrant to antibiotic treatment *in vivo* is *Mycobacterium tuberculosis*, which forms latent infections that persist over many years in the absence of overt signs of growth (*6*). Then there are bacterial

antibiotic persisters, a small subpopulation of non-growing or slowly growing cells that develop

in many bacterial infections and are tolerant to antibiotic treatment (*7-9*).

Studying non-growing bacteria is also important in environmental, industrial and other non-

medical contexts. For example, non-growing bacteria are excellent biocatalysts because they

can convert substrates that might be used for growth to value-added products. In past work we

have studied production of hydrogen gas, a biofuel, by the phototrophic α-proteobacterium

Rhodopseudomonas palustris. When illuminated to allow ATP synthesis by

photophosphorylation, non-growing cells diverted cell resources to hydrogen gas production and

generated 300% more hydrogen gas than growing cells (*10*). We also found that non-growing *R.*

palustris cells embedded in thin strips of latex film remained viable and produced hydrogen gas

for over four months (*11*).

These results prompted us to develop *R. palustris* as a model for studies of bacterial longevity,

and to ask the question: are there universal features of long-lived bacteria that we can uncover

from studies of this proteobacterium? Several attributes make *R. palustris* a good model for

addressing this question. First, it generates ATP and protonmotive force from light, even when it

is not growing (Figure 1). Also, *R. palustris* carries out photophosphorylation under anaerobic

conditions and does not generate oxygen, thus avoiding complications of oxidative stress during

growth-arrest. A Tn-seq study of growth-arrested *R. palustris* identified 116 genes, which we

call longevity genes, that are essential for long-term viability of non-growing cells, but are not

required for growth (*12*). Among these are genes that are conserved in bacteria and associated

with basic cellular processes including protein synthesis. Here, we carried out a physiological

characterization of *R. palustris* in its growth-arrested, but viable state. We found that active

translation and optimized ribosomes were critical for the longevity of *R. palustris,* as was

(p)ppGpp. This goes against the conventional description of non-growth as a dormant and

inactive state. We suggest that protein synthesis and other cellular activities are likely important

for many bacteria to maintain viability when they are not growing.

Results

<u>**Growth-arrested *R. palustris* cells do not appear to undergo cycles of death and**</u>

<u>**regrowth.**</u> A major barrier to studies of bacterial longevity in the laboratory is that most

heterotrophic bacteria, the best studied of which is *E. coli*, undergo cycles of death and growth

after entry into stationary phase (*1, 13*). Typically a fraction of cells survive long-term stationary

phase by feeding on nutrients released from dead cells. This leads to physiological and genetic

diversity in the population and there is strong selection pressure for mutations that enable

cannibalism (*14, 15*). In previous work we reported that populations of growth-arrested *R.*

palustris cells were nearly 100% viable and morphologically homogenous (*12*). To firmly

establish that cells were truly growth-arrested, rather than undergoing cycles of growth and

death, we introduced an unstable plasmid expressing a gentamycin (Gm) resistance gene into

wild-type *R. palustris*. When Gm is absent, the percentage of growing cells that retain this

plasmid would be expected to decrease (*16*). On the other hand, a non-growing culture would

be expected to retain the plasmid in Gm-free medium. When this strain (WT::pBBR[Gm]) was

inoculated into Gm-free medium, about 60% of the cells had lost the plasmid by the time they

reached stationary phase. When these cells were transferred into fresh Gm-free medium and

successively transferred several times over a period of 20 days after reaching stationary phase,

the plasmid continued to be lost. In contrast, growth-arrested cells retained the plasmid over the

same period of time (Figure 2a), suggesting that most or all of the cells were indeed in a non-

growing state.

In further support of the conclusion that growth-arrested *R. palustris* does not undergo cycles of

growth and death, we found that non-growing cells were tolerant to the cell wall synthesis

inhibitor ampicillin and the DNA synthesis inhibitor nalidixic acid at concentrations that

prevented growth (Figure 2b). However, we also found that the viabilities of non-growing cells

were compromised by protein synthesis inhibitors (Figure 2b and SI Figure 1). In all these cases

and throughout this study, growth arrest was imposed by providing cells a growth-limiting

amount of the carbon source acetate.

Long-lived, growth-arrested cells continue to synthesize proteins. Growth-arrested *R.*

palustris was sensitive to Gm, an aminoglycoside antibiotic that affects protein synthesis by

causing mistranslation, which results in accumulation of damaged proteins. We found that over

99% of growth-arrested cells lost viability over a few days when treated with Gm at a final

concentration of 50 µg/ml (Figure 2b). Although cells died relatively rapidly, they were not as

sensitive to Gm as were growing cells, which died within a few hours after exposure to this

antibiotic. Chloramphenicol (Cm) belongs to another class of antibiotic that targets ribosomes.

Although typically bacteriostatic, Cm was effective in killing growth-arrested *R. palustris* over a

prolonged period of time (Figure 2b and SI Figure 1). These results suggest that active

translation is important for *R. palustris* longevity. In support of the conclusion that growth-

arrested cells continue to synthesize proteins, proteomics experiments revealed that after 6

days and 20 days of growth-arrest, 289 and 525 proteins, respectively, were present at greater

than two-fold higher levels than in growing cells (Figure 3a and SI Table 1). Among these were

enzymes involved in the degradation of compounds, including benzoate and fatty acids (SI

Table 1 and 2), that are preferred carbon sources for *R. palustris*, suggesting that cells respond

to growth-arrest by increasing the translation of proteins that may aid in scavenging carbon. In

the period between 6 and 20 days post growth-arrest, 51 proteins of unknown function

increased in abundance. Some proteins (163 and 310 proteins at days 6 and 20 post growth-

arrest) were diminished in abundance by two-fold or more following growth-arrest (Figure 3a

and SI Table 1). This set included 53 out of 55 ribosomal proteins that were identified in our

proteomics dataset. On average, ribosome proteins were at 25-50% of their original levels after

6 days and 20 days of growth-arrest (Figure 3b). However, not all translation related proteins

were decreased in their abundances. For example, translation elongation factors made up ~5%

of the total protein from growing cells as well as growth-arrested cells. Similar patterns were

also observed for proteins involved in aminoacyl-tRNA biosynthesis (tRNA charging) and for two

previously identified longevity proteins YbeY and Era (Figure 3c).

Characterization of protein synthesis apparatus in long-lived cells. We isolated *R. palustris*

ribosomes by sucrose gradient centrifugation and found that the ribosome profile of growing *R.*

palustris cells included 30S, 50S and 70S populations, as is typical for growing bacteria (Figure

4a). There was also a small peak corresponding to a 100S population, which is likely a dormant

form of ribosomes, as has been seen in other bacteria (*17*). The ribosome profile of growth-

arrested *R. palustris* cells resembled that of growing cells (Figure 4a and 4b), but the quantity of

the 70S population was lower by day 25 post growth-arrest. Ribosome degradation is often

accompanied by the non-specific degradation of rRNA (*18-21*). In growing *R. palustris*, rRNA

made up about 70% of the total RNA (Figure 4c). After growth-arrest, rRNA accounted for about

50% of the total RNA, and it stayed at about this level for 120 d (Figure 4c). The ratio between

23S and 16S rRNA remained stable during this time (SI Figure 2), and little degradation was

observed (Figure 4c). The relative levels of aminoacylated (or charged) and free forms of tRNAs

are important for promoting and inhibiting translation (*22*). We used Northern blot analysis to

estimate the percentage of tryptophan-charged $tRNA_{trp}$ (Trp-$tRNA_{trp}$) relative to $tRNA_{trp}$. In

growing cells, ~60% of $tRNA_{trp}$ was in the charged Trp-$tRNA_{trp}$ form (Figure 4d and 4e). This

level is comparable to that reported in other bacteria (*23*). The percentage of Trp-$tRNA_{trp}$

increased to ~85% upon growth-arrest and was maintained at this level for ~20 days (Figure 4d

and 4e).

Two proteins identified as essential for *R. palustris* longevity are critical for ribosome

integrity in growth-arrested cells. Among the longevity genes that we have identified in *R.*

palustris (*12*), *ybeY* and *era* are predicted to encode proteins that interact with each other and

the S11 ribosome protein during ribosome maturation (*24-27*). *ybeY* or *era* deletion mutants

have compromised longevity but grow normally (*12*). Here we found that the ribosome profiles

of growing cells were similar between the wild-type and these mutants (Figure 5a). All had

distinctive 30S, 50S and 70S populations, although the 50S subunit was slightly more abundant

in the mutants. The ribosome profiles of growth-arrested *ΔybeY* and *Δera* mutant cells were

noticeably different from that of the wild-type, however. A prominent 50S subunit peak was seen

but the amounts of 30S subunits and 70S ribosomes were greatly diminished (Figure 5a).

As predicted based on studies of their *E. coli* homologs (*24-27*), *ΔybeY* and *Δera* strains were

defective in 16S rRNA maturation. An extra band immediately above the 16S rRNA was present

in gels of RNA from both mutants (SI Figure 2). Northern blot experiments confirmed that the

extra band was 17S rRNA, a precursor of 16S rRNA (Figure 5b). In growth-arrested cells, both

16S rRNA and 17S rRNA species were severely depleted in the mutated strains (Figure 5b, SI

Figure 2). This likely led to the defective ribosome production shown in Figure 5a, and the

compromised longevity of *R. palustris*. *ybeY* or *era* deletion mutants had wild-type levels of

charged tRNA-Trp (SI Figure 3), suggesting that tRNA aminoacylation is not directly coordinated

with the process of ribosome maturation.

As shown in Figure 3b, there was no significant change in the abundance of YbeY and Era

proteins in growth-arrested cells. Since the core ribosomal proteins were decreased 2 to 4-fold

in abundance (Figure 3b), the relative abundance of YbeY and Era was increased on a per

ribosome basis following growth-arrest. To test whether we could change the effects of YbeY on

longevity by modulating its relative abundance, we constructed and expressed a YbeY variant

predicted based on work in *E. coli*, to interact weakly with ribosomes. In *E. coli*, the interaction

between YbeY and ribosomes is partially disrupted by replacement of a conserved aspartate

with an arginine (*28*). When we made the corresponding YbeY$_{D118R}$ change in *R. palustris*, the

strain was compromised in its longevity to a similar degree as the *ΔybeY* mutant (Figure 6a).

When we overexpressed YbeY$_{D118R}$ *in trans* (*ΔybeY*::pBBP-*ybeY*$_{D118R}$) however, the longevity

defect was complemented (Figure 6a). The quantity of 16S rRNA was similar to that of the wild-

type, and no 17S rRNA was observed during growth-arrest (Figure 6b). Also, the ribosome

profile of the growth-arrested *ΔybeY*::pBBP-*ybeY*$_{D118R}$ complemented strain was comparable to

that of growth-arrested wild-type cells (Figure 6c).

The stringent response is important for longevity. Our results suggest that translation is

reduced but still maintained in growth-arrested cells. A well-known mechanism that contributes

to this is the stringent response mediated by guanosine polyphosphate [(p)ppGpp] (*29, 30*). In

many bacteria, the level of (p)ppGpp increases upon growth-arrest and this inhibits the

transcription of ribosomal genes, although the mechanism by which this is achieved varies

among bacteria (*29-31*). As with other bacteria, we detected little to no (p)ppGpp in

exponentially growing *R. palustris*. However, the level of (p)ppGpp increased sharply as growing

cells reached stationary phase and remained at relatively high levels after growth-arrest (Figure

7a).

Unlike *E. coli*, but like most other bacteria outside the γ- and β-proteobacteria, *R. palustris* has a

single bifunctional (p)ppGpp synthesis/hydrolysis *rsh* gene, here called *rsh*$_{rp}$, that has multiple

domains including a hydrolase and RelA_SpoT domain (Figure 7b). Our Tn-seq results

indicated that this gene (*rpa2693*) was essential for *R. palustris* longevity, but not for growth (*12,*

32, 33). However, we were unable to delete the full-length *rsh*$_{rp}$ gene. A subsequent inspection

of the locations of the Tn-insertions in this gene in our bank of Tn mutagenized cells revealed

that there were relatively few insertions in its RelA_SpoT domain (Figure 7b). When we made

domain specific deletions in Rsh$_{rp}$, we obtained a hydrolase domain deletion mutant (ΔRsh$_{rp64\text{-}192}$, Figure 7b) that produced very little (p)ppGpp during growth or growth-arrest. (Figure 7c).

This mutant grew with a generation time of 9.4 ± 0.6 h, compared to a wild-type 6.4 ± 0.2 h

generation time, a growth rate difference of about 30%. In contrast to this relatively modest

effect on the growth rate the ΔRsh$_{rp60\text{-}189}$ mutation had a severe effect on longevity (Figure 7d).

Discussion

We found that illuminated *R. palustris* responds to growth-arrest by reducing transcription of

ribosomal genes and making fewer ribosomes as is typical of most bacteria when they slow

their growth rate. Growth-arrested cells continued to synthesize proteins, some at elevated

levels relative to growing cells, over the period of 20 days that we followed protein synthesis by

proteomics (SI Table 1). We saw no evidence of significant rRNA degradation even 120 d post-

growth arrest (Figure 4c). The levels of proteins involved in tRNA charging and translation

elongation and were at similar levels in growing and growth-arrested cells (Figure 3c), which is

consistent with the notion from a previous report that elongation rates are maintained as

bacteria slow down their growth-rates even towards zero growth (*34*). In addition, a tRNA that

we followed was almost fully charged 20 d into growth-arrest (Figure 4d and 4e). Not only does

protein synthesis continue after growth-arrest, but it is essential for viability of non-growing cells

as indicated by their sensitivity to Gm and Cm (Figure 2b, SI Figure 1).

There are reports of continued protein synthesis in other species of bacteria in growth-arrest,

the more convincing of which come from studies in which production of fluorescent proteins in

single cells was used as a proxy for protein synthesis. Antibiotic persisters in early stages of

formation, as well as non-growing *E. coli* and *M. tuberculosis* have all been inferred to carry out

protein synthesis (*35-37*). These results suggest that maintenance of some level of protein

synthesis may be required for viability of many non-growing bacteria. From this it follows that it

would make sense to prioritize development of antibiotics that target protein synthesis in growth-

arrested cells. This will likely require understanding in detail what happens to bacterial

ribosomes following growth-arrest. Bacterial ribosomes are known to be differentially post-

translationally modified in stationary phase cells (*38*). Two longevity genes, *ybeY* and *era*, were

essential for normal ribosomes to form in non-growing *R. palustris* (Figure 3b). Era is a well-

studied bacterial GTPase (*27*), and genes for it and YbeY, are present and conserved in most

bacteria (*25*). Our data are consistent with studies of the corresponding *E coli* proteins showing

that YbeY and Era are involved in rRNA processing in *R. palustris*. Drawing on results from *E.*

coli (*28*) it is likely that the *R. palustris* proteins interact with each other as well as with ribosome

protein S11. When we looked at rRNAs formed by Δ*ybeY* and Δ*era* mutants that were actively

growing, it was apparent that growing cells and growth-arrested cells had the same defect in

17S rRNA processing (Figure 5b), but only the growth-arrested cells had a phenotype. This

highlights a feature of at least some *R. palustris* longevity proteins, which is that they are

functional in growing cells as well as in non-growing cells, but it is only when the additional

demands of growth-arrest on cellular functionality are imposed that essentiality becomes

apparent. Our work on Era and YbeY has led us to conclude that optimized ribosomes are a

feature of long-lived cells.

In early stationary phase and when in growth arrest, *R. palustris* synthesizes increasing

amounts of (p)ppGpp, indicating that cells mount a stringent response. The gene *rpa2693*,

which we call rsh_{rp}, is among the genes identified by Tn-seq to be essential for longevity (*12*). A

Δ$rsh_{rp60-189}$ mutant did not produce detectable (p)ppGpp and the viability of this mutant

decreased rapidly after growth-arrest (Figure 7c). We do not know why this deletion, which is in

the hydrolase domain of Rsh_{rp}, affected (p)ppGpp synthesis, but one idea is that it destabilized

or altered the structure of the protein so that its ability to synthesize (p)ppGpp was

compromised. In *R. palustris* the signal for (p)ppGpp synthesis appears to be nutrient limitation

or slowing growth rate. A future area of investigation is to understand why (p)ppGpp is essential

for *R. palustris* longevity and to identify genes and proteins whose activities may be modulated

by (p)ppGpp. (p)ppGpp has recently been shown to bind to Era from *Staphylococcus aureus*

and *E. coli* to inhibit its GTPase activity (*39-41*), and this may influence ribosome biogenesis

and integrity. There is good evidence that (p)ppGpp is important for development of antibiotic

persisters (*42*) and for long term survival of latent and growth-arrested *M. tuberculosis* (*43, 44*).

R. palustris is an extreme example of a bacterial species for which an entire population of non-

growing cells maintains viability for very long periods of time. It is likely that this is primarily due

to its ability to generate ATP by photophosphorylation when it is in growth arrest, as growth-

arrested cells incubated in dark do not survive well*(12)*. This feature of *R. palustris* makes it a

good model for understanding molecular mechanisms of longevity that can be extended to other

bacteria that remain alive in growth arrest, but only as heterogeneous populations. One

hypothesis is that varying ATP levels per cell underlie this heterogeneity. In addition to a

universal requirement for ATP to support viability of growth-arrested bacteria, we can say that

protein synthesis, optimized ribosomes and (p)ppGpp likely lie at the nexus of diverse strategies

used by non-growing bacteria to survive and even thrive.

Methods

Bacterial strains, growth and incubation conditions. *R. palustris* strain CGA009 was used

as the wild-type for this study. Anaerobic cultures were grown under illumination at 30℃, using

a defined media PM with 20 mM sodium acetate or 10mM sodium succinate as the carbon

source (*45*). The longevity of *R. palustris* cultures was determined by counting the colony-

forming units as described previously (*12*). In-frame deletions of *R. palustris* genes were created

by using suicide vector pJQ200SK as described before (*46*). For *in trans* expression in *R.*

palustris, the target genes were cloned into plasmid pBBPgdh and the plasmids were mobilized

into *R. palustris* by conjugation using *E. coli* strain S17-1 (*47*). Media were supplemented with

100 µg/ml of gentamicin for *in trans* complementation experiments.

Ribosome purification. Ribosomes from *R. palustris* cells were isolated following a method

adapted from a previous study (*48*). About 35 ml of culture was grown to the desired state and

poured over a 10-ml ice sheet (fast-cool). The ice sheet melted within 10 min and the cells were

collected by centrifugation at 4000 rpm for 15 min. The cell pellet was resuspended in 2 ml of

Buffer A (20mM HEPES-KOH, pH 7.5, 30mM NaCl, 8mM $MgCl_2$) with 1 µl/ml RNasin

(Promega). Three hundred microliter of 0.1 mm Zirconia/Silica beads (Biospec Products) were

added to every ml of sample. The samples were lysed with a Mini-Beadbeater-24 (Biospec

Products) at 3500 rpm for 1 min at 4℃, kept on ice for 1 min and the cycle was repeated 4

more times. The micro-beads and cell debris were removed by centrifugation at 15000 rpm for

10 min. The lysates were further clarified by spinning at 30,000 g in Beckman TL-1000

ultracentrifuge for 30 min, followed by pelleting ribosomes at 100,000 g for 4 hours. The pellet

was resuspended with Buffer A overnight on ice.

Sucrose gradient analysis. About five A_{260} units of ribosomes (less than 200 µl in volume)

were applied on top of 11 ml of a sucrose gradient (7-47%). The samples were centrifuged

through the gradient at 39,000 rpm for 4.5 h with a SW41 rotor, and then were separated with a

Brandel gradient pump. The sucrose gradient was made with Buffer A.

Ribosomal RNA analysis. Total RNA was purified from 3-5 ml of *R. palustris* culture using an

miRNeasy Mini Kit. The rRNAs were then analyzed with an RNA ScreenTape system (Agilent).

To probe 16S rRNA by Northern blot, 1 µg of total RNA was heated at 70℃ for 10 min and

immediately chilled on ice. The samples were then separated on a 1% agarose gel with 1X TBE

buffer. After electrophoresis, RNA was transferred to an Amersham Hybond-XL membrane (GE

Lifesciences) with a downward nucleic acid transfer system overnight using 20X SSC buffer. A

probe (5'-TCGTGCCTCAGCGTCAGTAATGGCCCAGTGA) labeled with DIG was used to

detect 16S rRNA following the manufacturer's instruction (DIG, Roche).

Charging analysis of tRNA$_{Trp}$. Total RNA was isolated with QIAzol lysis reagent according to

the manufacturer's instructions (Qiagen). The dried RNA pellet was dissolved and stored in 10

mM sodium acetate, pH 4.5. A deacylated control sample was prepared by incubating 10 µg of

total RNA with 1 mM EDTA and 100 mM Tris-HCl pH 9.0 at 42℃ for an hour (*49*). Northern blot

and acid-urea gel electrophoresis were performed essentially as described previously (*23, 49*).

Loading dye (0.01% xylene cyanol, 0.01% bromophenol blue, 8 M Urea, 1 mM EDTA and 10

mM sodium acetate, pH 5.0) was mixed with sample at 1:1 ratio. Samples were loaded on a

12% acid-urea gel of 20 cm and then were run at 15 W for 12 h at 4℃. Transfer of RNA to an

Amersham Hybond-XL membrane (GE Lifesciences) was performed in 0.5* TBE buffer at 650

mA for 30 min 4℃. The aminoacylated and the deacylated tRNA$_{Trp}$ were detected using a DIG-

labeled probe (5'-GGTTTTGGAGACCGGTGCTCTACCAATTGAG) following the manufacturer's

instruction (DIG, Roche).

Determination of cellular (p)ppGpp levels. *In vivo* (p)ppGpp labeling and detection were

performed as described with some modifications (*50*). Anaerobic *R. palustris* cultures were

grown in 1.5 ml PM medium with 100 µCi/ml of Na$_2$H^{32}PO$_4$ (PerkinElmer-NEX011001MC) added

at the time of inoculation. At the desired time points, 200-300 µl of culture was collected and

extracted with 8 M formic acid. Supernatants of the extracts were spotted on PEI-cellulose TLC

plates (Sigma-Aldrich) and separated with 1.5 M KH$_2$PO$_4$ (pH 3.4). The plates were then dried

and visualized by phosphor imaging. The level of (p)ppGpp was calculated by the ratio of

(ppGpp + ppGpp) / (GTP + ppGpp + pppGpp).

<u>Proteomic analysis.</u> Cells were lysed, digested with trypsin and analyzed by liquid

chromatography-tandem mass spectrometry as described in (*51*). Collected data were analyzed

using   MaxQuant software (v 1.5.5.1) (*32, 52*) for peptide/protein identification and

quantification. Identification of peptides was done by searching against the *R. palustris* ATCC

BAA-98 (strain CGA009) sequences from the Uniprot Knowledgebase (downloaded on August

30, 2016). The search was performed first with a larger parent mass tolerance (20 ppm) to

identify peptides and re-calibrate peptide masses. After re-calibration, the parent mass

tolerance was set to 4.5 ppm for a second round of search. During the searches oxidation of

methionine as a variable modification, and trypsin digestion was required in both peptide

termini. Identifications were filtered with 1% false-discovery rate at peptide-spectrum match and

protein levels. The function "matching between runs" was checked to detect peptide peaks in

LC-MS/MS runs that they were not fragmented. Label-free quantification (LFQ) values were

used to quantitative analysis. Proteins were considered differentially abundant with a p-value

≤0.05 by T.test considering two tails and equal variance.

Funding information

This work was supported by grant W911NF-15-1-0150 to CSH from the US Army Research

Office. SHP and ESN were supported by the U.S. Department of Energy, Office of Biological

and Environmental Research, Early Career Research Program. Mass spectrometry data was

generated in the Environmental Molecular Science Laboratory, a U.S. Department of Energy

(DOE) national scientific user facility at Pacific Northwest National Laboratory (PNNL) in

Richland, WA. Battelle operates PNNL for the DOE under contract DE-AC05-76RLO01830.

Acknowledgements

The authors thank Amy Schaefer, Kathryn Fixen, Yasuhiro Oda and Yanning Zheng for helpful

discussions, and Karl Weitz for technical assistance with the mass spectrometry analysis.

Abbreviations

Tn: transposon.

Author contribution

L.Y. and C.S.H. designed the research. L.Y. and H.M. performed the antibiotics assays with *R.*

palustris. L.Y. made the *R. palustris* mutants. L.Y. grew the *R. palustris* cells. S.H.P. and E.S.N.

performed proteomic analysis. L.Y. and D.R.M. carried out the *in vitro* analysis of ribosome. L.Y.

carried out the analysis of rRNA and tRNA. L.Y. performed the analysis of (p)ppGpp. All authors

analyzed the data. L.Y., E.S.N. and C.S.H. wrote the paper. All authors contributed to the

revision of the manuscript.

Competing interests

The authors declare no competing financial interests.

References

1.   Finkel, S. E. (2006) Long-term survival during stationary phase: evolution and the GASP
phenotype, *Nature reviews. Microbiology 4*, 113-120.
2.   Bergkessel, M., Basta, D. W., and Newman, D. K. (2016) The physiology of growth arrest:
uniting molecular and environmental microbiology, *Nature reviews. Microbiology 14*,
549-562.
3.   Wayne, L. G., and Sohaskey, C. D. (2001) Nonreplicating persistence of mycobacterium
tuberculosis, *Annual Review of Microbiology 55*, 139-163.
4.   Thieringer, H. A., Jones, P. G., and Inouye, M. (1998) Cold shock and adaptation,
BioEssays : news and reviews in molecular, cellular and developmental biology 20, 49-57.
5.   Hibbing, M. E., Fuqua, C., Parsek, M. R., and Peterson, S. B. (2010) Bacterial competition:
surviving and thriving in the microbial jungle, *Nature reviews. Microbiology 8*, 15-25.
6.   Ai, J. W., Ruan, Q. L., Liu, Q. H., and Zhang, W. H. (2016) Updates on the risk factors for
latent tuberculosis reactivation and their managements, *Emerg Microbes Infect 5*, e10.
7.   Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L., and Leibler, S. (2004) Bacterial
persistence as a phenotypic switch, *Science 305*, 1622-1625.

8.   Lewis, K. (2010) Persister cells, *Annual Review of Microbiology 64*, 357-372.

9.   Balaban, N. Q., Helaine, S., Lewis, K., Ackermann, M., Aldridge, B., Andersson, D. I.,
Brynildsen, M. P., Bumann, D., Camilli, A., Collins, J. J., Dehio, C., Fortune, S., Ghigo, J.-
M., Hardt, W.-D., Harms, A., Heinemann, M., Hung, D. T., Jenal, U., Levin, B. R., Michiels,
J., Storz, G., Tan, M.-W., Tenson, T., Van Melderen, L., and Zinkernagel, A. (2019)
Definitions and guidelines for research on antibiotic persistence, *Nature Reviews*
Microbiology.

10.   McKinlay, J. B., Oda, Y., Ruhl, M., Posto, A. L., Sauer, U., and Harwood, C. S. (2014) Non-
growing *Rhodopseudomonas palustris* increases the hydrogen gas yield from acetate by
shifting from the glyoxylate shunt to the tricarboxylic acid cycle, *The Journal of biological*
chemistry 289, 1960-1970.

11.   Gosse, J. L., Engel, B. J., Hui, J. C., Harwood, C. S., and Flickinger, M. C. (2010) Progress
toward a biomimetic leaf: 4,000 h of hydrogen production by coating-stabilized
nongrowing photosynthetic Rhodopseudomonas palustris, *Biotechnol Prog 26*, 907-918.

12.   Pechter, K. B., Yin, L., Oda, Y., Gallagher, L., Yang, J., Manoil, C., and Harwood, C. S.
(2017) Molecular Basis of Bacterial Longevity, *MBio 8*.

13.   Zambrano, M. M., Siegele, D. A., Almiron, M., Tormo, A., and Kolter, R. (1993) Microbial
competition: *Escherichia coli* mutants that take over stationary phase cultures, *Science*
259, 1757-1760.

14.   Zinser, E. R., and Kolter, R. (2004) *Escherichia coli* evolution during stationary phase,
Research in Microbiology 155, 328-336.

15.   Finkel, S. E., and Kolter, R. (1999) Evolution of microbial diversity during prolonged
starvation, *Proceedings of the National Academy of Sciences of the United States of*
America 96, 4023-4027.

16.   Gill, W. P., Harik, N. S., Whiddon, M. R., Liao, R. P., Mittler, J. E., and Sherman, D. R.
(2009) A replication clock for Mycobacterium tuberculosis, *Nat Med 15*, 211-214.

17.   Basu, A., and Yap, M. N. (2016) Ribosome hibernation factor promotes Staphylococcal
survival and differentially represses translation, *Nucleic Acids Research 44*, 4881-4893.

18.   Mandelstam, J., and Halvorson, H. (1960) Turnover of protein and nucleic acid in soluble
and ribosome fractions of non-growing Escherichia coli, *Biochimica et biophysica acta*
40, 43-49.

19.   Kaplan, R., and Apirion, D. (1975) The fate of ribosomes in Escherichia coli cells starved
for a carbon source, *The Journal of biological chemistry 250*, 1854-1863.

20.   Davis, B. D., Luger, S. M., and Tai, P. C. (1986) Role of ribosome degradation in the death
of starved Escherichia coli cells, *Journal of Bacteriology 166*, 439-445.

21.   Deutscher, M. P. (2003) Degradation of stable RNA in bacteria, *The Journal of biological*
chemistry 278, 45041-45044.

22.   Raina, M., and Ibba, M. (2014) tRNAs as regulators of biological processes, *Front Genet*
5.

23.   Bullwinkle, T. J., and Ibba, M. (2016) Translation quality control is critical for bacterial
responses to amino acid stress, *Proceedings of the National Academy of Sciences of the*
United States of America 113, 2252-2257.

24.   Jacob, A. I., Kohrer, C., Davies, B. W., RajBhandary, U. L., and Walker, G. C. (2013)
443   Conserved bacterial RNase YbeY plays key roles in 70S ribosome quality control and 16S
444   rRNA maturation, *Molecular Cell 49*, 427-438.

445   25.   Davies, B. W., Kohrer, C., Jacob, A. I., Simmons, L. A., Zhu, J., Aleman, L. M., Rajbhandary,
446   U. L., and Walker, G. C. (2010) Role of Escherichia coli YbeY, a highly conserved protein,
447   in rRNA processing, *Mol. Microbiol. 78*, 506-518.

448   26.   Inoue, K., Alsina, J., Chen, J., and Inouye, M. (2003) Suppression of defective ribosome
449   assembly in a rbfA deletion mutant by overexpression of Era, an essential GTPase in
450   Escherichia coli, *Mol. Microbiol. 48*, 1005-1016.

451   27.   Verstraeten, N., Fauvart, M., Versees, W., and Michiels, J. (2011) The universally
452   conserved prokaryotic GTPases, *Microbiology and molecular biology reviews : MMBR 75*,
453   507-542, second and third pages of table of contents.

454   28.   Vercruysse, M., Kohrer, C., Shen, Y., Proulx, S., Ghosal, A., Davies, B. W., RajBhandary, U.
455   L., and Walker, G. C. (2016) Identification of YbeY-Protein Interactions Involved in 16S
456   rRNA Maturation and Stress Regulation in Escherichia coli, *MBio 7*.

457   29.   Paul, B. J., Barker, M. M., Ross, W., Schneider, D. A., Webb, C., Foster, J. W., and Gourse,
458   R. L. (2004) DksA: a critical component of the transcription initiation machinery that
459   potentiates the regulation of rRNA promoters by ppGpp and the initiating NTP, *Cell 118*,
460   311-322.

461   30.   Krasny, L., and Gourse, R. L. (2004) An alternative strategy for bacterial ribosome
462   synthesis: Bacillus subtilis rRNA transcription regulation, *The EMBO journal 23*, 4473-
463   4483.

464   31.   Hauryliuk, V., Atkinson, G. C., Murakami, K. S., Tenson, T., and Gerdes, K. (2015) Recent
465   functional insights into the role of (p)ppGpp in bacterial physiology, *Nature reviews.*
466   *Microbiology 13*, 298-309.

467   32.   Yang, J., Yin, L., Lessner, F. H., Nakayasu, E. S., Payne, S. H., Fixen, K. R., Gallagher, L., and
468   Harwood, C. S. (2017) Genes essential for phototrophic growth by a purple
469   alphaproteobacterium, *Environmental Microbiology*.

470   33.   Pechter, K. B., Gallagher, L., Pyles, H., Manoil, C. S., and Harwood, C. S. (2015) Essential
471   Genome of the Metabolically Versatile Alphaproteobacterium *Rhodopseudomonas*
472   *palustris*, *Journal of Bacteriology 198*, 867-876.

473   34.   Dai, X., Zhu, M., Warren, M., Balakrishnan, R., Patsalo, V., Okano, H., Williamson, J. R.,
474   Fredrick, K., Wang, Y. P., and Hwa, T. (2016) Reduction of translating ribosomes enables
475   Escherichia coli to maintain elongation rates during slow growth, *Nat Microbiol 2*,
476   16231.

477   35.   Gefen, O., Fridman, O., Ronin, I., and Balaban, N. Q. (2014) Direct observation of single
478   stationary-phase bacteria reveals a surprisingly long period of constant protein
479   production activity, *Proceedings of the National Academy of Sciences of the United*
480   *States of America 111*, 556-561.

481   36.   Manina, G., Dhar, N., and McKinney, J. D. (2015) Stress and host immunity amplify
482   Mycobacterium tuberculosis phenotypic heterogeneity and induce nongrowing
483   metabolically active forms, *Cell Host Microbe 17*, 32-46.

484   37.   Gefen, O., Gabay, C., Mumcuoglu, M., Engel, G., and Balaban, N. Q. (2008) Single-cell
485   protein induction dynamics reveals a period of vulnerability to antibiotics in persister

486		bacteria, *Proceedings of the National Academy of Sciences of the United States of America 105*, 6145-6149.
487		
488	38.	Byrgazov, K., Vesper, O., and Moll, I. (2013) Ribosome heterogeneity: another level of complexity in bacterial translation regulation, *Curr Opin Microbiol 16*, 133-139.
489		
490	39.	Corrigan, R. M., Bellows, L. E., Wood, A., and Grundling, A. (2016) ppGpp negatively impacts ribosome assembly affecting growth and antimicrobial tolerance in Gram-positive bacteria, *Proceedings of the National Academy of Sciences of the United States of America 113*, E1710-1719.
491		
492		
493		
494	40.	Zhang, Y., Zbornikova, E., Rejman, D., and Gerdes, K. (2018) Novel (p)ppGpp Binding and Metabolizing Proteins of Escherichia coli, *MBio 9*.
495		
496	41.	Wang, B., Dai, P., Ding, D., Del Rosario, A., Grant, R. A., Pentelute, B. L., and Laub, M. T. (2019) Affinity-based capture and identification of protein effectors of the growth regulator ppGpp, *Nature chemical biology 15*, 141-150.
497		
498		
499	42.	Nguyen, D., Joshi-Datar, A., Lepine, F., Bauerle, E., Olakanmi, O., Beer, K., McKay, G., Siehnel, R., Schafhauser, J., Wang, Y., Britigan, B. E., and Singh, P. K. (2011) Active starvation responses mediate antibiotic tolerance in biofilms and nutrient-limited bacteria, *Science 334*, 982-986.
500		
501		
502		
503	43.	Primm, T. P., Andersen, S. J., Mizrahi, V., Avarbock, D., Rubin, H., and Barry, C. E., 3rd. (2000) The stringent response of Mycobacterium tuberculosis is required for long-term survival, *Journal of Bacteriology 182*, 4889-4898.
504		
505		
506	44.	Dutta, N. K., Klinkenberg, L. G., Vazquez, M. J., Segura-Carro, D., Colmenarejo, G., Ramon, F., Rodriguez-Miquel, B., Mata-Cantero, L., Porras-De Francisco, E., Chuang, Y. M., Rubin, H., Lee, J. J., Eoh, H., Bader, J. S., Perez-Herran, E., Mendoza-Losana, A., and Karakousis, P. C. (2019) Inhibiting the stringent response blocks Mycobacterium tuberculosis entry into quiescence and reduces persistence, *Sci Adv 5*, eaav2104.
507		
508		
509		
510		
511	45.	Fixen, K. R., Baker, A. W., Stojkovic, E. A., Beatty, J. T., and Harwood, C. S. (2014) Apo-bacteriophytochromes modulate bacterial photosynthesis in response to low light, *Proceedings of the National Academy of Sciences of the United States of America 111*, E237-244.
512		
513		
514		
515	46.	Quandt, J., and Hynes, M. F. (1993) Versatile suicide vectors which allow direct selection for gene replacement in gram-negative bacteria, *Gene 127*, 15-21.
516		
517	47.	McKinlay, J. B., and Harwood, C. S. (2010) Carbon dioxide fixation as a central redox cofactor recycling mechanism in bacteria, *Proceedings of the National Academy of Sciences of the United States of America 107*, 11669-11675.
518		
519		
520	48.	Lawrence, M. G., Shamsuzzaman, M., Kondopaka, M., Pascual, C., Zengel, J. M., and Lindahl, L. (2016) The extended loops of ribosomal proteins uL4 and uL22 of Escherichia coli contribute to ribosome assembly and protein translation, *Nucleic Acids Research 44*, 5798-5810.
521		
522		
523		
524	49.	Janssen, B. D., Diner, E. J., and Hayes, C. S. (2012) Analysis of aminoacyl- and peptidyl-tRNAs by gel electrophoresis, *Methods Mol Biol 905*, 291-309.
525		
526	50.	Fang, M., and Bauer, C. E. (2018) Regulation of stringent factor by branched-chain amino acids, *Proceedings of the National Academy of Sciences of the United States of America 115*, 6446-6451.
527		
528		

529   51.   Nakayasu, E. S., Burnet, M. C., Walukiewicz, H. E., Wilkins, C. S., Shukla, A. K., Brooks, S.,
530   Plutz, M. J., Lee, B. D., Schilling, B., Wolfe, A. J., Muller, S., Kirby, J. R., Rao, C. V., Cort, J.
531   R., and Payne, S. H. (2017) Ancient Regulatory Role of Lysine Acetylation in Central
532   Metabolism, *MBio 8*.
533   52.   Tyanova, S., Temu, T., and Cox, J. (2016) The MaxQuant computational platform for
534   mass spectrometry-based shotgun proteomics, *Nat Protoc 11*, 2301-2319.

Acetate

central
metabolism

CO$_2$

hv

e$^-$

H$^+$

H$^+$

Biosynthesis

ADP

ATP

NH$_4^+$

Figure 1. Metabolism of *R. palustris* during anaerobic growth in light. Acetate was used as the

carbon source and ammonium as the nitrogen source for growth in this study. *R. palustris*

generates ATP and protommotive force from light energy by cyclic photophosphorylation.

Energy generation and carbon use are independent from each other. In this study growth-arrest

occurred when *R. palustris* had used all available acetate. In this circumstance cells can

continue to generate ATP and protonmotive force from light.

576
577
Figure 2. Growth-arrested *R. palustris* cells do not appear to undergo cycles of death and
regrowth. a: Percentage of *R. palustris* cells that continue to maintain plasmid pBBR^{Gm} during
continuous growth (re-inoculated in medium with fresh carbon source repeatedly) or during
growth-arrest (no treatment after the initial carbon source was depleted). Gm-free medium was
used for this experiment. b: Growth-arrested *R. palustris* is sensitive to antibiotics targeting
ribosomes. Antibiotics are added five days after growth-arrest to independent cultures. Gm:
gentamycin. Cm: chloramphenicol. Nal: nalidixic acid. Ap: ampicillin. Error bars indicate the
standard deviation (n>3).

596
597

Figure 3. *R. palustris* protein abundances. Growing cells and growth-arrested cells are compared. a: Log$_2$-fold changes in abundances of all identified *R. palustris* proteins in growing cells (log), 6 days after growth-arrest (Day 6) and 20 days after growth arrest (Day 20) (n = 3, SI Table 1). b: The average relative quantities of ribosome proteins at day 6 (black, Day 6) and day 20 (red, Day 20) growth-arrest relative to the growing cells (log). The horizontal dashed line indicates the position of p = 0.05 (Student's *t*-test), and the vertical dashed lines indicate the position of two-fold abundance changes. Ribosome proteins decreased to ~50% and ~25% after 6 days and 20 days of growth-arrest, respectively. c: The average relative quantities of proteins involved in tRNA charging (KEGG map00970, SI Table 2), translation elongation factors (SI Table 2) and two previously identified longevity proteins [YbeY and Era, (*12*)]. The horizontal dashed line indicates the position of p = 0.05 (Student's *t*-test), and the vertical dashed lines indicate the position of two-fold abundance changes.

610
611
612

Figure 4. Translation is intact and active in growth-arrested *R. palustris*. a: The ribosome profile of cells at 6 days (Day 6) post growth-arrest is shown and compared to ribosomes from growing *R. palustris* (log). b: The ribosome profile of cells at 25 days (Day 25) post growth-arrest is shown and compared to ribosomes from growing *R. palustris* (log). c: Total RNA was purified and analyzed from growing and growth arrested *R. palustris* cells. The quantification of replicated experiments is shown in SI Figure 2. The standard is the internal standard of the RNA ScreenTape system used (Agilent). d: The charged and uncharged species of tRNA$_{trp}$ are shown here with a representative Northern blot. The aminoacylated (charged) and deacylated (uncharged) species of tRNA$_{trp}$ were separated with an acid urea gel and probed with a tRNA$_{trp}$ probe. e: Quantification of the charged and uncharged species of tRNA$_{trp}$ as analyzed in Figure 4d. Error bars represent the standard deviation (n>3).

628
629

Figure 5. YbeY and Era are important for ribosome maintenance during growth-arrest. a: Deletion of either *ybeY* or *era* had little effect on ribosomes from growing *R. palustris*, while the ribosomes from growth-arrested cultures of *ybeY* and *era* deletion mutants lacked 30S subunits. b: *R. palustris* total RNA was analyzed by Northern blot with a probe targeting 16S rRNA. In growing cells, both Δ*ybeY* and Δ*era* mutants accumulated 17S rRNA but maintained substantial levels of 16S rRNA. In growth-arrested cells, both 17S rRNA and 16S rRNA species were depleted.

Figure 6. Overexpression of YbeY$_{D118R}$ complemented the longevity defect of a Δ*ybeY* mutant. a: Longevity of different *R. palustris* strains. Deletion of the *ybeY* gene compromised the longevity of growth-arrested cells (Δ*ybeY*), and this was complemented by expressing this gene *in trans* (Δ*ybeY*::pBBP-YbeY) (*12*). A *ybeY*$_{D118R}$ mutant had a longevity defect similar to that of the Δ*ybeY* deletion mutant. Overexpression of *ybeY*$_{D118R}$ *in trans*, restored the longevity of a Δ*ybeY* mutant (Δ*ybeY*::pBBP-YbeY$_{D118R}$). Error bars indicate the standard deviation (n>3). b: *R. palustris* total RNA was analyzed by Northern blot with a probe targeting 16S rRNA. In the growth-arrested cells of Δ*ybeY*::pBBP-*ybeY*$_{D118R}$, only 16S rRNA appeared and no 17S rRNA was detected. c: The ribosomes from Δ*ybeY*::pBBP-YbeY$_{D118R}$ were analyzed on a sucrose gradient. In the growth-arrested cells, 30S, 50S and 70S ribosomes were clearly seen, though the abundance of the 50S subunit is higher than that of WT.

Figure 7. Growth-arrested cells with very low levels of (p)ppGpp have compromised longevity. a: The level of (p)ppGpp was low in exponentially growing *R. palustris* and increased during growth-arrest. b: The domain structure of Rsh*rp* (top). Locations and relative quantities of Tn insertions in Rsh*rp* (*12*). A ΔRsh*rp*64-192 mutant was made by deleting the sequence encoding the 64th-192th amino acids of Rsh*rp* from *R. palustris*. c: (p)ppGpp synthesis was severely compromised in the ΔRsh*rp*64-192 strain. The time points shown correspond to the time points shown in Figure 7a, representing early-mid log phase, mid-late log phase, entry into growth-arrest and day 5 following growth-arrest. d: The longevity of Δrsh*rp*64-192 was severely compromised. Less than 10% of cells remained viable after 2 weeks of growth-arrest. Error bars indicate the standard deviation (n>3).

677   **SI Figure 1.** The responses of *R. palustris* to various antibiotics at different growth stages.

678   Antibiotics are added to growing cultures or five days after growth-arrest. Cm: chloramphenicol.

679   Nal: nalidixic acid. Error bars indicate the standard deviation (n>3). For growing cells treated

680   with Nal, the number of viable cells was determined by A_{660} and a representative result is shown

681   (n > 3).

687 **SI Figure 2.** rRNA levels in different *R. palustris* strains at different growth-stages. The

688 quantities of 16S, 17S and 23S rRNA are determined using an RNA ScreenTape system

689 (Agilent) and normalized against the levels of 23S rRNA in growing cells of wild-type *R.*

690 *palustris*. An unpaired *t* test was performed with at least three replicates (n.s.: not significant; *:

691 $p < 0.05$; **: $p < 0.01$; ***: $p < 0.005$; ****: $p < 0.001$, as determined by Student's *t*-test).

705　**SI Figure 3.** Trp-tRNATrp remained highly charged in the growth-arrested cells of the strains of

706　Δ*ybeY* and Δ*era*.