1 **Bacterial longevity requires protein synthesis and a stringent response**

4 Liang Yin[a], Hongyu Ma[b], Ernesto S. Nakayasu[c], Samuel H. Payne[d], David R. Morris[e] and

5 Caroline S. Harwood[a†]

[a] Department of Microbiology, University of Washington, Seattle, USA.

[b] Department of Environmental Science and Engineering, Xi'an Jiaotong University, Xi'an, P.R.

China

[c] Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA, USA

[d] Department of Biology, Brigham Young University, Provo, UT, USA

[e] Department of Biochemistry, University of Washington, Seattle, USA.

Address correspondence to Caroline S. Harwood, csh5@uw.edu

Running title: Bacterial longevity

Keywords: *Rhodopseudomonas palustris*, ribosome, growth-arrest, translation, longevity,

stringent response

ABSTRACT

Gram-negative bacteria in infections, biofilms and industrial settings often stop growing due to

nutrient depletion, immune responses or environmental stresses. Bacteria in this state tend to

be tolerant to antibiotics and are often referred to as dormant. *Rhodopseudomonas palustris*, a

phototrophic α-proteobacterium, can remain fully viable for more than four months when growth

is arrested. Here, we show that protein synthesis, specific proteins involved in translation and a

stringent response are required for this remarkable longevity. Because it can generate ATP from

light during growth arrest, *R. palustris* is an extreme example of a bacterial species that will stay

alive for long periods of time as a relatively homogeneous population of cells and it is thus an

excellent model organism for studies of bacterial longevity. There is evidence that other Gram-

negative species also continue to synthesize proteins during growth arrest and that a stringent

response is required for their longevity as well. Our observations challenge the notion that

growth-arrested cells are necessarily dormant and metabolically inactive, and suggest that such

bacteria may have a level of metabolic activity that is higher than many would have assumed.

Our results also expand our mechanistic understanding of a crucial but understudied phase of

the bacterial life cycle.

IMPORTANCE

We are surrounded by bacteria; but they do not completely dominate our planet despite the

ability of many to grow extremely rapidly in the laboratory. This has been interpreted to mean

that bacteria in nature are often in a dormant state. We investigated life in growth arrest of

Rhodopseudomonas palustris, a proteobacterium that stays alive for months when it is not

growing. We found that cells were metabolically active; they continued to synthesize proteins

and mounted a stringent response, both of which were required for their longevity. Our results

suggest that long-lived bacteria are not necessarily inactive but have an active metabolism that

is well adjusted to life without growth.

INTRODUCTION. Bacteria in nature often exist in viable but non-growing states without forming

differentiated structures like spores (*1, 2*), but this crucial phase of the bacterial life cycle is

underexplored. Factors that limit cell replication include nutrient limitation, competition within

microbial communities and stresses like oxygen-depletion and extreme temperatures (*3-5*).

Several bacterial pathogens, including *Vibrio cholera*, *Pseudomonas aeruginosa* and

Burkholderia pseudomallei can survive for months and even years in a growth-arrested state in

distilled water, sterilized seawater or basal salts medium in a laboratory setting *(6-9)*. In keeping

with this, drinking water is a known reservoir for bacterial pathogens (*10*). Non-growing

pathogenic bacteria also occur in human infections and in this form present challenges for

treatment because antibiotics tend to target processes such as cell-wall biosynthesis or DNA

replication that are active in growing, but not non-growing cells. One example of a bacterium

that is recalcitrant to antibiotic treatment *in vivo* is *Mycobacterium tuberculosis*, which forms

latent infections that persist over many years in the absence of overt signs of growth (*11*). Then

there are bacterial antibiotic persisters, a small subpopulation of non-growing or slowly-growing

cells that develop in some bacterial infections and are tolerant to antibiotic treatment (*12-14*).

Studying non-growing bacteria is also important in environmental, industrial and other non-

medical contexts. For example, non-growing bacteria are excellent biocatalysts because they

can convert substrates that might be used for growth to value-added products. In past work we

have studied production of hydrogen gas, a biofuel, by the phototrophic α-proteobacterium

Rhodopseudomonas palustris. When illuminated to allow ATP synthesis by

photophosphorylation, non-growing cells diverted cell resources to hydrogen gas production and

generated 300% more hydrogen gas than growing cells (*15*). We subsequently determined

changes in metabolic fluxes and biomass composition that accounted for diversion of electrons

to hydrogen gas in growth-arrested cells (*15*). We also found that non-growing *R. palustris* cells

embedded in thin strips of latex film remained viable and produced hydrogen gas for over four

months (*16*).

These results prompted us to develop *R. palustris* as a model for studies of bacterial longevity,

and to ask the question: are there universal features of long-lived bacteria that we can uncover

from studies of this proteobacterium? Several attributes make *R. palustris* a good model for

addressing this question. First, it generates ATP and proton motive force from light, even when

it is not growing (Figure 1). Also, *R. palustris* carries out photophosphorylation under anaerobic

conditions and does not generate oxygen, thus avoiding complications of oxidative stress during

growth arrest. A Tn-seq study of growth-arrested *R. palustris* identified 116 genes, which we call

longevity genes, that are essential for long-term viability of non-growing cells, but are not

required for growth (*17*). Among these are genes that are conserved in bacteria and associated

with basic cellular processes including protein synthesis. Here, we carried out a physiological

characterization of *R. palustris* in its growth-arrested, but viable state. We found that active

translation and optimized ribosomes were critical for the longevity of *R. palustris,* as was

guanosine polyphosphate [(p)ppGpp]. This goes against the conventional description of non-

growth as state of metabolic inactivity often referred to as dormancy. We suggest that protein

synthesis and other cellular activities are likely important for many bacteria to maintain viability

when they are not growing.

RESULTS

Conditions of growth arrest. In this study *R. palustris* was grown anaerobically in minimal

salts medium in sealed glass tubes in light until growth arrest occurred due to carbon source

depletion (*17*). Cultures were incubated post-growth arrest in the same tubes in which they were

grown. Nitrogen was supplied in excess as ammonium and the carbon supplied was 20 mM

sodium acetate. We took as day 0 of growth arrest, the time at which cells stopped growing as

measured by a leveling off of the increase in the optical density (OD) of cultures. Cells grew to

twice the OD when we supplied them with 40 mM acetate, confirming that growth arrest

occurred due to carbon source limitation.

Characterization of cell size. Upon growth arrest due to starvation for nutrients many bacteria

decrease their size and change from a rod-shaped to coccid form (*6, 18-20*). Degradation of

ribosomes that accompanies growth arrest typically results in a loss of total cellular protein and

RNA, which likely contributes to cell shrinkage. To put our work on *R. palustris* into context with

the large body of work on starvation-survival of heterotrophic bacteria, we measured cell size

before and after growth arrest. As shown in Figure 2, cells in log phase and at 6 days post-

growth arrest do not differ appreciably in length or overall morphology.

Growth-arrested *R. palustris* cells do not appear to undergo cycles of death and

regrowth. To put the work reported here in context with another aspect of the starvation

literature, we note that depending on the conditions of growth arrest some bacteria, the best

studied of which is *E. coli*, undergo cycles of death and growth after entry into stationary phase

(*1, 21*). Typically a fraction of cells survive long-term stationary phase by feeding on nutrients

released from dead cells. This leads to heterogeneity among cells, making it difficult to carry out

meaningful studies of growth-arrested bacteria at the population level. Also genetic

heterogeneity develops in populations of growth-arrested bacteria as there is strong selection

pressure for mutations that enable cannibalism (*22, 23*). In previous work we reported that

populations of growth-arrested *R. palustris* cells were nearly 100% viable (*17*). To firmly

establish that cells were truly growth arrested, rather than undergoing cycles of growth and

death, we introduced an unstable plasmid expressing a gentamycin (Gm) resistance gene into

wild-type *R. palustris*. When Gm is absent, the percentage of growing cells that retain this

plasmid would be expected to decrease (*24*). On the other hand, a non-growing culture would

be expected to retain the plasmid in Gm-free medium. When this strain (WT::pBBR^{Gm}) was

inoculated into Gm-free medium, about 60% of the cells had lost the plasmid by the time they

reached stationary phase. When these cells were transferred into fresh Gm-free medium and

successively transferred several times over a period of 20 days after reaching stationary phase,

the plasmid continued to be lost. In contrast, growth-arrested cells retained the plasmid over the

same period of time (Figure 3a). This suggests that most or all of the cells were indeed in a non-

growing state. However, we cannot exclude that some small percentage of the cells underwent

cycles of growth and death.

In further support of the conclusion that growth-arrested *R. palustris* does not undergo major

cycles of growth and death, we found that non-growing cells were tolerant to the cell wall

synthesis inhibitor ampicillin and the DNA synthesis inhibitor nalidixic acid at concentrations that

prevented growth (Figure 3b and SI Figure 1). However, we also found that the viabilities of

non-growing cells were compromised by protein synthesis inhibitors (Figure 3b and SI Figure 1).

Long-lived, growth-arrested cells continue to synthesize proteins. Growth-arrested *R.*

palustris was sensitive to Gm, an aminoglycoside antibiotic that affects protein synthesis by

causing mistranslation, which results in accumulation of damaged proteins. We found that over

99% of growth-arrested cells lost viability over a few days when treated with Gm at a final

concentration of 50 µg/ml (Figure 3b). Although cells died relatively rapidly, they were not as

sensitive to Gm as growing cells, which died within a few hours after exposure to this antibiotic.

We noticed that the viability of growth-arrested cells exposed to Gm at 20 d post-growth arrest

tend to be higher than the number of viable cells at day 10 post-growth arrest (Figure 3b). It is

possible that some cells developed resistance to Gm or that the Gm supplied may have become

degraded, allowing for a small amount of growth on debris released from dying cells.

Chloramphenicol (Cm) belongs to another class of antibiotic that targets ribosomes. Although

typically bacteriostatic, Cm was effective in killing growth-arrested *R. palustris* over a prolonged

period of time (Figure 3b and SI Figure 1). These results suggest that active translation is

important for *R. palustris* longevity.

Proteomics experiments revealed that after 6 days and 20 days of growth arrest, 289 and 525

proteins, respectively, were present at greater than two-fold higher levels than in growing cells

(Figure 4a and SI Table 1). Among these were enzymes involved in the degradation of

compounds, including benzoate and fatty acids (SI Tables 1 and 2), that are preferred carbon

sources for *R. palustris*, suggesting that cells respond to growth arrest by increasing the

translation of proteins that may aid in scavenging carbon. In the period between 6 and 20 days

post growth arrest, 51 proteins of unknown function increased in abundance. Some proteins

(163 and 310 proteins at days 6 and 20 post growth arrest) were diminished in abundance by

two-fold or more following growth-arrest (Figure 4a and SI Table 1). This set included 53 out of

55 ribosomal proteins that were identified in our proteomics dataset. On average, ribosome

proteins were at approximately 50% and 25% of their original levels after 6 days and 20 days of

growth arrest (Figure 4b and 4c). We calculated the relative mass of these proteins per cell and

from this calculated that ribosomes comprised 25.7% of the total protein mass in growing cells,

which was reduced to 13.1% (49% reduction) and 6.8% (74% reduction) after 6 and 20 days of

growth arrest (Figure 4c, SI Table 3). Not all translation related proteins were decreased in their

abundances. For example, translation elongation factors made up ~5% of the total protein from

growing cells as well as growth-arrested cells (Figures 4d and 4e, SI Table 3). Similar patterns

were also observed for proteins involved in aminoacyl-tRNA biosynthesis (tRNA charging) and

for two previously identified longevity proteins YbeY and Era (*17*) that are involved in ribosome

maturation (Figure 4d).

Finally, to directly examine protein synthesis in growth-arrested *R. palustris*, we constructed a

LacZ-reporter driven by a P_{hirl} promoter that can be induced with phenylacetyl-homoserine

lactone (PA-HSL) (SI Figure 2). After 6 days of growth arrest, PA-HSL was added to the culture

and samples were collected after 2 additional days of incubation. The negative control without

added signal showed a base level of LacZ expression. Compared to this, LacZ acitvity was

~70% higher when the PA-HSL signal was added to induce lacZ expression (SI Figure 2). This

observation directly demonstrates that *R. palustris* has the ability to synthesize proteins during

growth arrest.

Characterization of protein synthesis apparatus in long-lived cells. We isolated *R. palustris*

ribosomes by sucrose gradient centrifugation and found that the ribosome profile of growing *R.*

palustris cells included 30S, 50S and 70S populations, as is typical for growing bacteria (Figure

5a). There was also a small peak corresponding to a 100S population, which is likely a dormant

form of ribosomes, as has been seen in other bacteria (*25, 26*). The ribosome profile of growth-

arrested *R. palustris* cells resembled that of growing cells (Figure 5a and 5b), but the quantity of

the 70S population was lower at day 6 and even lower at day 25 post growth arrest, suggesting

that cells degraded some fraction of their ribosomes after they stopped growing. This is

consistent with our proteomics data (Figure 4b) and fits the notion that the total rate of protein

synthesis is expected to be much lower in growth-arrested cells compared to growing cells.

Ribosome degradation is often accompanied by the non-specific degradation of rRNA (*27-30*).

However, even after 120 d of growth arrest, cells retained intact species of 23S and 16S rRNA,

indicating that cells continuously repaired and replaced ribosomes albeit at probably a low rate

after they stopped growing (Figure 5c, SI Figure 3). The relative levels of aminoacylated (or

charged) and free forms of tRNAs are important for promoting and inhibiting translation (*31*). We

used Northern blot analysis to estimate the percentage of tryptophan-charged $tRNA_{trp}$ (Trp-

$tRNA_{trp}$) relative to $tRNA_{trp}$. In growing cells, ~60% of $tRNA_{trp}$ was in the charged Trp-$tRNA_{trp}$

form (Figure 5d and 5e). This level is comparable to that reported in other bacteria (*32*). The

percentage of Trp-tRNA$_{trp}$ increased to ~85% upon growth arrest and was maintained at this

level for ~20 days (Figure 5d and 5e).

Two proteins identified as essential for *R. palustris* longevity are critical for ribosome

integrity in growth-arrested cells. Among the longevity genes that we have identified in *R.*

palustris (*17*), *ybeY* and *era* are predicted to encode proteins that interact with each other and

the S11 ribosome protein during ribosome maturation (*33-37*). *ybeY* or *era* deletion mutants

have compromised longevity but grow normally (*17*). Here we found that the ribosome profiles

of growing cells were similar between the wild type and these mutants (Figure 6a). All had

distinctive 30S, 50S and 70S populations, although the 50S subunit was slightly more abundant

in the mutants. The ribosome profiles of growth-arrested *ΔybeY* and *Δera* mutant cells were

noticeably different from that of the wild type, however. A prominent 50S subunit peak was seen

but the amounts of 30S subunits and 70S ribosomes were greatly diminished (Figure 6a).

As predicted based on studies of their *E. coli* homologs (*33-36*), *ΔybeY* and *Δera* strains were

defective in 16S rRNA maturation. An extra band immediately above the 16S rRNA was present

in gels of RNA from both mutants (SI Figure 3). Northern blot experiments confirmed that the

extra band was 17S rRNA, a precursor of 16S rRNA (Figure 6b). In growth-arrested cells, both

16S rRNA and 17S rRNA species were severely depleted in the mutated strains (Figure 6b and

SI Figure 3). This is likely connected to the defective 30S ribosome subunit production shown in

Figure 6a, and the compromised longevity of *R. palustris*. We do not know why the 17S and 16S

rRNAs were so much lower in abundance in growth-arrested vs. growing cells (Figure 6b and SI

Figure 3). It is possible that factors associated with these mutations coupled with the stress of

growth arrest lead to their degradation. *ybeY* or *era* deletion mutants had wild-type levels of

charged tRNA-Trp (SI Figure 4), suggesting that tRNA aminoacylation is not directly coordinated

with the process of ribosome maturation.

As shown in Figure 4b, there was no significant change in the abundance of YbeY and Era

proteins in growth-arrested cells. Since the core ribosomal proteins were decreased 2 to 4-fold

in abundance (Figure 4b), the relative abundance of YbeY and Era was increased on a per

ribosome basis following growth arrest. To test whether we could change the effects of YbeY on

longevity by modulating its relative abundance, we constructed and expressed a YbeY variant

predicted based on work in *E. coli*, to interact weakly with ribosomes. In *E. coli*, the interaction

between YbeY and ribosomes is partially disrupted by replacement of a conserved aspartate

with an arginine (*37*). When we made the corresponding YbeY$_{D118R}$ change in *R. palustris*, the

strain was compromised in its longevity to a similar degree as the *ΔybeY* mutant (Figure 7a).

When we overexpressed YbeY$_{D118R}$ *in trans* (*ΔybeY*::pBBP-*ybeY*$_{D118R}$) however, the longevity

defect was complemented (Figure 7a). The quantity of 16S rRNA was similar to that of the wild-

type, and no 17S rRNA was observed during growth arrest (Figure 7b). The ribosome profile of

the growth-arrested *ΔybeY*::pBBP-*ybeY*$_{D118R}$ -complemented strain differed from that of growth-

arrested wild-type cells, especially in the quantities of 50S and 70S ribosomes (Figure 7c).

However, growth-arrested cells could apparently tolerate these differences when challenged in

longevity assays (Figure 7a). These results suggest that it may be helpful for growth-arrested

cells to have relatively more YbeY protein compared to growing cells and that this promotes 30S

ribosome subunit assembly in this condition.

The stringent response is important for longevity. Our results suggest that translation is

reduced but still maintained in growth-arrested cells. A well-known mechanism that contributes

to this is the stringent response mediated by guanosine polyphosphate [(p)ppGpp] (*38, 39*). In

many bacteria, the level of (p)ppGpp increases upon growth arrest and this inhibits the

transcription of ribosomal genes, although the mechanism by which this is achieved varies

among bacteria (*38-40*). As with other bacteria, we detected little to no (p)ppGpp in

exponentially growing *R. palustris*. However, the level of (p)ppGpp increased sharply as growing

cells reached stationary phase and remained at relatively high levels after growth arrest (Figure

8a).

Unlike *E. coli*, but like most other bacteria outside the γ- and β-proteobacteria, *R. palustris* has a

single bifunctional (p)ppGpp synthesis/hydrolysis *rsh* gene, here called rsh_{rp}, that has multiple

domains including a hydrolase (HDc) and RelA_SpoT domain (Figure 8b). Our Tn-seq results

indicated that this gene (*rpa2693*) was essential for *R. palustris* longevity, but not for growth (*17,*

41, 42). However, we were unable to delete the full-length rsh_{rp} gene. A subsequent inspection

of the locations of the Tn-insertions in this gene in our bank of Tn mutagenized cells revealed

that there were relatively few insertions in its RelA_SpoT domain (Figure 8b). When we made

domain specific deletions in Rsh$_{rp}$, we obtained a hydrolase domain deletion mutant (ΔRsh$_{rp64-}$

$_{192}$, Figure 8b) that produced very little (p)ppGpp during growth or growth arrest. (Figure 8c). We

do not know why this deletion affected (p)ppGpp synthesis in this way, but one idea is that it

destabilized or altered the structure of the protein so that its ability to synthesize (p)ppGpp was

compromised. The ΔRsh$_{rp64-192}$ mutant grew with a generation time of 9.4 ± 0.6 h, compared to a

wild type 6.4 ± 0.2 h generation time, a growth rate difference of about 30%. In contrast to this

relatively modest effect on the growth rate, the ΔRsh$_{rp60-189}$ mutation had a severe effect on

longevity. As shown in Figure 8d, the mutant survived poorly in growth arrest.

DISCUSSION

We found that illuminated *R. palustris* responds to growth arrest by reducing ribosome

abundance as is typical of most bacteria when they slow their growth rate. Growth-arrested cells

continued to synthesize proteins, some at elevated levels relative to growing cells, over the

period of 20 days that we followed protein composition by proteomics (SI Table 1) and it is likely

that they repaired and replaced some portion of their original ribosome inventory over this time

period. The levels of most proteins involved in tRNA charging and translation elongation and

were at similar levels in growing and growth-arrested cells (Figure 4d and SI Table 2), which is

consistent with the notion from a previous report that elongation rates are maintained as

bacteria slow down their growth-rates even towards zero growth (*43*). In addition, a tRNA that

we followed was almost fully charged 20 d into growth arrest (Figure 5d and 5e). Not only does

protein synthesis continue after growth arrest, but it is essential for viability of non-growing cells

as indicated by their sensitivity to Gm and Cm (Figure 3b, SI Figure 1).

There are reports of continued protein synthesis in other species of bacteria in growth arrest,

the more convincing of which come from studies in which production of fluorescent proteins in

single cells was used as a proxy for protein synthesis. Antibiotic persisters in early stages of

formation, as well as non-growing *E. coli*, *Mycobacterium* and *Salmonella* have all been inferred

to carry out protein synthesis (*44-47*). These results suggest that maintenance of some level of

protein synthesis may be required for viability of many non-growing bacteria. From this it follows

that it would make sense to prioritize development of antibiotics that target protein synthesis in

growth-arrested cells. This will likely require understanding in detail what happens to bacterial

ribosomes following growth arrest. Bacterial ribosomes are known to be differentially post-

translationally modified in stationary phase cells (*48*). Two longevity genes, *ybeY* and *era*, were

essential for normal ribosomes to form in non-growing *R. palustris* (Figure 6a), and their

encoded proteins increased abundance on a per ribosome basis during growth arrest (Figure

4b). Era is a well-studied bacterial GTPase (*36*), and genes for it and YbeY, are present and

conserved in most bacteria (*34*). Our data are consistent with studies of the corresponding *E*

coli proteins showing that YbeY and Era are involved in rRNA processing in *R. palustris*. When

we looked at rRNAs formed by Δ*ybeY* and Δ*era* mutants that were actively growing, it was

apparent that growing cells and growth-arrested cells had the same defect in 17S rRNA

processing (Figure 6b), but only the growth-arrested cells had a physiological phenotype. This

highlights a feature of at least some *R. palustris* longevity proteins, which is that they are

functional in growing cells as well as in non-growing cells, but it is only when the additional

demands of growth arrest on cellular functionality are imposed that essentiality becomes

apparent in growth arrest. Our work on Era and YbeY has led us to conclude that optimized

ribosomes are a feature of long-lived cells.

In early stationary phase and when in growth arrest, *R. palustris* synthesizes increasing

amounts of (p)ppGpp, indicating that cells mount a stringent response. The gene *rpa2693*,

which we call rsh_{rp}, is among the genes identified by Tn-seq to be essential for longevity (*17*). A

$\Delta rsh_{rp60\text{-}189}$ mutant did not produce detectable (p)ppGpp and the viability of this mutant

decreased rapidly after growth arrest (Figure 8c and 8d). There is good evidence that (p)ppGpp

is also important for development of antibiotic persisters (*49*) and for long term survival of latent

and growth-arrested *M. tuberculosis* (*50, 51*). A great deal still needs to be learned about how

(p)ppGpp signaling is activated in growth-arrested *R. palustris*. Studies of (p)ppGpp in different

species of bacteria have generally not followed the levels of this nucleotide deep into stationary

phase. *E. coli* has been shown to have elevated levels of (p)ppGpp only in early stationary

phase, with (p)ppGpp levels returning to baseline later in stationary phase. This is very different

from the situation in *R. palustris* where high levels of (p)ppGpp are maintained for at least five

days post-growth arrest. Uncharged tRNAs are understood to be the classic signal for activation

of RelA and up-regulation of (p)ppGpp synthesis in *E. coli* and other gamma proteobacteria

(*40*). But in *R. palustris*, tRNA$_{\text{Trp}}$ charging is even higher in growth arrest than during growth

(Figure 5d and 5e). It could be that one or several tRNA species other than tRNA$_{\text{Trp}}$ become

uncharged during carbon starvation, which could potentially trigger (p)ppGpp synthesis in a

fashion similar to the case of *E. coli*. However, it is also known for that the exact triggers of

(p)ppGpp synthesis can vary between species (*40, 52, 53*). For example, amino acid starvation

that typically leads to uncharged tRNA species is insufficient to trigger (p)ppGpp accumulation

in *Rhizobium meliloti (54, 55)*, *Caulobacter crescentus* (*56*) and *Rhodobacter sphaeroides* (*57*).

A nitrogen-related phosphotransferase system (PTS^{NTR}) has been found to be important for

stimulating (p)ppGpp production in *C. crescentus* in response to nitrogen limitation (*58, 59*), but

the mechanism by which carbon limitation, the condition studied here, may stimulate (p)ppGpp

accumulation in α-proteobacteria remains unclear *(56)*.

Other future areas of investigation are to understand why (p)ppGpp is essential for *R. palustris*

longevity and to identify its target genes and proteins. Based on work with other bacteria it is

very likely that (p)ppGpp interacts with RNA polymerase to inhibit transcription of genes

encoding ribosomal RNAs and proteins (*38-40*). But this would need to be done in a very

controlled fashion because some new ribosome biogenesis is very likely to be required for *R.*

palustris longevity in growth arrest. (p)ppGpp has recently been shown to bind to Era from

Staphylococcus aureus and *E. coli* to inhibit its GTPase activity (*60-62*), and a similar effect on

Era from *R. palustris* might influence the biogenesis and integrity of its ribosomes.

Virtually all bacteria probably stop growing at one time or other due to nutrient limitation in their

native niches and it is also likely that that they employ diverse strategies to stay alive. However,

there also may be evolutionarily conserved core survival strategies that all bacteria use. We

suggest that optimized ribosomes, a stringent response and protein synthesis are part of this

core strategy   The primary reason for the extreme longevity of *R. palustris* is likely its ability to

continuously generate ATP by photophosphorylation when it is in growth arrest, because

growth-arrested cells incubated in dark do not survive nearly as well *(17)*. All bacteria need ATP

to stay alive. There is plenty of evidence in the literature suggesting that ATP, obtained by

scavenging resources from the environment or from degradation of internal energy stores or cell

biomass is necessary to support the viability of diverse non-spore forming growth-arrested

bacteria (*20*). A recent study also suggested that varying ATP levels per cell might underlie the

heterogeneity in populations and lead to non-growing persisters (*63*). Protein synthesis and

(p)ppGpp synthesis are core features that underlie the extraordinary longevity of *R. palustris*.

Both of these processes are highly energy-consuming and crucial for all bacteria. To understand

how *R. palustris* integrates these processes and utilizes its ATP during growth-arrest could be a

key to understand how non-growing bacteria in general stay alive.

METHODS

Bacterial strains, growth and incubation conditions. *R. palustris* strain CGA009 was used

as the wild-type for this study. Anaerobic cultures were grown under illumination at 30 □C, using

a defined media PM with 20 mM sodium acetate or 10 mM sodium succinate as the carbon

source (*64*). The longevity of *R. palustris* cultures was determined by counting colony-forming

units as described previously (*17*). In-frame deletions of *R. palustris* genes were created by

using suicide vector pJQ200SK as described before (*65*). For *in trans* expression in *R. palustris*,

the target genes were cloned into plasmid pBBPgdh and the plasmids were mobilized into *R.*

palustris by conjugation using *E. coli* strain S17-1 (*66*). Media were supplemented with 100

µg/ml of gentamicin for *in trans* complementation experiments.

Construction of inducible LacZ-reporter and LacZ assays. To construct the inducible LacZ-

reporter, we assembled a *lacZ* fragment and a *hirR*-P_{hirl} fragment (*67*) into a pBBP plasmid

backbone (*66*) (SI Figure 2). The promoter P_{hirl} responds specifically to PA-HSL signal (*67*). At

day 6 of *R. palustris* growth arrest, PA-HSL was added into the culture at final concentration of 1

µM. After an additional ~48 h of incubation, 5 ml of culture was collected and resuspended in 2

ml of Z-buffer (60 mM Na_2HPO_4, 40 mM NaH_2PO_4, 12.5 mM NaCl, 1 mM MgSO4 and 40mM 2-

mercaptoethanol, pH 7.0). The samples were sonicated at a frequency of 20~40 kHz for 3

cycles (20 s sonication and 20 s cooling on ice). Cell lysis was cleared by centrifugation at 4

$\square$C, 15,000 rpm for 15 min, and the supernatant was collected for subsequent β-galactosidase

assays. To start the reaction, 200 μl supernatant was mixed with 40 μl of 4 mg/ml ONPG at 30

$\square$C, and Abs_{420nm} was monitored using a Biotek H1 plate-reader. The level of LacZ was

calculated as the V_{max} of the reaction normalized by total protein concentration of the

supernatant.

Ribosome purification. Ribosomes from *R. palustris* cells were isolated following a method

adapted from a previous study (*68*). About 35 ml of culture was grown to the desired state and

poured over a 10-ml ice sheet (fast-cool). The ice sheet melted within 10 min and the cells were

collected by centrifugation at 4000 rpm for 15 min. The cell pellet was resuspended in 2 ml of

Buffer A (20mM HEPES-KOH, pH 7.5, 30mM NaCl, 8mM $MgCl_2$) with 1 μl/ml RNasin

(Promega). Three hundred microliter of 0.1 mm Zirconia/Silica beads (Biospec Products) were

added to every ml of sample. The samples were lysed with a Mini-Beadbeater-24 (Biospec

Products) at 3500 rpm for 1 min at 4 $\square$C, kept on ice for 1 min and the cycle was repeated 4

more times. The micro-beads and cell debris were removed by centrifugation at 15000 rpm for

10 min. The lysates were further clarified by spinning at 30,000 g in Beckman TL-1000

ultracentrifuge for 30 min, followed by pelleting ribosomes at 100,000 g for 4 hours. The pellet

was resuspended with Buffer A overnight on ice.

Sucrose gradient analysis. About five A_{260} units of ribosomes (less than 200 μl in volume)

were applied on top of 11 ml of a sucrose gradient (7-47%). The samples were centrifuged

through the gradient at 39,000 rpm for 4.5 h with a SW41 rotor, and then were separated with a

Brandel gradient pump. The sucrose gradient was made with Buffer A.

Ribosomal RNA analysis. Total RNA was purified from 3-5 ml of *R. palustris* culture using an

miRNeasy Mini Kit. The rRNAs were then analyzed with an RNA ScreenTape system (Agilent).

To probe 16S rRNA by Northern blot, 1 µg of total RNA was heated at 70 □C for 10 min and

immediately chilled on ice. The samples were then separated on a 1% agarose gel with 1X TBE

buffer. After electrophoresis, RNA was transferred to an Amersham Hybond-XL membrane (GE

Lifesciences) with a downward nucleic acid transfer system overnight using 20X SSC buffer. A

probe (5'-TCGTGCCTCAGCGTCAGTAATGGCCCAGTGA) labeled with DIG was used to

detect 16S rRNA following the manufacturer's instruction (DIG, Roche).

Charging analysis of tRNA$_{Trp}$. Total RNA was isolated with QIAzol lysis reagent according to

the manufacturer's instructions (Qiagen). The dried RNA pellet was dissolved and stored in 10

mM sodium acetate, pH 4.5. A deacylated control sample was prepared by incubating 10 µg of

total RNA with 1 mM EDTA and 100 mM Tris-HCl pH 9.0 at 42 □C for an hour (*69*). Northern

blot and acid-urea gel electrophoresis were performed essentially as described previously (*32,*

69). Loading dye (0.01% xylene cyanol, 0.01% bromophenol blue, 8 M Urea, 1 mM EDTA and

10 mM sodium acetate, pH 5.0) was mixed with sample at 1:1 ratio. Samples were loaded on a

12% acid-urea gel of 20 cm and then were run at 15 W for 12 h at 4 □C. Transfer of RNA to an

Amersham Hybond-XL membrane (GE Lifesciences) was performed in 0.5* TBE buffer at 650

mA for 30 min 4 □C. The aminoacylated and the deacylated tRNA$_{Trp}$ were detected using a

DIG-labeled probe (5'-GGTTTTGGAGACCGGTGCTCTACCAATTGAG) following the

manufacturer's instruction (DIG, Roche).

Determination of cellular (p)ppGpp levels. *In vivo* (p)ppGpp labeling and detection were

performed as described with some modifications (*70*). Anaerobic *R. palustris* cultures were

grown in 1.5 ml PM medium with 100 µCi/ml of Na$_2$H^{32}PO$_4$ (PerkinElmer-NEX011001MC) added

at the time of inoculation. At the desired time points, 200-300 µl of culture was collected and

extracted with 8 M formic acid. Supernatants of the extracts were spotted on PEI-cellulose TLC

plates (Sigma-Aldrich) and separated with 1.5 M KH_2PO_4 (pH 3.4). The plates were then dried

and visualized by phosphor imaging. The level of (p)ppGpp was calculated by the ratio of

(ppGpp + pppGpp) / (GTP + ppGpp + pppGpp).

Proteomic analysis. Biological triplicates of cells were lysed, digested with trypsin and

analyzed by liquid chromatography-tandem mass spectrometry as described in (*71*). Collected

data were analyzed using MaxQuant software (v 1.5.5.1) (*41, 72*) for peptide/protein

identification and quantification. Identification of peptides was done by searching against the *R.*

palustris ATCC BAA-98 (strain CGA009) sequences from the Uniprot Knowledgebase

(downloaded on August 30, 2016). The search was performed first with a larger parent mass

tolerance (20 ppm) to identify peptides and re-calibrate peptide masses. After re-calibration, the

parent mass tolerance was set to 4.5 ppm for a second round of search. During the searches

oxidation of methionine as a variable modification, and trypsin digestion was required in both

peptide termini. Identifications were filtered with 1% false-discovery rate at peptide-spectrum

match and protein levels. The function "matching between runs" was checked to detect peptide

peaks in LC-MS/MS runs that they were not fragmented. Label-free quantification (LFQ) values

were used for quantitative analysis. This method considers only the most reliably detected

peptides and normalizes according to the total signal of the run. It is, therefore, ideal for

comparing individual proteins in different samples (*73*). For absolute quantification, intensity-

based absolute quantification (iBAQ) (*74*) measurements of individual proteins were

sequentially normalized by the protein molecular mass and total mass in the sample, resulting in

relative protein mass compared to the entire cell proteome. Proteins were considered

differentially abundant with a p-value ≤0.05 by T.test considering two tails and equal variance.

The mass spectometry data have been deposited to the ProteomeXchange Consortium via the

PRIDE partner repository with the dataset identifier PXD013729.

Funding information

This work was supported by grant W911NF-15-1-0150 to CSH from the US Army Research Office. SHP and ESN were supported by the U.S. Department of Energy, Office of Biological and Environmental Research, Early Career Research Program.

Acknowledgements

The authors thank Amy Schaefer, Kathryn Fixen, Yasuhiro Oda and Yanning Zheng for helpful discussions, and Karl Weitz for technical assistance with the mass spectrometry analysis. Mass spectrometry data was generated in the Environmental Molecular Science Laboratory, a U.S. Department of Energy (DOE) national scientific user facility at Pacific Northwest National Laboratory (PNNL) in Richland, WA. Battelle operates PNNL for the DOE under contract DE-AC05-76RLO01830.

Abbreviations

Tn: transposon.

Author contribution

L.Y. and C.S.H. designed the research. L.Y. and H.M. performed the antibiotics assays with *R. palustris*. L.Y. made the *R. palustris* mutants. L.Y. grew the *R. palustris* cells. S.H.P. and E.S.N. performed proteomic analysis. L.Y. and D.R.M. carried out the *in vitro* analysis of ribosome. L.Y. carried out the analysis of rRNA and tRNA. L.Y. performed the analysis of (p)ppGpp. All authors analyzed the data. L.Y., E.S.N. and C.S.H. wrote the paper. All authors contributed to the revision of the manuscript.

Competing interests

The authors declare no competing financial interests.

References

1.   Finkel, S. E. (2006) Long-term survival during stationary phase: evolution and the GASP
phenotype, *Nature reviews. Microbiology 4*, 113-120.
2.   Bergkessel, M., Basta, D. W., and Newman, D. K. (2016) The physiology of growth arrest:
uniting molecular and environmental microbiology, *Nature reviews. Microbiology 14*,
549-562.
3.   Wayne, L. G., and Sohaskey, C. D. (2001) Nonreplicating persistence of mycobacterium
tuberculosis, *Annual Review of Microbiology 55*, 139-163.
4.   Thieringer, H. A., Jones, P. G., and Inouye, M. (1998) Cold shock and adaptation,
BioEssays : news and reviews in molecular, cellular and developmental biology 20, 49-57.
5.   Hibbing, M. E., Fuqua, C., Parsek, M. R., and Peterson, S. B. (2010) Bacterial competition:
surviving and thriving in the microbial jungle, *Nature reviews. Microbiology 8*, 15-25.
6.   Hood, M. A., Guckert, J. B., White, D. C., and Deck, F. (1986) Effect of nutrient
deprivation on lipid, carbohydrate, DNA, RNA, and protein levels in Vibrio cholerae, *Appl
Environ Microbiol 52*, 788-793.
7.   Moore, R. A., Tuanyok, A., and Woods, D. E. (2008) Survival of Burkholderia
pseudomallei in water, *BMC Res Notes 1*, 11.
8.   Lewenza, S., Abboud, J., Poon, K., Kobryn, M., Humplik, I., Bell, J. R., Mardan, L., and
Reckseidler-Zenteno, S. (2018) Pseudomonas aeruginosa displays a dormancy
phenotype during long-term survival in water, *PLoS One 13*, e0198384.
9.   Machreki, Y., Kouidhi, B., Machreki, S., Chaieb, K., and Saenz, Y. (2019) Analysis of a long
term starved Pseudomonas aeruginosa ATCC27853 in seawater microcosms, *Microb
Pathog 134*, 103595.
10.   Szewzyk, U., Szewzyk, R., Manz, W., and Schleifer, K. H. (2000) Microbiological safety of
drinking water, *Annual Review of Microbiology 54*, 81-127.
11.   Ai, J. W., Ruan, Q. L., Liu, Q. H., and Zhang, W. H. (2016) Updates on the risk factors for
latent tuberculosis reactivation and their managements, *Emerg Microbes Infect 5*, e10.
12.   Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L., and Leibler, S. (2004) Bacterial
persistence as a phenotypic switch, *Science 305*, 1622-1625.
13.   Lewis, K. (2010) Persister cells, *Annual Review of Microbiology 64*, 357-372.
14.   Balaban, N. Q., Helaine, S., Lewis, K., Ackermann, M., Aldridge, B., Andersson, D. I.,
Brynildsen, M. P., Bumann, D., Camilli, A., Collins, J. J., Dehio, C., Fortune, S., Ghigo, J.-
M., Hardt, W.-D., Harms, A., Heinemann, M., Hung, D. T., Jenal, U., Levin, B. R., Michiels,
J., Storz, G., Tan, M.-W., Tenson, T., Van Melderen, L., and Zinkernagel, A. (2019)
Definitions and guidelines for research on antibiotic persistence, *Nature Reviews
Microbiology*.
15.   McKinlay, J. B., Oda, Y., Ruhl, M., Posto, A. L., Sauer, U., and Harwood, C. S. (2014) Non-
growing *Rhodopseudomonas palustris* increases the hydrogen gas yield from acetate by
shifting from the glyoxylate shunt to the tricarboxylic acid cycle, *The Journal of biological
chemistry 289*, 1960-1970.

16. Gosse, J. L., Engel, B. J., Hui, J. C., Harwood, C. S., and Flickinger, M. C. (2010) Progress
toward a biomimetic leaf: 4,000 h of hydrogen production by coating-stabilized
nongrowing photosynthetic Rhodopseudomonas palustris, *Biotechnol Prog 26*, 907-918.
17. Pechter, K. B., Yin, L., Oda, Y., Gallagher, L., Yang, J., Manoil, C., and Harwood, C. S.
(2017) Molecular Basis of Bacterial Longevity, *MBio 8*.
18. Sun, J., Gao, X., Qun, J., Du, X., Bi, K., Zhang, X., and Lin, L. (2016) Comparative analysis
of the survival and gene expression of pathogenic strains Vibrio harveyi after starvation,
FEMS Microbiology Letters 363.
19. Amy, P. S., and Morita, R. Y. (1983) Starvation-survival patterns of sixteen freshly
isolated open-ocean bacteria, *Appl Environ Microbiol 45*, 1109-1115.
20. Lever, M. A., Rogers, K. L., Lloyd, K. G., Overmann, J., Schink, B., Thauer, R. K., Hoehler, T.
M., and Jorgensen, B. B. (2015) Life under extreme energy limitation: a synthesis of
laboratory- and field-based investigations, *FEMS Microbiol Rev 39*, 688-728.
21. Zambrano, M. M., Siegele, D. A., Almiron, M., Tormo, A., and Kolter, R. (1993) Microbial
competition: *Escherichia coli* mutants that take over stationary phase cultures, *Science*
259, 1757-1760.
22. Zinser, E. R., and Kolter, R. (2004) *Escherichia coli* evolution during stationary phase,
Research in Microbiology 155, 328-336.
23. Finkel, S. E., and Kolter, R. (1999) Evolution of microbial diversity during prolonged
starvation, *Proceedings of the National Academy of Sciences of the United States of*
America 96, 4023-4027.
24. Gill, W. P., Harik, N. S., Whiddon, M. R., Liao, R. P., Mittler, J. E., and Sherman, D. R.
(2009) A replication clock for Mycobacterium tuberculosis, *Nat Med 15*, 211-214.
25. Basu, A., and Yap, M. N. (2016) Ribosome hibernation factor promotes Staphylococcal
survival and differentially represses translation, *Nucleic Acids Research 44*, 4881-4893.
26. Yoshida, H., and Wada, A. (2014) The 100S ribosome: ribosomal hibernation induced by
stress, *Wiley Interdiscip Rev RNA 5*, 723-732.
27. Mandelstam, J., and Halvorson, H. (1960) Turnover of protein and nucleic acid in soluble
and ribosome fractions of non-growing Escherichia coli, *Biochimica et biophysica acta*
40, 43-49.
28. Kaplan, R., and Apirion, D. (1975) The fate of ribosomes in Escherichia coli cells starved
for a carbon source, *The Journal of biological chemistry 250*, 1854-1863.
29. Davis, B. D., Luger, S. M., and Tai, P. C. (1986) Role of ribosome degradation in the death
of starved Escherichia coli cells, *Journal of Bacteriology 166*, 439-445.
30. Deutscher, M. P. (2003) Degradation of stable RNA in bacteria, *The Journal of biological*
chemistry 278, 45041-45044.
31. Raina, M., and Ibba, M. (2014) tRNAs as regulators of biological processes, *Front Genet*
5.
32. Bullwinkle, T. J., and Ibba, M. (2016) Translation quality control is critical for bacterial
responses to amino acid stress, *Proceedings of the National Academy of Sciences of the*
United States of America 113, 2252-2257.
33. Jacob, A. I., Kohrer, C., Davies, B. W., RajBhandary, U. L., and Walker, G. C. (2013)
Conserved bacterial RNase YbeY plays key roles in 70S ribosome quality control and 16S
rRNA maturation, *Molecular Cell 49*, 427-438.

34. Davies, B. W., Kohrer, C., Jacob, A. I., Simmons, L. A., Zhu, J., Aleman, L. M., Rajbhandary,
U. L., and Walker, G. C. (2010) Role of Escherichia coli YbeY, a highly conserved protein,
in rRNA processing, *Mol. Microbiol. 78*, 506-518.
35. Inoue, K., Alsina, J., Chen, J., and Inouye, M. (2003) Suppression of defective ribosome
assembly in a rbfA deletion mutant by overexpression of Era, an essential GTPase in
Escherichia coli, *Mol. Microbiol. 48*, 1005-1016.
36. Verstraeten, N., Fauvart, M., Versees, W., and Michiels, J. (2011) The universally
conserved prokaryotic GTPases, *Microbiology and molecular biology reviews : MMBR 75*,
507-542, second and third pages of table of contents.
37. Vercruysse, M., Kohrer, C., Shen, Y., Proulx, S., Ghosal, A., Davies, B. W., RajBhandary, U.
L., and Walker, G. C. (2016) Identification of YbeY-Protein Interactions Involved in 16S
rRNA Maturation and Stress Regulation in Escherichia coli, *MBio 7*.
38. Paul, B. J., Barker, M. M., Ross, W., Schneider, D. A., Webb, C., Foster, J. W., and Gourse,
R. L. (2004) DksA: a critical component of the transcription initiation machinery that
potentiates the regulation of rRNA promoters by ppGpp and the initiating NTP, *Cell 118*,
311-322.
39. Krasny, L., and Gourse, R. L. (2004) An alternative strategy for bacterial ribosome
synthesis: Bacillus subtilis rRNA transcription regulation, *The EMBO journal 23*, 4473-
4483.
40. Hauryliuk, V., Atkinson, G. C., Murakami, K. S., Tenson, T., and Gerdes, K. (2015) Recent
functional insights into the role of (p)ppGpp in bacterial physiology, *Nature reviews.*
Microbiology 13, 298-309.
41. Yang, J., Yin, L., Lessner, F. H., Nakayasu, E. S., Payne, S. H., Fixen, K. R., Gallagher, L., and
Harwood, C. S. (2017) Genes essential for phototrophic growth by a purple
alphaproteobacterium, *Environmental Microbiology*.
42. Pechter, K. B., Gallagher, L., Pyles, H., Manoil, C. S., and Harwood, C. S. (2015) Essential
Genome of the Metabolically Versatile Alphaproteobacterium *Rhodopseudomonas*
palustris, *Journal of Bacteriology 198*, 867-876.
43. Dai, X., Zhu, M., Warren, M., Balakrishnan, R., Patsalo, V., Okano, H., Williamson, J. R.,
Fredrick, K., Wang, Y. P., and Hwa, T. (2016) Reduction of translating ribosomes enables
Escherichia coli to maintain elongation rates during slow growth, *Nat Microbiol 2*,
16231.
44. Gefen, O., Fridman, O., Ronin, I., and Balaban, N. Q. (2014) Direct observation of single
stationary-phase bacteria reveals a surprisingly long period of constant protein
production activity, *Proceedings of the National Academy of Sciences of the United*
States of America 111, 556-561.
45. Manina, G., Dhar, N., and McKinney, J. D. (2015) Stress and host immunity amplify
Mycobacterium tuberculosis phenotypic heterogeneity and induce nongrowing
metabolically active forms, *Cell Host Microbe 17*, 32-46.
46. Gefen, O., Gabay, C., Mumcuoglu, M., Engel, G., and Balaban, N. Q. (2008) Single-cell
protein induction dynamics reveals a period of vulnerability to antibiotics in persister
bacteria, *Proceedings of the National Academy of Sciences of the United States of*
America 105, 6145-6149.

47.   Stapels, D. A. C., Hill, P. W. S., Westermann, A. J., Fisher, R. A., Thurston, T. L., Saliba, A.
E., Blommestein, I., Vogel, J., and Helaine, S. (2018) Salmonella persisters undermine
host immune defenses during antibiotic treatment, *Science 362*, 1156-1160.
48.   Byrgazov, K., Vesper, O., and Moll, I. (2013) Ribosome heterogeneity: another level of
complexity in bacterial translation regulation, *Curr Opin Microbiol 16*, 133-139.
49.   Nguyen, D., Joshi-Datar, A., Lepine, F., Bauerle, E., Olakanmi, O., Beer, K., McKay, G.,
Siehnel, R., Schafhauser, J., Wang, Y., Britigan, B. E., and Singh, P. K. (2011) Active
starvation responses mediate antibiotic tolerance in biofilms and nutrient-limited
bacteria, *Science 334*, 982-986.
50.   Primm, T. P., Andersen, S. J., Mizrahi, V., Avarbock, D., Rubin, H., and Barry, C. E., 3rd.
(2000) The stringent response of Mycobacterium tuberculosis is required for long-term
survival, *Journal of Bacteriology 182*, 4889-4898.
51.   Dutta, N. K., Klinkenberg, L. G., Vazquez, M. J., Segura-Carro, D., Colmenarejo, G.,
Ramon, F., Rodriguez-Miquel, B., Mata-Cantero, L., Porras-De Francisco, E., Chuang, Y.
M., Rubin, H., Lee, J. J., Eoh, H., Bader, J. S., Perez-Herran, E., Mendoza-Losana, A., and
Karakousis, P. C. (2019) Inhibiting the stringent response blocks Mycobacterium
tuberculosis entry into quiescence and reduces persistence, *Sci Adv 5*, eaav2104.
52.   Cashel, M., Gentry, D. R., Hernandez, V. J., and Vinella, D. (1996) The stringent response,
In *Escherichia coli and Salmonella: cellular and molecular biology* (Neidhardt, F. C., Ed.),
pp 1458-1496, American Society for Microbiology Press, Washington, DC.
53.   Boutte, C. C., and Crosson, S. (2013) Bacterial lifestyle shapes stringent response
activation, *Trends Microbiol 21*, 174-180.
54.   Belitsky, B., and Kari, C. (1982) Absence of accumulation of ppGpp and RNA during
amino acid starvation in Rhizobium meliloti, *The Journal of biological chemistry 257*,
4677-4679.
55.   Krol, E., and Becker, A. (2011) ppGpp in Sinorhizobium meliloti: biosynthesis in response
to sudden nutritional downshifts and modulation of the transcriptome, *Mol. Microbiol.*
81, 1233-1254.
56.   Boutte, C. C., and Crosson, S. (2011) The complex logic of stringent response regulation
in Caulobacter crescentus: starvation signalling in an oligotrophic environment, *Mol.*
Microbiol. 80, 695-714.
57.   Acosta, R., and Lueking, D. R. (1987) Stringency in the absence of ppGpp accumulation in
Rhodobacter sphaeroides, *Journal of Bacteriology 169*, 908-912.
58.   Ronneau, S., Caballero-Montes, J., Coppine, J., Mayard, A., Garcia-Pino, A., and Hallez, R.
(2019) Regulation of (p)ppGpp hydrolysis by a conserved archetypal regulatory domain,
Nucleic Acids Research 47, 843-854.
59.   Ronneau, S., Petit, K., De Bolle, X., and Hallez, R. (2016) Phosphotransferase-dependent
accumulation of (p)ppGpp in response to glutamine deprivation in Caulobacter
crescentus, *Nat Commun 7*, 11423.
60.   Corrigan, R. M., Bellows, L. E., Wood, A., and Grundling, A. (2016) ppGpp negatively
impacts ribosome assembly affecting growth and antimicrobial tolerance in Gram-
positive bacteria, *Proceedings of the National Academy of Sciences of the United States*
of America 113, E1710-1719.

61.   Zhang, Y., Zbornikova, E., Rejman, D., and Gerdes, K. (2018) Novel (p)ppGpp Binding and
675   Metabolizing Proteins of Escherichia coli, *MBio 9*.
676   62.   Wang, B., Dai, P., Ding, D., Del Rosario, A., Grant, R. A., Pentelute, B. L., and Laub, M. T.
677   (2019) Affinity-based capture and identification of protein effectors of the growth
678   regulator ppGpp, *Nature chemical biology 15*, 141-150.
679   63.   Shan, Y., Brown Gandt, A., Rowe, S. E., Deisinger, J. P., Conlon, B. P., and Lewis, K. (2017)
680   ATP-Dependent Persister Formation in Escherichia coli, *MBio 8*.
681   64.   Fixen, K. R., Baker, A. W., Stojkovic, E. A., Beatty, J. T., and Harwood, C. S. (2014) Apo-
682   bacteriophytochromes modulate bacterial photosynthesis in response to low light,
683   *Proceedings of the National Academy of Sciences of the United States of America 111*,
684   E237-244.
685   65.   Quandt, J., and Hynes, M. F. (1993) Versatile suicide vectors which allow direct selection
686   for gene replacement in gram-negative bacteria, *Gene 127*, 15-21.
687   66.   McKinlay, J. B., and Harwood, C. S. (2010) Carbon dioxide fixation as a central redox
688   cofactor recycling mechanism in bacteria, *Proceedings of the National Academy of
689   Sciences of the United States of America 107*, 11669-11675.
690   67.   Liao, L., Schaefer, A. L., Coutinho, B. G., Brown, P. J. B., and Greenberg, E. P. (2018) An
691   aryl-homoserine lactone quorum-sensing signal produced by a dimorphic prosthecate
692   bacterium, *Proceedings of the National Academy of Sciences of the United States of
693   America 115*, 7587-7592.
694   68.   Lawrence, M. G., Shamsuzzaman, M., Kondopaka, M., Pascual, C., Zengel, J. M., and
695   Lindahl, L. (2016) The extended loops of ribosomal proteins uL4 and uL22 of Escherichia
696   coli contribute to ribosome assembly and protein translation, *Nucleic Acids Research 44*,
697   5798-5810.
698   69.   Janssen, B. D., Diner, E. J., and Hayes, C. S. (2012) Analysis of aminoacyl- and peptidyl-
699   tRNAs by gel electrophoresis, *Methods Mol Biol 905*, 291-309.
700   70.   Fang, M., and Bauer, C. E. (2018) Regulation of stringent factor by branched-chain amino
701   acids, *Proceedings of the National Academy of Sciences of the United States of America
702   115*, 6446-6451.
703   71.   Nakayasu, E. S., Burnet, M. C., Walukiewicz, H. E., Wilkins, C. S., Shukla, A. K., Brooks, S.,
704   Plutz, M. J., Lee, B. D., Schilling, B., Wolfe, A. J., Muller, S., Kirby, J. R., Rao, C. V., Cort, J.
705   R., and Payne, S. H. (2017) Ancient Regulatory Role of Lysine Acetylation in Central
706   Metabolism, *MBio 8*.
707   72.   Tyanova, S., Temu, T., and Cox, J. (2016) The MaxQuant computational platform for
708   mass spectrometry-based shotgun proteomics, *Nat Protoc 11*, 2301-2319.
709   73.   Cox, J., Hein, M. Y., Luber, C. A., Paron, I., Nagaraj, N., and Mann, M. (2014) Accurate
710   proteome-wide label-free quantification by delayed normalization and maximal peptide
711   ratio extraction, termed MaxLFQ, *Molecular & cellular proteomics : MCP 13*, 2513-2526.
712   74.   Schwanhausser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., and
713   Selbach, M. (2011) Global quantification of mammalian gene expression control, *Nature
714   473*, 337-342.
715
716

Figure 1. Metabolism of *R. palustris* during anaerobic growth in light. Acetate was used as the carbon source and ammonium as the nitrogen source for growth in this study. *R. palustris* generates ATP and proton motive force from light energy by cyclic photophosphorylation. Energy generation and carbon use are independent from each other. In this study growth arrest occurred when *R. palustris* had used all available acetate. In this circumstance cells can continue to generate ATP and proton motive force from light.

725
726
727
728

729
730

5μm

growing growth arrest D6

(μm)	Cell length	Min.	Max.
growing	2.5±0.6	1.2	3.8
growth arrest D6	2.6±0.6	1.1	4.4

731
732

734 **Figure 2.** Comparison of lengths of *R. palustris* cells from growing cultures and from cultures at

735 day 6 post growth arrest. Cells were photographed at 100X magnification with a Nikon Eclipse

736 80i digital microscope (n > 50).

Figure 3. Growth-arrested *R. palustris* cells do not appear to undergo cycles of death and
regrowth. a: Percentage of *R. palustris* cells that continue to maintain plasmid pBBR^Gm during
continuous growth (re-inoculated in medium with fresh carbon source repeatedly) or during
growth arrest (no treatment after the initial carbon source was depleted). Gm-free medium was
used for this experiment. b: Growth-arrested *R. palustris* is sensitive to antibiotics targeting
ribosomes. Antibiotics are added five days after growth arrest to independent cultures. Gm:
gentamycin. Cm: chloramphenicol. Nal: nalidixic acid. Ap: ampicillin. Error bars indicate the
standard deviation (n>3).

767
768

Figure 4. *R. palustris* protein abundances. Growing cells and growth-arrested cells are
compared. a: Log$_2$-fold changes in abundances of differentially abundant (p ≤ 0.05 by Student's
t-test) *R. palustris* proteins in growing cells (log), 6 days after growth arrest (Day 6) and 20 days
after growth arrest (Day 20). The proteome profiles of three samples (n = 3) for each condition
are shown (SI Table 1). b: The average relative quantities of 53 identified ribosome proteins at
day 6 (black, Day 6) and day 20 (red, Day 20) growth arrest relative to the growing cells (log).
The horizontal dashed line indicates the position of p = 0.05 (Student's *t*-test), and the vertical
dashed lines indicate the position of two-fold abundance changes. c: Ribosome proteins
compared in growing cells (log) and at day 6 and day 20 post growth arrest as % mass faction
of the proteome (SI Table 3). Error bars represent standard deviation. d: The average relative
quantities of selected proteins at day 6 (black, Day 6) and day 20 (red, Day 20) growth arrest
relative to the growing cells (log). The horizontal dashed line indicates the position of p = 0.05
(Student's *t*-test), and the vertical dashed lines indicate the position of two-fold abundance
changes. This panel includes proteins involved in tRNA charging (KEGG map00970, SI Table
2), translation elongation factors (SI Table 2) and two previously identified longevity proteins

(YbeY and Era). e: Translation elongation factors (SI Table 3) compared in growing cells (log)

and at day 6 and day 20 post growth arrest as % mass fraction of the proteome. Error bars

represent standard deviation.

Figure 5. Translation is intact and active in growth-arrested *R. palustris.* a: The ribosome profile
of cells at 6 days (Day 6) post growth arrest is shown and compared to ribosomes from growing
R. palustris (log). b: The ribosome profile of cells at 25 days (Day 25) post growth arrest is
shown and compared to ribosomes from growing *R. palustris* (log). c: Total RNA was purified
and analyzed from growing and growth-arrested *R. palustris* cells. The same amount of RNA
was loaded in each lane. The quantification of replicated experiments is shown in SI Figure 3.
The standard is the internal standard of the RNA ScreenTape system used (Agilent). d: The
charged and uncharged species of tRNA$_{trp}$ are shown here with a representative Northern blot.
The aminoacylated (charged) and deacylated (uncharged) species of tRNA$_{trp}$ were separated
with an acid urea gel and probed with a tRNA$_{trp}$ probe. e: Quantification of the charged and
uncharged species of tRNA$_{trp}$ as analyzed in Figure 5d. Error bars represent the standard
deviation (n>3).

Figure 6. YbeY and Era are important for ribosome maintenance during growth arrest. a:

Deletion of either *ybeY* or *era* had little effect on ribosomes from growing *R. palustris*, while the

ribosomes from growth-arrested cultures of *ybeY* and *era* deletion mutants lacked 30S subunits.

b: *R. palustris* total RNA was analyzed by Northern blot with a probe targeting 16S rRNA. In

growing cells, both Δ*ybeY* and Δ*era* mutants accumulated 17S rRNA but maintained substantial

levels of 16S rRNA. In growth-arrested cells, both 17S rRNA and 16S rRNA species were

depleted.

Figure 7. Overexpression of YbeY$_{D118R}$ complemented the longevity defect of a Δ*ybeY* mutant.
a: Longevity of different *R. palustris* strains. Deletion of the *ybeY* gene compromised the
longevity of growth-arrested cells (Δ*ybeY*), and this was complemented by expressing this gene
in trans (Δ*ybeY*::pBBP-YbeY) (*17*). A *ybeY*$_{D118R}$ mutant had a longevity defect similar to that of
the Δ*ybeY* deletion mutant. Overexpression of *ybeY*$_{D118R}$ *in trans*, restored the longevity of a
Δ*ybeY* mutant (Δ*ybeY*::pBBP-YbeY$_{D118R}$). Error bars indicate the standard deviation (n>3). b: *R.*
palustris total RNA was analyzed by Northern blot with a probe targeting 16S rRNA. In the
growth-arrested cells of Δ*ybeY*::pBBP-*ybeY*$_{D118R}$, only 16S rRNA appeared and no 17S rRNA
was detected. c: The ribosomes from Δ*ybeY*::pBBP-YbeY$_{D118R}$ were analyzed on a sucrose
gradient. In the growth-arrested cells, 30S, 50S and 70S ribosomes were clearly seen, though
the abundance of the 50S subunit is higher than that of WT.

838
839

Figure 8. Growth-arrested cells with very low levels of (p)ppGpp have compromised longevity.

a: The level of (p)ppGpp was low in exponentially growing *R. palustris* and increased during

growth-arrest. b: The domain structure of Rsh$_{rp}$ (top). Locations and relative quantities of Tn

insertions in Rsh$_{rp}$ (*17*). A ΔRsh$_{rp64-192}$ mutant was made by deleting the sequence encoding the

64th-192th amino acids of Rsh$_{rp}$ from *R. palustris.* c: (p)ppGpp synthesis was severely

compromised in the ΔRsh$_{rp64-192}$ strain. The time points shown correspond to the time points

shown in Figure 8a, representing early-mid log phase, mid-late log phase, entry into growth-

arrest and day 5 following growth-arrest. d: The longevity of Δrsh$_{rp64-192}$ was severely

compromised. Less than 10% of cells remained viable after 2 weeks of growth-arrest. Error bars

indicate the standard deviation (n>3).

850
851

SI Figure and Table Legends

SI Figure 1. The responses of *R. palustris* to various antibiotics at different growth stages.

Antibiotics are added to growing cultures or five days after growth-arrest. Cm: chloramphenicol.

Nal: nalidixic acid. Error bars indicate the standard deviation (n>3). For growing cells treated

with Nal, the number of viable cells was determined by A_{660} and a representative result is shown

(n > 3).

SI Figure 2. The *hirR* gene is expressed with a GDH (*rpa0944*) promoter and LacZ protein is

expressed with P_{hirl} promoter. *R. palustris* culture was grown to Day 6 of growth-arrest and the

PA-HSL signal was added into the culture. After two days, the samples were collected and the

level of LacZ was analyzed as stated in the Method session. With the signal added, the LacZ

level was ~70% higher comparing to the negative control.

SI Figure 3. rRNA levels in different *R. palustris* strains at different growth-stages. The

quantities of 16S, 17S and 23S rRNA are determined using an RNA ScreenTape system

(Agilent) and normalized against the levels of 23S rRNA in growing cells of wild-type *R.*

palustris. An unpaired *t* test was performed with at least three replicates (n.s.: not significant; *:

$p < 0.05$; **: $p < 0.01$; ***: $p < 0.005$; ****: $p < 0.001$, as determined by Student's *t*-test).

SI Figure 4. Trp-tRNATrp remained highly charged in the growth-arrested cells of Δ*ybeY* and

Δ*era* strains.

SI Table 1. Proteome of R. palustris in log phase and at 6 days and 20 days post growth arrest.

SI Table 2. Categories (KEGG) of proteins with changes in abundances in growth-arrested

compared to growing *R. palustris*.

SI Table 3. Relative mass % of individual *R. palustris* proteins compared to the whole cell

proteome in log phase cells and in cells at 6 days and 20 days post growth arrest.