The female urinary microbiota in relation to the reproductive tract microbiota

Chen Chen[1,2,5,*], Lilan Hao[1,2,*], Weixia Wei[3,4,*], Fei Li[1], Liju Song[1,2], Xiaowei Zhang[1,2], Juanjuan Dai[3,4], Zhuye Jie[1,2,6], Jiandong Li[1,2], Xiaolei Song[1], Zirong Wang[1], Zhe Zhang[1,2], Liping Zeng[3,4], Hui Du[3,4], Huiru Tang[3,4], Huanming Yang[1,7], Jian Wang[1,7], Karsten Kristiansen[1,5], Xun Xu[1,2], Ruifang Wu[4,5,†], Huijue Jia[1,2,6,8,†]

[1]BGI-Shenzhen, Shenzhen 518083, China.

[2]China National Genebank, BGI-Shenzhen, Shenzhen 518120, China.

[3]Peking University Shenzhen Hospital, Shenzhen 518036, China.

[4]Shenzhen Key Laboratory on Technology for Early Diagnosis of Major Gynecological diseases, Shenzhen, PR China.

[5]Department of Biology, Ole MaalØes Vej 5, University of Copenhagen, Copenhagen, Denmark.

[6]Shenzhen Key Laboratory of Human Commensal Microorganisms and Health Research, BGI-Shenzhen, Shenzhen, China.

[7]James D. Watson Institute of Genome Sciences, Hangzhou, China.

[8]Macau University of Science and Technology, Taipa, Macau 999078, China.

[*]These authors contributed equally to this work.

[†]Correspondence should be addressed to H. J. (jiahuijue@genomics.cn) or R. W. (wurf100@126.com).

E-mails and ORCIDs:

Chen Chen*. Email: chenchen1@genomics.cn, 0000-0001-6788-2894

Lilan Hao*. Email: haolilan@genomics.cn

Fei Li. Email: lifei1@genomics.cn, 0000-0002-8142-1100

Liju Song. Email: songliju@genomics.cn

Xiaowei Zhang. Email: zhangxiaowei@genomics.cn

Zhuye Jie. Email: jiezhuye@genomics.cn

Jiandong Li. Email: lijiandong@genomics.cn

Xiaolei Song. Email: songxiaolei@genomics.cn

Zirong Wang. Email: wangzirong@genomics.cn

Zhe Zhang. Email: zhangzhe4@cngb.org

Huanming Yang. Email: yanghm_old@genomics.cn

Jian Wang Email: wangjian@genomics.cn

Karsten Kristiansen. Email: kk@bio.ku.dk, 0000-0002-6024-0917

Xun Xu Email: xuxun@genomics.cn

Huijue Jia†. Email: jiahuijue@genomics.cn, 0000-0002-3592-126X

Weixia Wei*. Email: weixwei_2013@126.com

Juanjuan Dai. Email: djjsd001@126.com

Liping Zeng. Email: zengliping@163.com

Hui Du. Email: duhui_107108@163.com

Huiru Tang. Email: 13828857955@126.com

Ruifang Wu†. Email: wurf100@126.com

Abstract

Human urine is traditionally considered to be sterile, and whether the urine harbours distinct microbial communities has been a matter of debate. The potential link between female urine and reproductive tract microbial communities is currently not clear.

Here we collected the urine samples from 147 Chinese women of reproductive age, and explored the nature of colonization by 16S rRNA gene amplicon sequencing, real-time qPCR and live bacteria culture. To demonstrate utility intra-individual Spearman's correlation was used to explore the relationship between urine and multi-sites of the reproductive tract. PERMANOVA was also performed to explore potential correlations between the lifestyle and various clinical factors and urinary bacterial communities. Our data demonstrated distinct bacterial communities in urine, indicative of a non-sterile environment. Types of diverse, *Streptococcus*-dominated, and *Lactobacillus*-dominated were the three most common types in the cohort. Detailed comparison of the urinary microbiota to the multi-sites of reproductive tract microbiota demonstrated the urinary microbiota was more similar to the microbiota in the cervix and uterine cavity instead of vagina in the same women.

Our data demonstrates the potential connectivity of the microbiota in the female urogenital system and provided insight into the exploration of urethra and genital tract diseases.

Data Description

Purpose of data acquisition

The role of microbiota played in the vaginal environment has received a lot of attention over the past decade, while the female upper reproductive tract was traditionally believed to be sterile and mostly studied in the context of infections or incontinence [1]. Recently, Chen et al. has determined that the female upper reproductive tract is colonised by a microbiota that can be associated with lifestyle, pregnancy history and gynaecological diseases. Like the female upper reproductive tract [2], the sterile hypothesis of urine has also been overturned by emerging evidences that indicates the existence of microorganisms in the urinary tract by cultural or sequencing approaches [3,4]. A recent study by Thomas-White *et al.* (2018), using an expanded quantitative urine culture in combination with whole-genome sequencing, has isolated and genome

sequenced 149 bacterial strains from catheterized urine of both symptomatic and

asymptomatic peri-menopausal women, and also showed highly similar strains of

commensal bacteria in both the bladder and vagina of the same individual [5].

Gottschick *et al.* (2017) analysed the urinary microbiota of 189 individuals using 16S

rRNA gene amplicon sequencing, and suggested the urethra and bladder can harbor

microbial communities distinct from the vagina [6]. However, the relationship between

female urine microbiota and the upper reproductive tract microbiota has so for not been

studied.

Here we present a dataset of the urinary microbiota for a relatively large cohort of 147

women of reproductive age. Together with our recently published peritoneal fluid,

uterine and vaginal samples from the same individuals, this data shows that although

with greater amounts of *Lactobacillus* and *Streptococcus*, the urinary microbiota is

more similar to the microbiota of the cervix and uterine cavity, in accordance with the

anatomical opening of the bladder. Together with a wealth of collected metadata, we

demonstrate this data is useful to explore the potential of the urinary microbiota for

clinical diagnosis.

Methods

Sample collection

In this study, a total of 147 reproductive age women (age 22-48) were recruited by

Peking University Shenzhen Hospital (**Supplementary Table 1**). All participants were

reproductive age women who underwent hysteroscopy and/or laparoscopy for

conditions without infections, such as hysteromyoma, adenomyosis, endometriosis, or

salpingemphraxis. Subjects with other related diseases, such as vaginal inflammation,

severe pelvic adhesion, endocrine or autoimmune disorders were removed. Pregnant

women, breastfeeding women and menstrual women at sampling were also excluded.

None of the subjects received any hormone treatments, antibiotics or vaginal

medications within a month of sampling. In addition, no cervical treatment was

performed within the previous 7 days, no vaginal douching was performed within 5

days, and no sexual activity was performed within at least 2 days.

137 self-sampling morning mid-stream urine samples were collected between

December 2013 and July 2014 prior to the surgery (**Table S1**), and then stored at -80°C

until transported on dry ice to BGI-Shenzhen for sequencing. The samples from an

additional 10 women were collected for validation purposes by a doctor during the

surgery in July 2017. For each operation, a urine catheter was inserted into the urethra

to collect mid-stream urine, and same amount of sterile saline which collected through

the urine catheter was set as control samples. The samples were then placed at 4°C,

transported to BGI-Shenzhen, and processed within 6 hours. A portion of each sample

was used for culturing live bacteria and the rest was used for sequencing.

DNA extraction of urine samples

Genomic DNA extraction was carried out following the steps: TE buffer, lysozyme

solution, proteinase K, 10% SDS were added to each urine sample, and vortexed to mix.

Sample was then incubated at 55 °C in a shaking water bath for 120 min to effect

complete lysis. Added phenol-chloroform-isoamylalcohol (25:24:1) to the supernatant

and vortexed to mix well. Then the supernatant was transferred to a nuclease-free

microfuge tube, added chloroform-isoamylalcohol (24:1) and vortexed to mix well.

After the supernatant was transferred to a nuclease-free microfuge tube, sodium acetate,

glycogen and cold isopropanol were added to the supernatant, and then the mixture was

incubated at -20 °C for the night. 70% ethanol was used to wash the precipitate twice.

TE buffer was then added to resuspend the precipitate. Extracts were treated with

DNase-free RNase to eliminate RNA contamination. DNA quantity was determined

using NanoDrop spectrophotometer, Qubit Fluorometer (with the Quant-iTTMdsDNA

BR Assay Kit), and gel electrophoresis.[7]

16S rRNA amplicon sequencing

The primers 515F and 907R were utilized for PCR amplification of the hypervariable

regions V4-V5 of the bacterial 16S rRNA gene. The 907R primer include a unique

barcoded fusion. The primers were as follow: 515F:

5'-GTGCCAGCMGCCGCGGTAA- 3' and 907R: 5'- CCGTCAATTCMTTTRAGT

-3', where M denotes A or C and R denotes purine. The condition of PCR amplification

was as follows: 3 min of denaturation at 94°C, followed by 25 cycles of 45 s at 94 °C

(denaturing), 60 s at 50 °C (annealing), 90 s at 72 °C (elongation), with a final

elongation for 10 min at 72 °C. The amplification products were purified by the

AxyPrep^{TM} Mag PCR Clean-Up Kit (Axygen, USA). The amplicon libraries were

constructed with an Ion Plus Fragment Library Kit (Thermo Fisher Scientific Inc.) [8],

then sequenced by Ion PGM^{TM} Sequencer with Ion 318^{TM} Chip v2 with a read length of

400bp (Thermo Fisher Scientific Inc., Ion PGM™ Hi□Q™ OT2 Kit, Cat.No: A27739;

Ion PGM™ Hi-Q ™ Sequencing Kit, Cat.No: A25592)[9].

Processing of sequencing reads

The raw sequencing reads were firstly subjected to Mothur (Mothur,

RRID:SCR_011947;V1.33.3) [10] for filtering of the low-quality reads utilizing the

following criteria: 1) reads with less than 200bp; 2) reads not matching the degenerated

PCR primers for up to two errors; 3) reads with an average quality score less than 25. A

total of 8,812,607 reads, with an average of 57,225 reads per sample (a minimum of

1,113 reads and a maximum of 194,564 reads) were obtained. Subsequently, the

sequences with identity greater than 97% were clustered into Operational Taxonomic

Units (OTUs) using QIIME (QIIME, RRID:SCR_008249; V1.8.0) uclust programme

[11], where each cluster was thought of as representing a species. The seed sequences

of each OTU were aligned against the Greengene reference sequences (gg_13_8_otus)

for annotation using the UCLUST taxonomy assigner.

We also calculated the Unifrac distance using QIIME based on taxonomic abundance

profiles at the OTU level [11].

PERMANOVA on the influence of phenotypes

160 Permutational multivariate analysis of variance (PERMANOVA) was used to assess

161 the effect of different covariate based on the relative abundances of OTUs of the

162 samples [12,13] using Bray-Curtis and UniFrac distance and 9999 permutations from

163 vegan package(vegan, RRID:SCR_011950) in R [13,14].

Real-time quantitative PCR

We quantified the four *Lactobacillus* species, including *L.iners*, *L jensenii*, *L. crispatus*

and *L. gasseri* using the modified real-time quantitative PCR [15]. SYBR Premix Ex

Taq GC (TAKARA) was applied to run on StepOnePlus Real-time PCR System (Life

Technologies). PCR reaction mixture contained 10 μl of 2×SYBR Premix Ex Taq GC,

0.2 μM forward primer, 0.2 μM reverse primer, 1.6 μl of DNA sample and 8.2 μl

ultrapure water to make up the final reaction volume of 20 μl. Each run included a

standard curve and all samples were amplified in triplicate. Ultrapure water was

included as blank control template.

To construct the standard curves, the sequencing-confirmed plasmids of four species

were used after quantification with Qubit Fluorometer and serial 10-fold dilutions. The

amplification efficiency was (100±10) % and linearity values were all ≥0.99.

Bacterial culturing

The urine samples and controls from additional 10 subjects were cultivated in the

laboratory by spreading 100 μl of sample on the different agars with 5% horse blood,

such as PYG agar (DSMZ 104 medium), BHI agar and EG agar. The plates were

incubated in both aerobic and anaerobic conditions at 37 °C for 72 hours. To keep the

medium anaerobically during culture, resazurin and cysteine-HCl were added as

reducing agents. The genomic DNA of the isolates was extracted by the Bacterial DNA

Kit (OMEGA), and then 16S rRNA gene amplification by the universal primers

27F/1492R were performed [16]. The amplicons were purified and sent for Sanger

sequencing. The generated sequences were then submitted to BLAST on the

EzBioCloud [17] for identification.

Preliminary analysis and validation

Microbiota composition of the urine

To explore the urinary microbiota in this dataset, morning midstream urine (UR) was

self-collected prior to surgery from an exploratory study cohort of 137 Chinese women

recruited (median age 31.6, range 22-48). As with our previous vagino-uterine

microbiota study [2], all volunteers had conditions that were not known to involve

infections (**Table S1**). From 95 women in the cohort, six locations within the female

reproductive tract, including lower third of vagina (CL), posterior fornix (CU), cervical

mucus (CV), endometrium (ET), left and right fallopian tubes (FLL and FRL), and

peritoneal fluid (PF) were also sampled. Their vagino-uterine microbiota information

has been published previously [2]. After 16S rRNA gene amplicon sequencing, the

sequencing reads were pre-processed for quality control and filtering, then clustered

into operational taxonomic units (OTUs) (**Methods**, **Table S2, S3**).

Due to the anatomical structures, voided urine samples from women were considered to

be easily contaminated by the microbiota from the surrounding vulvovaginal region

[18]. Most vaginal communities (88%) in this cohort were dominated by one genus that

constitutes >50% relative abundance in the individuals. In contrast, the urinary

microbiota in this study showed more heterogeneity. 56.93% of the cohort harboured a

diverse type, which was represented by a significant part of the bacteria, including

Streptococcus, *Lactobacillus*, *Pseudomonas*, *Staphylococcus*, *Acinetobacter* and

Vagococcus, while dominated (>50%) by none of them (**Fig. 1**). In addition, 22.63% of

the women harboured over 50% *Streptococcus*, and 13.87% of the women

contained >50% *Lactobacillus* (**Fig. 1a, Fig. 1b**). Rare subtypes such as *Enterococcus*

(2.19%), Bifidobacteriaceae (1.46%), *Prevotella* (0.73%), Enterobacteriaceae (0.73%),

Coriobacteriaceae (0.73%), and *Veillonella* (0.73%) were also detected in this cohort

(**Fig. 1a, Fig. 1b**). Notably, the median relative abundance of *Lactobacillus*,

Pseudomonas and *Acinetobacter* in the urine samples were more similar to the uterus

samples (**Fig. 1c**) [2]. At the phylum level, urinary microbiota was dominated by

Firmicutes and Proteobacteria (**Fig. 1c**).

Intra-individual similarity in the urine- reproductive tract microbiota

To further assess the microbiota relationship between the urine and the six positions of the reproductive tract, we computed the intra-individual correlation of microbial profile between the urine and different sites of the reproductive tract and clustered the individuals into 4 groups (Spearman correlation coefficient, **Fig. 2a, Table S4**). Interestingly, the microbiota of group 3, which accounted for 41% of the cohort, showed the significance of correlation with urine samples increased from CL to CV, ET and PF gradually. In contrast, 9% women in group 1 presented a reverse trend. In group 2 (22%) and group 4 (27%), there appeared to be a weak relationship between the urine and reproductive tract. Taken together, we observed the most similar distribution of microbiota between urine and CV/ET (**Fig. 2b**). The principal coordinate analyses (PCoA) of the weighted and unweighted intra-individual UniFrac distance further corroborated our results that there is an intra-individual similarity of the microbiota between the urine and the upper sites of reproductive tract, especially the junction sites (CV and ET) (**Fig. 3**).

Lifestyle and clinical factors influencing the urinary microbiota

The human microbiome is dynamic and highly affected by its host environment. Age, menstrual cycle, benign conditions such as adenomyosis, and infertility due to endometriosis have previously been reported to shape the vagino-uterine microbiota [2]. With our comprehensive collection of demographic and baseline clinical characteristics from women of reproductive age (**Table S1**), such variations in the urinary microbiota can be explored in this dataset. Urinary microbial composition was significantly associated with these factors, such as age, surgical history, abortion, vaginal deliveries, experience of given birth (multipara vs. nullipara), infertility due to endometriosis and hysteromyoma (PERMANOVA, $P < 0.05$, $q < 0.05$, **Table 1**). Although the urinary microbiota also correlated with some other factors, such as menstrual phase, contraception, endometriosis, pelvic adhesiolysis and anemia, the statistical significance was not achieved after controlling for multiple testing (PERMANOVA, $P < 0.05$ but $q > 0.05$, **Table 1**). The initial results here indicate a close link between the urinary microbiota with the general and diseased physiological conditions, and can be taken further with further exploration of this data.

Potential uses

As the large-scale cohort for studying the female urinary microbiota, our data provide a

useful baseline and reference dataset in women of reproductive age. We also explored

the association between the composition of urinary microbiota and that of reproductive

tract microbiota. It is valuable to note that a higher intra-individual compositional

similarity of the microbiota between urine and cervical canal/uterus instead of the

vagina, which indicate that sampling of midstream urine (the least invasive and the

easiest way) could be potentially used to survey the micro-environment of the cervical

canal and uterus in the general population. This is of relevance in relation to the

demonstrated associations between the urinary microbiota and various uterine-related

diseases such as hysteromyoma, and infertility due to endometriosis. Our data provide a

reference for clinical diagnosis, and warrants further detailed exploration. Thus, we

hope this dataset will be the beginning of a new round of accelerated discoveries,

including a novel scientific explanation of the uterine-related diseases by longitudinal

studies on the microbiota of urinary and reproductive tract.

Declarations
265

Ethics approval and consent to participate

The study was approved by the Institutional Review Board of BGI-Shenzhen and

Peking University Shenzhen Hospital. All participants gave written informed consent

prior to their recruitment into the study.

Availability of supporting data

The sequence reads generated by 16S rRNA gene amplicon sequencing have been

deposited in both the European Nucleotide Archive with the accession number

PRJEB29341 and the CNSA (https://db.cngb.org/cnsa/) of CNGB database with

accession code CNP0000166. The sequences of bacterial isolates have been deposited

in the European Nucleotide Archive with the accession number PRJEB36743

Acknowledgments

The study was supported by the Shenzhen Municipal Government of China

(JCYJ20160229172757249 and JCYJ20170817145523036) and Shenzhen Peacock

Plan (No. KQTD20150330171505310). The authors really appreciate colleagues at

BGI-Shenzhen for DNA extraction, library construction, and sequencing.

Author contributions

Competing interests

There were no competing financial interests.

References

1. Whiteside S a., Razvi H, Dave S, Reid G, Burton JP. The microbiome of the urinary

tract - A role beyond infection. Nat. Rev. Urol. Nature Publishing Group; 2015. p.

81–90.

2. Chen C, Song X, Wei W, Zhong H, Dai J, Lan Z, et al. The microbiota continuum

along the female reproductive tract and its relation to uterine-related diseases. Nat

Commun [Internet]. Springer US; 2017;8. Available from:

http://dx.doi.org/10.1038/s41467-017-00901-0

3. Siddiqui H, Nederbragt AJ, Lagesen K, Jeansson SL, Jakobsen KS. Assessing

diversity of the female urine microbiota by high throughput sequencing of 16S rDNA

amplicons. BMC Microbiol. BioMed Central; 2011;11:244.

4. Wolfe AJ, Toh E, Shibata N, Rong R, Kenton K, FitzGerald MP, et al. Evidence of

uncultivated bacteria in the adult female bladder. J Clin Microbiol. American Society

for Microbiology; 2012;50:1376–83.

5. Thomas-White K, Forster SC, Kumar N, Van Kuiken M, Putonti C, Stares MD, et al.

Culturing of female bladder bacteria reveals an interconnected urogenital microbiota.

Nat Commun. Nature Publishing Group; 2018;9:1557.

6. Gottschick C, Deng ZL, Vital M, Masur C, Abels C, Pieper DH, et al. The urinary

microbiota of men and women and its changes in women during bacterial vaginosis and

antibiotic treatment. Microbiome. BioMed Central; 2017;5:99.

7. DNA extraction for human microbe samples. Available from:

http://dx.doi.org/10.17504/protocols.io.bcmriu56

8. Prepare Amplicon Libraries without Fragmentation Using the Ion Plus Fragment

Library Kit. Available from:

https://assets.thermofisher.com/TFS-Assets/LSG/manuals/MAN0006846_PrepAmpli

conLibr_using_IonPlusFragLibraryKit_UB.pdf

9. Ion Personal Genome Machine^{TM} (PGM^{TM}) System REFERENCE GUIDE.

Available from:

https://assets.thermofisher.com/TFS-Assets/LSG/manuals/MAN0009783_IonPGM_R

efGuide.pdf

10. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al.

Introducing mothur: Open-source, platform-independent, community-supported

software for describing and comparing microbial communities. Appl Environ

Microbiol. 2009;75:7537–41.

11. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et

al. QIIME allows analysis of high-throughput community sequencing data. Nat

Methods. 2010;7:335–6.

12. Anderson MJ. A new method for non-parametric multivariate analysis of variance.

Aust Ecol. 2001;26:32–46.

13. Feng Q, Liang S, Jia H, Stadlmayr A, Tang L, Lan Z, et al. Gut microbiome

development along the colorectal adenoma-carcinoma sequence. Nat Commun

[Internet]. 2015;6:6528. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/25758642

14. Zapala MA, Schork NJ. Multivariate regression analysis of distance matrices for

testing associations between gene expression patterns and related variables. Proc Natl

Acad Sci U S A. 2006;103:19430–5.

15. De Backer E, Verhelst R, Verstraelen H, Alqumber M a, Burton JP, Tagg JR, et al.

Quantitative determination by real-time PCR of four vaginal Lactobacillus species,

Gardnerella vaginalis and Atopobium vaginae indicates an inverse relationship

between L. gasseri and L. iners. BMC Microbiol. 2007;7:115.

16. Augustinos AA, Kyritsis GA, Papadopoulos NT, Abd-Alla AMM, C??ceres C,

Bourtzis K. Exploitation of the medfly gut microbiota for the enhancement of sterile

insect technique: Use of Enterobacter sp. in larval diet-based probiotic applications.

PLoS One. 2015;10:1–17.

17. EzBioCloud [Internet]. Available from: https://www.ezbiocloud.net

18. Karstens L, Asquith M, Caruso V, Rosenbaum JT, Braun J, Gregory T, et al. HHS

Public Access. 2019;15:735–49.

Figure legends

Fig. 1. Urinary microbiota of the initial cohort of 137 Chinese reproductive-age women. (a) The relative abundances of genera detected in each individual are shown in the bar chart. The dendrogram is a result of a centroid linage hierarchical clustering based on Euclidean distances between the microbial composition proportion of urinary bacterial communities. (b) The ratio of different urinary microbiota types. The genus whose relative abundance account for >50% in an individual is selected as an identified type. The genera all account <50% of the microbiota in an individual are identified as diverse type. (c) Pie chart for the urinary microbial genera according to the median relative abundance. Genera that took up less than 1% of the microbiota were labelled together as 'others'. The outer-ring indicates the distribution of microbiota in phylum level (**Table S1**).

Fig. 2. Similarity of the urine-reproductive tract microbiota within individuals.

(**a**) Heatmap for the intra-individual Spearman's correlation coefficient between urine and different body sites of reproductive tract (Table S4). Samples derive from the initial cohort of 95 Chinese reproductive-age women, who collected both the urine and reproductive tract samples. As the number of samples from fallopian tubes (FLL, FRL) is too small, the correlations between urine and fallopian tubes are not shown. The dendrogram is a result of a centroid linage hierarchical clustering based on Euclidean distances between the intra-individual Spearman's correlation coefficient of different body sites. The colored squares illustrate the subtypes of urinary microbiome. (**b**) Spearman's correlation coefficient between urine and different body sites of reproductive tract. Wilcoxon ranked sum test is used to calculate the difference. Boxes denote the interquartile range (IQR) between the first and third quartiles (25th and 75th percentiles, respectively), and the line inside the boxes denote the median. The whiskers denote the lowest and highest values within 1.5 times the IQR from the first

and third quartiles, respectively. An asterisk denotes p <0.05, two asterisks denote p <0.01, three asterisks denote p <0.001.

Figure 3. PCoA on the samples based on Unweighted-UniFrac (a) and Weighted-UniFrac (b) distances. Samples were taken from UR, CL, CU and CV before operation, and from ET and PF during operation. Samples derive from the initial cohort of 137 Chinese reproductive-age women. Each dot represents one sample (n□=□94 CL, 95 CU, 95 CV, 80 ET, 93 PF, 9 FLL, 10 FRL and 137 UR).