1

Niche complementarity among pollinators increases

community-level plant reproductive success

Ainhoa Magrach[1,2], Francisco P. Molina[3], Ignasi Bartomeus[3]

Short running title: Network structure effects on ecosystem functioning

[1]Basque Centre for Climate Change-BC3, Edif. Sede 1, 1º, Parque Científico UPV-EHU,

Barrio Sarriena s/n, 48940, Leioa, Spain

[2]IKERBASQUE, Basque Foundation for Science, María Díaz de Haro 3, 48013, Bilbao,

Spain

[3]Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, Isla de la

Cartuja, 41092, Sevilla, Spain

Type of article: Letter

Words in Abstract: 170 words

Words in main text (excluding abstract, acknowledgements, references, table and figure

legends): 5,302

Number of references: 60

Number of figures, tables and text boxes: 7

Corresponding author: ainhoa.magrach@bc3research.org, nacho.bartomeus@gmail.com

Keywords: nestedness, niche complementary, fruit set, pollination, plant-pollinator

interactions

Abstract

Declines in pollinator diversity and abundance have been reported across different

regions, with implications for the reproductive success of plant species. However,

research has focused primarily on pairwise plant-pollinator interactions, largely

overlooking community-level dynamics. Here, we present one of the first efforts linking

pollinator visitation to plant reproduction from a community-wide perspective using a

well-replicated dataset encompassing 16 well-resolved plant-pollinator networks and data

on reproductive success for 19 plant species from Mediterranean shrub ecosystems. We

find that models including simple visitation metrics are sufficient to explain the

variability in reproductive success observed. However, insights into the mechanisms

through which differences in pollinator diversity translate into changes in reproductive

success require additional information on network structure. Specifically, we find a

positive effect of increasing niche complementarity between pollinators on plant

reproductive success. This shows that maintaining communities with a diversity of

species but also of functions is paramount to preserving natural ecosystems.

Introduction

Pollinators provide key services to plants by facilitating pollen flow between individuals. The recent declining trends found for some pollinator species in some regions of the planet (Potts *et al.* 2010, Bartomeus *et al.* 2019) have led many researchers to focus on the functional impacts of these changes in pollinator diversity, with a major focus being placed on the consequences for plant reproductive success (Biesmeijer *et al.* 2006).

Many research efforts have targeted the reproductive success of individual plant species (Albrecht *et al.* 2012; Thomson 2019), and used relatively simple visitation metrics (e.g., the number of pollinator species that visit a plant or the number of visits they perform) to explain the differences observed across different plant individuals. Contrastingly, community-level analyses remain scarce (Bennett *et al.* 2018). Yet plants and pollinators do not interact in isolation, but rather are embedded within larger networks of interactions encompassing other plant and pollinator species. We are thus missing an important part of the picture, which includes the direct interactions between the whole ensemble of plants and pollinators, but also the indirect interactions between species within one guild (e.g., plants) through their shared resources (Pauw 2013; Carvalheiro *et al.* 2014; Lázaro *et al.* 2014; Mayfield & Stouffer 2017; Johnson & Bronstein 2019). Understanding how changes in pollinator diversity and interaction structure affect whole community functioning is thus a major challenge that requires attention.

The few pollination studies that have analysed the effects of changing pollinator diversity for reproductive success at the community level have done so using mainly experimental setups. As an example, a study that experimentally recreated a plant

community with 9 plant species and differing levels of pollinator diversity across

different enclosures, found that not only pollinator species diversity had an effect for

average reproductive success, but that plant-pollinator interaction structure also had an

important effect (Fründ *et al.* 2013). In particular, these authors found that niche

complementarity between pollinators, in terms of plant species and temperature coverage

(a measure of the overlap in the use of plant resources and optimum temperature activity)

had a positive effect for average seed set at the community level (Fründ *et al.* 2013). This

provides added information on the effects of biodiversity for ecosystem functioning,

suggesting that not only the diversity of species present, but also the diversity of roles and

ways in which a community is structured, are determinant factors.

Indeed, theoretical research has long suggested that the structure of multitrophic

communities has an effect for ecosystem functioning (reviewed in Thompson *et al.*

2012). This line of research, rooted in niche theory and revamped by food-web studies

(MacArthur & Levins 1967; May & MacArthur 1972, Tilman 1982, Godoy *et al* 2018),

has greatly advanced theory, but these ideas have not yet been tested using empirical data

(but see Poisot *et al.* 2013). Specifically, a major knowledge gap resides in understanding

which aspects of structure determine which aspects of function (Thompson *et al.* 2012).

This is because although a network perspective has promised to encapsulate complex

ecological mechanisms occurring at the community level – such as indirect interactions

(Holt 1977, Abrams *et al* 1998) or niche overlap (Woodward & Hildrew 2002)- less

attention has been given to the ways in which these mechanisms relate to observed

ecosystem processes (Blüthgen 2010). In contrast, we are now at a point in which there is

considerable understanding on the attributes characterizing mutualistic interaction

networks (Bascompte & Jordano 2007). Especially, in the case of pollination, we have

ample knowledge on the attributes that shape these mutualistic interactions at the

community level. Amongst them is the prevalence of nested structures, i.e., arrangements

where specialist species interact with a subset of the species that generalists interact with

(Bascompte *et al* 2003) and which is thought to promote species diversity (Bastolla *et al*

2009); or the relatively high extent of complementary specialization at the community

scale, which may be directly related to key ecosystem functions (Blüthgen & Klein

2011). However, the mechanisms by which these attributes affect plant reproduction

remain to be understood (Winfree 2013). The time is thus ripe to use the existing

knowledge around plant-pollinator network structures to explore the relationship between

network structure and ecosystem functioning empirically, with special emphasis being

placed on the underlying ecological mechanisms that drive these relationships.

Here, we present one of the first efforts linking pollinator visitation and plant

reproductive success at the community level using empirical data on plant-pollinator

interaction networks and plant reproductive success. To this end, we use a well-replicated

dataset encompassing 16 well-resolved plant-pollinator interaction networks coupled with

data on the reproductive success of 19 plant species recorded in Mediterranean shrub

ecosystems. Our study focuses on understanding whether adding information on selected

interaction network structure indices to previously used simple visitation metrics (e.g.,

the number and diversity of pollinator species visiting a plant species) aids in better

explaining the differences observed in community-wide reproductive success. In doing

so, we conducted our analyses focusing on reproductive success at two different levels:

(i) at the species level by considering the effect of the position of a focal species within

the larger network and its impact on its individual reproductive success, and (ii) at the site

level, by evaluating how attributes that describe the whole site might affect average

values of reproductive success. Specifically, our study focuses on how the interplay

between niche complementarity and redundancy determines reproductive success. Plant

reproductive success requires the delivery of conspecific pollen and thus of a certain

degree of niche complementarity (Blüthgen & Klein 2011). Yet, greater values of

nestedness, which imply redundancy in species functions, are thought to promote species

diversity (Bastolla *et al.* 2009) and stability (Thébault & Fontaine 2010) within plant-

pollinator networks. At present, we do not know how either of these network

characteristics affects the functions performed by pollinators. Finally, in addition to

average values, we also evaluate whether network structure helps explain differences in

equity in reproductive success across species within a community, as a measure of

evenness in the pollination service delivered.

Our results suggest that models including information on simple visitation metrics

alone are good in explaining the variability observed in reproductive success. However,

insights into the mechanisms through which differences in pollinator diversity translate

into changes in reproductive success require additional information on network structure,

notably information on the complementarity between the functions performed and the

niches occupied by different pollinator species. Specifically, we find a positive effect of

increasing niche complementarity between pollinators on plant reproductive success.

Methods

Plant pollinator interactions

Our study was conducted in SW Spain within the area of influence of Doñana National

Park, i.e., within the limits of the Natural Space of Doñana as defined by the local

government (Junta de Andalucía, Fig. 1). All sites were located within similar elevations

(ranging from 50 to 150 m a.s.l.), similar habitat and soil types, and presented similar

plant composition (plant mean Sørensen beta-diversity among sites = 0.41), reducing

potential confounding factors. Here, we surveyed 16 Mediterranean woodland patches

with an average distance of 7 km between them (min= 3 km, max= 46.5 km). Each site

was surveyed 7 times during the flowering season of 2015 (from February to May)

following a 100-m x 2 m transect for 30 mins. Along each transect, we identified all plant

species and recorded all the floral visitors that landed on their flowers and touched the

plant´s reproductive parts during each 30-min period. Only floral visitors (from now on

referred to as pollinators) that could not be identified in the field were captured, stored

and identified in the laboratory by FPM and experts in the different taxonomic groups

(see acknowledgements). In addition, at each round we conducted 3 minutes of focal

observations recording all floral visitors observed on 3 plant individuals per species

belonging to the 19 most common (based on previous surveys) plant species across the

study area (mean $\pm$ SD: 6.25 $\pm$ 1.73 species per site). Furthermore, we included some

interactions between plant and pollinator individuals that were not observed during the

sampling but that were opportunistically recorded immediately before or after the

sampling periods, as some of these interactions are difficult to document and might be

important to define network structure (Jordano 2016). These opportunistic interactions

represented 22.96% of all interactions recorded. All surveys were done under similar

weather conditions, avoiding windy or rainy days. Surveys were done during mornings

and afternoons with the sampling order being established randomly.

Plant reproductive success

Within each site, we marked between 3 and 12 individuals (mean $\pm$ SD: 6.49 $\pm$ 2.37,

Table S2) belonging to 1 to 6 plant species (mean $\pm$ SD: 4.06 $\pm$ 1.69, Table S1),

depending on the availability and presence of flowers during the sampling events. For

each individual, at the end of the season, we recorded fruit set (i.e. the proportion of

flowers that set fruit), the average number of seeds per fruit and the average fruit and

seed weight per fruit (1-36 fruits subsampled; mean $\pm$ SD: 11.17 $\pm$ 6.85, Table S3). Our

survey included a total of 19 different plant species across our 16 sites. Plants species

were selected based on their availability, with sampling being focused on the most

abundant plant species. The values at the species level were then averaged per site to

calculate unique reproductive success measures at the site level. All plant species depend

on pollinators to maximize their reproduction (Table S4).

Data analyses

In order to evaluate the completeness of our sampling of the pollinator and plant

community as well as that of their interactions, we estimated the asymptotic number of

species present (Chao *et al.* 2009), a non-parametric estimator of species richness for

abundance data. This estimator includes non-detected species and allowed us to calculate

the proportion detected with our original sampling data. We used Chao 1 asymptotic

species richness estimators (Chao *et al.* 2009) and estimated the richness of pollinators,

plants and plant–pollinator links accumulated as sampling effort increased up to 100%

sampling coverage using package iNEXT (Hsieh *et al.* 2016) within the R environment

(R Development Core Team 2011). We then extracted the values covered by our

sampling.

In order to analyse how differences in network structure might affect plant

reproductive success, we constructed plant-pollinator interaction networks by pooling the

data for the 7 rounds of sampling. We thus obtained one interaction network per site,

representing the number of individuals of different pollinator species recorded visiting

each different plant species. For each network, we then proceeded to extract a series of

relevant network metrics at the species and site levels.

In addition, we checked for potential spatial autocorrelation in our data by means

of Mantel correlograms. Autocorrelation values were low for all variables included in our

analyses (Figure S1) and hence we treat each site as independent in our analysis.

Species-level network analysis

At the species level, we focused on attributes defining the position of a focal plant species

within the larger community. As such, we considered two metrics providing

complementary non-redundant information: (i) average niche overlap in terms of

pollinators between a focal plant species and each of the other plant species in the

community, which estimates the potential indirect interactions between different plant

species through shared resources (in this case pollinators), and (ii) centrality, which

depicts the importance of the role played by a plant species within the larger community

(as resource for a large number of pollinator species) and its contribution to network

cohesiveness.

Niche overlap was calculated as the average overlap in pollinator species visiting

a focal plant and each of the other plants in the community using the Morisita overlap

index, a measure of similarity between two sets of data (Zhang 2016). As a measure of

centrality, we used weighted closeness centrality, which represents the number of shortest

paths going through a focal plant based on a unipartite projection of the bipartite plant-

pollinator network using a weighted representation of the network (Dormann *et al.* 2009).

With this method, links between plant species represent shared pollinator species.

Site-level network analysis

At the site level, we followed the same logic as the one presented at the species level.

Thus, we also calculated two network metrics providing complementary non-redundant

information. In this case we focused on (i) nestedness and (ii) pollinator niche

complementarity.

Nestedness is the property by which specialists interact with a subset of the

species that generalists interact with (Bascompte *et al.* 2003). Although there is an

ongoing debate in the literature, some studies have found that nested networks are more

stable and resilient to perturbations because nestedness promotes a greater diversity by

minimizing competition among species in a community (Bastolla *et al.* 2009). However,

many network attributes vary with network size and complexity (Blüthgen *et al.* 2006). In

the case of nestedness, we know it can be affected by network size and connectance

(Song *et al.* 2017). An approach that is often used to correct for this, is to use null models

and to compare null-model corrected nestedness values across different networks.

However, this approach has been recently shown to present the same issues, as z-scores

also change with network size and connectance (Song *et al.* 2017). We thus followed the

advice provided by Song et al. (2017) by using a normalized value of the widely-used

nestedness metric NODF (Almeida-Neto & Ulrich 2011), $NODF_c$. This normalized value

is calculated as $NODF_c = NODF_n/(C * log(S))$, where C is connectance and S is

network size. $NODF_n$ is calculated as $NODF/max(NODF)$, which is independent of

network size and thus comparable across different networks (Song *et al* 2017). To

calculate max(NODF) we used a recently corrected version of the algorithm (Simmons *et*

al 2019) in all but three sites, where the condition that the number of links>number of

species was not met and thus precluded us from using this new version. Results did not

change qualitatively when using the uncorrected version of the algorithm for all sites.

Niche complementarity metrics are important because plant reproductive success

depends on the delivery of conspecific pollen and thus of a certain level of specialization

or niche divergence (reviewed in Brosi 2016). To calculate niche complementarity, we

used a community-level measure defined as the total branch length of a functional

dendrogram based on qualitative differences measured using a Euclidean distance in

visitor assemblages between plants (Petchey & Gaston 2007; Devoto *et al.* 2012). All

network metrics were calculated using package bipartite (Dormann *et al.* 2009).

All of these metrics were calculated using all the data as well as for the subset of

the data excluding interactions observed outside of sampling periods. Differences

between results are minimal for both and thus we will only present results for the analysis

using the full dataset (see Table S12A-H for results removing observations out of

transect).

Statistical analyses

In order to evaluate whether adding information on network structure improves our

ability to explain differences in reproductive success - both at the species and the site

level - we used generalized linear (GLMs) and generalized linear mixed models

(GLMMs) respectively. In both cases (species and site-level models) we fit two types of

models: (i) model 1, that only included simple visitation metrics and (ii) model 2 that

additionally included information on network structure. These models are meant to be

additive, so that the network metrics included are intended to complement rather than

substitute the simple metrics traditionally used.

At the species level, response variables included the fruit set for different

individuals of each species analyzed using a binomial distribution and the average

number of seeds per fruit, the average fruit and seed weight fitted using normal

distributions. The number of seeds per fruit was centered and scaled (i.e., we subtracted

column means and divided by standard deviation) to allow meaningful comparisons

across species with contrasting life histories. As explanatory variables, model 1 included

pollinator richness, and the total number of visits received by each plant species; while

model 2 added the two network attributes calculated at the species level: average plant

niche overlap and centrality. For both models, we included plant species nested within

site and site as random effects to account for the non-independence of several individuals

measured per species and site.

At the site level, we upscaled our species-level analyses. As response variables we

had the average reproductive success per site (i.e., average fruit set analyzed using a

binomial distribution, average number of seeds per fruit and average fruit and seed

weight using a normal distribution). We thus had a single value per site and no random

effects are needed. In this case, model 1 included total pollinator richness and total

pollinator abundance (i.e. number of visits received by all plants within the community)

as explanatory variables. Model 2, in turn, added information on network structure by

including nestedness and pollinator niche complementarity as explanatory variables.

265   Average values of reproductive success at the site level can be driven to a large

extent by a single plant species. Yet, what will determine the persistence of a diverse

plant community, is the presence of some sort of "equity" or evenness in reproductive

success across the whole community. We therefore calculated a measure of equity in

reproductive success at the site level as the proportion of species with normalized

(between 0 and 1) average fruit set values that were above the 50^{th} percentile. As any

selected threshold is arbitrary, we repeated this using the 25^{th} and 75^{th} percentile

thresholds (Byrnes *et al* 2014). We then used the same framework as that used for species

and site-level analyses and fit the same models 1 and 2 GLMs but using equity in

reproductive success as the response variable and fitting a binomial distribution.

275   In all cases, we used variance inflation factors to check for collinearity between

explanatory variables. Additionally, we ran residual diagnostics to check if model

assumptions were met. Then, we used the Akaike Information Criterion (AIC) to

compare model performance and complexity. Whenever the difference between the AIC

of both models was < 2 ($\Delta AIC < 2$), we considered that both models were equally good

(Burnham *et al.* 2011). All predictor variables were standardized prior to analysis. For

every model we also calculate the R^2 value, using the approximation suggested for

generalized mixed models when necessary (Nakagawa *et al* 2017).

Finally, we tested whether the importance of network structure in explaining

differences in equity in reproductive success within communities increases with the

number of plant species being considered. We expect that when only one plant species is

considered, then the importance of network structure will be negligible, while we expect

this importance to increase as more plant species are considered (up to a maximum

number of 6 species which is the maximum we have measured in our study at a particular

site).

To test this, we ran a simple simulation in which the number of species

considered increased at each step and for each step we re-calculated equity in

reproductive success. Instead of drawing plant species randomly for each step, we tested

all possible combinations for each plant number level and network, as the number of

combinations is small (e.g. for n = 3 plant selected out of 6 there is only 20 possible

combinations). Then, we tested if the relationship between equity in reproductive success

and functional complementarity (given its importance in determining differences in

reproductive success, see Results section) changes as a function of the number of plants

considered within our simulated communities. To this end, for each level of species

number considered, we randomly selected one of the generated equity values across each

of the 16 communities and regressed these 16 values against our network level predictor

and extracted the model slope estimates. We repeated this process 1,000 times and

averaged all slope estimates. We expect that the more plants considered, the larger the

resulting average estimates will be. Note that we only interpret the mean effects, as the

variance among different plant number of species considered depends on the initial

number of possible combinations.

Results

Within our sampling we recorded 1,472 plant-pollinator interactions involving 277

pollinator species and 57 plant species. Within the pollinator community the distribution

of individuals in different orders was: 87.84% Hymenoptera, 6.78% Diptera, 4.05%

Coleoptera and 1.09% Lepidoptera.

Our sampling completeness analyses revealed that with our survey we were able

to capture 18-62% of pollinator species (average = 35%), 47-98% for plant species

(average = 78%) and 13-41% for plant-pollinator links (average = 27%), in line with that

obtained with other studies (e.g., Chacoff *et al.* 2012, Fig. S2). Our values of sampling

completeness are slightly smaller in the case of pollinators, probably as a consequence of

the great diversity found in the Mediterranean region and within our study area in

particular, a hotspot of insect diversity (Nieto *et al.* 2014). In addition, the fact that we

include an extra effort to capture rare interactions observed outside of our main sampling

might also increase the number of singletons which directly affect richness estimates.

Species-level analyses

At the species level, in the case of fruit set, our results show that model 2 shows the best

fit to our data (lowest AIC value), and fixed effects explains 4% of the variability

observed. In this case, we find a positive effect of a network structure metric, the

centrality of a focal plant within the overall network on its fruit set (Table 1, Fig. 2A, Fig.

S3).

For the average number of seeds per fruit, our results show again that model 2

shows the best fit, with fixed effects explaining 4% of the variability observed in our

data. In this case, we find a positive effect of the niche overlap between plant species on

the number of seeds produced (Table 1B, Fig. 2B, Fig. S3).

For all other measures of reproductive success considered (i.e., fruit and seed

weight), model 1 showed the best fit. However, none of the variables included within our

model explain the differences observed (Tables S5-S6, Fig. S3).

Site-level analyses

At the site level, we find different patterns for fruit set and the number of seeds per fruit

as compared to those for fruit and seed weight. In the case of fruit set and the number of

seeds per fruit, we find that both model 1 and 2 are equally good in describing the

differences observed when penalizing for model complexity (i.e., $\Delta AIC < 2$; Burnham *et*

al 2011). This suggests model 2 is a good model despite its added complexity, and

actually shows a substantially better predictive ability than model 1 ($R^2 = 0.59$ for model

2 versus 0.47 for model 1 in the case of fruit set and $R^2 = 0.52$ for model 2 versus 0.31

for model 1 in the case of the number of seeds per fruit) and therefore we will comment

results for this model only. In particular, we find that both fruit set and the number of

seeds per fruit are positively related to niche complementarity between pollinators (Table

2, Fig. 3). Additionally, we find a negative effect of site-level pollinator richness on

average fruit set (Table 2A, Fig. 3, Fig. S4).

Contrastingly, in the case of weight variables (fruit and seed weight), in both

cases we find that the best model is model 1, i.e., that only including simple visitation

metrics ($R^2 = 0.29$ in the case of fruit weight and 0.51 in the case of seed weight). Here,

we find a consistent positive effect of site-level pollinator richness for both weight

descriptors (Tables S7A-S8A, Fig. 4, Fig. S4). This effect is maintained even after

removing a site that has a particularly large pollinator richness value (Tables S7B-S8B,

Fig. S5, Fig. S4).

Equity in fruitset

When evaluating the effect of differences in community composition and network

structure for equity in reproductive success across the different species within a

community we find that model 1 is the best model for all the thresholds considered (50^{th},

25^{th} and 75^{th} percentiles). However, none of the variables considered are able to explain

differences observed in equity across sites (Tables S9, S10, S11).

359   Within our simulation evaluating the effect of niche complementarity on equity in

reproductive success as more plants within the community are considered, we find that

the effect of complementarity becomes more important as the reproductive success of

more species is considered (Fig. 5). This importance seems to reach some sort of plateau

at 6 species. However, this should be further evaluated, as this is the maximum number of

species simultaneously observed in a community for our study, which precludes us from

simulating further numbers of species.

Discussion

The existence of relationships between interaction network structure and ecosystem

function have been long hypothesized, yet, the specific mechanisms by which structure

influences function have remained elusive until now (Thompson *et al.* 2012). Our results

show that different aspects of network structure affect different dimensions of ecosystem

functioning. In particular, we find that the centrality of a plant species within a

community, which measures the number of connections it receives from other species in

the community, has a positive effect for its fruit set. At the site level, we find that greater

values of niche complementarity between pollinators result in larger average fruit sets

and number of seeds per fruit.

One of the first conclusions we can extract from the fact that in most cases both of

the models we considered (i.e., the simple model based on visitation metrics and the more

complex one including network structure metrics) were equally good, is that the added

complexity of measuring the full network of interactions may not pay off for rapid

assessments. Hence, simple visitation metrics, such as pollinator richness, might be

enough to describe general patterns (Garibaldi *et al.* 2013, 2015). Yet, adding network

level information may inform us of the potential ecological mechanisms underlying the

processes driving the observed patterns.

Consistent with previous experimental (Fontaine *et al.* 2005; Fründ *et al.* 2013),

theoretical (Pauw 2013), and empirical studies (Valdovinos *et al.* 2016, Poisot *et al.*

2013), we find that niche complementarity is key in determining differences in

reproductive outputs. Indeed, we find that communities where there is less overlap in the

niches occupied by pollinator species had greater values of reproductive success, both

greater fruit set and larger numbers of seeds per fruit. This therefore reflects the fact that

reproductive success in plant species requires the delivery of conspecific pollen and thus

of a certain degree of specialization amongst pollinator species on a particular plant

resource in order to avoid the negative effects of inter-specific pollen deposition (e.g.,

pollen loss, Flanagan *et al.* 2009) or interference with conspecific pollen (Morales &

Traveset 2008). However, we also find that some level of redundancy in these functions

is needed as revealed by the positive effect of plant niche overlap on the number of seeds

per fruit at the species level.

In our study, we did not find an effect of nestedness for reproductive success in

any case. This metric, widely used across network analysis, and which is deemed to

stabilize natural communities (Bastolla *et al.* 2009), does not seem to play a direct role in

ecosystem function measured as plant reproductive success. However, our study is

limited to a maximum of six common plant species per community, and including more

species, including rare species, might reveal different patterns, in which nestedness and

the redundancy it implies might play a more important role.

Site-level plant reproductive success measured as average fruit or seed set across

all the species considered, is an important part of the functions delivered by pollinators to

plants. However, these average values might be masking a great deal of variability

amongst plant species, and thus a nuanced view of the effect of pollinators on whole-

plant ensembles is needed. This can be captured by the effect of pollinators on equity in

reproductive success across plant species. This aspect ensures that reproductive success is

equally distributed amongst a larger number of species, thus contributing to the

maintenance of greater species diversity values in natural populations. Indeed, we know

that plant species diversity within a community is largely driven by different types of

direct and indirect interactions including those amongst plant species (e.g., resource

competition, Goldberg & Barton 1992, or facilitation, Bruno *et al.* 2003), as well as those

defining antagonistic (e.g., involving pathogens, Bagchi *et al.* 2010), or mutualistic

interactions (e.g, pollinators, Benadi *et al.* 2013; Lanuza *et al.* 2018). However,

equitability in reproductive success across species is seldom taken into account, despite

its importance in maintaining genetic diversity and ensuring the resilience of populations

to further change.

In the case of equity, we did not find a strong effect of either simple visitation or

network structure metrics. However, the results of our simulation on the importance of

network structure as the number of plant species considered increases, shows us that this

effect increases when more than four plant species are considered. This implies that if we

were able to measure reproductive success for all the plant species in all the communities

(which is not feasible given constraints in sampling effort), we might find that the effects

of network structure on equity might be more prevalent.

One of the unexpected results of our analyses is the strong negative effect of

pollinator richness for fruit set at the site level. An explanation to this might be the fact

that pollinator richness here includes all the pollinators recorded during our sampling

efforts, i.e., it includes species that do not pollinate some of the species whose

reproductive success was measured. More complex communities with more pollinators,

but also with more plant species (Pearson correlation between plant and pollinator

richness = 0.42 in our case) may require stabilizing mechanisms that reduce the

competition exerted by the dominant plant species. A way to reduce the competition

exerted by these dominant species, which are precisely those evaluated in this study, is by

reducing their reproductive success (Lanuza *et al* 2018, Stavert *et al* 2019). These ideas

open the door to exploring the positive or negative effects of the complete pollinator

community on full plant species coexistence, which may be determined by density-

dependence effects (Benadi & Pauw 2018). In our case, while fruit set is negatively

related to pollinator richness, it is important to note that fruit and seed weight show the

opposite relationship, indicating that this density-dependent effect might only be limiting

fruit quantity and not fruit quality. Thus, taking into account the densities of co-flowering

plant species may be the next step (Vanbergen *et al.* 2014).

Our study illustrates the complexity of linking network structure to ecosystem

function empirically, because measuring both structure and function is challenging. For

example, there is an ongoing debate as to which network metrics better reflect classic

ecological mechanisms, such as niche partitioning or competition (Delmas *et al* 2018).

Here, we focus on testing two specific hypotheses, but other structural properties can be

explored when more data becomes available. Furthermore, the structure of plant-

pollinators networks is dynamic due to ecological and evolutionary reasons, but so far,

we are only able to characterize it for single snap-shots. Moreover, different aspects of

functioning may be important, such as the presence of non-linear relationships or the

need to consider the functioning of both trophic levels (Godoy *et al* 2018). In terms of

plant reproductive success and the functions performed by pollinators we can measure

different aspects, ranging from pollen deposition (the direct pollinator function), to its

final effects on plant fitness. Here, we focus on an intermediate stage including fruit

quantity and quality, which is of clear ecological importance.

In summary, our findings show that the analysis of natural communities of

interacting species using network analysis not only represents an ideal way of visualizing

and grasping the complexity present within these communities. Rather, it also represents

a manner of mechanistically understanding differences observed across the reproductive

success of individuals and/or species while linking them to potential ecological

mechanisms.

Data accessibility

All the data used is available

Acknowledgements

The authors would like to thank Oscar Aguado for identifying pollinator species. We would also like to acknowledge the efforts of the editor Cédric Gaucherel, as well as those of Michael Lattorff, Nicolas Deguines, Alberto Pascual, and three anonymous reviewers who provided very helpful comments and suggestions in a previous version of this manuscript. AM received funding from a Juan de la Cierva (IJCI-2014-22558) and Ikerbasque fellowships. IB acknowledges funding from MSC-PCIG14-GA-2013-631653 BeeFun Project. We thank Doñana's Singular Scientific-Technical Infrastructure (ICTS-RBD) for access to the park.

479 **Table 1.** Results of GLMM showing the effect of simple visitation and network structure
480 metrics on A) species-level fruit set and B) average number of seeds per fruit based on
481 best model selected. Bold letters indicate variables with large effects.
482

A) Fruit set	Estimate	Std.Error	z.value
(Intercept)	1.72	0.21	8.16
Pollinator richness	-0.01	0.21	-0.07
Total number of visits	0.14	0.25	0.57
Centrality	**0.46**	**0.25**	**1.81**
Plant niche overlap	0.05	0.24	0.20

483
484

B) Seeds per fruit	Estimate	Std..Error	t.value
(Intercept)	0.05	0.14	0.36
Pollinator richness	0.14	0.15	0.90
Total number of visits	-0.01	0.16	-0.04
Centrality	-0.15	0.15	-1.00
Plant niche overlap	**0.22**	**0.17**	**1.32**

487 **Table 2.** Results of GLM showing effect of simple visitation and network structure

488 metrics on A) site-level average fruit set and B) site-level average number of seeds per

489 fruit based on best model selected. Bold letters indicate variables with large effects.

A) Fruit set	Estimate	Std. Error	z value
(Intercept)	**1.22**	**0.13**	**9.18**
Pollinator richness	**-0.75**	**0.17**	**-4.35**
Total number of visits	-0.16	0.16	-1.02
Nestedness	0.11	0.16	0.72
Pollinator niche complementarity	**0.29**	**0.18**	**1.57**

B) Seeds per fruit	Estimate	Std. Error	t value
(Intercept)	45.37	8.55	5.31
Pollinator richness	3.26	12.18	0.27
Total number of visits	8.38	9.99	0.84
Nestedness	-10.94	10.21	-1.07
Pollinator niche complementarity	**29.51**	**13.31**	**2.21**

 Figure legends

Figure 1. Map showing location of 16 Mediterranean woodland patches where plant-

pollinator interactions were surveyed from February to May 2015. Inset shows location

of study area within SW Spain.

Figure 2. Partial residual plots showing the effect of the single predictor which best

explains the variability in species-level reproductive success. A) Shows the effect of plant

species centrality on the fruit set of each of the plant species considered and B) shows the

effect of plant niche overlap on the average number of seeds per fruit. Dots represent

each of the individuals sampled for each species within each site.

Figure 3. Partial residual plots showing the effect of the single predictor which best

explains the variability in site-level reproductive success. A) Shows the effect of

pollinator richness, and B) of niche complementarity among pollinator species on site-

level average fruit set. C) Shows the effect of niche complementarity among pollinator

species on the average number of seeds per fruit at the site level. Dots represent average

values of fruit set at the level of the community for all plant species considered (N=16

sites).

Figure 4. Partial residual plots showing the effect of pollinator richness on site-level

average A) fruit and B) seed weight. Dots represent values for each site (N=16 sites).

Figure 5. Results of simulation evaluating the importance of niche complementarity in

determining differences in equity in reproductive across communities harboring from one

to six species. Points represent average values across 1,000 simulated combinations.

Figure 1.

Plant species
- *Anchusa azurea*
- *Asphodelus fistulosus*
- *Astragalus lusitanicus*
- *Cistus albidus*
- *Cistus crispus*
- *Cistus ladanifer*
- *Cistus libanotis*
- *Cistus monspeliensis*
- *Cistus salvifolius*
- *Halimium commutatum*
- *Halimium halimifolium*
- *Lavandula pedunculata*
- *Lavandula stoechas*
- *Phlomis purpurea*
- *Rosmarinus officinalis*
- *Spartium junceum*
- *Teucrium fruticans*
- *Ulex australis*

520
521
522

Figure 2.

526
527

Figure 3.

Figure 4.

Figure 5.

References

Abrams, P.A., Holt, R.D. & Roth, J.D. (1998). Apparent Competition or Apparent

Mutualism? Shared Predation When Populations Cycle. *Ecology*, 79, 201–212.

Albrecht, M., Schmid, B., Hautier, Y. & Müller, C.B. (2012). Diverse pollinator

communities enhance plant reproductive success. *Proceedings of the Royal Society B:*

Biological Sciences, 279, 4845 LP–4852.

Almeida-Neto, M. & Ulrich, W. (2011). A straightforward computational approach for

measuring nestedness using quantitative matrices. *Environ Model Softw*, 26, 173–178.

Bagchi, R., Swinfield, T., Gallery, R.E., Lewis, O.T., Gripenberg, S. & Narayan, L. *et al.*

(2010). Testing the Janzen-Connell mechanism: pathogens cause overcompensating

density dependence in a tropical tree. *Ecology Letters*, 13, 1262–1269.

Bartomeus, I., Stavert, J., Ward, D. & Aguado, O. (2019). Historical collections as a tool

for assessing the global pollination crisis. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, 374, 20170389.

Bascompte, J. & Jordano, P. (2007). Plant-Animal Mutualistic Networks: The

Architecture of Biodiversity. *Annu Rev Ecol Evol S*, 38, 567–593.

Bascompte, J., Jordano, P., Melián, C.J. & Olesen, J.M. (2003). The nested assembly of

plant–animal mutualistic networks. *P. Natl. Acad. Sci. USA*, 100, 9383–9387.

Bastolla, U., Fortuna, M.A., Pascual-Garcia, A., Ferrera, A., Luque, B. & Bascompte, J.

(2009). The architecture of mutualistic networks minimizes competition and increases

biodiversity. *Nature*, 458, 1018–1020.

Benadi, G., Blüthgen, N., Hovestadt, T. & Poethke, H.-J. (2013). When can plant-

pollinator interactions promote plant diversity? *Am Nat*, 182, 131–146.

Benadi, G. & Pauw, A. (2018) Frequency dependence of pollination visitation rates

suggests that pollination niches can allow plant species coexistence. *Journal of Ecology*,

106, 1892-1901.

Bennett, J.M., Thompson, A., Goia, I., Feldmann, R., Ştefan, V. & Bogdan, A. *et al.*

(2018). A review of European studies on pollination networks and pollen limitation, and

a case study designed to fill in a gap. *AoB PLANTS*, 10, ply068–ply068.

Biesmeijer, J.C., Roberts, S.P., Reemer, M., Ohlemuller, R., Edwards, M. & Peeters, T. *et*

al. (2006). Parallel declines in pollinators and insect-pollinated plants in Britain and the

Netherlands. *Science*, 313, 351–354.

Blüthgen, N. (2010). Why network analysis is often disconnected from community

ecology: A critique and an ecologist's guide. *Basic and Applied Ecology*, 11, 185–195.

Blüthgen, N. & Klein, A.-M. (2011). Functional complementarity and specialisation: The

role of biodiversity in plant–pollinator interactions. *Basic and Applied Ecology*, 12, 282–

291.

Blüthgen, N., Menzel, F. & Blüthgen, N. (2006). Measuring specialization in species

interaction networks. *BMC Ecology*, 6, 1–12.

Brosi, B.J. (2016). Pollinator specialization: from the individual to the community. *New*

Phytologist, 210, 1190–1194.

Bruno, J.F., Stachowicz, J.J. & Bertness, M.D. (2003). Inclusion of facilitation into

ecological theory. *Trends in Ecology & Evolution*, 18, 119–125.

Burnham, K.P., Anderson, D. & Huyvaert, K.P. (2011). AIC model selection and

multimodel inference in behavioral ecology: some background, observations, and

comparisons. *Behav Ecol Sociobiol*, 65, 23–35.

Carvalheiro, L.G., Biesmeijer, J.C., Benadi, G., Fründ, J., Stang, M. & Bartomeus, I. *et*

al. (2014). The potential for indirect effects between co-flowering plants via shared

pollinators depends on resource abundance, accessibility and relatedness. *Ecology*

Letters, 17, 1389–1399.

Chacoff, N.P., Vázquez, D.P., Lomáscolo, S.B., Stevani, E.L., Dorado, J. & Padrón, B.

(2012). Evaluating sampling completeness in a desert plant-pollinator network. *Journal*

of Animal Ecology, 81, 190–200.

Chao, A., Colwell, R.K., Lin, C. & Gotelli, N.J. (2009). Sufficient sampling for

asymptotic minimum species richness estimators. *Ecology*, 90, 1125–1133.

Delmas, E., Besson, M., Brice, M.-H., Burkle, L.A., Dalla Riva, G.V. & Fortin, M.-J. *et*

al. (2019). Analysing ecological networks of species interactions. *Biological Reviews*, 94,

16–36.

Devoto, M., Bailey, S., Craze, P. & Memmott, J. (2012). Understanding and planning

ecological restoration of plant–pollinator networks. *Ecology Letters*, 15, 319–328.

Dormann, C.F., Frund, J., Bluthgen, N. & Gruber, B. (2009). Indices, graphs and null

models: analyzing bipartite ecological networks. *The Open Ecology Journal*, 2, 7–24.

Flanagan, R.J., Mitchell, R.J., Knutowski, D. & Karron, J.D. (2009). Interspecific

pollinator movements reduce pollen deposition and seed production in Mimulus ringens

(Phrymaceae). *American Journal of Botany*, 96, 809–815.

Fontaine, C., Dajoz, I., Meriguet, J. & Loreau, M. (2005). Functional Diversity of Plant–

Pollinator Interaction Webs Enhances the Persistence of Plant Communities. *PLOS*

Biology, 4, e1.

Fründ, J., Dormann, C.F., Holzschuh, A. & Tscharntke, T. (2013). Bee diversity effects

on pollination depend on functional complementarity and niche shifts. *Ecology*, 94,

2042–2054.

Garibaldi, L., Carvalheiro, L.G., Leonhardt, S.D., Aizen, M.A., Blaauw, B.R. & Isaacs,

R. *et al.* (2015). From research to action: enhancing crop yield through wild pollinators.

Front Ecol Environ, 12, 439–447.

Garibaldi, L.A., Steffan-Dewenter, I., Winfree, R., Aizen, M.A., Bommarco, R. &

Cunningham, S.A. *et al.* (2013). Wild Pollinators Enhance Fruit Set of Crops Regardless

of Honey Bee Abundance. *Science*, 339, 1608–1611.

Godoy, O., Bartomeus, I., Rohr, R.P. & Saavedra, S. (2018). Towards the Integration of

Niche and Network Theories. *Trends in Ecology & Evolution*, 33, 287–300.

Goldberg, D.E. & Barton, A.M. (1992). Patterns and Consequences of Interspecific

Competition in Natural Communities: A Review of Field Experiments with Plants. *The*

American Naturalist, 139, 771–801.

Holt, R.D. (1977). Predation, apparent competition, and the structure of prey

communities. *Theoretical Population Biology*, 12, 197–229.

Hsieh, T.C., Ma, K.H. & Chao, A. (2016). iNEXT: iNterpolation and EXTrapolation for

species diversity. R package version 2.0.8.

Johnson, C.A. & Bronstein, J.L. (2019). Coexistence and competitive exclusion in

mutualism. *Ecology*, 100, e02708.

Jordano, P. (2016) Sampling networks of ecological interactions. *Functional ecology*, 30,

1183-1893.

Lanuza, J.B., Bartomeus, I. & Godoy, O. (2018). Opposing effects of floral visitors and

soil conditions on the determinants of competitive outcomes maintain species diversity in

heterogeneous landscapes. *Ecology Letters*, 21, 865–874.

Lázaro, A., Lundgren, R. & Totland, Ø. (2014). Experimental reduction of pollinator

visitation modifies plant–plant interactions for pollination. *Oikos*, 123, 1037–1048.

Macarthur, R. & Levins, R. (1967). The Limiting Similarity, Convergence, and

Divergence of Coexisting Species. *The American Naturalist*, 101, 377–385.

May, R.M. & MacArthur, R.H. (1972). Niche Overlap as a Function of Environmental

Variability. *Proceedings of the National Academy of Sciences*, 69, 1109 LP–1113.

Mayfield, M.M. & Stouffer, D.B. (2017). Higher-order interactions capture unexplained

complexity in diverse communities. *Nature Ecology &Amp; Evolution*, 1, 62.

Morales, C.L. & Traveset, A. (2008). Interspecific Pollen Transfer: Magnitude,

Prevalence and Consequences for Plant Fitness. *Critical Reviews in Plant Sciences*, 27,

221–238.

Nakagawa, S., Johnson, P.C.D. & Schielzeth, H. (2017). The coefficient of determination

R2 and intra-class correlation coefficient from generalized linear mixed-effects models

revisited and expanded. *Journal of The Royal Society Interface*, 14, 20170213.

Nieto, A., Roberts, S., Kemp, J., Rasmont, P., Kuhlmann, M. & García Criado, M. *et al.*

(2014). European Red List of Bees.

Pauw, A. (2013) Can pollination niches facilitate plant coexistence? *Trends in Ecology &*

Evolution, 28, 30-37.

Petchey, O.L. & Gaston, K.J. (2007). Dendrograms and measuring functional diversity.

Oikos, 116, 1422–1426.

Poisot, T., Mouquet, N. & Gravel, D. (2013). Trophic complementarity drives the

biodiversity–ecosystem functioning relationship in food webs. *Ecology Letters*, 16, 853–

861.

Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O. & Kunin, W.E.

(2010). Global pollinator declines: trends, impacts and drivers. *Trends Ecol Evol*, 25,

345–353.

R Development Core Team, R. (2011). R: A Language and Environment for Statistical

Computing. R foundation for statistical computing.

Simmons, B.I., Hoeppke, C. & Sutherland, W.J. (2019). Beware greedy algorithms.

Journal of Animal Ecology, 88, 804–807.

Song, C., Rohr, R.P. & Saavedra, S. (2017). Why are some plant–pollinator networks

more nested than others? *Journal of Animal Ecology*, 86, 1417–1424.

Stavert, J.R., Bartomeus, I., Beggs, J.R., Gaskett, A.C. & Pattemore, D.E. (2019). Plant

species dominance increases pollination complementarity and plant reproductive

function. *Ecology*, 0, e02749.

Thébault, E. & Fontaine, C. (2010). Stability of Ecological Communities and the

Architecture of Mutualistic and Trophic Networks. *Science*, 329, 853–856.

Thompson, R.M., Brose, U., Dunne, J.A., Hall Jr., R.O., Hladyz, S. & Kitching, R.L. *et*

al. (2012). Food webs: reconciling the structure and function of biodiversity. *Trends in*

Ecology & Evolution, 27, 689–697.

Thomson, D.M. (2019). Effects of long-term variation in pollinator abundance and

diversity on reproduction of a generalist plant. *Journal of Ecology*, 0.

Valdovinos, F.S., Brosi, B.J., Briggs, H.M., Moisset de Espanés, P., Ramos-Jiliberto, R.

& Martinez, N.D. (2016). Niche partitioning due to adaptive foraging reverses effects of

nestedness and connectance on pollination network stability. *Ecology Letters*, 19, 1277–

1286.

Vanbergen, A., Woodcock, B., Gray, A., Grant, F., Telford, A., Lambdon, P., Chapman,

D.S., Pywell, R.F., Heard, M.S. & Cavers, S. (2014) Grazing alters insect visitation

networks and plant mating systems. *Functional Ecology*, 28, 178-189.

Winfree, R. (2013). Global change, biodiversity, and ecosystem services: What can we

learn from studies of pollination? *Basic and Applied Ecology*, 14, 453–460.

Woodward, G. & Hildrew, A.G. (2002). Body-size determinants of niche overlap and

intraguild predation within a complex food web. *Journal of Animal Ecology*, 71, 1063–

1074.

Zhang, J. (2016). Package spaa: SPecies Association Analysis.