Probing the polarity of spontaneous GABAergic synaptic transmission in vivo

Olivier Dubanet[1], Arnaldo Ferreira[1], Andreas Frick[1], Hajime Hirase[2], Anna Beyeler[1], Xavier
Leinekugel[1]*

1. University of Bordeaux, INSERM U1215, Neurocentre Magendie, 33077 Bordeaux, France
2. Center for Translational Neuromedicine, University of Copenhagen
*corresponding author : xavier@arcadi.eu

9
10
11
12

Abstract

Neuronal chloride gradient directly affects the polarity of GABAergic transmission, a major factor of
network dysfunction in a variety of brain pathologies. But the influence of $[Cl^-]i$ on functional
GABAergic transmission critically depends on the very dynamic combination of membrane potential,
conductance and occurrence of other synaptic inputs. Taking this complexity into account implies
measuring the postsynaptic responses to spontaneously occurring GABAergic events, in vivo, without
interfering with neuronal $[Cl^-]i$. We have overcome this challenge by combining extracellular detection
of unitary inhibitory postsynaptic field-potentials (fIPSPs) and silicon probe recording of the firing
activity of multiple individually identified neurons. Probing the polarity of spontaneous, ongoing
GABAergic transmission across the CA3 hippocampal circuit in vivo with single cell resolution, our
results support depolarizing actions of perisomatic GABAergic transmission and time-locked excitation
of CA3 pyramidal neurons in acute and chronic mouse models of epilepsy, but little contribution of
excitatory GABA to epileptogenesis.

Introduction

GABAergic inhibition relies on ionotropic GABA$_A$ and metabotropic GABA$_B$ receptors. The fast and time-locked inhibition considered essential for neuronal coding is the one mediated by GABA$_A$ receptors (GABA$_A$ R), permeable to chloride ions (Cl$^-$). In normal conditions, neurons maintain a low intracellular concentration of chloride ([Cl$^-$]i), so that the reversal potential of Cl$^-$ mediated currents is more negative than the resting potential[1]. GABA release therefore generates an influx of Cl$^-$, with an hyperpolarizing effect on membrane potential, and inhibition of neuronal discharge.

Recent publications have proposed the implication of defective inhibition in the etiology of various pathologies including epilepsy, depression, schizophrenia, Down Syndrome or Autism Spectrum Disorders [2-9]. In vitro and in vivo experiments have suggested that defective function of the K-Cl co-transporter KCC2, that would normally extrude Cl$^-$ from the cell, was responsible for a reversed polarity of GABAergic synaptic transmission from inhibitory to excitatory, participating in epileptogenesis in animal models [10-20] as well as in human tissue resected from epileptic patients [21-23]. But one major difficulty in the study of GABAergic inhibition is to avoid artefactual interference with the native Cl$^-$ gradient, a main determinant of GABA$_A$ R mediated inhibitory function. Perforated-patch clamp recordings with the compound antibiotic gramicidin provide electrical access to the cell through the formation of pores in the membrane, permeable to cations but impermeant to Cl$^-$. Perforated-patch recordings with gramicidin have been successfully used in vitro to record GABAergic synaptic currents, while keeping intracellular Cl$^-$ unaffected by the pipette solution[10,11,16,24]. But previous work has also demonstrated that Cl$^-$ gradient can be affected by the slicing procedure [25-27], questioning the claims for excitatory GABAergic transmission based on in vitro recordings [26].

In vivo treatment with the diuretic bumetanide, which also reduces the accumulation of Cl$^-$ ions in neurons through the inhibition of the Cl$^-$ intruder NKCC1, was found to reduce behavioural abnormalities in rat and mouse models of Autism Spectrum Disorders[5] and seizures in animal models of epilepsy [19,28-32]. But bumetanide has a limited bio-availability in the brain due to poor blood brain barrier penetration[33-35], and it is finally unclear if its action in vivo was indeed due to the restoration of neuronal Cl$^-$ gradient and inhibitory GABAergic transmission.

Direct investigation of the polarity of GABAergic transmission in vivo puts serious constraints in terms of experimental access, and the evaluation of responses to spontaneous synaptic GABA$_A$ R activation is challenging. The ideal method would fully preserve the exact spatio-temporal interactions between membrane potential, conductance and ongoing synaptic inputs (both excitatory and inhibitory), which natural dynamics are highly complex and yet critical for the net inhibitory or excitatory effect of GABAergic transmission[18,24,36-39]. Adapting to in vivo conditions the extracellular recording of unitary GABAergic postsynaptic potentials (field IPSPs) previously described in vitro[40-42], we could evaluate the polarity of perisomatic GABAergic signaling during spontaneous activity in the intact brain. As an exemple application, we report direct evidence for depolarizing actions of perisomatic GABAergic transmission and time-locked excitation of CA3 pyramidal neurons in acute and chronic mouse models

64 of epilepsy, but little contribution of excitatory GABA to epileptogenesis. We believe that this

65 demonstration will pave the way for the investigation of Cl⁻ homeostasis and inhibitory function in vivo,

66 in physiological conditions and in situations in which alteration or even reversal of GABAergic

67 transmission is hypothesized to occur. It also constitutes an invaluable tool to quantify the actual in vivo

68 efficacy of drugs designed to modulate Cl⁻ homeostasis and restore physiological GABAergic inhibition,

69 thereby meeting high clinical and therapeutical expectations.

72 **Results**

74 **Extracellular recording of perisomatic field-IPSPs and time-locked inhibition of CA3 pyramidal**

75 **cell firing in vivo.**

76  Non-invasive recording of the dynamics of GABAergic synaptic transmission in vivo is a

77 challenge. Previous work in vitro has suggested that perisomatic inhibitory postsynaptic signals

78 originating from individual interneurons (basket and chandelier cells) could be recorded as extracellular

79 field potentials (fIPSPs) from their target pyramidal-cell population[40-42]. This approach offers several

80 main advantages. First, extracellular recordings preserve native [Cl⁻]i and other cytoplasmic factors

81 critical for GABAergic transmission[43]. Second, the polarity of the field events indicates the direction of

82 the current elicited in the target population of pyramidal cells, thereby addressing the question of

83 depolarizing vs hyperpolarizing polarity of GABAergic postsynaptic signaling. And third, combined

84 with units recording and spike sorting of neuronal discharges, extracellular detection of population

85 fIPSPs provides a time reference to investigate the net effect of GABAergic perisomatic synaptic signals

86 on the firing of the target neurons with single cell resolution, allowing a direct quantification of the

87 efficacy of inhibition or excitation provided by perisomatic GABAergic inputs.

88  In order to evaluate the efficiency and polarity of perisomatic GABAergic transmission in the

89 hippocampus in vivo, we have therefore inserted silicon probes into the dorsal CA3 pyramidal layer of

90 urethane-anesthetized mice, recording spontaneous local field potentials and unitary neuronal activity.

91 As illustrated in Fig. 1, spontaneous activity from the CA3 pyramidal layer included action potentials

92 (units) and positive local field events with a profile similar to that of fIPSPs previously described in

93 vitro as the GABA$_A$ receptor mediated postsynaptic potentials originating from PV interneurons[42].

94 Accordingly, these events were characterized by a fast rise time (2.01±0.53ms, n=8 mice) and a decay

95 time constant increased by the GABA$_A$ modulator diazepam (2mg/kg, IP) from 3.65±0.05ms to

96 4.19±0.06ms (n=3 mice, p<0.01, two-tailed paired t-test, t = 9.13, DF = 2). As observed in vitro and

97 expected from perisomatically projecting interneurons (basket and chandelier cells), in vivo fIPSPs

98 provided very powerful time-locked inhibition of local pyramidal cells. Accordingly, multi-unit

99 pyramidal cell activity was almost fully interrupted by fIPSPs, as observed in the traces shown in Fig.

100 1b and in the trough that follows fIPSPs in the peri-event time histogram (PETH) shown in Fig. 1d-e.

Beside the peak firing that precedes fIPSPs and probably reflects the recruitment of perisomatic

inhibition by pyramidal cells discharge[42], 79.05±14.93% of the 139 putative pyramidal cells obtained

from 8 mice were significantly inhibited by fIPSPs, resulting in a population firing reduced by

92.3±4.5%, returning to baseline values after 9.75±2.8ms (n=8 mice).

Perisomatic fIPSPs specifically originate from PV interneurons and provide homeostatic

redistribution of individual pyramidal cells firing rates in the CA3 hippocampal network.

In order to verify that fIPSPs actually originate from PV interneurons, we recorded the extracellular

field-potential responses to the optogenetic activation of PV interneurons. PV-Cre mice were injected

with the DIO hChR2 (E123T/T159C)-EYFP construct in the CA3 region in order to express the cationic

actuator ChR2 specifically in PV interneurons. PV immuno-staining of the infected mice showed

excellent specificity, with 96 to 100% (98.3±2%, n=3 mice) of the infected (GFP-positive) cells being

also PV-positive (Supplementary Fig. 1a). As illustrated in Fig. 2, brief (2ms) local optogenetic

activation of PV interneurons did evoke reliable fIPSPs, together with powerfull time-locked inhibition

of local pyramidal cells (n=33 pyramidal cells from 5 mice). PV interneuron firing is therefore sufficient

to evoke fIPSPs. In order to know whether other types of interneurons also significantly contributed to

the generation of perisomatic fIPSPs during spontaneous CA3 activity, PV-Cre mice were injected with

the FLEX-rev::PSAML141F,Y115F:GlyR-IRES-GFP construct to express the inhibitory

pharmacogenetic actuator PSAM-GlyR specifically in PV interneurons. PV immuno-staining of the

infected mice showed that in addition to high specifity (99±1%, n=3 mice), 98±1 % (n=3 mice) of the

PV-positive cells in the region of recording were also GFP-positive (Supplementary Fig. 1b), indicating

that most PV interneurons expressed PSAM-GlyR. As illustrated in Fig. 3g-h, the pharmacogenetic

inhibition of PV interneurons induced by local injection of the exogenous ligand PSEM-89S very

efficiently suppressed spontaneous fIPSPs, from 6.58±3.50 to 0.45±0.66 events/s (n= 5 mice, p<0.05,

two-sided Wilcoxon Mann-Whitney Test), suggesting that the recorded fIPSPs originated

predominantly from PV interneurons.

Even though pharmacogenetic silencing of PV interneurons in the anesthetized mouse did not

increase the global firing rate of the CA3 pyramidal cell population (Fig. 3i, 1.19±2.72Hz in control vs

0.96±1.04Hz in PSEM-89S, n=5 mice, NS, two-sided Wilcoxon Mann-Whitney Test), it did affect

individual cells firing. As illustrated in Supplementary Fig. 2, the correlation coefficient between

individual cells firing rates in consecutive epochs of 10min indicated a remarkable stability, both in

baseline (r=0.987, p<0.001) and after PV interneuron silencing (r=0.79, p<0.001). On the other hand, as

illustrated in Fig. 3i, the firing rates before vs after PSEM-89S were hardly correlated (r=0.396, p<0.05),

indicating that PV interneuron silencing induced a redistribution of firing rates among pyramidal

neurons within the CA3 circuit.

Depolarizing and excitatory perisomatic GABAergic transmission in acute seizures in the drug-free mouse.

The reversal potential of Cl-mediated GABAergic currents (E-GABA) is controlled by the passive movement and active transport of Cl^- through various ionic channels and transporters in the plasma membrane. During spontaneous activity, Cl^- ions flowing through $GABA_A$ channels tend to shift the reversal potential of GABAergic currents toward membrane potential. E-GABA is therefore a very dynamic process that depends on ongoing fluctuations of membrane potential and GABAergic conductances. During seizures, GABA is massively released by interneurons while pyramidal cells are depolarized, therefore getting loaded with Cl^- until extrusion mechanisms restore an hyperpolarizing Cl^- gradient. It is therefore expected that due to the time course of Cl^- extrusion, high Cl^- load and excitatory GABA might participate in a deficit of inhibition immediately after a seizure[16,44]. However, the time dynamics of Cl^- balance through intrusion and extrusion systems that has been studied in vitro largely remains to be characterized in vivo. As illustrated in Fig. 4, taking advantage of fIPSPs as a non invasive index of the polarity of GABAergic currents in the CA3 pyramidal population, we have investigated the dynamics of perisomatic inhibition during acute interictal and ictal seizures in unanesthetized mice. In order to monitor both spontaneous and evoked fPISPs, PV-Cre mice were infected with the DIO-hChR2-EYFP construct in the CA3 region. Light stimulation with a locally positioned optical fiber provided evoked fIPSPs by direct stimulation of PV interneurons. Acute seizures were induced by a brief and focal intra-hippocampal injection of the convulsive agent bicuculline in the contralateral hemisphere. Electrophysiological recordings displayed both interictal activity and hippocampal ictal seizures, recognized as complex events lasting 30 to 40 seconds (39.25 ±2.5s, n=6 seizures from 4 mice), even though the animal did not show behavioural convulsions. While in comparison to baseline, fIPSPs were not significantly affected between epileptic discharges, they transiently appeared with a reversed polarity after the termination of ictal seizures, returning to normal polarity within 182 to 434 seconds after the termination of the seizure event (219.76±128.56s, n=6 seizures from 4 mice), suggesting transiently reversed polarity of GABAergic transmission after ictal seizures, but not after acute interictal discharges. Moreover, reversed fIPSPs entrained the discharge of postsynaptic neurons, as indicated by the peak in the peri-event histograms between reversed fIPSPs and multi-unit activity, suggesting not only depolarizing but also excitatory action of perisomatic GABAergic synaptic transmission. No more fIPSP-related excitatory interactions were seen after the recovery of the polarity of fIPSPs. These results suggest that an ictal seizure can entrain transient depolarizing and excitatory actions of perisomatic GABAergic transmission in the CA3 circuit.

Discrete excitatory perisomatic GABAergic transmission in a chronic model of epilepsy.

Acute ictal seizures, as induced experimentally by the injection of the convulsive agent kainic acid (KA), can turn into chronic epilepsy after a silent period of up to several weeks. Previous work on another model of epilepsy has suggested that reversed (excitatory) GABAergic transmission might be

involved in the transition between acute seizures and chronic epilepsy[11,14,31]. This condition is of high

clinical relevance because incident occurrence of isolated seizures in human is known to be a main factor

for the development of chronic epilepsy after a latent period that can last from months to several years[45].

Identifying the cellular mechanisms responsible for this transition would therefore potentially offer

opportunities for preventive treatment against epilepsy.

We have evaluated the dynamics of perisomatic inhibition in the CA3 hippocampal region of

anesthetized KA-treated mice during the latent period, one week after KA injection in the contralateral

hippocampus[46]. Interictal discharges were present in all KA-treated mice, but in none of the control

animals (Supplementary Fig. 3). Among all identified fIPSPs, none appeared of reversed polarity,

whether in control or in KA-treated mice, suggesting that GABAergic transmission remained globally

hyperpolarizing at the network level. Nevertheless, because the polarity of fIPSPs represents the average

response of all the local pyramidal targets of the discharging presynaptic interneuron(s), the possibility

remained that a subpopulation of pyramidal cells might be depolarized or even excited by GABAergic

transmission in spite of a globally hyperpolarizing fIPSP population response. We therefore evaluated

the response of individual neurons to spontaneous fIPSPs, and indeed identified a minority of neurons

(n=4 out of 78 putative pyramidal cells from 3 KA-treated mice, among a total of 195 putative pyramidal

cells from 8 KA-treated mice) that fired action potentials in response to GABAergic synaptic inputs.

Individual examples of putative pyramidal cells displaying the three distinct types of observed responses

to spontaneous fIPSPs (time locked excitation, time-locked inhibition, or no clear change in net firing

rate) are illustrated in Fig. 5. Therefore, analyzing single cell responses to fIPSPs revealed an

heterogenity of GABAergic alterations among KA-injected animals. These observations suggest that

fIPSPs in vivo may be useful to investigate whether the extent of reversed GABAergic transmission at

various delays after an acute seizure episode is involved in the distinct etiological outcome among

individual animals, differentially affected by the initial insult.

Discussion

The possibility that altered Cl⁻ homeostasis and functionally reversed GABAergic transmission

(from inhibitiory to excitatory) might be involved in major pathologies such as epilepsy, autism

spectrum disorders or schizophrenia, has recently raised considerable interest[2-5,10-15,19,21,22,29,30,47,48].

However, a direct assessment of this hypothesis has been hindered by the technical difficulty of probing

endogenous GABAergic synaptic function in vivo, because it implies to minimize perturbation of

cytoplasmic content, membrane potential and natural synaptic GABAergic activity[43]. The difficulties

reside in both the recording methods and the activation of GABAergic transmission. Neuronal [Cl⁻]i is

very labile and even intracellular recordings with sharp electrodes, which limit cytoplasmic wash-out,

do not preserve membrane integrity, potentially altering resting potential and endogenous Cl⁻ gradient.

Patch clamp (cell-attached) recordings are challenging in vivo, yield information about a single neuron
at a time, and potentially affect neuronal resting potential[49], a major parameter for the function of
GABAergic transmission. Imaging techniques (eg of calcium or chloride) are advantageous in terms of
spatial resolution, but do not offer the time resolution of electrophysiological recordings and get invasive
when applied to deep structures. Calcium imaging, in addition, does not detect depolarizations that do
not reach the threshold of calcium channel activation, and hardly discriminates between depolarizing
and excitatory effects that can both result in calcium increases, for example during a shunting
depolarization not associated with action potential firing[50], or $GABA_A$-induced calcium increases not
mediated by changes in membrane potential[51]. As for the activation of $GABA_A$-R, neither the exogenous
applications of GABAergic agonists, nor the electrical or optogenetic activation of GABAergic inputs
used in previous studies[18,26,36,37,50,52,53], respect the dynamics of spontaneous synaptic transmission. For
example, boosting Cl^- extrusion through KCC2 was recently shown to nicely prevent the prolongation
of seizures induced by the optogenetic activation of PV interneurons, but had no effect on spontaneous,
ongoing seizures[18]. Therefore, in spite of a reversed Cl^- gradient and GABAergic polarity with important
functional consequences if activated optogenetically, there is no evidence that excitatory GABAergic
transmission functionally participated in spontaneous ongoing seizure activity. Another caveat is that
different types of interneurons interact with each other, such as the mutually inhibiting PV and SOM-
expressing interneurons [38]. Experimental manipulation of specific interneurons (eg PV) may therefore
also influence seizure expression indirectly by the modulation of pyramidal cells' GABAergic inputs
from other interneurons. This illustrates the potential confound of interpreting the effects of GABAergic
activation at the network level rather than getting a direct readout of monosynaptic, spontaneous
GABAergic transmission.

On the other hand, previous work in vitro suggested that the direct postsynaptic response of
hippocampal pyramidal cells to their spontaneously occurring perisomatic GABAergic inputs could be
recorded extracellularly, as field-Inhibitory Postsynaptic Potentials (fIPSPs), and that the polarity of
fIPSPs could be used as an index of Cl^- gradient and polarity of GABAergic transmission. Indeed,
perisomatically projecting interneurons (basket and chandelier cells) have a densely arborised axon
specifically confined to the pyramidal cell layer, and contact a large proportion of pyramidal cells within
a restricted projection area[54,55]. Therefore, each action potential of the interneuron triggers the nearly
synchronous release of GABA on hundreds of target pyramidal cells, which postsynaptic responses
(IPSPs) sum up and can be readily identified from a local extracellular electrode[40-42], with a polarity
reflecting that of the Cl^- gradient[40,41]. This approach is well suited to preserving endogenous $[Cl^-]i$, can
be performed in vivo, and opens the possibility to resolve the dynamics of Cl^- gradient and polarity of
GABAergic transmission over the course of ongoing synaptic activity. A strong limitation however is
that fIPSPs are the average response of multiple pyramidal cells and therefore lack the resolution of
individual cells, so that changes in GABAergic polarity could be detected if they affect the whole
neuronal population but may be missed if affecting only a subset of cells. Moreover, as mentioned

before, GABAergic transmission involves both voltage and conductance changes, so that reversed [Cl$^-$
]i and depolarizing GABA carry a complex combination of excitatory (through depolarization) and
inhibitory (through increased conductance) influences[24,39,50,56-58]. As a consequence, determining the net
postsynaptic effect of GABAergic transmission requires a direct readout of postsynaptic firing activity.
For all these reasons, we now propose that the method of choice to probe Cl$^-$ gradient and GABAergic
synaptic function in vivo lies in combined multi-electrode recording and spike-sorting methods. While
fIPSPs provide the timing and global (average) polarity of spontaneous perisomatic GABAergic
synaptic events, the distributions (peri-event time histograms) of the spikes of individually identified
neurons provide the net functional effect of GABAergic transmission, with single cell resolution.

In line with previous in vitro evidence for the direct involvement of basket and chandelier cells in
the generation of fIPSPs[40-42], we provide direct evidence that in vivo fIPSPs originate mostly from PV
interneurons, because they were elicited by the optogenetic activation of PV interneuronal firing, and
largely eliminated by the pharmacogenetic inhibition of PV interneuronal firing. Perisomatic inhibition
is considered to be functionally optimal for the control of neuronal discharge, with synapses strategically
located on the soma and close to the site of action potential initiation[59,60]. PV basket and chandelier cells
are thus considered to play a major role in the control of pyramidal cells firing through time-locked
inhibition, presumably preventing excessive firing within the hippocampal circuit. However, in spite of
the critical importance of excitatory-inhibitory interactions in the recurrent CA3 circuit, the literature is
strikingly missing direct reports of the dynamics of excitatory-inhibitory interactions in this neuronal
circuit, presumably because of the technical difficulties to identify perisomatic inhibitory events in vivo.
Our experimental results confirm that perisomatic GABAergic transmission provides very powerful
time-locked inhibition of most CA3 pyramidal neurons. An interesting observation is that the blockade
of PV interneuron firing was responsible for a redistribution of pyramidal cell activity, but was not
accompanied by any major quantitative change of activity at the network level. This suggests that
perisomatic inhibition may be involved in the control of the timing of pyramidal cells discharge rather
than quantitative control of the network discharge. Previous reports are compatible with this hypothesis.
In a previous study conducted in vitro, we have reported that fIPSPs were rather poorly recruited by
network activity, suggesting that perisomatic inhibition should be more efficient in shaping the timing
of spike flow than in limiting excitatory runaway recruitment and preventing the generation of hyper-
synchronized discharges. Previous report that transgenic mice with specifically disrupted glutamatergic
inputs to parvalbumin-positive interneurons displayed hippocampo-dependent spatial memory
impairment but no epileptic phenotype are also in support of this interpretation[61,62].

But one major advantage offered by the extracellular recording of fIPSPs is the possibility to
evaluate the efficiency, and even the polarity, of perisomatic GABAergic transmission in a variety of
conditions, including pathological conditions in which GABAergic transmission is suspected to be
affected, or even reversed, due to altered neuronal Cl$^-$ homeostasis. Cl$^-$ gradient is dependent on both
recent neuronal activity involving Cl$^-$ currents and the dynamics of pumps and transporters involved in

maintaining physiological Cl⁻ gradient. Our experiments have tested the possibility to detect the consequences of both acute seizures and chronic epilepsy on Cl⁻ regulation and GABAergic transmission. We report that transient hypersynchronized discharges induce a global reversal of neuronal Cl⁻ and polarity of GABAergic transmission, shifting from inhibitory to excitatory for a duration of several tens of seconds before returning to normal polarity and efficiency, but only in the post-ictal period during which neurons (including GABAergic interneurons) are silent, which does not support a contribution of excitatory GABA to the seizures. While this reversal of GABAergic polarity is readily seen as a reversed polarity of fIPSPs, we also report a more subtle effect during the course of epileptogenesis in the KA model of chronic epilepsy. One week after KA injection, we observed that perisomatic GABAergic transmission provided time-locked excitation of pyramidal neurons in the hippocampus, but restricted to a minor subset of neurons, and we did not observe a reversed polarity of fIPSPs. These results suggest that excitatory GABAergic transmission may have only a minor contribution to hippocampal epileptogenesis in contralerally bicuculline-induced acute seizures or during the latent period in the mouse KA model of epilepsy, as recently suggested in the subicular region during the development of kindling-epilepsy [19]. Nevertheless, subtle alterations in the regulation of Cl⁻ homeostasis and GABAergic transmission already operate in the hippocampal circuit during the latent period that precedes the establishment of chronic epilepsy. The functional consequences of defective GABAergic transmission in a minority of neurons, both in terms of coding and circuit dynamics, and in other periods, other models of epilepsy, and other pathologies, remain to be investigated. Independently of seizure generation, reversed GABAergic transmission might significantly affect neuronal processing and information coding.

The extracellular detection of perisomatic IPSPs therefore provides an invaluable tool to evaluate Cl⁻ homeostasis and inhibitory function, in physiological conditions and in situations in which alteration or even reversal of GABAergic transmission is hypothesized to occur. This approach opens new possibilities to evaluate the relevance of the excitatory GABA hypothesis in physiological development, autism spectrum disorders, schizophrenia, various forms of epilepsy and the time course of epileptogenesis, as well as to evaluate the potential of pharmacological interference with neuronal Cl⁻ gradient as an effective therapeutic strategy against major brain diseases.

Acknowledgements. We thank Roustem Khazipov, Rosa Cossart, Valérie Crepel, Tarek Deeb and Isabel Del Pino for useful comments and discussions on a previous version of the manuscript, Delphine Gonzales, Nathalie Aubailly, and all the personnel of the Animal Facility of the NeuroCentre Magendie for animal care, Katy Lecorf for her help regarding virus preparation, the UMR5293 vector core facility for producing AAV vectors, and the laboratory of S.M. Sternson (Howard Hughes Medical Institute) for providing the chemical compound PSEM-89S.

Funding. This work was performed thanks to the following funding sources: INSERM, CNRS (XL),

Région Nouvelle Aquitaine (XL, AB), the Japanese Society for the Promotion of Science (JSPS, XL,

HH), the Brain and Behavior Research Foundation (AB), Agence Nationale pour la Recherche (ANR,

XL), the Fondation Française pour la Recherche sur les Epilepsies (FFRE, XL), the French Ministry of

Research and Education (OD). The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript.

Competing interests. The authors have declared that no competing interests exist.

Author contributions. XL and HH performed pilot experiments and initiated the study. XL conceived

and designed the experiments. OD and XL planned the study. OD, A. Ferreira, AB and XL performed

the experiments. OD and XL analyzed the data. A. Frick contributed with lab and breeding space, and a

fraction of running expenses. OD, AB and XL wrote the paper.

Data availability. The data shown in the paper will be made available upon reasonable request to the

corresponding author.

Code availability. The custom code used for analysis will be made available upon reasonable request

to the corresponding author.

References

1. Blaesse P, Airaksinen MS, Rivera C, Kaila K. Cation-Chloride Cotransporters and Neuronal Function. *Neuron* **61**, 820-838 (2009).

2. Ben-Ari Y. NKCC1 Chloride Importer Antagonists Attenuate Many Neurological and Psychiatric Disorders. *Trends in Neurosciences* **40**, 536-554 (2017).

3. Doyon N, Vinay L, Prescott Steven A, De Koninck Y. Chloride Regulation: A Dynamic Equilibrium Crucial for Synaptic Inhibition. *Neuron* **89**, 1157-1172 (2016).

4. Deidda G, Parrini M, Naskar S, Bozarth IF, Contestabile A, Cancedda L. Reversing excitatory GABAAR signaling restores synaptic plasticity and memory in a mouse model of Down syndrome. *Nature Medecine* **21**, 318-326 (2015).

5. Tyzio R, *et al.* Oxytocin-mediated GABA inhibition during delivery attenuates autism pathogenesis in rodent offspring. *Science* **343**, 675-679 (2014).

6. Lopez-Pigozzi D, *et al.* Altered Oscillatory Dynamics of CA1 Parvalbumin Basket Cells during Theta–Gamma Rhythmopathies of Temporal Lobe Epilepsy. *eNeuro* **3**, ENEURO.0284-0216.2016 (2016).

7. Valero M, *et al.* Mechanisms for Selective Single-Cell Reactivation during Offline Sharp-Wave Ripples and Their Distortion by Fast Ripples. *Neuron* **94**, 1234-1247.e1237 (2017).

8. Menendez de la Prida L, Trevelyan AJ. Cellular mechanisms of high frequency oscillations in epilepsy: On the diverse sources of pathological activities. *Epilepsy Research* **97**, 308-317 (2011).

9. Cellot G, Cherubini E. GABAergic Signaling as Therapeutic Target for Autism Spectrum Disorders. *Frontiers in Pediatrics* **2**, (2014).

10. Barmashenko G, Hefft S, Aertsen A, Kirschstein T, Köhling R. Positive shifts of the GABAA receptor reversal potential due to altered chloride homeostasis is widespread after status epilepticus. *Epilepsia* **52**, 1570-1578 (2011).

11. Pathak HR, *et al.* Disrupted Dentate Granule Cell Chloride Regulation Enhances Synaptic Excitability during Development of Temporal Lobe Epilepsy. *The Journal of Neuroscience* **27**, 14012-14022 (2007).

12. Nardou R, *et al.* Neuronal chloride accumulation and excitatory GABA underlie aggravation of neonatal epileptiform activities by phenobarbital. *Brain* **134**, 987-1002 (2011).

13. Moore YE, Kelley MR, Brandon NJ, Deeb TZ, Moss SJ. Seizing Control of KCC2: A New Therapeutic Target for Epilepsy. *Trends in Neurosciences* **40**, 555-571 (2017).

14. Bragin DE, Sanderson JL, Peterson S, Connor JA, Müller WS. Development of epileptiform excitability in the deep entorhinal cortex after status epilepticus. *European Journal of Neuroscience* **30**, 611-624 (2009).

372   15.   Chen L, *et al.* KCC2 downregulation facilitates epileptic seizures. *Scientific Reports* **7**, 156
373   (2017).

374   16.   Ellender TJ, Raimondo JV, Irkle A, Lamsa KP, Akerman CJ. Excitatory Effects of Parvalbumin-
375   Expressing Interneurons Maintain Hippocampal Epileptiform Activity via Synchronous
376   Afterdischarges. *The Journal of Neuroscience* **34**, 15208-15222 (2014).

377   17.   Di Cristo G, Awad PN, Hamidi S, Avoli M. KCC2, epileptiform synchronization, and epileptic
378   disorders. *Progress in Neurobiology* **162**, 1-16 (2018).

379   18.   Magloire V, Cornford J, Lieb A, Kullmann DM, Pavlov I. KCC2 overexpression prevents the
380   paradoxical seizure-promoting action of somatic inhibition. *Nature Communications* **10**, 1225
381   (2019).

382   19.   Wang Y, *et al.* Depolarized GABAergic Signaling in Subicular Microcircuits Mediates
383   Generalized Seizure in Temporal Lobe Epilepsy. *Neuron* **95**, 92-105.e105 (2017).

384   20.   Riekki R, Pavlov I, Tornberg J, Lauri SE, Airaksinen MS, Taira T. Altered Synaptic Dynamics
385   and Hippocampal Excitability but Normal Long-Term Plasticity in Mice Lacking
386   Hyperpolarizing GABAA Receptor-Mediated Inhibition in CA1 Pyramidal Neurons. *Journal of*
387   *Neurophysiology* **99**, 3075-3089 (2008).

388   21.   Cohen I, Navarro V, Clemenceau S, Baulac M, Miles R. On the Origin of Interictal Activity in
389   Human Temporal Lobe Epilepsy in Vitro. *Science* **298**, 1418-1421 (2002).

390   22.   Huberfeld G, *et al.* Perturbed Chloride Homeostasis and GABAergic Signaling in Human
391   Temporal Lobe Epilepsy. *The Journal of Neuroscience* **27**, 9866-9873 (2007).

392   23.   Pallud J, *et al.* Cortical GABAergic excitation contributes to epileptic activities around human
393   glioma. *Science Translational Medicine* **6**, 244ra289-244ra289 (2014).

394   24.   Gulledge AT, Stuart GJ. Excitatory actions of GABA in the cortex. *Neuron* **37**, 299-309 (2003).

395   25.   van der Pol AN, Obrietan K, Chen G. Excitatory Actions of GABA after Neuronal Trauma. *The*
396   *Journal of Neuroscience* **16**, 4283-4292 (1996).

397   26.   Valeeva G, Tressard T, Mukhtarov M, Baude A, Khazipov R. An Optogenetic Approach for
398   Investigation of Excitatory and Inhibitory Network GABA Actions in Mice Expressing
399   Channelrhodopsin-2 in GABAergic Neurons. *The Journal of Neuroscience* **36**, 5961-5973
400   (2016).

401   27.   Dzhala V, Valeeva G, Glykys J, Khazipov R, Staley K. Traumatic alterations in GABA
402   signaling disrupt hippocampal network activity in the developing brain. *JNeurosci* **32**, 4017-
403   4031 (2012).

404   28.   Dzhala VI, *et al.* NKCC1 transporter facilitates seizures in the developing brain. *Nature*
405   *Medicine* **11**, 1205 (2005).

406   29.   Kahle KT, Staley KJ. The bumetanide-sensitive Na-K-2Cl cotransporter NKCC1 as a potential
407   target of a novel mechanism-based treatment strategy for neonatal seizures. *Neurosurgical*
408   *Focus* **25**, E22 (2008).

30. Marguet SL*, et al.* Treatment during a vulnerable developmental period rescues a genetic
epilepsy. *Nature Medecine* **21**, 1436-1444 (2015).

31. Kourdougli N*, et al.* Depolarizing γ-aminobutyric acid contributes to glutamatergic network
rewiring in epilepsy. *Annals of Neurology* **81**, 251-265 (2017).

32. Brandt C, Nozadze M, Heuchert N, Rattka M, Löscher W. Disease-Modifying Effects of
Phenobarbital and the NKCC1 Inhibitor Bumetanide in the Pilocarpine Model of Temporal
Lobe Epilepsy. *The Journal of Neuroscience* **30**, 8602-8612 (2010).

33. Römermann K*, et al.* Multiple blood-brain barrier transport mechanisms limit bumetanide
accumulation, and therapeutic potential, in the mammalian brain. *Neuropharmacology* **117**,
182-194 (2017).

34. Donovan MD, Schellekens H, Boylan GB, Cryan JF, Griffin BT. In vitro bidirectional
permeability studies identify pharmacokinetic limitations of NKCC1 inhibitor bumetanide.
European Journal of Pharmacology **770**, 117-125 (2016).

35. Puskarjov M, Kahle KT, Ruusuvuori E, Kaila K. Pharmacotherapeutic targeting of cation-
chloride cotransporters in neonatal seizures. *Epilepsia* **55**, 806-818 (2014).

36. Astorga G*, et al.* An excitatory GABA loop operating in vivo. *Frontiers in Cellular
Neuroscience* **9**, (2015).

37. Assaf F, Schiller Y. The antiepileptic and ictogenic effects of optogenetic neurostimulation of
PV-expressing interneurons. *Journal of Neurophysiology* **116**, 1694-1704 (2016).

38. Lovett-Barron M*, et al.* Regulation of neuronal input transformations by tunable dendritic
inhibition. *NatNeurosci* **15**, 423-423 (2012).

39. Lu T, Trussell LO. Mixed excitatory and inhibitory GABA-mediated transmission in chick
cochlear nucleus. *J Physiol* **535**, 125-131 (2001).

40. Glickfeld LL, Roberts JD, Somogyi P, Scanziani M. Interneurons hyperpolarize pyramidal cells
along their entire somatodendritic axis. *Nature Neuroscience* **12**, 21-23 (2009).

41. Bazelot M, Dinocourt C, Cohen I, Miles R. Unitary inhibitory field potentials in the CA3 region
of rat hippocampus. *J Physiol* **588**, 2077-2090 (2010).

42. Beyeler A, Retailleau A, Molter C, Mehidi A, Szabadics J, Leinekugel X. Recruitment of
Perisomatic Inhibition during Spontaneous Hippocampal Activity. *PLoSOne* **8**, e66509 (2013).

43. Glykys J*, et al.* Local Impermeant Anions Establish the Neuronal Chloride Concentration.
Science **343**, 670-675 (2014).

44. Fujiwara-Tsukamoto Y, Isomura Y, Nambu A, Takada M. Excitatory GABA input directly
drives seizure-like rhythmic synchronization in mature hippocampal CA1 pyramidal cells.
Neuroscience **119**, 265-275 (2003).

45. French JA*, et al.* Characteristics of medial temporal lobe epilepsy: I. Results of history and
physical examination. *Annals of Neurology* **34**, 774-780 (1993).

46.   Riban V, Bouilleret V, Pham-Lê BT, Fritschy JM, Marescaux C, Depaulis A. Evolution of
hippocampal epileptic activity during the development of hippocampal sclerosis in a mouse
model of temporal lobe epilepsy. *Neuroscience* **112**, 101-111 (2002).

47.   Löscher W, Puskarjov M, Kaila K. Cation-chloride cotransporters NKCC1 and KCC2 as
potential targets for novel antiepileptic and antiepileptogenic treatments. *Neuropharmacology*
69, 62-74 (2013).

48.   MacKenzie G, O'Toole KK, Moss SJ, Maguire J. Compromised GABAergic inhibition
contributes to tumor-associated epilepsy. *Epilepsy Research* **126**, 185-196 (2016).

49.   Chavas J, Marty A. Coexistence of Excitatory and Inhibitory GABA Synapses in the Cerebellar
Interneuron Network. *The Journal of Neuroscience* **23**, 2019-2031 (2003).

50.   Kirmse K, Kummer M, Kovalchuk Y, Witte OW, Garaschuk O, Holthoff K. GABA depolarizes
immature neurons and inhibits network activity in the neonatal neocortex in vivo. *Nature*
Communications **6**, 7750 (2015).

51.   Chavas J, Forero ME, Collin T, Llano I, Marty A. Osmotic tension as a possible link between
GABA(A) receptor activation and intracellular calcium elevation. *Neuron* **44**, 701-713 (2004).

52.   Minlebaev M, Valeeva G, Tcheremiskine V, Coustillier G, Khazipov R. Cell-attached
recordings of responses evoked by photorelease of GABA in the immature cortical neurons.
Frontiers in Cellular Neuroscience **7**, (2013).

53.   Krook-Magnuson E, Armstrong C, Oijala M, Soltesz I. On-demand optogenetic control of
spontaneous seizures in temporal lobe epilepsy. *Nature Communications* **4**, 1376 (2013).

54.   Sik A, Penttonen M, Ylinen A, Buzsaki G. Hippocampal CA1 interneurons: an in vivo
intracellular labeling study. *The Journal of Neuroscience* **15**, 6651-6665 (1995).

55.   Buhl EH, Han ZS, Lorinczi Z, Stezhka VV, Karnup SV, Somogyi P. Physiological properties
of anatomically identified axo-axonic cells in the rat hippocampus. *Journal of Neurophysiology*
71, 1289-1307 (1994).

56.   Khalilov I, Minlebaev M, Mukhtarov M, Khazipov R. Dynamic Changes from Depolarizing to
Hyperpolarizing GABAergic Actions during Giant Depolarizing Potentials in the Neonatal Rat
Hippocampus. *The Journal of Neuroscience* **35**, 12635-12642 (2015).

57.   Staley KJ, Mody I. Shunting of excitatory input to dentate gyrus granule cells by a depolarizing
GABAA receptor-mediated postsynaptic conductance. *Journal of Neurophysiology* **68**, 197-212
(1992).

58.   Tyzio R*, et al.* Postnatal changes in somatic gamma-aminobutyric acid signalling in the rat
hippocampus. *European Journal of Neuroscience* **27**, 2515-2528 (2008).

59.   Somogyi P, Klausberger T. Defined types of cortical interneurone structure space and spike
timing in the hippocampus. *J Physiol* **562**, 9-26 (2005).

60.   Miles R, Toth K, Gulyas AI, Hajos N, Freund TF. Differences between somatic and dendritic
inhibition in the hippocampus. *Neuron* **16**, 815-823 (1996).

61.   Racz A, Ponomarenko AA, Fuchs EC, Monyer H. Augmented hippocampal ripple oscillations

in mice with reduced fast excitation onto parvalbumin-positive cells. *The Journal of*

Neuroscience **29**, 2563-2568 (2009).

62.   Fuchs EC*, et al.* Recruitment of parvalbumin-positive interneurons determines hippocampal

function and associated behavior. *Neuron* **53**, 591-604 (2007).

Figure 1: Spontaneous extracellular field inhibitory post-synaptic potentials (fIPSPs) in vivo

(a) Schematic of electrode track and histological verification of recording location within the CA3 hippocampal region (**DiI-labeled silicon probe in red**, **DAPI staining in blue**).

(b) Exemple trace of spontaneous wide-band (0.1Hz-9KHz) field-recording activity in the CA3 pyramidal layer, displaying 4 channels (10-20µm distance between recording sites) of the same shank of a 2-shanks Neuronexus Buzsaki16 silicon probe. Field-events of positive polarity (upward deflections), fIPSPs (peaks, **dashed blue lines)**. Fast downward deflections, action potentials (AP), also labeled below as raster display (**red ticks,** multi-unit activity). Note the transient interruption of neuronal firing after fIPSPs.

(c) Comparison of fIPSP decay (left, superimposed average traces normalized on amplitude, right, fIPSPs decay times, mean± SD, ** p<0.01) in control (**black**) and in presence of diazepam (**grey**). Note prolonged fIPSP decay under diazepam.

(d) Peri-event time histogram (mean and SD, n=8 WT mice, time bin 1 ms) between all the spikes discharged by putative pyramidal cells and fIPSPs (reference, peak-time), shown as average trace on top at the same time scale and aligned on peak (**dotted blue line**). Baseline mean and 2SD are respectively shown as **plain and dotted horizontal grey lines**. Note increased pyramidal cell discharge in the 10 ms prior to fIPSP, and the strong inhibition (trough, max inhibition) that follows. **Inset**, schematic illustration of the recurrent excitatory/inhibitory loop: **(1)** the collective discharge of interconnected pyramidal cells activates the discharge of local interneurons, resulting **(2)** in perisomatic fIPSP and inhibition of pyramidal population activity.

(e) Diagram illustrating firing rates (mean ± SD, same 8 mice as in D; baseline level = 100%, calculated from -50 ms to -30 ms) at **peak** (bin of maximum firing in the 10ms prior to fIPSPs), **trough** (bin of minimum firing in the 4ms post-fIPSPs), and **recovery** (last of post-fIPSPs time bin below baseline - 2SD) time points. **Inset**, proportion of pyramidal cells (n=139 from 8 mice) with a significant post-fIPSPs trough (ie inhibited vs not-inhibited neurons).

Figure 2: In vivo fIPSPs generated by optogenetic activation of PV interneurons

(a) Schematic representation of electrode and optic fiber positions in PV-Cre mice with AAV-driven ChR2 and GFP expression in CA3 PV interneurons.

(b-e) Confocal stacks of GFP-positive (**green**) and PV-positive (**red**) immunostaining in the hippocampal CA3 region. (**b**, **scale bar**, 100μm) GFP immuno staining. (**c-e**, **scale bar**, 50μm), higher magnification of the CA3a region delimited by the **white rectangle** in B). (**C**) GFP immuno-staining. (**d**) PV immunostaining. (**e**) Overlay. Note the co-localization on the 4 identified PV interneurons (arrow-heads).

(f) Peri-event time histogram (mean ± SD, n=5 mice, time bin 1 ms) between all the spikes discharged by putative pyramidal cells and optogenetically evoked (laser ON 2ms, **blue bar**) fIPSPs (reference), shown as superimposed (**grey**) and average (**black**) traces (n=60 from a single representative recording) on top, aligned and at the same time scale. Note the powerful time-locked inhibition of pyramidal cell firing in response to fIPSPs evoked by the optogenetic stimulation of PV interneurons.

Figure 3: Inhibition of fIPSPs by pharmacogenetic inactivation of PV interneurons

(a) Schematic representation of electrode and injection site in PV-Cre mice with AAV-driven PSAM-GlyR and GFP expression in CA3 PV interneurons.

(b-e) Confocal stacks of GFP-positive (**green**) and PV-positive (**red**) immunostaining in the hippocampal CA3 region. (**b**, **scale bar**, 100μm) GFP immuno staining. (**c-e**, **scale bar**, 50μm), higher magnification of the CA3a region delimited by the **white rectangle** in B). **(c)** GFP immuno-staining. **(d)** PV immunostaining. **(e)** Overlay. Note the co-localization of PV and PSAM-GlyR immuno-staining on the 3 labeled interneurons (**arrow-heads**).

(f) Example traces of spontaneous wide-band (0.1Hz-9KHz) field-recording activity in the CA3 pyramidal layer, in control (left, **baseline**) and in presence of the exogenous ligand PSEM-89S (right, **PSEM**). Note the drastic reduction in fIPSPs expression (*).

(g-h) Summary plots (mean ± SD, n=5 mice) of fIPSP frequency **(g)** and pyramidal population firing rate **(h)** in control (baseline) and in presence of PSEM (* p<0.05).

(i) Individual pyramidal cells firing rates (n=36 putative pyramidal cells from 3 mice) in control ("**before**" is the 10min period that immediately preceeds PSEM-89S injection) and in presence of PSEM ("**after**" period is from 5 to 15min after PSEM-89S injection). **Black line**, bissector (y=x). **Red line**, linear regression line. Note that the pharmacogenetic blockade of PV interneurons with PSAM-GlyR redistributed individual firing rates without any significant change on global population firing activity.

Figure 4: Transiently reversed fIPSPs and perisomatic GABAergic synaptic transmission following acute seizures

(a) Schematic representation of electrode and optic fiber positions in the left hippocampus of PV-Cre mice with AAV-driven ChR2 expression in CA3 PV interneurons, and contralateral injection of the convulsivant bicuculline.

(b) Exemple trace (wide-band LFP) of a seizure as acute response to contralateral bicuculline injection.

(c) Time course of individual fIPSPs amplitude (**black dots**, spontaneous, **blue dots**, evoked by optogenetic activation of PV interneurons), relative to acute seizures induced by the contralateral injection of bicuculline (**vertical grey bar**, 3 seizures from the same animal, aligned by their ending time). **Insets**, representative individual field responses (optogenetically evoked fIPSPs; **horizontal blue bars**, laser ON) in control (**pre-seizure**), during the period right after the seizure (**reversed polarity**), or after recovery (**post-seizure**). Note transiently evoked spiking activity induced by fIPSPs of reversed polarity right after the seizure and progressive recovery of fIPSPs amplitude.

(d) Peri-event time histogram (mean and SD, time bin 1ms) between all the spikes discharged by putative pyramidal cells (n=4 mice) and spontaneous (in **black**, n=31682 events) or optogenetically evoked (in **blue**, n=2756 events) fIPSPs (reference) from the pre- and post-seizure periods, shown as average traces on top (spontaneous, **black**; evoked, **blue**) at the same time scale and aligned on peak.

(e) Peri-event time histogram (mean and SD, time bin 1ms) between multi-unit activity and optogenetically evoked fIPSPs (reference) in the period right after the seizure (n=57 fIPSPs from 6 seizures from 4 mice), shown as average traces on top at the same time scale and aligned on peak. Note the clear time-locked excitation of neuronal firing and reversed polarity of the fIPSPs evoked by the optogenetic stimulation of PV interneurons in the period right after the seizure.

Figure 5: Normal fIPSP polarity but GABA-mediated time-locked excitation in head-fixed KA-treated mice
(a) CA3 neuronal activity was recorded from urethane-anesthetized WT mice one week after contralateral, intra-hippocampal KA injection.
(b) Exemple trace (wide-band LFP) of interictal burst (interictal spikes, **blue arrow-heads**).
(c) Exemple traces of spontaneous CA3 pyramidal layer activity (left) and superimposed and averaged traces of fIPSPs (right), in control (top) and KA (below) conditions. Note the presence of normal polarity, frequent fIPSPs, in both conditions.
(d-f) Individual examples of putative pyramidal cells displaying the three distinct types of observed responses to spontaneous fIPSPs in KA-treated mice (time locked excitation **(d)**, time-locked inhibition **(e)**, or no clear change in net firing rate (**f, non responding**)). **Upper left**, average waveform from the 8 channels of the recording electrode. **Upper right**, spike autocorrelogram. **Middle**, raster display of spike firing aligned on fIPSP ocurrence (**vertical blue line**, fIPSP peak). **Lower plot**, normalized peri-event time histogram (time bin, 1ms) of the neuron discharge (AP count) relative to fIPSP peak (reference, **vertical blue line**). **Inset (d)**, similar peri-event time histogram but ignoring action potential bursts (ie all spikes with preceding inter spike interval < 10ms omitted from the histogram).