Enhanced Disease Susceptibility 5 (EDS5) is required for *N*-hydroxy pipecolic acid formation

Dmitrij Rekhter[1], Lennart Mohnike[1], Kirstin Feussner[1,2], Krzysztof Zienkiewicz[1,2], Yuelin Zhang[3], Ivo Feussner[1,2,4,*]

[1] University of Goettingen, Albrecht-von-Haller-Institute for Plant Sciences, Department of Plant Biochemistry, D-37077 Goettingen, Germany.

[2] University of Goettingen, Goettingen Center for Molecular Biosciences (GZMB), Service Unit for Metabolomics and Lipidomics, D-37077 Goettingen, Germany.

[3] University of British Columbia, Department of Botany, Vancouver, BC V6T 1Z4, Canada.

[4] University of Goettingen, Goettingen Center for Molecular Biosciences (GZMB), Department of Plant Biochemistry, D-37077 Goettingen, Germany.

*Correspondence to Ivo Feussner: ifeussn@uni-goettingen.de.

Abstract

Pipecolic acid is essential for the establishment of systemic acquired resistance in plants. It is synthesized in the plastid and further processed in the cytosol to its active form *N*-hydroxy pipecolic acid. Here we provide strong evidence that the exporter Enhanced Disease Susceptibility 5 is not only required for the biosynthesis of salicylic acid, but also of *N*-hydroxy pipecolic acid, suggesting that it represents a convergent point of plant immunity.

Plants have to adapt to biotic and abiotic challenges in nature. In order to cope with these risks, they produce signaling molecules which are critical for the activation of their defense system[1]. The accumulation of salicylic acid (SA) and pipecolic acid (Pip) at the site of infection as well as in systemic

tissues is a key event for establishing the immune response against biotrophic pathogens[2]. The first

step in SA biosynthesis is the conversion of chorismic acid (CA) to isochorismic acid (IC) by

Isochorismate Synthase 1 (ICS1) in plastids[3]. We recently showed that Enhanced Disease Susceptibility

5 (EDS5) is essential for the export of IC from plastids into the cytosol[4]. IC is further processed to

isochorismate-9-glutamate by avrPphB Susceptible 3 (PBS3) that subsequently decomposes to SA in a

non-enzymatic process. Loss of any of these three genes leads to a drastic reduction of pathogen-

induced SA production and to enhanced disease susceptibility[5,6].

Pip has been shown to be as crucial as SA for plant immunity[7]. Its biosynthesis occurs in plastids starting

from lysine[8,9]. First, the α-aminotransferase AGD2-like Defense Response Protein 1 (ALD1) catalyzes

the formation of ε-amino α-keto caproic acid, which spontaneously cyclizes to Δ^1-piperideine-2-

carboxylic acid (P2C) in solution. The ketimine reductase SAR-Deficient 4 (SARD4) catalyzes

subsequently the formation of Pip from P2C[10,11]. Pip-based signaling relies on Flavin-dependent

Monooxygenase 1 (FMO1), which is responsible for the *N*-hydroxylation of Pip to yield *N*-hydroxy

pipecolic acid (NHP)[12,13]. This newly discovered compound was proposed to be a crucial regulator of

systemic acquired resistance (SAR)[12]. Although the exact subcellular localization of FMO1 has not been

determined yet, studies of other FMOs strongly suggest a localization on the cytoplasmic face of the

endoplasmic reticulum[14], implying the need for a transporter from the site of Pip biosynthesis to the

location where it is further processed. Both SA and NHP can be further glycosylated to form SA-β-

glucoside (SAG), SA-glucose ester (SGE)[15] and NHP-glycoside (NHP-*O*Glc)[13], respectively. Glycosylation

has been proposed to inactivate plant hormone signaling and is typically facilitated by cytosolic UDP-

dependent glycosyltransferases (UGTs)[16].

Not only pathogen infection, but also abiotic stresses like ozone or UV-C treatment stimulate the

biosynthesis of SA and SAG[17], leading to comparable changes in gene expression[18]. In order to have a

fast and reproducible experimental setup for Pip synthesis, we examined the possibility to induce Pip

production by UV-C stress. Indeed, Pip accumulates over time in *Arabidopsis thaliana* leaves in a similar

course of time (Fig. 1a), as it was described for *Pseudomonas syringae* infection before[12]. Beside Pip,

we also observed increased amounts of NHP and NHP-*O*Glc (of which the MS/MS fragmentation patterns are depicted in Supplementary Fig. 1)[13]. This indicates that Pip oxidation and glycosylation also occur after UV-C treatment. To exclude that the observed synthesis of Pip and NHP-*O*Glc is activated by UV-C-triggered SA accumulation, we examined Pip and NHP-*O*Glc contents in the SA-deficient mutants, *eds5, pbs3* (Fig. 1b-e) and Salicylic acid Induction Deficient 2 (*sid2),* harboring a mutation in the *ICS1* gene (Supplementary Figure 2a and b)*.* As expected, the mutant lines, but not wild type plants, lack both SA and SAG 24 hours after being exposed to UV-C for 20 min (Fig. 1b and c). On the other hand Pip and NHP-*O*Glc accumulation was observed only in wild type, *pbs3* and *sid2*, but surprisingly not in *eds5* plants (Fig. 1d and e). Similar to pathogen-treated plants[12], UV-C stress-induced Pip-biosynthesis and -catabolism do not depend on the presence of SA in UV-C treated plants. The absence of Pip and NHP-*O*Glc in *eds5* raised the question whether EDS5 is responsible for the export of not only the SA precursor IC, but also of Pip. Recently, it was shown that *fmo1* mutants are impaired in the hydroxylation of Pip to NHP. Instead, *fmo1* accumulates large amounts of Pip upon UV-C irradiation (Supplementary Fig. 3), again in a similar extent as described for pathogen assays[12]. In line with FMO1 functioning downstream of Pip synthesis, these results corroborate that EDS5 acts upstream of FMO1 by exporting Pip to the cytosol.

The assumed block in Pip export did not result in an enrichment of Pip in *eds5* plants, despite the induction of Pip biosynthesis on the transcriptional level: Both *SARD4* and *EDS5* were strongly upregulated upon UV-C treatment in the SA deficient *sid2* mutant and wild type plants (Supplementary Figure 2c and d), supporting our metabolite profiling results. In analogy, we observed increased levels of IC in *pbs3*, but not in *eds5* mutant plants upon UV-C treatment (Supplementary Fig. 4). It is likely that a diversion of the metabolic flux prevents a harmful accumulation of both Pip and IC in plastids[19]. Pip may either inhibit its own biosynthesis via a feedback loop or feed into the lysine degradation pathway towards the Krebs cycle[20]. IC on the other hand might be channeled towards the synthesis of aromatic amino acids.

In order to exclude that the observed absence of NHP and NHP-*O*Glc in *eds5* mutants is due to their

inability to accumulate SA, we tested whether external SA supply could induce Pip and NHP-*O*Glc

production in *eds5* and *pbs3* plants. As reported, SA treatment triggers the biosynthesis of Pip and

NHP-*O*Glc in wild type plants (Fig. 2a)[7]. SA biosynthesis is impaired in the *pbs3* mutant, but drenching

the soil with SA still initiated Pip biosynthesis in these plants (Fig. 2a). However, as with UV-C

treatment, external SA also did not lead to Pip accumulation in *eds5* mutants. This is consistent with

EDS5 representing a plastidial exporter of Pip independent of SA. When Pip irrigation was used to

trigger *in planta* Pip biosynthesis, we detected increased levels of the downstream product NHP-*O*Glc

in wild type plants (Fig. 2b). In *eds5* and the Pip biosynthesis mutant *sard4*, the levels of NHP-*O*Glc are

reduced by more than 60%. This suggests that in the mutants only the external Pip was metabolized

further, whereas *de novo* Pip biosynthesis did not occur. The ability of *eds5* plants to convert

exogenously applied Pip into NHP-*O*Glc supports FMO1 being active outside of plastids and not

affected by the loss of function of *EDS5*.

Taken together, we show that UV-C treatment is sufficient to induce the production of Pip and its

metabolites. This process does not require the presence of SA and can thus be used to study the SA-

independent branch of plant defense signaling. Moreover, we identified here a previously unknown

connection between Pip and EDS5, a protein that, so far, was only recognized for its involvement in SA

biosynthesis[17]. Most likely EDS5 is also responsible for the export of Pip from plastids into the cytosol,

where it is further processed by FMO1. This spatial separation of Pip biosynthesis and the site of

activation to NHP adds an additional layer for regulation. Our study implies that EDS5 serves as the

central hub in the biosynthesis of two major defense signaling molecules, SA and NHP (Fig. 2c). It is

surprising that no pathogen effector has been found to target EDS5 and thereby exploit this key point

in plant immunity.

Figures

Figure 1 UV-C treatment of *Arabidopsis thaliana* leads to an accumulation of the signaling

compounds SA and Pip and their corresponding glycosides. a, Relative amounts (counts per second)

of pipecolic acid (Pip, white bars) and its downstream products *N*-hydroxy pipecolic acid (NHP, grey

bars) and *N*-hydroxy pipecolic acid glycoside (NHP-*O*Glc, black bars) in *Arabidopsis* wild type leaves

(Col-0) at different time points after UV-C light treatment. **b-e,** Relative amounts of salicylic acid (SA)

and its glycoside SAG (nmol g^{-1} leaf fresh weight [f.w.]) and of Pip and its glycoside NHP-*O*Glc (counts

per second) in leaves of wild type (Col-0), respectively *eds5*-3 and *pbs3*-1 mutant plants 24 hours after

UV-C treatment in comparison to untreated plants. Bars represent the mean ± STD of three biological

replicates. Statistical differences among replicates are labeled with different letters ($P < 0.05$, one-way

ANOVA and post hoc Tukey's Test; n = 3). The experiments were repeated twice with similar results.

Figure 2 Root application of either SA or Pip induces Pip biosynthesis and processing in leaves. a,

Relative amounts (counts per second) of pipecolic acid (Pip) in leaves of wild type (Col-0) *eds5*-3 and

pbs3-1 plants 24 hours after soil drenching with water (-) or SA (+). **b,** Relative amounts (counts per

second) of *N*-hydroxy pipecolic acid glycoside (NHP-*O*Glc) in leaves of wild type (Col-0), *eds5*-3 and

sard4-5 plants 24 hours after soil drenching with water (-) or Pip (+). Bars represent the mean ± STD of

three biological replicates. Statistical differences among replicates are labeled with different letters (P

< 0.05, one-way ANOVA and post hoc Tukey's Test; n = 3). The experiment was repeated twice with

similar results. **c,** A working model depicting EDS5 as the central hub in the biosynthesis of the signaling

compounds SA and NHP. The induction of these pathways was facilitated here by UV-C light treatment.

Methods

Plant material. *Arabidopsis* plants were grown in a chamber at 22 °C with a 16 h light period and 60% relative humidity for 4-5 weeks. For our experiments, we use *Arabidopsis* ecotype Col-0 and the following mutants in this background: *eds5-3*[21], *pbs3-1*[6], *sard4-5*[10], and fmo1-1[12], which were described previously.

UV-C and soil drench treatment. For the UV-C treatment, we followed previous protocol[17]. In short, 4-5 week old *Arabidopsis* plants were exposed to UV-C light (254 nm) for 20 min at 50 cm distance to the lamp (TUV T8 30W, Philips) for the induction of SA and Pip biosynthesis. For the treatment with SA and Pip, we followed previously described protocols[7,21]. 4-5 week old plants, were soil drenched with either 10 mL water, 10 mL of a 5 mM Pip solution (P45850, Sigma) or 10 mL of a 5 mM SA solution (S5922, Sigma), equals to 50 µM final concentration. Samples were collected 24 hours after treatment and metabolites were extracted and analyzed as described earlier[10,22]. The deviation of exact mass to accurate mass for Pip, NHP and NHP-*O*Glc was less than ±2 mDa in the untargeted metabolite analysis. SA and SAG were quantified based on internal D4-SA standard (C/D/N Isotopes Inc., Pointe-Claire, Canada). The NHP standard was chemical synthesized as described in Hartmann et al., 2018[12]. The MS/MS spectra of NHP and NHP-*O*Glc corresponds to the results from Chen et al., 2018[13].

Quantitative real-time PCR analysis. To analyze the expression of *SARD4* (At5g52810) and *EDS5* (AT4G39030) after UV-C treatment, total RNA was isolated from frozen leaves with the Spectrum™ Plant Total RNA Kit (Sigma) following the manufacturer's instruction. One microgram of RNA was treated with DNase I (Thermo Fisher Scientific) and cDNA was synthesized using Revert Aid H Minus Reverse Transcriptase (Thermo Fisher Scientific). Quantitative RT-PCR was performed using Takyon No ROX SYBR Mastermix blue dTTP (Kaneka Eurogentec) in reaction volume 20 µl. The gene ACTIN8 (At1g49240) was used as a control. Each reaction was performed with material from plants harvested

154 from three independent samples in iQ5 real time detection system (Bio-Rad). Primers are depicted in

155 Supplementary Table 1.

Acknowledgments

We are grateful to Ulf Diedrichsen and Brigitte Worbs for the synthesis of the NHP authentic standard

and to Amelie Kelly for critical reading the manuscript. This work was supported by the DFG (IRTG 2172

"PRoTECT" program of the Göttingen Graduate Center of Neurosciences, Biophysics, and Molecular

Biosciences.) to D.R., L.M. and I.F.; I.F. was additionally supported by the DFG (ZUK 45/2010 and INST

186/822-1). Y.Z. was supported by Natural Sciences and Engineering Research Council of Canada,

Canada Foundation for Innovation, and British Columbia Knowledge Development Fund.

Author contributions

D.R., K.F. and I.F. conceived and designed the experiments. D.R., L.M., and K.Z. performed the

experiments. D.R., K.F., Y.Z., and I.F. wrote the article. **Competing interests:** Authors declare no

competing interests. **Data and materials availability:** All data is available in the main text or the

supplementary materials. The authors responsible for distribution of materials integral to the findings

presented in this article are: Ivo Feussner (ifeussn@uni-goettingen.de) and Yuelin Zhang

(yuelin.zhang@ubc.ca).

References

[1] Fu, Z. Q. & Dong, X. Systemic acquired resistance: Turning local infection into global defense. *Annu.*
Rev. Plant Biol. **64**, 839-863, doi:10.1146/annurev-arplant-042811-105606 (2013).
[2] Bernsdorff, F. *et al.* Pipecolic acid orchestrates plant systemic acquired resistance and defense
priming via salicylic acid-dependent and -independent pathways. *Plant Cell* **28**, 102-129,
doi:10.1105/tpc.15.00496 (2016).
[3] Garcion, C. *et al.* Characterization and biological function of the ISOCHORISMATE SYNTHASE2 gene
of Arabidopsis. *Plant Physiol.* **147**, 1279-1287, doi:10.1104/pp.108.119420 (2008).
[4] Rekhter, D. *et al.* PBS3 is the missing link in plant-specific isochorismate-derived salicylic acid
biosynthesis. *bioRxiv*, 600692, doi:10.1101/600692 (2019).
[5] Nawrath, C. & Métraux, J. P. Salicylic acid induction-deficient mutants of Arabidopsis express *PR-2*
and *PR-5* and accumulate high levels of camalexin after pathogen inoculation. *Plant Cell* **11**, 1393-
1404, doi:10.1105/tpc.11.8.1393 (1999).
[6] Nobuta, K. *et al.* The GH3 acyl adenylase family member PBS3 regulates salicylic acid-dependent
defense responses in Arabidopsis. *Plant Physiol.* **144**, 1144-1156, doi:10.1104/pp.107.097691
(2007).
[7] Navarova, H., Bernsdorff, F., Döring, A.-C. & Zeier, J. Pipecolic acid, an endogenous mediator of
defense amplification and priming, is a critical regulator of inducible plant immunity. *Plant Cell* **24**,
5123-5141, doi:10.1105/tpc.112.103564 (2012).

8   Song, J. T., Lu, H. & Greenberg, J. T. Divergent roles in *Arabidopsis thaliana* development and
defense of two homologous genes, *ABERRANT GROWTH AND DEATH2* and *AGD2-LIKE DEFENSE*
RESPONSE PROTEIN1, encoding novel aminotransferases. *Plant Cell* **16**, 353-366,
doi:10.1105/tpc.019372 (2004).
9   Sharma, S., Shinde, S. & Verslues, P. E. Functional characterization of an ornithine cyclodeaminase-
like protein of *Arabidopsis thaliana*. *BMC Plant Biol* **13**, 182, doi:10.1186/1471-2229-13-182 (2013).
10 Ding, P. *et al.* Characterization of a pipecolic acid biosynthesis pathway required for systemic
acquired resistance. *Plant Cell* **28**, 2603-2615, doi:10.1105/tpc.16.00486 (2016).
11 Hartmann, M. *et al.* Biochemical principles and functional aspects of pipecolic acid biosynthesis in
plant immunity. *Plant Physiol.* **174**, 124-153, doi:10.1104/pp.17.00222 (2017).
12 Hartmann, M. *et al.* Flavin monooxygenase-generated *N*-hydroxypipecolic acid is a critical element
of plant systemic immunity. *Cell* **173**, 456-469, doi:10.1016/j.cell.2018.02.049 (2018).
13 Chen, Y.-C. *et al.* *N*-hydroxy-pipecolic acid is a mobile metabolite that induces systemic disease
resistance in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **115**, E4920-E4929,
doi:10.1073/pnas.1805291115 (2018).
14 Zhang, M. & Robertus, J. D. Molecular cloning and characterization of a full-length flavin-dependent
monooxygenase from yeast. *Arch. Biochem. Biophys.* **403**, 277-283, doi:10.1016/S0003-
9861(02)00237-0 (2002).
15 Song, J. T. Induction of a salicylic acid glucosyltransferase, AtSGT1, is an early disease response in
Arabidopsis thaliana. *Molecules and Cells* **22**, 233-238 (2006).
16 Li, Y., Baldauf, S., Lim, E.-K. & Bowles, D. J. Phylogenetic analysis of the UDP-glycosyltransferase
multigene family of *Arabidopsis thaliana*. *J. Biol. Chem.* **276**, 4338-4343,
doi:10.1074/jbc.M007447200 (2001).
17 Serrano, M. *et al.* Export of salicylic acid from the chloroplast requires the multidrug and toxin
extrusion-like transporter EDS5. *Plant Physiol.* **162**, 1815-1821, doi:10.1104/pp.113.218156 (2013).
18 Kilian, J. *et al.* The AtGenExpress global stress expression data set: protocols, evaluation and model
data analysis of UV-B light, drought and cold stress responses. *Plant J.* **50**, 347-363,
doi:10.1111/j.1365-313X.2007.03052.x (2007).
19 Widhalm, J. R. *et al.* Identification of a plastidial phenylalanine exporter that influences flux
distribution through the phenylalanine biosynthetic network. *Nat. Commun.* **6**, 8142,
doi:10.1038/ncomms9142 (2015).
20 Kirma, M., Araújo, W. L., Fernie, A. R. & Galili, G. The multifaceted role of aspartate-family amino
acids in plant metabolism. *J. Exp. Bot.* **63**, 4995-5001, doi:10.1093/jxb/ers119 (2012).
21 Nawrath, C. EDS5, an essential component of salicylic acid-dependent signaling for disease
resistance in Arabidopsis, is a member of the MATE transporter family. *Plant Cell* **14**, 275-286,
doi:10.1105/tpc.010376 (2002).
22 Kusch, S. *et al.* Arabidopsis *mlo3* mutant plants exhibit spontaneous callose deposition and signs of
early leaf senescence. *bioRxiv*, 558122, doi:10.1101/558122 (2019).