The plastidial exporter Enhanced Disease Susceptibility 5 is required for the biosynthesis of *N*-hydroxy pipecolic acid

Short title: EDS5 is required for NHP biosynthesis

Dmitrij Rekhter[1], Lennart Mohnike[1], Kirstin Feussner[1,2], Krzysztof Zienkiewicz[1,2], Yuelin Zhang[3], Ivo Feussner[1,2,4,*]

[1] University of Goettingen, Albrecht-von-Haller-Institute for Plant Sciences, Department of Plant Biochemistry, D-37077 Goettingen, Germany.

[2] University of Goettingen, Goettingen Center for Molecular Biosciences (GZMB), Service Unit for Metabolomics and Lipidomics, D-37077 Goettingen, Germany.

[3] University of British Columbia, Department of Botany, Vancouver, BC V6T 1Z4, Canada.

[4] University of Goettingen, Goettingen Center for Molecular Biosciences (GZMB), Department of Plant Biochemistry, D-37077 Goettingen, Germany.

*Correspondence to Ivo Feussner: ifeussn@uni-goettingen.de.

Abstract

Pipecolic acid is essential for the establishment of systemic acquired resistance in plants. It is synthesized in the plastid and further processed in the cytosol to its active form *N*-hydroxy pipecolic acid. Here we provide strong evidence that the exporter Enhanced Disease Susceptibility 5 is required for the biosynthesis of not only salicylic acid, but also *N*-hydroxy pipecolic acid, suggesting that it represents a convergent point of plant immunity.

Introduction

Plants face numerous biotic and abiotic challenges in nature. In order to cope with these threats, they produce a variety of metabolites. These small molecules are critical for the activation of their defense system[1]. The accumulation of salicylic acid (SA) and pipecolic acid (Pip) at the site of infection as well as in systemic tissues is a key event for the successful immune response against biotrophic pathogens[2]. The first step in SA biosynthesis, the conversion of chorismic acid (CA) to isochorismic acid (IC) by Isochorismate Synthase 1 (ICS1) in plastids[3]. We previously showed that Enhanced Disease Susceptibility 5 (EDS5) is essential for the export of IC from plastids into the cytosol. IC is further processed to isochorismate-9-glutamate by avrPphB Susceptible 3 (PBS3) that subsequently decomposes to SA in a non-enzymatical process[4]. Loss of any of these three genes leads to a drastic reduction of pathogen-induced SA production and to enhanced disease susceptibility[5,6].

Pip has been shown to be equally crucial for plant immunity as SA[7]. Its biosynthesis occurs in plastids starting from lysine[8,9]. First, the α-aminotransferase AGD2-like Defense Response Protein 1 (ALD1) catalyzes the formation of ε-amino α-keto caproic acid, which spontaneously cyclizes to Δ^1-piperideine-2-carboxylic acid (P2C) in solution. The ketimine reductase SAR-Deficient 4 (SARD4) catalyzes subsequently the formation of Pip from P2C[10]. Pip-based signaling relies on Flavin-dependent Monooxygenase 1 (FMO1), which is responsible for the *N*-hydroxylation of Pip to yield *N*-hydroxy pipecolic acid (NHP)[11,12]. This newly discovered compound was proposed to be a crucial regulator of

systemic acquired resistance (SAR)[11]. Although the exact subcellular localization of FMO1 has not been

determined yet, studies of other FMOs strongly suggest a localization on the cytoplasmic face of the

endoplasmic reticulum[13], implying the need for a transporter from the site of Pip biosynthesis to the

location where it is further processed. Both SA and NHP can be further glycosylated to form SA-β-

glucoside (SAG), SA-glucoseester (SGE)[14] and NHP-glycoside (NHP-OGlc)[12], respectively. Glycosylation

has been proposed to inactivate plant hormone signaling and is typically facilitated by cytosolic UDP-

dependent glycosyltransferases (UGTs)[15].

Results and discussion

Not only pathogenic infection, but also abiotic stresses like ozone or UV-C treatment stimulate the

biosynthesis of SA and SAG[16], leading to similar changes in gene expression[17]. In order to have a fast

and reproducible test system for Pip synthesis, we examined the possibility to induce Pip production

by UV-C stress. Indeed, Pip accumulates over time in *Arabidopsis thaliana* leaves in a similar course of

time (Fig. 1a), as it was described for *Pseudomonas syringae* infection before[11]. Beside Pip, we also

observed increased amounts of NHP and NHP-OGlc (of which the MS/MS fragmentation patterns are

depicted in Fig. S1)[12]. This indicates that Pip oxidation and glycosylation similarly occur after UV-C

treatment. To exclude that the observed synthesis of Pip and NHP-OGlc is activated by UV-C-triggered

SA accumulation, we examined Pip and NHP-OGlc contents in the SA-deficient mutants, *eds5, pbs3*

(Fig. 1b-e) and Salicylic acid Induction Deficient 2 (*sid2),* harboring a mutation in the *ICS1* gene (Fig. S2a

and b)*.* As expected, the mutant lines, but not wild type plants, lack both SA and SAG 24 hours after

being exposed to UV-C for 20 min (Fig. 1b and c). On the other hand Pip and NHP-OGlc accumulation

was observed only in wild type, *pbs3* and *sid2*, but surprisingly not in *eds5* plants (Fig. 1d and e). Similar

to pathogen-treated plants[11], UV-C stress- induced Pip-biosynthesis and -processing do not depend on

the presence of SA in UV-C treated plants. The absence of Pip and NHP-OGlc in *eds5* raised the question

whether EDS5 is responsible for the export of not only the SA precursor IC, but also of Pip. Recently, it

was shown that *fmo1* mutants are impaired in the hydroxylation of Pip to NHP. Instead, they

accumulate large amounts of Pip upon UV-C irradiation (Fig. S3), again in a similar extent as described

for pathogen assays[11]. In line with FMO1 functioning downstream of Pip synthesis, these results

corroborate that EDS5 acts upstream of FMO1 by exporting Pip to the cytosol.

The assumed block in Pip export did not result in an enrichment of Pip in *eds5* plants, despite the

induction of Pip biosynthesis on the transcriptional level: Both *SARD4* and *EDS5* were strongly

upregulated upon UV-C treatment in the SA deficient *sid2* mutant and wild type plants (Fig. S2c and

d), supporting our metabolite profiling results. In analogy, we observed increased levels of IC in *pbs3*,

but not in *eds5* mutant plants upon UV-C treatment (Fig. S4). It is likely that a diversion of the metabolic

flux prevents a harmful accumulation of both Pip and IC in plastids[18]. Pip may either inhibit its own

biosynthesis via a feedback loop or feed into the lysine degradation pathway towards the Krebs cycle[19].

IC on the other hand might be channeled towards the synthesis of aromatic amino acids.

In order to exclude that the observed absence of NHP and NHP-*O*Glc in *eds5* mutants is due to their

inability to accumulate SA, we tested whether external SA supply could induce Pip and NHP-*O*Glc

production in *eds5* and *pbs3* plants. As reported, SA treatment triggers the biosynthesis of Pip and

NHP-*O*Glc in wild type plants (Fig. 2a)[7]. SA biosynthesis is impaired in the *pbs3* mutant, but drenching

the soil with SA still initiated Pip biosynthesis in these plants (Fig. 2a). However, as with UV-C

treatment, external SA also did not lead to Pip accumulation in *eds5* mutants. This is consistent with

EDS5 representing a plastidial exporter of Pip independent of SA. When Pip irrigation was used to

trigger *in planta* Pip biosynthesis, we detected increased levels of the downstream product NHP-*O*Glc

in wild type plants (Fig. 2b). In *eds5* and the Pip biosynthesis mutant *sard4*, the levels of NHP-*O*Glc are

reduced by more than 60%. This suggests that in the mutants only the external Pip was metabolized

further, whereas *de novo* Pip biosynthesis did not occur. The ability of *eds5* plants to convert

exogenously applied Pip into NHP-*O*Glc supports FMO1 being active outside of plastids and not

affected by the loss of function of *EDS5*.

Taken together, we show that UV-C treatment is sufficient to induce the production of Pip and its

metabolites. This process does not require the presence of SA and can thus be used to study the SA-

independent branch of plant defense signaling. Moreover, we identified here a previously unknown

connection between Pip and *EDS5*, a gene that, so far, was only recognized for its involvement in SA

biosynthesis[16]. Most likely EDS5 is also responsible for the export of Pip from plastids into the cytosol,

where it is further processed by FMO1. This spatial separation of Pip biosynthesis and site of activation

adds an additional layer of regulation. Our study implies that EDS5 serves as the central hub in the

biosynthesis of two major defense signaling molecules, SA and NHP (Fig. 2c). It is surprising that no

pathogen effector has been found to target EDS5 and thereby exploit this key point in plant immunity.

Figures

Figure 1 UV-C treatment of *Arabidopsis thaliana* leads to an accumulation of the signaling compounds SA and Pip and their corresponding glycosides. a, Levels in counts per second of Pip (white bars) and its downstream products NHP (grey bars) and NHP-*O*Glc (black bars) in *Arabidopsis* wild type leaves (Col-0) at different time points after UV-C light treatment. **b-e,** Levels of SA and its

glycoside SAG (nmol g^{-1} leaf fresh weight [f.w.]) and levels of Pip and its glycoside NHP-*O*Glc (counts

per second) in leaves of wild type (Col-0), respectively *eds5*-3 and *pbs3*-1 mutant plants 24 hours after

UV-C treatment in comparison to untreated plants. Bars represent the mean ± STD of three biological

replicates. Statistical differences among replicates are labeled with different letters ($P < 0.05$, one-way

ANOVA and post hoc Tukey's Test; n = 3). The experiments were repeated twice with similar results.

Figure 2 Root application of either SA or Pip induces Pip biosynthesis and processing in leaves. a,

Levels of Pip (counts per second [cps]) in leaves of wild type (Col-0) *eds5*-3 and *pbs3*-1 plants 24 hours

after soil drenching with water (-) or SA (+). **b,** Levels of NHP-*O*Glc (counts per second [cps]) in leaves

of wild type (Col-0), *eds5*-3 and *sard4-5* plants 24 hours after soil drenching with water (-) or Pip (+).

Bars represent the mean ± STD of three biological replicates. Statistical differences among replicates

are labeled with different letters (P < 0.05, one-way ANOVA and post hoc Tukey's Test; n = 3). The

experiment was repeated twice with similar results. **c,** A working model depicting EDS5 as the central

hub in the biosynthesis of the signaling compounds SA and NHP. The induction of these pathways was

facilitated here by UV-C light treatment.

Methods

Plant material. *Arabidopsis* plants were grown in a chamber at 22 °C with a 16 h light period and 60% relative humidity for 4-5 weeks. For our experiments, we use *Arabidopsis* ecotype Col-0 and the following mutants in this background: *eds5-3*[20], *pbs3-1*[6], *sard4-5*[10], and fmo1-1[11], which were described previously.

UV-C and soil drench treatment. For the UV-C treatment, we followed previous protocol[16]. In short, 4-5 week old *Arabidopsis* plants were exposed to UV-C light (254 nm) for 20 min at 50 cm distance to the lamp (TUV T8 30W, Philips) for the induction of SA and Pip biosynthesis. For the treatment with SA and Pip, we followed previously described protocols[7,20]. 4-5 week old plants, were soil drenched with either 10 mL water, 10 mL of a 5 mM Pip solution (P45850, Sigma) or 10 mL of a 5 mM SA solution (S5922, Sigma), equals to 50 µM final concentration. Samples were collected 24 hours after treatment and metabolites were extracted and analyzed as described earlier[10,21]. The deviation of exact mass to accurate mass for Pip, NHP and NHP-*O*Glc was less than ±2 mDa in the untargeted metabolite analysis. SA and SAG were quantified based on internal D4-SA standard (C/D/N Isotopes Inc., Pointe-Claire, Canada). The NHP standard was chemical synthesized as described in Hartmann et al., 2018[11]. The MS/MS spectra of NHP and NHP-*O*Glc corresponds to the results from Chen et al., 2018[12].

Quantitative real-time PCR analysis. To analyze the expression of *SARD4* (At5g52810) and *EDS5* (AT4G39030) after UV-C treatment, total RNA was isolated from frozen leaves with the Spectrum™ Plant Total RNA Kit (Sigma) following the manufacturer's instruction. One microgram of RNA was treated with DNase I (Thermo Fisher Scientific) and cDNA was synthesized using Revert Aid H Minus Reverse Transcriptase (Thermo Fisher Scientific). Quantitative RT-PCR was performed using Takyon No ROX SYBR Mastermix blue dTTP (Kaneka Eurogentec) in reaction volume 20 µl. The gene ACTIN8 (At1g49240) was used as a control. Each reaction was performed with material from plants harvested

160 from three independent samples in iQ5 real time detection system (Bio-Rad). Primers are depicted in

161 Supplementary Table 1.

Acknowledgments

We are grateful to Ulf Diedrichsen and Brigitte Worbs for the synthesis of the NHP authentical

standard. This work was supported by the DFG (IRTG 2172 "PRoTECT" program of the Göttingen

Graduate Center of Neurosciences, Biophysics, and Molecular Biosciences.) to D.R., L.M. and I.F.; I.F.

was additionally supported by DFG excellence initiative (ZUK 45/2010 and INST 186/822-1). Y.Z. was

supported by Natural Sciences and Engineering Research Council of Canada, Canada Foundation for

Innovation, and British Columbia Knowledge Development Fund.

Author contributions

D.R., K.F. and I.F. conceived and designed the experiments. D.R., L.M., and K.Z. performed the

experiments. D.R., K.F., Y.Z., and I.F. wrote the article. **Competing interests:** Authors declare no

competing interests. **Data and materials availability:** All data is available in the main text or the

supplementary materials. The authors responsible for distribution of materials integral to the findings

presented in this article are: Ivo Feussner (ifeussn@uni-goettingen.de) and Yuelin Zhang

(yuelin.zhang@ubc.ca).

1   Fu, Z. Q. & Dong, X. Systemic acquired resistance: Turning local infection into global defense.
Annu. Rev. Plant Biol. **64**, 839-863, doi:10.1146/annurev-arplant-042811-105606 (2013).
2   Bernsdorff, F. *et al.* Pipecolic acid orchestrates plant systemic acquired resistance and defense
priming via salicylic acid-dependent and -independent pathways. *Plant Cell* **28**, 102-129,
doi:10.1105/tpc.15.00496 (2016).
3   Garcion, C. *et al.* Characterization and biological function of the ISOCHORISMATE SYNTHASE2
gene of Arabidopsis. *Plant Physiol.* **147**, 1279-1287, doi:10.1104/pp.108.119420 (2008).
4   Rekhter, D. *et al.* PBS3 is the missing link in plant-specific isochorismate-derived salicylic acid
biosynthesis. *bioRxiv*, 600692, doi:10.1101/600692 (2019).
5   Nawrath, C. & Métraux, J. P. Salicylic acid induction-deficient mutants of Arabidopsis express
PR-2 and *PR-5* and accumulate high levels of camalexin after pathogen inoculation. *Plant Cell*
11, 1393-1404, doi:10.1105/tpc.11.8.1393 (1999).
6   Nobuta, K. *et al.* The GH3 acyl adenylase family member PBS3 regulates salicylic acid-
dependent   defense   responses   in   Arabidopsis.   *Plant   Physiol.*   **144**,   1144-1156,
doi:10.1104/pp.107.097691 (2007).
7   Navarova, H., Bernsdorff, F., Döring, A.-C. & Zeier, J. Pipecolic acid, an endogenous mediator
of defense amplification and priming, is a critical regulator of inducible plant immunity. *Plant
Cell* **24**, 5123-5141, doi:10.1105/tpc.112.103564 (2012).
8   Song, J. T., Lu, H. & Greenberg, J. T. Divergent roles in *Arabidopsis thaliana* development and
defense of two homologous genes, *ABERRANT GROWTH AND DEATH2* and *AGD2-LIKE
DEFENSE RESPONSE PROTEIN1*, encoding novel aminotransferases. *Plant Cell* **16**, 353-366,
doi:10.1105/tpc.019372 (2004).
9   Sharma, S., Shinde, S. & Verslues, P. E. Functional characterization of an ornithine
cyclodeaminase-like protein of *Arabidopsis thaliana*. *BMC Plant Biol* **13**, 182,
doi:10.1186/1471-2229-13-182 (2013).
10   Ding, P. *et al.* Characterization of a pipecolic acid biosynthesis pathway required for systemic
acquired resistance. *Plant Cell* **28**, 2603-2615, doi:10.1105/tpc.16.00486 (2016).
11   Hartmann, M. *et al.* Flavin monooxygenase-generated *N*-hydroxypipecolic acid is a critical
element of plant systemic immunity. *Cell* **173**, 456-469, doi:10.1016/j.cell.2018.02.049 (2018).
12   Chen, Y.-C. *et al.* *N*-hydroxy-pipecolic acid is a mobile metabolite that induces systemic disease
resistance   in   *Arabidopsis*.   *Proc.   Natl.   Acad.   Sci.   USA*   **115**,   E4920-E4929,
doi:10.1073/pnas.1805291115 (2018).
13   Zhang, M. & Robertus, J. D. Molecular cloning and characterization of a full-length flavin-
dependent monooxygenase from yeast. *Arch. Biochem. Biophys.* **403**, 277-283,
doi:10.1016/S0003-9861(02)00237-0 (2002).
14   Song, J. T. Induction of a salicylic acid glucosyltransferase, AtSGT1, is an early disease response
in *Arabidopsis thaliana*. *Molecules and Cells* **22**, 233-238 (2006).
15   Li, Y., Baldauf, S., Lim, E.-K. & Bowles, D. J. Phylogenetic analysis of the UDP-glycosyltransferase
multigene   family   of   *Arabidopsis   thaliana*.   *J.   Biol.   Chem.*   **276**,   4338-4343,
doi:10.1074/jbc.M007447200 (2001).
16   Serrano, M. *et al.* Export of salicylic acid from the chloroplast requires the multidrug and toxin
extrusion-like transporter EDS5. *Plant Physiol.* **162**, 1815-1821, doi:10.1104/pp.113.218156
(2013).
17   Kilian, J. *et al.* The AtGenExpress global stress expression data set: protocols, evaluation and
model data analysis of UV-B light, drought and cold stress responses. *Plant J.* **50**, 347-363,
doi:10.1111/j.1365-313X.2007.03052.x (2007).
18   Widhalm, J. R. *et al.* Identification of a plastidial phenylalanine exporter that influences flux
distribution through the phenylalanine biosynthetic network. *Nat. Commun.* **6**, 8142,
doi:10.1038/ncomms9142 (2015).
19   Kirma, M., Araújo, W. L., Fernie, A. R. & Galili, G. The multifaceted role of aspartate-family
amino acids in plant metabolism. *J. Exp. Bot.* **63**, 4995-5001, doi:10.1093/jxb/ers119 (2012).

20   Nawrath, C. EDS5, an essential component of salicylic acid-dependent signaling for disease
231   resistance in Arabidopsis, is a member of the MATE transporter family. *Plant Cell* **14**, 275-286,
232   doi:10.1105/tpc.010376 (2002).
233   21   Kusch, S. *et al.* Arabidopsis *mlo3* mutant plants exhibit spontaneous callose deposition and
234   signs of early leaf senescence. *bioRxiv*, 558122, doi:10.1101/558122 (2019).