Title: Postpartum breast cancer progression is driven by semaphorin 7a mediated invasion and survival

Running title: PPBC is driven by SEMA7A-mediated invasion and survival

Sarah E Tarullo[1,2], Ryan C Hill[3], Kirk Hansen[3], Fariba Behbod[4], Virginia F Borges[1,2,5], Andrew C Nelson[6], and Traci R Lyons[1,2,5]*

Affiliations:

[1]Department of Medicine, Division of Medical Oncology, CU Anschutz Medical Campus, Aurora, CO 80045

[2]Young Women's BC Translational Program, CU Anschutz Medical Campus, Aurora, CO 80045

[3]Department of Biochemistry and Molecular Genetics, CU Anschutz Medical Campus, Aurora, CO 80045

[4]Division of Cancer and Developmental Biology, University of Kansas Medical Center, Kansas City, KS 66160

[5]University of Colorado Cancer Center, Aurora, CO 80045

[6]Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN 55455

***Corresponding Author:** Traci R Lyons (303-724-3885; traci.lyons@cuanschutz.edu)

Conflict of Interest: The authors have declared that no conflict of interest exists.

ABSTRACT:

Young women diagnosed with breast cancer (BC) have poor prognosis due to

increased rates of metastasis. Additionally, women within 10 years of most recent

childbirth at diagnosis are ~3 times more likely to develop metastasis than age and

stage matched nulliparous women. We define these cases as postpartum BC (PPBC)

and propose that the unique biology of the postpartum mammary gland drives tumor

progression. Our published results revealed roles for SEMA7A in breast tumor cell

growth, motility, invasion, and tumor associated-lymphangiogenesis, all of which are

also increased in pre-clinical models of PPBC. However, whether SEMA7A drives

progression in PPBC remains largely unexplored. Our results presented herein show

that silencing of SEMA7A decreases tumor growth in a model of PPBC while

overexpression is sufficient to increase growth in nulliparous hosts. Further, we show

that SEMA7A promotes multiple known drivers of PPBC progression including tumor

associated COX-2 expression and fibroblast-mediated collagen deposition in the tumor

microenvironment. Additionally, we show for the first time that SEMA7A expressing cells

deposit fibronectin to promote tumor cell survival. Finally, we show that co-expression of

SEMA7A/COX-2/FN predicts for poor prognosis in breast cancer patient cohorts. These

studies suggest SEMA7A as a key mediator of BC progression and that targeting

SEMA7A may open avenues for novel therapeutic strategies.

INTRODUCTION:

Postpartum breast cancer (PPBC), or breast cancers (BC) diagnosed within 5-10 years of last childbirth, are ~three times more likely to become metastatic [1-3]. Specifically, PPBC patients exhibit distant metastasis free five-year survival (DMFS) rates as low as 70% [1], which are further decreased to 50% after ten years [2]. Additionally, PPBC may account over half of BCs diagnosed in women aged <45 [1]. In a pre-clinical model of PPBC a non-metastatic BC cell line becomes invasive and metastatic upon orthotopic implantation at the onset of postpartum mammary gland involution [4]. Postpartum/post-lactational mammary involution returns the gland to the pre-pregnant state; we and others have shown that programs associated with postpartum involution are similar to tumor-promotional microenvironments [1, 4-9]. Since the MCF10DCIS model initially resembles ductal carcinoma in situ (DCIS), which progresses to ER/PR/HER2 negative invasive ductal carcinoma (IDC) [10], we utilized this model to show accelerated tumor growth and progression to IDC in postpartum hosts [4]. This progression was driven by collagen deposition and expression of cyclooxygenase-2 (COX-2), both of which were required for tumor cell invasion. Additionally, a weakly tumorigenic breast epithelial cell line, HMLE-Ras[lo], was similarly promoted via host driven mechanisms [11]. More recently, expression of a neuronal guidance molecule, Semaphorin 7a (SEMA7A), was observed in mouse mammary epithelium during postpartum involution and in the tumors that outgrew after implantation during involution [6].

Semaphorins are characterized for their roles during development, however and have reported roles in multiple cancer types [12-16] and SEMA7A expression is

emerging as poor prognostic indicator [6, 16-20]. SEMA7A can promote cell-

autonomous signaling when it remains bound to the cell via its

glycosylphosphatidylinositol (GPI) membrane link or non-cell-autonomous signaling

when shed via cleavage into the extracellular environment. Here, we show that

SEMA7A protein is expressed in DCIS from BC patients, is necessary for postpartum

tumor progression in our pre-clinical model, and sufficient to drive tumor progression in

nulliparous hosts. We also demonstrate that shed SEMA7A drives collagen deposition

in the tumor microenvironment (TME) via upregulation of collagen I mRNA in fibroblasts,

which promotes expression of COX-2 and invasion. Furthermore, we propose a cell-

autonomous pro-invasive and survival role for SEMA7A that is mediated through

fibronectin (FN), epithelial-to-mesenchymal transition (EMT) and downstream pro-

survival signaling via phosphorylation of AKT. Additionally, we show that SEMA7A

expressing cells exhibit enhanced metastatic capabilities. Finally, our results we

demonstrate that a gene signature of SEMA7A, COX-2, and FN1 predicts for poor

prognosis for BC patients suggest that SEMA7A merits further studies to develop a

novel therapeutic for BC patients.

RESULTS

SEMA7A promotes postpartum tumor progression in a mouse model and is

expressed in DCIS from patients

The MCF10DCIS model is ideal for monitoring early events in the metastatic

cascade, such as the transition from in situ to invasive. To examine whether SEMA7A

drives DCIS progression in postpartum/post-lactational hosts, we orthotopically injected

MCF10DCIS cells stably expressing a SEMA7A targeted shRNA (SEMA7A-KD) or

nontargeting control (Crtl) [18] at involution day 1 (SFigure1A). We observed that

silencing of SEMA7A is sufficient to decrease tumor growth in postpartum hosts despite

some of the tumors regaining expression of SEMA7A protein (Figure1A; SFigure1B).

Harvested tumors (H&E stained sections) were scored for invasion at 5 weeks post-

injection, when the majority of postpartum tumors are normally invasive (SFigure1C),

and 33% of the SEMA7A-KD were invasive compared to 86% in the controls. Further,

56% of tumors in the SEMA7A-KD group maintained evidence of DCIS versus 12.5%

controls (Figure1B). Finally, 11% of tumors in the KD group were DCIS with

microinvasion compared to 0% of controls. Collagen-mediated upregulation of COX-2 is

a dominant feature that drives invasion in postpartum hosts in the MCF10DCIS model

[4]. Consistent with a role for SEMA7A in collagen/COX-2 dependent invasion in

postpartum hosts, we observe that fibrillar collagen deposition, by Massan's trichrome

stain, and COX-2 expression, by immunohistochemistry (IHC) are significantly

decreased in SEMA7A-KD tumors (Figure1C&D).

To determine whether SEMA7A expression was higher in DCIS from postpartum

patients, we performed IHC on DCIS lesions and adjacent normal breast tissue from

women in our BC cohort (Figure2A) and quantitated medium + strong staining to show

that DCIS lesions express higher levels of SEMA7A than normal breast tissue

(Figure2B,SFigure2A). Then, when separated by parity status, we observe increased

SEMA7A in DCIS lesions in both nulliparous and postpartum patients, with a trend

toward highest expression in postpartum patients, suggesting that SEMA7A expression

may have relevance to all patients with DCIS, but may be particularly relevant to

postpartum patients (Figure2B,SFigure2B). Consistent with this, paired analysis

between normal and DCIS within each patient revealed that the majority of patients in

both the nulliparous and postpartum groups exhibited increased SEMA7A expression in

DCIS compared to normal (Figure 2C&D). We confirmed this observation by showing

similar data on SEMA7A expression in a tissue microarray consisting of additional

normal breast and matched patient samples of DCIS using kidney and placenta as

negative and positive controls, respectively (Figure2E-G,SFigure2C). Finally, we also

show that SEMA7A mRNA levels are increased in DCIS compared to normal in the

METABRIC [21] dataset and that SEMA7A is in the top 11% of upregulated genes in

DCIS compared normal (p=0.002)(SFigure2D). Taken together, our results suggest that

SEMA7A may represent a general mediator of DCIS growth and invasion in both parous

and nulliparous women.

To test our hypothesis that SEMA7A expression is sufficient to drive DCIS

progression, we injected MCF10DCIS cells that overexpress SEMA7A protein

(SEMA7A-OE), along with controls, into a separate cohort of nulliparous hosts; we

observed accelerated growth of SEMA7A-OE tumors (Figure3A). These tumors were

similarly scored for invasion, but at three weeks post-injection, when tumors are

normally DCIS in nulliparous hosts (SFigure1C). While, control tumors were all DCIS, as

expected, greater than 50% of the SEMA7A-OE tumors had microinvasion and/or were

mixed IDC+DCIS (Figure3B;SFigure 3). Additionally, levels of collagen and COX-2,

which are normally very low in DCIS tumors in nulliparous hosts, are higher in the

SEMA7A-OE tumors, which is consistent with a role for SEMA7A in promoting

production (Figure3C&D).

SEMA7A promotes tumor cell invasion via matrix deposition and acquisition of

mesenchymal phenotypes

To model SEMA7A dependent invasion in vitro, we utilized a 3D organoid model

with SEMA7A-OE cells suspended in Matrigel or Matrigel+collagen [4]. We found that

SEMA7A overexpression was not sufficient to drive increased invasion on Matrigel

alone but sufficient when collagen I was present in the matrix (Figure4A&B). We then

validated the requirement for both collagen and SEMA7A in invasion via knockdown in

the MDA-MB-231 cell line, which requires collagen for 3D organoid formation [22]

(SFigure4A, Figure4C). Since fibroblasts in the TME produce collagen, we then stained

for alpha-smooth muscle actin (αSMA), a marker of activated fibroblasts, but did not

observe differences in fibroblast infiltration (SFigure4). These results suggest that

SEMA7A does not recruit tumor infiltrating fibroblasts.

Alternatively, to determine whether SEMA7A expressing tumor cells promote

collagen production by fibroblasts via shedding of SEMA7A, we induced serum starved

fibroblasts with conditioned media from our SEMA7A-OE tumor cells, alongside

conditioned media from control cells and TGFβ as positive control and examined

expression of the *COL1A1* mRNA by qPCR. We focused on *COL1A1* because collagen

I is the most abundant collagen in the mammary gland [23]. Conditioned medias from

SEMA7A OE cells induced *COL1A1* expression in fibroblasts to higher levels than

observed in control or TGFβ treated fibroblasts (Figure4D). Then, we treated fibroblasts

with purified SEMA7A to show specific induction of COL1A1 gene expression

(Figure4E). To understand additional SEMA7A mediated changes to the TME, we

performed an unbiased mass spectrometry analysis of conditioned medias from ex vivo

tumors derived from SEMA7A-KD or control cells. When we restricted our analysis to

ECM peptides of human origin we observed downregulation of several ECM associated

molecules with SEMA7A-KD (SFigure4A), but focused on the significant decrease in

fibronectin (FN) (Figure5A) after immunostaining of our tumors confirmed decreased FN

expression with SEMA7A-KD (Figure5B&C). As FN is frequently associated with EMT,

we examined whether SEMA7A expression is also associated with additional

mesenchymal markers. We observed SEMA7A dependent expression of matrix

remodeling enzyme, matrix metalloproteinase-2, vimentin, and S100A4, as well as other

members of the S100 family of proteins (Figure5D-F;SFigure4B-E), which are all

secreted by mesenchymal-like cells [24]. To confirm SEMA7A-dependent

mesenchymal-like phenotypes we measured cell aspect ratios with the prediction that

epithelial-like cells, due to their cuboidal morphology, would exhibit a ratio of ~1 and

ratios >1 would be indicative of mesenchymal-like cells. Our results reveal that the

mesenchymal-like MDA-MB-231 cells, exhibit decreased average aspect ratios with

SEMA7A-KD (Figure5G). Concordantly, SEMA7A-OE in the MCF10DCIS cells, which

are less mesenchymal-like, resulted in increased aspect ratios (Figure5H). Furthermore,

immunoblot analysis of MCF10DCIS SEMA7A-OE cells reveals decreased E-cadherin

and increased vimentin expression, markers of epithelial and mesenchymal cells,

respectively (SFigure5F). Together, our results suggest that SEMA7A promotes cellular

invasion via both cell-autonomous and non-cell-autonomous mechanisms of matrix

deposition and remodeling.

SEMA7A promotes cell survival via fibronectin, AKT and COX-2

We also observed decreased tumor growth of SEMA7A-KD tumors in vivo but did

not see significant changes in proliferation marker Ki67 (SFigure5A). However, we did

observe a trend toward increased cleaved-caspase 3, a marker of apoptosis, in

SEMA7A-KD tumors with a corresponding decrease in SEMA7A-OE tumors

(SFigure5B&C). We also observed increased cell death in SEMA7A-KD cells in culture

via a luminescent assay for caspase activity, which we validated in the MDA-MB-231

cell line (Figure6A&B). Conversely, decreased cell death was observed in the SEMA7A

OE cell in both cell lines (SFigure5D, SFigure5E&F). Additionally, analysis of cell death

in real time confirmed that SEMA7A-KD MDA-MB-231 cells exhibit increases in cell

death via activation of apoptotic signaling (Figure6C). One known pro-survival

mechanism co-opted by tumors to block activation of caspase cleavage is activation of

pro-survival kinase AKT [25] and we observed decreased levels of phosphorylated AKT

via immunoblot for pS473 in SEMA7A-KD and a corresponding increase in levels in

SEMA7A OE cells (Figure6D&SFigure5G). Interestingly, FN signals via integrins, which

can activate downstream pro-survival pathways, including AKT [26]. To determine

whether add-back of FN could rescue SEMA7A-KD cells from cell death, we KD plated

cells FN-, laminin- or collagen-coated plates with control cells on tissue culture plastic

for reference. We observed increased cell death in all SEMA7A-KD conditions except

when FN was present where we observed levels of cell death were more similar to the

Ctrl cells (Figure6E). Thus, we suggest that SEMA7A promotes AKT mediated survival

by increasing FN deposition and therefore cellular attachment.

Mesenchymal-like tumor cells, known to have high levels of FN, are not as

dependent on matrix attachment for survival and can also exhibit resistance to normal

programs that mediate cell death including survival in detached or anchorage

independent conditions [27]. An initial step of the metastatic cascade involves cell

detachment from ECM to facilitate local invasion and access to the vasculature; then,

cells in circulation must survive in matrix detached conditions prior to extravasating to

seed metastatic sites. To assess a role for SEMA7A in promoting survival in circulation,

we forced cellular detachment in vitro and measured cleaved-caspase 3/7 to show that

SEMA7A-KD increases cell death while SEMA7A-OE promotes cell survival in detached

conditions (Figure6F&G, SFigure5H&I). Interestingly, fibronectin was not sufficient to

rescue cell death in suspension suggesting that fibronectin deposition is necessary

(data not shown). However, since activation of AKT is a known mediator of COX-2

expression [28] we also examined the ability of COX-2 KD cells to survive in detached

conditions. Similar to our results in the SEMA7A-KD cells we observe that COX-2

knockdown cells exhibit increased cell death in detached conditions, which was not

observed in attached conditions (SFigure5J) suggesting that SEMA7A mediated

upregulation of AKT signaling, and subsequent expression of COX-2, support cell

survival in detached conditions such as those encountered during local invasion and in

circulation (Figure 6H).

SEMA7A, COX-2, FN and metastatic potential

To assess whether SEMA7A promotes survival in circulation in vivo, we

performed tail vein injections of MDA-MB-231 SEMA7A-KD or Ctrl cells (SFigure6A)

and assessed for pulmonary metastasis. Our results reveal a decrease in both the

number and average size of metastatic lesions per lung with SEMA7A KD (Figure7A)

suggesting that SEMA7A plays a role in survival in circulation, as well as a possible role

in seeding and outgrowth of BC metastasis. We have previously published that

SEMA7A expression is upregulated in IDC compared to normal and the worst overall

survival (OS) was observed in SEMA7A+ER-BC in the METABRIC dataset [18];

however, in SEMA7A+ER- BCs we do not observe decreased distant metastasis free

survival (DMFS) via KmPlot analysis (SFigure6B). Similarly, although transcripts for

COX-2 and FN1 are also upregulated in BC (STable1), neither COX-2 or FN1 are

associated with decreased DMFS in ER-BC (SFigure6E&F). However, co-expression of

COX-2, SEMA7A, and FN1 increases risk for metastasis in ER-BC patients, with 5 year

DMFS rates approaching 70% by both KmPlot and GOBO (Gene expression-based

Outcome for Breast cancer Online or GOBO) analysis (Figure7B&C) [1]. We also

observed that co-expression of our 3 gene signature significantly associates with

decreased DMFS in basal and PAM50 basal subtype tumors using GOBO (SFigure6G-

I). Finally, co-expression of SEMA7A with both FN and COX-2 is observed in BCs in the

TCGA dataset (Figure6D&E).

DISCUSSION

All women experience a transient increase in risk for developing breast cancer after each completed pregnancy [4, 29, 30]. Additionally, patients diagnosed with breast cancer within ten years postpartum comprise nearly 50% of all BCs in women <40 in both US and Norwegian cohorts and these PPBC patients are at high risk for metastatic spread [1, 31]. A recent publication by Welch and Hurst defines the hallmarks of metastasis as motility and invasion, modulation of the microenvironment, plasticity, and colonization [32]. Here, we identify roles for SEMA7A in promotion of metastasis using preclinical models regardless of parity status. Our results suggest that SEMA7A drives one described mechanism of tumor progression pertinent to postpartum women, which is collagen mediated upregulation of COX-2 that drives motility and invasion of DCIS cells. However, our results also suggest that this mechanism of invasion can be driven by SEMA7A in both nulliparous and postpartum hosts suggesting it is not unique to parous patients. Further, we extend our observations and identify a novel cell-autonomous mechanism by which SEMA7A mediates tumor cell survival via stimulation of FN production and activation of pro-survival kinase AKT regardless of the host parity status. We also show that SEMA7A supports tumor cell invasion by stably altering cells to a more mesenchymal phenotype, which promotes cell survival in anchorage-independent conditions to allow for colonization of distant organs. Thus, we suggest that SEMA7A plays a key role in multiple steps toward progression to metastatic disease. We support this claim by showing a that co-expression of COX-2, SEMA7A, and FN correlates with distant metastasis formation in BC patients. We propose a model whereby SEMA7A signaling supports metastatic progression (Figure8).

SEMA7A was first identified on lymphocytes and originally designated CDw108

[33]. CDw108 was renamed SEMA7A due to its structural similarities with members of

the Semaphorin family of proteins best known for their roles in neuronal guidance [34].

SEMA7A is the only semaphorin linked to the membrane via a GPI anchor; as such, it

can be cleaved to result in shedding of SEMA7A into the extracellular environment,

which was shown to promote β1-integrin dependent inflammation and fibrosis [35-41].

During fibrosis and in response to TGFβ, SEMA7A activates AKT signaling, which

results in upregulation of collagen and FN in a β1-integrin dependent manner [42].

While our current studies do not explore a role for β1-integrin, we and others have

shown that SEMA7A-β1-integrin binding promotes tumor growth, EMT,

migration/invasion, metastasis, and neo-vasculogenesis [17, 19, 20]. Herein, we

hypothesize that SEMA7A mediated tumor cell production of FN activates integrin

mediated PI3K signaling leading to activation of AKT and cell survival [26, 43]. We also

show that SEMA7A promotes mesenchymal phenotypes and tumor cell invasion, which

could may be mediated by FN engagement of $\alpha_v\beta_1$-integrin and downstream activation

of Slug resulting in transcription of mesenchymal genes [44, 45]. In support of this

hypothesis, SEMA7A is linked to EMT in a murine model of BC where TGFβ fails to

induce EMT in the absence of SEMA7A [17]. Additional studies to determine how

SEMA7A and FN contribute to EMT in our model are necessary to fully understand this

mechanism in BC progression.

Since upregulation of EMT pathways such as Snail, Slug, and Zeb1 are

associated with decreased sensitivity to chemotherapeutic drugs and targeted tyrosine

kinase inhibitors (TKIs) [17, 46-55], it is also possible that SEMA7A may promote

metastasis by conferring resistance to current therapies. In support of this, SEMA7A

promotes resistance to EGFR targeted therapies in lung cancer [13]. Additional studies

will explore the role of SEMA7A in resistance and/or susceptibility to current therapies in

BC. Moreover, one major limitation of our current study is investigation of this

mechanism only in ER- models, which is supported, in part, by our patient dataset

studies showing that SEMA7A expression invokes a higher risk for metastasis in ER-

breast cancers compared to all breast cancer. However, since ER+ breast cancers

comprise the largest percentage of breast cancers in women, additional studies in ER+

breast cancers must be performed. Interestingly, estrogen and/or progesterone promote

SEMA7A expression in the hypothalamus, suggesting that hormones may drive

elevated levels of SEMA7A in BCs [56]. Furthermore, in hormone-receptor positive

(HR+) BCs both AKT and FN promote tamoxifen resistance [57-64] and Semaphorin

4C, which is structurally similar to SEMA7A, drives hormonal-independence in HR+ BCs

[65]. Thus, we are also exploring the relationship between SEMA7A expression and

tumor progression in ER+ models [66].

We also demonstrate that SEMA7A is also involved in a non-cell-autonomous

signaling axis via induction of fibroblast production of collagen, which then promotes

COX-2 expression in the tumor cell [4]. Collagen has been characterized for multiple

roles in promoting tumor progression as well as increased breast density, which can

also increase risk for developing BC [8, 67, 68]. Additionally, a single pregnancy is

sufficient to convert the high collagen content of the postpartum breast into pro-

tumorigenic collagen [69] and activated fibroblasts, which deposit collagen, are present

during postpartum involution [4, 8, 23]. Thus, our previous studies showing that

SEMA7A is expressed in the postpartum mammary epithelium during involution

SEMA7A, coupled with those presented herein, suggest that SEMA7A could also play a

role in the increased risk for developing BC after pregnancy via increasing collagen

deposition [1, 4, 31, 67, 68]. COX-2 is also well characterized for multiple roles in tumor

initiation and progression [70, 71]. While our results suggest that SEMA7A stimulates

collagen production by fibroblasts to result in tumor cell expression of COX-2 we still do

not fully understand this mechanism. Additionally, our current studies have not

addressed whether COX-2 activity is increased in tumor cells on collagen. Our data also

suggest SEMA7A activation of AKT may play a role in promoting survival via

upregulation of COX-2. Further studies are needed to understand the molecular

mechanisms underlying SEMA7A and COX-2 signaling, regulation, and their roles in

promoting tumor cell invasion and survival.

Currently there are no prevention strategies, targeted therapies, or specific

treatment options for women diagnosed postpartum or for women with high tumor

expression of SEMA7A. If our proposed SEMA7A mediated mechanism of progression

is also dependent on COX-2 activity, as is suggested by our previous studies [4, 7], the

addition of a COX-2 inhibitor to current treatment regimens could have efficacy in

postpartum or SEMA7A+ BC patients. We also propose that SEMA7A expression could

predict progression or itself be a potential therapeutic target for BC patients. If direct

targeting of SEMA7A is not feasible, SEMA7A activates downstream targets such as

FAK, Src, and ERK [16, 72], for which targeted therapies are available. Although these

targeted inhibitors have been largely unsuccessful in BC, SEMA7A could serve as a

predictive biomarker for patients who may benefit. Since metastases are the leading

cause of BC related deaths and are largely untreatable, these novel anti-metastatic

treatment strategies should be explored.

342
MATERIALS AND METHODS

Cell Culture

MCF10DCIS and MDA-MB-231 were cultured in 2D and 3D cultures as previously

described [4, 18, 22]. MCF10DCIS cells and previously described shCOX-2 derivatives

[73] were obtained from K. Polyak and A. Marusyk (Harvard University, Cambridge,

MA). MDA-MB-231 cells were obtained from P. Schedin (Oregon Heath and Sciences

University, Portland OR). HLF-1 cells were gifted from M. Fini (CU Anschutz Medical

Campus, Denver, CO). Cells were validated by the DNA sequencing core at the CU

Anschutz Medical Campus and identified to be a pure population of their respective cell

lines. Cells were regularly tested for mycoplasma throughout studies. shRNA silencing

was achieved using shRNA SEMA7A targeting plasmids (SABiosciences, Frederik, MD,

and Functional Genomics Facility at CU Anschutz Medical Campus, Denver, CO) and

confirmed via qPCR and Western blot analysis. Overexpression plasmid (SEMA7A-Fc)

was a generous gift from R. Medzhitov (Yale University, New Haven, CT). Control

plasmid (pcDNA3.1) was obtained from H. Ford (CU Anschutz Medical Campus,

Denver, CO). All other overexpression plasmids (p304-V5-Blasticidin and V5-SEMA7A)

were obtained from the Functional Genomics Core at the CU Anschutz Medical Campus

and overexpression was confirmed via qPCR and Western blot analysis. Purified

SEMA7A was isolated from MDA-MB-231 cells engineered to overexpress SEMA7A-Fc

in collaboration with the Protein Purification/MoAB/Tissue culture core at the CU

Anschutz Medical Campus. Cells were forced into suspension by coating plates with 12

mg/ml poly-HEMA (poly (2-hydroxyethyl methacrylate), Sigma, St. Louis, MO) prior to

plating.

Tissue microarray

367 Tissue microarrays containing normal and DCIS samples were prepared from biopsy

368 tissue following placement in preservation media (LiforCell, Lifeblood Medical, Inc.) and

369 storage at 4ºC, as previously described [74].

qPCR

RNA was isolated and qPCR were performed as previously described, with GAPDH and

RPS18 as reference genes [18]. Primers for SEMA7A, COL1A1, COX-2 were obtained

from Bio-Rad (Bio-Rad PrimePCR, Hercules, CA). Other primers were designed to be

intron spanning with the following sequences: GAPDH (forward:

CAAGAGCACAAGAGGAA GAGAG, reverse: CTACATGGCAACTGTGAGGAG) and

RPS18 (forward: GCGAGTACTCAACACCAACA, reverse:

GCTAGGACCTGGCTGTATTT).

Immunoblot analysis

379 Western blots were performed as previously described [18]. Antibody information is

380 provided in STable2.

Animal model

The MCF10DCIS model was utilized as previously described [4, 18]. Briefly, 6-8-week-

old female SCID Hairless Outbread mice from Charles River were bred and, after birth,

pup numbers were normalized to 6-8 pups per dam. After 10-13 days of lactation, pups

were removed to initiate involution (Day 0). Subsequently, injections of 250K

MCF10DCIS controls and cells with shSEMA7A were initiated one day post-weaning

(involution day 1) bilaterally into the #4 mammary fat pads. For SEMA7A

overexpressing studies, 6-8-week-old Nude athymic (nulliparous) from Charles River

were utilized because they are more cost-effective than SCID mice and breeding is not

necessary for studies in nulliparous hosts. Tumors were measured twice weekly. For

metastasis studies, nude mice were injected with $1x10^6$ cells into the tail vein, monitored

for weight loss and sacrificed 3 weeks post-injection.

Histologic analysis

Mammary glands with intact tumor were prepared for immunohistochemistry as

previously described [4, 7]. Hematoxylin and eosin stained sections were examined by a

board-certified anatomic pathologist (ACN) as scored as follows: 0-lesions which

contained only well-devolved DCIS structures with clearly defined basement

membranes and no evidence of microinvasion; 1-lesions that contained extensive DCIS

with identifiable micro-invasive foci; 2-lesions that contained significant areas of sheet-

like invasive tumor growth and mixed with areas of DCIS; 3-lesions that contained

entirely invasive tumor with rare to absent DCIS remnants. The pathologist was blinded

to study group by the randomization of animal numbers in each group.

Immunohistochemistry and Immunofluorescence

For FN and COX-2 400X images were taken of intact tumor and quantitated using

ImageJ software. For SEMA7A, cleaved caspase-3, and trichrome, stain quantification

of total tumor area (necrotic and stromal areas removed) and percent positive stain or

stain intensity was performed using ImageScope Aperio Analysis software (Leica,

Buffalo Grove, IL). Areas for quantification were annotated using Aperio analysis tools

and percent weak, medium, and strong determined using the color-deconvolution

algorithm. For COX-2 and FN analysis, areas for quantification were isolated from the

surrounding stroma and percent positive calculated as area with positive stain (positive

pixels) using Image J and divided by total area (total pixels) and multiplied by

100.Immunofluorescent images were obtained using 400X magnification on OLYMPUS

microscope. Antibody information is provided in STable 2.

In vitro cell death assay

Cell death was analyzed using Caspase-Glo 3/7 Assay (Promega, Madison, WI),

according to the manufacturer's instructions.

Analysis of publicly available datasets

Km plotter was queried for BC, and SEMA7A, PTGS2 (COX-2), and FN1 using the

multigene classifier mean centered option for distant metastasis free survival (DMFS)

[75]. ER- status was determined from ESR1 gene expression data. Ma breast cancer

dataset was queried for SEMA7A, PTGS2 (COX-2), and FN1 using Gene expression-

based Outcome for Breast Cancer Online (GOBO). TCGA was queried for co-

expression of SEMA7A and COX-2 or FN1 using CBioPortal.

Mass spectrometry analysis

Small tumor sections (~1mm) were placed on gelatin sponges (Novartis Animal Heath,

Greensboro, NC, USA) in serum-free media as previously described [76]. After 48

hours, tumor conditioned media was collected ~ 30 µg of total protein digested utilizing

the filter-aided sample digestion (FASP) protocol as previously described according to

MAIPE standards [77]. Briefly, samples were reduced, alkylated, and enzymatically

digested with trypsin. Resulting peptides were concentrated and de-salted by solid

phase extraction utilizing in-house made stage tips made with Sytrene Divinyl Benzene

disks (Empore™). Liquid chromatography tandem mass spectrometry (LC-MS/MS) was

performed on a Thermo nanoEasy LC II coupled to a Q Exactive HF. MS acquisition

parameters are detailed previously [78]. Raw files were searched with Proteome

Discoverer 2.2 against the Mus Musculus, Homo Sapiens, and Bos Taurus uniprotKB

database in Mascot. Precursor mass tolerance was set to +/- 10 ppm and MS/MS

fragment ion tolerance of +/- 25 ppm. Trypsin specificity was selected allowing for 1

missed cleavage. Variable modifications include Met oxidation, proline hydroxylation,

protein N-terminal acetylation, peptide N-terminal pyroglutamic acid formation, and a

fixed modification of Cys carbamidomethylation.

Search results were visualized using Metaboanalyst v4.0 [79] and gene ontology

mapping was done using PANTHER [80]. Data were prepared according to MIAPE

standards and will be made available upon publication.

Experimental replicates, sample size and statistical analyses.

All in vitro studies were performed in biological triplicates. For animal studies, we chose

the number of mice/group/time-point, based on power calculations form previous and

pilot studies to achieve at least 80% power (β) with α=0.05. All animal studies were

replicated twice with representative or pooled data shown. Unpaired and paired t-tests,

ANOVA, and Kaplan Meier statistical analyses were performed in GraphPad Prism,

assuming independent samples and normal distributions. Analyses for Figure 2 were

done using one-tailed t-tests, as our results from Figure 1 would predict significant

differences between groups. All other analyses were done using two-tailed tests. Only

p-values of less than 0.05 were considered significant. Aside from unexpended death or

cell contamination, no samples or animals were excluded from our analysis. For IHC

analysis of tumors, outliers were removed if they were significant by the ROUT (Q=1%)

test.

Study Approval

Prior to resection, patients provided written informed consent under an IRB-approved

protocol according to the guidelines of their respective institutions and conducted in

compliance with HIPPA regulations. All animal studies were approved by the IACUC of

the CU Anschutz Medical Campus, protocol number B106017(06)1E.

Author Contributions

SET and TRL conceived and designed the study. SET performed all in vitro and in vivo studies. VFB and FB were responsible for regulatory oversight of human tissue acquisition and providing cases for IHC analysis. RCH and KH were responsible for all mass spectrometry experiments and associated data analysis. ACN was responsible for analyzing and scoring all tumor for invasion. SET and TRL were responsible for hypothesis development, conceptual design, data analysis and data interpretation. SET and TRL wrote the manuscript with all authors providing critical evaluation.

Acknowledgements

MCF10DCIS cells were obtained from K. Polyak and A. Marusyk (Harvard University, Cambridge, MA). MDA-MB-231 cells were obtained from P. Schedin (Oregon Heath and Sciences University, Portland OR). HLF-1 cells were gifted from M. Fini (CU Anschutz Medical Campus, Denver, CO). H. Ford for the pcDNA3.1 vector (CU Anschutz Medical Campus, Denver, CO).; and R. Medzhitov (Yale University, New Haven, CT) for the SEMA7A-Fc overexpression vector. We also acknowledge V. Wessells, A. Elder, L. Crump, T. Wallace, C. Young, A. Stoller, M. Kobritz, and C. Hoang for technical support and advice. This work was supported by the American Cancer Society (RSG 16-171-01-CSM), and NIH/NCI (R01CA211696-01A1) to T. Lyons; NIH/CCTSI/CTSA (TL1 TR001081) to S. Tarullo.

Supplementary Information is available on *Oncogene's* website.

References

485   References
486

487   1   Callihan EB, Gao D, Jindal S, Lyons TR, Manthey E, Edgerton S *et al*.
488   Postpartum diagnosis demonstrates a high risk for metastasis and merits an
489   expanded definition of pregnancy-associated breast cancer. *Breast cancer*
490   *research and treatment* 2013; 138: 549-559.
491

492   2   Erica T. Goddard SB, Troy Schedin, Sonali Jindal, Jeremy Johnston, Ethan
493   Cabral, Traci R. Lyons, Motomi Mori, Pepper J. Schedin, Virginia F. Borges.
494   Defining the impact of a postpartum diagnosis on metastasis and the clinical
495   features underlying risk: A young women's breast cancer cohort study. *Jama*
496   *Network* 2018; In press.
497

498   3   Nichols HB, Schoemaker MJ, Cai J, Xu J, Wright LB, Brook MN *et al*. Breast
499   Cancer Risk After Recent Childbirth: A Pooled Analysis of 15 Prospective
500   Studies. *Ann Intern Med* 2018.
501

502   4   Lyons TR, O'Brien J, Borges VF, Conklin MW, Keely PJ, Eliceiri KW *et al*.
503   Postpartum mammary gland involution drives progression of ductal carcinoma in
504   situ through collagen and COX-2. *Nature medicine* 2011; 17: 1109-1115.
505

506   5   Slepicka PF, Cyrill SL, Dos Santos CO. Pregnancy and Breast Cancer: Pathways
507   to Understand Risk and Prevention. *Trends Mol Med* 2019.
508

509   6   Elder AM, Tamburini BAJ, Crump LS, Black SA, Wessells VM, Schedin PJ *et al*.
510   Semaphorin 7A Promotes Macrophage-Mediated Lymphatic Remodeling during
511   Postpartum Mammary Gland Involution and in Breast Cancer. *Cancer research*
512   2018; 78: 6473-6485.
513

514   7   Lyons TR, Borges VF, Betts CB, Guo Q, Kapoor P, Martinson HA *et al*.
515   Cyclooxygenase-2-dependent lymphangiogenesis promotes nodal metastasis of
516   postpartum breast cancer. *The Journal of clinical investigation* 2014; 124: 3901-
517   3912.
518

519   8   Guo Q, Minnier J, Burchard J, Chiotti K, Spellman P, Schedin P. Physiologically
520   activated mammary fibroblasts promote postpartum mammary cancer. *JCI*
521   *Insight* 2017; 2: e89206.
522

523   9   Asztalos S, Gann PH, Hayes MK, Nonn L, Beam CA, Dai Y *et al*. Gene
524   expression patterns in the human breast after pregnancy. *Cancer Prev Res*
525   *(Phila)* 2010; 3: 301-311.
526

527   10   Miller FR, Santner SJ, Tait L, Dawson PJ. MCF10DCIS.com xenograft model of
528   human comedo ductal carcinoma in situ. *Journal of the National Cancer Institute*
529   2000; 92: 1185-1186.
530

531   11   Gupta PB, Proia D, Cingoz O, Weremowicz J, Naber SP, Weinberg RA *et al*.
532   Systemic stromal effects of estrogen promote the growth of estrogen receptor-
533   negative cancers. *Cancer research* 2007; 67: 2062-2071.

534
535   12   Rehman M, Tamagnone L. Semaphorins in cancer: biological mechanisms and
536   therapeutic approaches. *Semin Cell Dev Biol* 2013; 24: 179-189.

537
538   13   Kinehara Y, Nagatomo I, Koyama S, Ito D, Nojima S, Kurebayashi R *et al*.
539   Semaphorin 7A promotes EGFR-TKI resistance in EGFR mutant lung
540   adenocarcinoma cells. *JCI Insight* 2018; 3.

541
542   14   Ma B, Herzog EL, Lee CG, Peng X, Lee CM, Chen X *et al*. Role of chitinase 3-
543   like-1 and semaphorin 7a in pulmonary melanoma metastasis. *Cancer research*
544   2015; 75: 487-496.

545
546   15   Saito T, Kasamatsu A, Ogawara K, Miyamoto I, Saito K, Iyoda M *et al*.
547   Semaphorin7A promotion of tumoral growth and metastasis in human oral cancer
548   by regulation of g1 cell cycle and matrix metalloproteases: Possible contribution
549   to tumoral angiogenesis. *PloS one* 2015; 10: 1-20.

550
551   16   Scott Ga, McClelland La, Fricke AF, Fender A. Plexin C1, A Receptor for
552   Semaphorin 7A, Inactivates Cofilin and Is a Potential Tumor Suppressor for
553   Melanoma Progression. *Journal of Investigative Dermatology* 2009; 129: 954-
554   963.

555
556   17   Allegra M, Zaragkoulias A, Vorgia E, Ioannou M, Litos G, Beug H *et al*.
557   Semaphorin-7a reverses the ERF-induced inhibition of EMT in Ras-dependent
558   mouse mammary epithelial cells. *Molecular biology of the cell* 2012; 23: 3873-
559   3881.

560
561   18   Black SA, Nelson AC, Gurule NJ, Futscher BW, Lyons TR. Semaphorin 7a exerts
562   pleiotropic effects to promote breast tumor progression. *Oncogene* 2016; 35:
563   5170-5178.

564
565   19   Garcia-Areas R, Libreros S, Amat S, Keating P, Carrio R, Robinson P *et al*.
566   Semaphorin7A promotes tumor growth and exerts a pro-angiogenic effect in
567   macrophages of mammary tumor-bearing mice. *Frontiers in physiology* 2014; 5:
568   17.

569
20   Garcia-Areas R, Libreros S, Simoes M, Castro-Silva C, Gazaniga N, Amat S *et*
al. Suppression of tumor-derived Semaphorin 7A and genetic ablation of host-
derived Semaphorin 7A impairs tumor progression in a murine model of
advanced breast carcinoma. *Int J Oncol* 2017; 51: 1395-1404.

21   Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ *et al*. The
genomic and transcriptomic architecture of 2,000 breast tumours reveals novel
subgroups. *Nature* 2012; 486: 346-352.
578

579   22   Jacob A, Jing J, Lee J, Schedin P, Gilbert SM, Peden AA *et al*. Rab40b regulates
580   trafficking of MMP2 and MMP9 during invadopodia formation and invasion of
581   breast cancer cells. *Journal of cell science* 2013; 126: 4647-4658.
582

583   23   Maller O, Hansen KC, Lyons TR, Acerbi I, Weaver VM, Prekeris R *et al*. Collagen
584   architecture in pregnancy-induced protection from breast cancer. *Journal of cell*
585   *science* 2013; 126: 4108-4110.
586

587   24   Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. *The*
588   *Journal of clinical investigation* 2009; 119: 1420-1428.
589

590   25   Balmanno K, Cook SJ. Tumour cell survival signalling by the ERK1/2 pathway.
591   *Cell death and differentiation* (Review) 2008; 16: 368.
592

593   26   Reddig PJ, Juliano RL. Clinging to life: cell to matrix adhesion and cell survival.
594   *Cancer Metastasis Rev* 2005; 24: 425-439.
595

596   27   Tiwari N, Gheldof A, Tatari M, Christofori G. EMT as the ultimate survival
597   mechanism of cancer cells. *Seminars in Cancer Biology* 2012; 22: 194-207.
598

599   28   St-Germain ME, Gagnon V, Parent S, Asselin E. Regulation of COX-2 protein
600   expression by Akt in endometrial cancer cells is mediated through NF-
601   kappaB/IkappaB pathway. *Mol Cancer* 2004; 3: 7.
602

603   29   Schedin P. Pregnancy-associated breast cancer and metastasis. *Nature reviews*
604   *Cancer* 2006; 6: 281-291.
605

606   30   Albrektsen G, Heuch I, Hansen S, Kvale G. Breast cancer risk by age at birth,
607   time since birth and time intervals between births: exploring interaction effects.
608   *British journal of cancer* 2005; 92: 167-175.
609

610   31   Goddard ET, Bassale S, Schedin T, Jindal S, Johnston J, Cabral E *et al*.
611   Association Between Postpartum Breast Cancer Diagnosis and Metastasis and
612   the Clinical Features Underlying Risk. *JAMA Netw Open* 2019; 2: e186997.
613

614   32   Welch DRA-Ohoo, Hurst DRA-Ohoo. Defining the Hallmarks of Metastasis. LID -
615   10.1158/0008-5472.CAN-19-0458 [doi].
616

617   33   Cerny J, Stockinger H, Horejsi V. Noncovalent associations of T lymphocyte
618   surface proteins. *Eur J Immunol* 1996; 26: 2335-2343.
619

Zhou Y, Gunput RA, Pasterkamp RJ. Semaphorin signaling: progress made and promises ahead. *Trends in biochemical sciences* 2008; 33: 161-170.

Liu H, Juo ZS, Shim AH, Focia PJ, Chen X, Garcia KC *et al*. Structural basis of semaphorin-plexin recognition and viral mimicry from Sema7A and A39R complexes with PlexinC1. *Cell* 2010; 142: 749-761.

Fong KP, Barry C, Tran AN, Traxler Ea, Wannemacher KM, Tang HY *et al*. Deciphering the human platelet sheddome. *Blood* 2011; 117: 15-27.

Jaimes Y, Gras C, Goudeva L, Buchholz S, Eiz-Vesper B, Seltsam A *et al*. Semaphorin 7A inhibits platelet production from CD34+ progenitor cells. *Journal of thrombosis and haemostasis : JTH* 2012; 10: 1100-1108.

Xie J, Wang H. Semaphorin 7A as a potential immune regulator and promising therapeutic target in rheumatoid arthritis. *Arthritis Res Ther* 2017; 19: 10.

Kopp MA, Brommer B, Gatzemeier N, Schwab JM, Pruss H. Spinal cord injury induces differential expression of the profibrotic semaphorin 7A in the developing and mature glial scar. *Glia* 2010; 58: 1748-1756.

Gan Y, Reilkoff R, Peng X, Russell T, Chen Q, Mathai SK *et al*. Role of semaphorin 7a signaling in transforming growth factor beta1-induced lung fibrosis and scleroderma-related interstitial lung disease. *Arthritis and rheumatism* 2011; 63: 2484-2494.

Reilkoff RA, Peng H, Murray LA, Peng X, Russell T, Montgomery R *et al*. Semaphorin 7a+ regulatory T cells are associated with progressive idiopathic pulmonary fibrosis and are implicated in transforming growth factor-beta1-induced pulmonary fibrosis. *American journal of respiratory and critical care medicine* 2013; 187: 180-188.

Kang H-R, Lee CG, Homer RJ, Elias Ja. Semaphorin 7A plays a critical role in TGF-beta1-induced pulmonary fibrosis. *The Journal of experimental medicine* 2007; 204: 1083-1093.

Gilmore AP, Owens TW, Foster FM, Lindsay J. How adhesion signals reach a mitochondrial conclusion--ECM regulation of apoptosis. *Current opinion in cell biology* 2009; 21: 654-661.

Knowles LM, Gurski LA, Engel C, Gnarra JR, Maranchie JK, Pilch J. Integrin αvβ3 and fibronectin upregulate Slug in cancer cells to promote clot invasion and metastasis. *Cancer research* 2013; 73: 6175-6184.

Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal transition. *Nature reviews Molecular cell biology* 2014; 15: 178-196.

667 46   Chao Y, Wu Q, Acquafondata M, Dhir R, Wells A. Partial mesenchymal to
668   epithelial reverting transition in breast and prostate cancer metastases. *Cancer*
669   *microenvironment : official journal of the International Cancer Microenvironment*
670   *Society* 2012; 5: 19-28.
671

672 47   Smith BN, Bhowmick NA. Role of EMT in Metastasis and Therapy Resistance. *J*
673   *Clin Med* 2016; 5.
674

675 48   Nurwidya F, Murakami A, Takahashi F, Takahashi K. Molecular mechanisms
676   contributing to resistance to tyrosine kinase-targeted therapy for non-small cell
677   lung cancer. *Cancer biology & medicine* 2012; 9: 18-22.
678

679 49   Nurwidya F, Takahashi F, Murakami A, Takahashi K. Epithelial mesenchymal
680   transition in drug resistance and metastasis of lung cancer. *Cancer research and*
681   *treatment : official journal of Korean Cancer Association* 2012; 44: 151-156.
682

683 50   Uchikado Y, Okumura H, Ishigami S, Setoyama T, Matsumoto M, Owaki T *et al*.
684   Increased Slug and decreased E-cadherin expression is related to poor
685   prognosis in patients with gastric cancer. *Gastric cancer : official journal of the*
686   *International Gastric Cancer Association and the Japanese Gastric Cancer*
687   *Association* 2011; 14: 41-49.
688

689 51   Deep G, Jain AK, Ramteke A, Ting H, Vijendra KC, Gangar SC *et al*. SNAI1 is
690   critical for the aggressiveness of prostate cancer cells with low E-cadherin. *Mol*
691   *Cancer* 2014; 13: 37.
692

693 52   McKeithen D, Graham T, Chung LW, Odero-Marah V. Snail transcription factor
694   regulates neuroendocrine differentiation in LNCaP prostate cancer cells. *Prostate*
695   2010; 70: 982-992.
696

697 53   Cheng GZ, Chan J, Wang Q, Zhang W, Sun CD, Wang LH. Twist
698   transcriptionally up-regulates AKT2 in breast cancer cells leading to increased
699   migration, invasion, and resistance to paclitaxel. *Cancer research* 2007; 67:
700   1979-1987.
701

702 54   Baritaki S, Yeung K, Palladino M, Berenson J, Bonavida B. Pivotal roles of snail
703   inhibition and RKIP induction by the proteasome inhibitor NPI-0052 in tumor cell
704   chemoimmunosensitization. *Cancer research* 2009; 69: 8376-8385.
705

706 55   Vesuna F, Lisok A, Kimble B, Domek J, Kato Y, van der Groep P *et al*. Twist
707   contributes to hormone resistance in breast cancer by downregulating estrogen
708   receptor-alpha. *Oncogene* 2012; 31: 3223-3234.
709

710   56   Parkash J, Messina A, Langlet F, Cimino I, Loyens A, Mazur D *et al*.
711   Semaphorin7A regulates neuroglial plasticity in the adult hypothalamic median
712   eminence. *Nature communications* 2015; 6: 6385.
713

714   57   Eke I, Storch K, Krause M, Cordes N. Cetuximab attenuates its cytotoxic and
715   radiosensitizing potential by inducing fibronectin biosynthesis. *Cancer research*
716   2013; 73: 5869-5879.
717

718   58   Pontiggia O, Sampayo R, Raffo D, Motter A, Xu R, Bissell MJ *et al*. The tumor
719   microenvironment modulates tamoxifen resistance in breast cancer: a role for
720   soluble stromal factors and fibronectin through beta1 integrin. *Breast cancer*
721   *research and treatment* 2012; 133: 459-471.
722

723   59   Han SW, Roman J. Fibronectin induces cell proliferation and inhibits apoptosis in
724   human bronchial epithelial cells: pro-oncogenic effects mediated by PI3-kinase
725   and NF-kappa B. *Oncogene* 2006; 25: 4341-4349.
726

727   60   Hattar R, Maller O, McDaniel S, Hansen KC, Hedman KJ, Lyons TR *et al*.
728   Tamoxifen induces pleiotrophic changes in mammary stroma resulting in
729   extracellular matrix that suppresses transformed phenotypes. *Breast cancer*
730   *research : BCR* 2009; 11: R5.
731

732   61   Cheng JQ, Jiang X, Fraser M, Li M, Dan HC, Sun M *et al*. Role of X-linked
733   inhibitor of apoptosis protein in chemoresistance in ovarian cancer: possible
734   involvement of the phosphoinositide-3 kinase/Akt pathway. *Drug resistance*
735   *updates : reviews and commentaries in antimicrobial and anticancer*
736   *chemotherapy* 2002; 5: 131-146.
737

738   62   Fraser M, Leung BM, Yan X, Dan HC, Cheng JQ, Tsang BK. p53 is a
739   determinant of X-linked inhibitor of apoptosis protein/Akt-mediated
740   chemoresistance in human ovarian cancer cells. *Cancer research* 2003; 63:
741   7081-7088.
742

743   63   Knuefermann C, Lu Y, Liu B, Jin W, Liang K, Wu L *et al*. HER2/PI-3K/Akt
744   activation leads to a multidrug resistance in human breast adenocarcinoma cells.
745   *Oncogene* 2003; 22: 3205-3212.
746

747   64   Li J, Feng Q, Kim JM, Schneiderman D, Liston P, Li M *et al*. Human ovarian
748   cancer and cisplatin resistance: possible role of inhibitor of apoptosis proteins.
749   *Endocrinology* 2001; 142: 370-380.
750

751   65   Gurrapu S, Pupo E, Franzolin G, Lanzetti L, Tamagnone LA-Ohoo.
752   Sema4C/PlexinB2 signaling controls breast cancer cell growth, hormonal
753   dependence and tumorigenic potential.
754

755 66 Crump LS, Wyatt G, Porter WW, Richer J, Lyons TR. Hormonal regulation of
756 Semaphorin 7a in ER+ breast cancer drives therapeutic resistance. *bioRxiv*
757 2019: 650135.
758
759 67 Provenzano PP, Eliceiri KW, Campbell JM, Inman DR, White JG, Keely PJ.
760 Collagen reorganization at the tumor-stromal interface facilitates local invasion.
761 *BMC medicine* 2006; 4: 38.
762
763 68 Provenzano PP, Inman DR, Eliceiri KW, Knittel JG, Yan L, Rueden CT *et al*.
764 Collagen density promotes mammary tumor initiation and progression. *BMC*
765 *medicine* 2008; 6: 11-11.
766
767 69 Slocum E, Craig A, Villanueva A, Germain D. Parity predisposes breasts to the
768 oncogenic action of PAPP-A and activation of the collagen receptor DDR2.
769
770 70 Ristimäki A, Sivula A, Lundin J, Ristima A, Lundin M, Salminen T *et al*.
771 Prognostic Significance of Elevated Cyclooxygenase-2 Expression in Breast
772 Cancer Advances in Brief Prognostic Significance of Elevated Cyclooxygenase-2
773 Expression in Breast Cancer 12002: 632-635.
774
775 71 Denkert C, Winzer K-J, Müller B-M, Weichert W, Pest S, Köbel M *et al*. Elevated
776 expression of cyclooxygenase-2 is a negative prognostic factor for disease free
777 survival and overall survival in patients with breast carcinoma. *Cancer* 2003; 97:
778 2978-2987.
779
780 72 Pasterkamp RJ, Peschon JJ, Spriggs MK, Kolodkin AL. Semaphorin 7A
781 promotes axon outgrowth through integrins and MAPKs. *Nature* 2003; 424: 398-
782 405.
783
784 73 Hu M, Peluffo G, Chen H, Gelman R, Schnitt S, Polyak K. Role of COX-2 in
785 epithelial-stromal cell interactions and progression of ductal carcinoma in situ of
786 the breast. *Proceedings of the National Academy of Sciences of the United*
787 *States of America* 2009; 106: 3372-3377.
788
789 74 Valdez KE, Fan F Fau - Smith W, Smith W Fau - Allred DC, Allred Dc Fau -
790 Medina D, Medina D Fau - Behbod F, Behbod F. Human primary ductal
791 carcinoma in situ (DCIS) subtype-specific pathology is preserved in a mouse
792 intraductal (MIND) xenograft model.
793
794 75 Nagy A, Lanczky A, Menyhart O, Gyorffy B. Author Correction: Validation of
795 miRNA prognostic power in hepatocellular carcinoma using expression data of
796 independent datasets. *Scientific reports* 2018; 8: 11515.
797
798 76 Dean JL, McClendon aK, Hickey TE, Butler LM, Tilley WD, Witkiewicz AK *et al*.
799 Therapeutic response to CDK4/6 inhibition in breast cancer defined by ex vivo
800 analyses of human tumors. *Cell cycle* 2012; 11: 2756-2761.

801
802   77   Barrett AS, Wither MJ, Hill RC, Dzieciatkowska M, D'Alessandro A, Reisz JA *et*
803   *al*. Hydroxylamine Chemical Digestion for Insoluble Extracellular Matrix
804   Characterization. *Journal of proteome research* 2017; 16: 4177-4184.
805
806   78   Reisz JA, Nemkov T, Dzieciatkowska M, Culp-Hill R, Stefanoni D, Hill RC *et al*.
807   Methylation of protein aspartates and deamidated asparagines as a function of
808   blood bank storage and oxidative stress in human red blood cells. *Transfusion*
809   2018; 58: 2978-2991.
810
811   79   Chong J, Xia J. MetaboAnalystR: an R package for flexible and reproducible
812   analysis of metabolomics data. *Bioinformatics* 2018; 34: 4313-4314.
813
814   80   Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14:
815   more genomes, a new PANTHER GO-slim and improvements in enrichment
816   analysis tools. *Nucleic acids research* 2019; 47: D419-d426.
817
818

Figure 1. **SEMA7A promotes growth and invasion in PPBC**. **A.** Tumor volumes for

control (Ctrl) or shSEMA7A (SEMA7A-KD) MCF10DCIS cells in postpartum hosts

(n=12/group), inset: representative immunoblot for SEMA7A and GAPDH. **B.** Tumors

from Ctrl and SEMA7A-KD from postpartum hosts scored for invasion (n=12/group). **C.**

Trichrome stained quantification (top) and representative images (bottom) of collagen of

tumors from Ctrl and SEMA7A-KD tumors from postpartum hosts, scale bars 50μm

(n=12/group) . **D.** IHC quantification (top) and representative images (bottom) of IHC for

COX-2 of tumors from Ctrl and SEMA7A-KD tumors from postpartum hosts, scale bars

50μm (n=12/group) (*p<0.05, **p<0.01, ***p<0.0005, ****p<0.0001, t-test).

Figure 2. SEMA7A expression is increased in DCIS patient samples. A.

Representative images of normal (top) or DCIS lesions (bottom) from postpartum

patients from the CU Anschutz Medical Campus Young Women's BC (YWBC) stained

by IHC for SEMA7A, with quantification mask to the right, blue=negative; yellow=weak;

orange=medium; red=strong; scale bars 50 μm. **B.** Quantification of SEMA7A IHC of

normal or DCIS patient samples from the CU Anschutz YWBC cohort (n=103), then

separated by parity status, nulliparous (n=52) or postpartum (n=51). **C**. Quantification of

SEMA7A IHC from nulliparous patients by patient, paired t-test. **D.** Quantification of

SEMA7A IHC from postpartum patients by patient, paired t-test. **E.** Representative

images of SEMA7A IHC stain from tissue array of samples, scale bars 50μm. **F.**

Quantification of SEMA7A IHC stain from **C**; normal (n=32) or DCIS (n=30). **G.**

Quantification of SEMA7A IHC by patient, paired t-test. (*p<0.05, **p<0.01, ****p<0.001)

Figure 3. **SEMA7A expression is sufficient to drive tumor growth and invasion**. **A**.

Tumor volumes for control (NUL Ctrl) or SEMA7A overexpressing (NUL SEMA7A OE)

in nulliparous hosts (n=10/group), inset: representative immunoblot for SEMA7A and

GAPDH. **B**. Tumors from NUL Ctrl and NUL SEMA7A OE scored for invasion

(n=10/group). **C**. Trichrome stained quantification (top) and representative images

(bottom) of collagen of tumors from NUL Ctrl and NUL SEMA7A OE tumors, scale bars

50μm **D.** IHC quantification (top) and representative images (bottom) of IHC for COX-2

of tumors from NUL Ctrl and NUL SEMA7A OE tumors, scale bars 50μm (*$p<0.05$,

**$p<0.01$, ***$p<0.0005$, ****$p<0.0001$, t-test).

Figure 4. **SEMA7A promotes invasion via fibroblast mediated collagen deposition.**

A. Ctrl or SEMA7A OE MCF10DCIS cells embedded in matrigel scored for invasion. **B.**

Ctrl or SEMA7A OE MCF10DCIS cells embedded in matrigel plus 20% collagen. **C**.

Control (Ctrl) or shSEMA7A (KD) MDA-MB-231 cells in matrigel plus 25% collagen

scored for invasion. **D**. Quantitative RT-PCR (q-PCR) for COL1A1 in fibroblasts in

serum free media (SFM) treated with 10 ng/mL TGFβ or conditioned media (CM) from

control (Ctrl) or SEMA7A overexpressing (SEMA7A OE) MDA-MB-231 cell lines. **E**.

qPCR for COLA1A in fibroblasts in SFM treated with 75 ng/μL purified SEMA7A.

(*$p<0.05$, **$p<0.01$,***$p<0.005$, t-test).

Figure 5. **SEMA7A promotes mesenchymal protein expression and phenotypes**.

A. Global proteomics analysis of secreted proteins reveals decreased fibronectin (FN) in

MCF10DCIS shSEMA7A (SEMA7A-KD) tumors compared to control (Ctrl) ex vivo

(n=10-11/group). **B&C**. IHC for FN in MCF10DCIS Ctrl and SEMA7A-KD tumors (B),

and MCF10DCIS Ctrl and SEMA7A overexpressing (SEMA7A OE) tumors, scale bar

50μm. **D-F**. Additional decreases in mesenchymal or mesenchymal associated proteins

observed by proteomics in shSEMA7A (SEMA7A-KD) tumors ex vivo. **G**.

Immunofluorescence for F-actin in MDA-MB-231 Ctrl or KD cells and quantification for

cell aspect ratio (length/width). **H**. Immunofluorescence for F-actin in MCF10DCIS Ctrl

or SEMA7A OE cells and quantification for cell aspect ratio (length/width). (*p<0.05,

**p<0.01, ***p<0.005, ****p<0.001, t-test)

<u>**Figure 6**</u>. **SEMA7A promotes cell survival via fibronectin. A&B**. Fold change of

cleaved caspase 3/7 activity in control (Ctrl) and shSEMA7A (KD) in MCF10DCIS (**A**) or

MDA-MB-231 (**B**) cell lines. **C.** Cleaved caspase 3/7 activity measured over time in

MDA-MB-231 Ctrl or KD cells. **D.** Representative immunoblot for phospho-AKT (S473),

total AKT, or GAPDH in Ctrl or KD, quantified to the right. **E**. Cleaved caspase 3/7

activity control (Ctrl) or shSEMA7A (KD) cells plated on tissue culture plastic (TC),

laminin (LAM), collagen I (COL) or fibronectin (FN). **F&G.** Cleaved caspase 3/7 activity

MCF10DCIS (**F**) or MDA-MB-231 (**G**) Ctrl or KD cells in forced suspension. **H.** Cleaved

caspase 3/7 activity in MCF10DCIS Ctrl or shCOX-2 (KD1/KD2) cell lines in forced

suspension.

<u>**Figure 7.**</u> **SEMA7A drives metastatic seeding and poor prognosis in patients**. **A.**

Average area of metastasis in lungs of mice after tail vein injection with MDA-MB-231

control (Ctrl) or shSEMA7A (KD1, KD2) cells (n=5/group), representative images of

lungs to the right. **B.** Kaplan-Meier analysis of ER- BCs using Km plotter for SEMA7A,

COX-2 and FN mRNA expression for distant metastasis free survival (DMFS) (n=228).

C. Kaplan-Meier analysis of ER-BCs in the Ma Breast Cancer dataset using GOBO for

SEMA7A, COX-2 and FN mRNA expression for distant metastasis free survival (DMFS)

(n=320) **D&E.** Co-expression analysis of the TCGA BC provisional cohort using

CBioPortal for all BCs (n=1108). (*$p<0.05$, **$p<0.01$,***$p<0.005$ ****$p<0.001$)**.**

<u>**Figure 8**</u>**. Model depicting SEMA7A mediated invasion and cell survival.** Our

current data suggests SEMA7A promotes fibroblast mediated collagen deposition,

resulting in COX-2 expression and tumor cell invasion. Our results also suggest that

SEMA7A can promote cell survival through SEMA7A mediated fibronectin expression.

Figure 1

Figure 2

Figure 3

A

SEMA7A — 75kDa
GAPDH — 37kDa
Ctrl OE

Tumor volume (mm³)

** ** **

NUL Ctrl
NUL SEMA7A OE

Weeks

B

100%
NUL Ctrl

44.4%
33.3%
22.2% *
NUL SEMA7A OE

■ DCIS ■ IDC+DCIS ■ DCIS+microinvasion

C

% Collagen +

***

NUL Ctrl   NUL SEMA7A OE

D

% positive COX-2 pixels

**

NUL Ctrl   NUL SEMA7A OE

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8