Enhanced biodegradation of naphthalene by *Pseudomonas* sp. consortium

immobilized in calcium alginate beads

Kunal Dutta[1], Sergey Shityakov[2], Ibrahim Khalifa[3,4], Saroj Ballav[1], Debarati Jana[1],
Tuhin Manna[1], Monalisha Karmakar[1], Priyanka Raul[1], Kartik Chandra Guchhait[1] and
Chandradipa Ghosh[1,*]

[1]Microbiology and Immunology Laboratory, Department of Human Physiology with
Community Health, Vidyasagar University, Midnapore -721102, West Bengal, India.

[2]Department of Anaesthesia and Critical Care, University of Würzburg, 97080
Würzburg, Germany.

[3]Food Technology Department, Faculty of Agriculture, 13736 Moshtohor, Benha
University, Egypt.

[4]College of Food Science and Technology, Huazhong Agricultural University, Wuhan
430070, PR China.
16
17
18 *Corresponding author:*
19   Prof. Chandradipa Ghosh
20   Microbiology and Immunology Laboratory
21   Department of Human Physiology with Community Health
22   Vidyasagar University
23   Midnapore - 721102
24   West Bengal,
25   INDIA
26   Email : ch_ghosh@mail.vidyasagar.ac.in
27   Phone : +91 3222 276554/555/557 Extn. 450
28   Fax   : +91 3222 275329
29
30
Kunal Dutta   : orcid.org/0000-0002-0818-8787

Sergey Shityakov   : orcid.org/0000-0002-6953-9771

Ibrahim Khalifa   : orcid.org/0000-0002-7648-2961

Abstract

Polycyclic aromatic hydrocarbons (PAHs) belong to a large group of organic

pollutant which considers as a potential health hazard to living beings. Herein,

naphthalene biodegradation potential by free and immobilized *Pseudomonas putida*

strain KD10 and *Pseudomonas* sp. consortium were studied. Additionally, naphthalene

1, 2-dioxygenase (*nah*Ac) was sequenced and analyzed, which reveals two altered

amino acid residues. However, the altered amino acid residues are not present in the

vicinity of the active site. The gas-phase binding free energy (ΔG_{London}) of the mutant

variant of naphthalene 1, 2-dioxygenase was -7.10 kcal mol^{-1} which closely resembles

the wild type variant. Naphthalene biodegradation rate by *Pseudomonas putida* strain

KD10 was 79.12 mg L^{-1} day^{-1} and it was significantly elevated up to 123 mg L^{-1} day^{-1}

by the immobilized *Pseudomonas* sp. consortium. The half-life ($t_{1/2}$) for naphthalene

biodegradation was 3.1 days with the inhibition constant (k_i), substrate saturation

constant (k_s) and maximum specific degradation rate constant (q_{max}) of 1268 mg L^{-1},

395.5 mg L^{-1} and 0.65 h^{-1}, respectively, for the *Pseudomonas putida* strain KD10.

However, the t$_{1/2}$ value was significantly reduced to 2 days along with k_i, k_s and q_{max}

values of 1475 mg L^{-1}, 298.8 mg L^{-1} and 0.71 h^{-1}, respectively, by the immobilized

Pseudomonas sp. consortium. The GC-MS data suggest that KD10 might follow D-

gluconic acid mediated meta-cleavage pathway of catechol biodegradation. It is

concluded that naphthalene biodegradation performance by immobilized *Pseudomonas*

sp. consortium was superior to free or immobilized *Pseudomonas putida* KD10.

Microbial consortium immobilization could be a useful tool for water quality

management and environmental remediation.

Keywords:

Pseudomonas sp., petroleum wastes, biodegradation, cell immobilization, mutant

naphthalene 1, 2-dioxygenase, rigid-flexible molecular docking.

Highlights

• Superior naphthalene biodegradation by *Pseudomonas* sp. consortium
immobilized in calcium alginate beads.

• A common mutation prone amino acid stretch inside chain A of naphthalene 1,
2-dioxygenase has been identified.

• A new naphthalene biodegradation pathway by *Pseudomonas putida* strain
KD10 has been proposed.

 1. Introduction

 Polycyclic aromatic hydrocarbons (PAHs) are considered as the potential health

 hazard for living beings (Kumari et al., 2018). Environmental agencies including US-

 EPA, European Union, Environment-Canada, registered PAH as the priority pollutants

 that require immediate human intervention (Wang et al., 2018). The physicochemical

 properties of PAH make them a major contributor for soil and groundwater

 contamination through bio-magnification (Norris, 2017). Moreover, according to the

 Environmental Health Hazard Assessment, U.S.A., naphthalene (NAP) concentration

 beyond 170 ppb is not safe drinking (Bruce et al., 1998). Different orthodox and

 expensive techniques for environmental remediation, *viz.,* incineration, gasification,

 plasma-gasification have been replaced by green-technologies such as bioremediation,

 phytoremediation, nanoremediation, *etc.* (Thomé et al., 2018). Bioremediation is

 considered as the most cost-effective and eco-friendly oil spill management technique

 (Wilson and Jones, 1993). However, while considering the vast volume of a mobile

 open water system, bioremediation stumble upon several limiting factors such as the

 low local concentration of the effective microorganisms, loss of active microorganisms,

 etc. (Chen et al., 2017). Conversely, cell immobilization provides several advantages

 such as, it helps to retain high local concentration of the effective microorganisms, keep

 intact bacterial cell membrane stability and can be stored for future reuse (Bhardwaj et

 al., 2000). Overall, cell immobilization can markedly improve the stability and

 efficiency of the bioremediation process (Mrozik and Piotrowska-Seget, 2010; Tyagi et

 al., 2011). Cell immobilization using calcium alginate beads (CABs) is a convenient

 option where maximum cells remain viable and can tolerate high concentration of the

 toxicant (Lee and Heo, 2000). Moreover, calcium alginate is nontoxic to the bacterial

cell and it has low production cost which facilities easy reuse (Bhardwaj et al., 2000).

Biodegradation of a toxicant (complex nutrient for bacteria) by the microbial

consortium can efficiently enhance biodegradation rate (Kumari et al., 2018). This rate

enhancement could be achieved by several ways such as, different biodegradation

pathways of each individual bacterium (Dutta et al., 2018), a metabolic intermediates of

one bacteria may act as the starting material of other bacteria (Surkatti and El-Naas,

2018), different genetic makeup (Woyke et al., 2006), synergetic effects of different

microbial species, (Ghazali et al., 2004) or by synthesising different variant of catalytic

enzymes. Bacterial cells in the microbial consortium can also combine their metabolic

capabilities to utilize the common complex nutrient (Gilbert et al., 2003).

Biodegradation of complex hydrocarbon mixture by the microbial consortium offers a

combination of diverse enzymes which promotes the biodegradation processes

(Wongwilaiwalin et al., 2010). However, microbial consortium immobilization for

biodegradation of PAHs was not studied previously. The aim of the present study is to

evaluate the naphthalene biodegradation performance by *Pseudomonas putida* strain

KD10 and *Pseudomonas* sp. consortium as a free and immobilized format. Previous

study showed that *Pseudomonas putida* strain KD9 cells decrease in size and shape

from rod to sphere during naphthalene biodegradation and their specific growth rate was

also little slower (Dutta et al., 2018). Keeping this fact in the mind, both morphological

types of bacteria were applied in this present set of study. Additionally, the naphthalene

1, 2-dioxygenase (*nah*Ac) was sequenced and analyzed.

2. Materials and methods

2.1. Chemicals

Naphthalene was purchased from Sigma-Aldrich chemicals Pvt. Ltd. (USA) and all other chemicals used for media preparation were procured from HiMedia Laboratory (Mumbai, India). GC-MS and HPLC grade solvents were procured from Fisher Scientific (Mumbai, India). Sodium alginate (CAS No. 9005-38-3) of medium viscosity was purchased from Merck Pvt. Ltd. (USA).

2.2. Microorganisms, growth media, growth condition and consortium preparation

Soil samples were collected from petroleum refinery waste sites near Indian Oil (Haldia, West Bengal, India). The enrichment isolation and strain identification were carried out according to the standard protocol described previously (Dutta et al., 2017). Carbon deficient minimum medium (CSM), with a pH of 7.1, was used to cultivate the bacteria. Naphthalene was used as a sole source of carbon and energy in CSM with following compositions: 0.2 g L^{-1} $MgSO_4$, $7H_2O$; 0.08 g L^{-1} Ca $(NO_3)_2$, $4H_2O$; 0.005 g L^{-1} $FeSO_4$, $7H_2O$; 4.8 g L^{-1}, K_2HPO_4; 1.2 g L^{-1} KH_2PO_4. *Pseudomonas putida* strain KD6 (KX786159.1) and *Pseudomonas putida* strain KD9 (KX786158.1) and the newly isolated *Pseudomonas putida* strain KD10 (KX786157.1) were used to prepare the blend of *Pseudomonas* sp. consortium. The consortium was maintained in Luria-Bertani broth at 31°C with 150 rpm in order to grow the bacterial cell in its normal size and shape. Alternatively, bacterial cells were grown in CSM with naphthalene a as sole source of carbon and energy to obtain altered morphological variant.

2.3. Gene sequence analysis

2.3.1. Detection of naphthalene 1, 2-dioxygenase and catechol 2, 3-dioxygenase

The conventional polymerase chain reaction (PCR) for naphthalene 1, 2-dioxygenase (*nah*Ac) and catechol 2, 3-dioxygenase (*nah*H) using specific primers (Table S1) were performed. Additionally, PCR product of *nah*Ac was sequenced and analyzed according to the methods described previously (Dutta et al., 2017).

2.3.2. Clustering and phylogenetic analysis

The evolutionary distance of naphthalene 1, 2-dioxygenase among different bacterial species was analyzed using BLOSUM weighted matrix followed by pairwise distance computation using MEGA (v7.0) (Kumar et al., 2016). The distance matrix was then clustered using R (Team, 2013) to create the cladogram and heatmap. The phylogenetic position of the isolated *Pseudomonas putida* strain KD10 was analyzed using the previous method (Dutta et al., 2017).

2.3.3. Molecular docking

The rigid-body molecular docking was conducted using Auto Dock (v4.2.1) (Morris et al., 1998). Briefly, the center grid dimensions were set to 20.271×61.989×87.168 with a grid spacing of 0.375 Å. The virtual screening was

repeated for 10 times with the unaltered docking parameters having 2.0 Å cluster

tolerance. Additionally, the rigid-flexible molecular docking was performed using

Molecular Operation Environment (Chemical Computing Group, Montreal Inc.,

Canada). The latter scoring function was employed to identify the most favourable

docked poses and to estimate the binding affinity of the protein-ligand complexes. The

non-covalent interactions were analyzed using the previous method (Salentin, S., et al.,

2015).

2.4. Enzyme kinetic assay

The enzyme kinetic parameters of the naphthalene 1, 2-dioxygenase $_{I250,\ V256}$ was

performed using the cell-free extract of the *Pseudomonas putida* strain KD10, grown in

250 mL of CSM with naphthalene (500 mg L^{-1}) as a sole source of carbon and energy as

described by previously (Dutta et al., 2017).

2.4. Detection of solvent efflux pumps system

The detection of solvent efflux pump system (*srp*ABC) in *Pseudomonas putida*

strain KD10 was performed using conventional polymerase chain reaction in a thermal

cycler (Mastercycler® nexus gradient, Eppendrof, (Germany). Standard reaction

mixtures were prepared using forward, reverse primers (Table S1) as described

previously (Dutta et al., 2018).

2.5. *Cell immobilization*

2.5.1. Cell immobilization in calcium alginate beads

Pseudomonas putida strain KD10 and *Pseudomonas* sp. consortium were

immobilized according to the standard protocol described previously (Daâssi et al.,

2014). Briefly as sodium alginate was dissolved in 0.9 wt. % NaCl (1 gm in 40 mL 0.9

wt. % NaCl) for 24 h and sterilized by autoclaving (121°C for 15 min). Two grams of

bacterial cell mass was added to 8 ml of NaCl solution and again added to the sterile

alginate solution. The mixture was then gently vortex for complete homogenization and

extruded dropwise through a hypodermic syringe into chilled sterile $CaCl_2$ solution

(Figure 2). The beads were hardened in the same solution at room temperature with

gentle stirring for 1 h. Finally, the beads were washed several times with 0.9 wt. %

NaCl to remove excess calcium ions and free cells. The beads have an average diameter

of 0.5 mm and stored at 4°C. Sterile beads (without microorganisms) were used to

monitor the abiotic loss of naphthalene. Sodium alginate of medium viscosity

(≥2,000 cP) was used to prepare the calcium alginate beads (CABs).

2.5.2. Cell viability count

The viable cell enumeration in the CABs was performed by using a protocol

described previously (Usha et al., 2010). In brief, CABs was washed in saline and keep

submerged for 10 min (for saline soaking). Following soaking, the CABs were shaken

with glass beads for 15 min and 1 g CABs were homogenized in 9 ml saline. The saline

was then used for viable cell enumeration onto a nutrient agar plate. Plating was done

with this treated saline by series dilution method up to 10^{-5} dilutions on nutrient agar

plates. Plating was also done for the initial CFU count. For every dilution, 10 µl of the

solution was plated. Plating was done by the pour plate method. Plates were incubated

at 37°C for 24 h. (Jain and Pradeep, 2005).

2.6. Scanning electron microscopic (SEM) study

Samples for SEM analysis were prepared by a protocol described elsewhere

(Dutta et al., 2018). The dried samples were coated in a sputter coater (Quorum-

SC7620) under vacuum with a thin gold layer right before SEM analysis using a

scanning electron microscope (Zeiss, EVO 18, Germany) with an accelerating voltage

of 5 kV.

2.7. Biodegradation kinetics

2.7.1. Biodegradation of naphthalene in the liquid medium

Naphthalene biodegradation study was conducted by the protocol described

previously (Dutta et al., 2017). The conical flasks were incubated at 31°C with 150 rpm

and uninoculated conical flask were used as control. Culture medium was collected at

regular interval of 72 h for degradation and growth kinetic study. Additionally, the

effect of initial concentration of naphthalene (150-2500 mg L^{-1}) was studied with

different immobilized systems.

2.7.2. Chemical analysis

Naphthalene biodegradation was analyzed by using the 1260 infinity series HPLC

system (Agilent, Santa Clara, CA, USA) equipped with Zorbax SB-C18 reversed-phase

column (4.6 × 12.5 mm, 5 µm). The analysis of naphthalene concentrations was

conducted using isocratic elution conditions with the mobile phase 80:20 (v/v)

methanol: water at a flow rate of 1 ml min^{-1}. The detection was performed at 254 nm

according to the protocol described in the literature (Dutta et al., 2017). Additionally,

the metabolic intermediates of naphthalene biodegradation were analyzed using the GC-

MS system (GC Trace GC Ultra, MS-Polarisq, Thermo Scientific India Pvt. Ltd)

equipped with a capillary column (TR-WaxMS, 30 m × 0.25 mm [ID] × 0.25 µm film

thickness) by the protocol described elsewhere (Dutta et al., 2017). The entire analysis

was performed in electron ionization, at full scan mode. The metabolite identification

was based on the mass spectra comparison using the NIST Mass Spectral library (v2.0,

2008).

2.8. Data analysis

The first-order degradation kinetics model was used to estimate the residual

naphthalene in CSM using equations 1, the algorithms as expressed determine the half-

life ($t_{1/2}$) values of naphthalene in CSM. The substrate inhibition kinetic parameters

were calculated using equation 2.

$$C_t = C_0 \times e^{-kt}$$
[1]

$$q = q_{max} \frac{S}{K_s + S + S^2/K_i} \qquad [2]$$

280   The effect of SCS on growth pattern was measured by calculating the difference

281   between optical densities at 600_{nm} and expressed by slight extension (Equation 4) of the

282   Gomperz's sigmoid growth fit equation (Equation 3).

$$y = ae^{-\exp(-k(x - x_c))} \qquad [3]$$

$$\Delta X_c = X_{c2} - X_{c1} \qquad [4]$$

286   The growth pattern change by SCS was considered as CFU shift (Equation 5). Where,

287   X_{c2} = Optical density at X_{c2} and X_{c1} = Optical density at X_{c1}.

$$CFU_{shift} = \Delta X_c \qquad [5]$$

The bacterial growth kinetics were analyzed by applying the Gomperz's model using

Levenberg-Marquardt algorithm in Origin® 2016 (California, USA) and the degradation

kinetics were analyzed by using Graphpad Prism® 6.01 (San Diego, CA, USA).

3. Results and discussion

3.1. Isolation, identification and growth patterns of NAP degrading bacterial strain

Pseudomonas putida strain KD10 was isolated from petroleum refinery waste

with its distinguished colony morphology on CSM agar plate (Figure S1a, b). The PCR

product of 16S rRNA gene of the strain KD10 was sequenced. Moreover, the multiple

sequence alignment followed by phylogenetic assessment suggests that the strain

belongs to the *Pseudomonas putida* group and it showed 99% sequence similarity with

the previously deposited sequences of *Pseudomonas putida* (Figure S2). The 16S rRNA

gene sequence was deposited at NCBI (https://www.ncbi.nlm.nih.gov/) under the

GenBank accession number KX786157.1.

Naphthalene biodegradation by the *Pseudomonas putida* KD10 was

preliminarily confirmed by the catechol test (Figure S1C) followed by growth pattern

analysis in CSM with naphthalene as a sole source of carbon and energy. Additionally,

growth pattern with 0.5 gm % sucrose as a secondary carbon supplement (SCS) was

compared with growth pattern of non-supplemented CSM. Gompertz's model fit

(Equation 3) was used to analyze the growth curve (Table S4).

3.2. Pseudomonas sp. *consortium, growth pattern and NAP biodegradation*

The growth pattern of the *Pseudomonas* sp. consortium, as shown in Figure 1

was optimized at 31°C and pH 7.1 which reveal successful cooperation among

Pseudomonas putida strain KD6, KD9 and KD10. Further, the growth curve model

indicates a significant increase in total biomass of the *Pseudomonas* sp. consortium and

they also showed a high concentration of naphthalene tolerance potential.

3.3. Gene sequence and phylogenetic analysis

Naphthalene 1, 2-dioxygenase (*nah*Ac), encoded by *Pseudomonas putida* strain

KD10 have only two point mutations at I250, and V256 which were replaced by

methionine and glycine respectively (Figure 3A, S6). However, additional mutations

were found in the same chain A at K200, A210, E264, M284 and N334 by replacing

glutamic acid, glycine, aspartic acid, isoleucine, respectively (Dutta et al. 2017, 2018).

The three dimensional structural analysis suggests that the mutations at 200, 210, 284

and 334 causes a little structural change in comparison to the wild type variant of

naphthalene 1, 2-dioxygenase (Figure 3B). Conversely, the amino acids stretch at the

close proximity of the active site residues exactly from 248 to 266 showed a structural

mismatch among all three mutant and wild type variants of naphthalene 1, 2-

dioxygenase (Figure 3B). The active site of an enzyme tends to evolve fast to attain its

maximum performance and functionality in a particular environment (James and

Tawfik, 2003).

The alteration of amino acids bases from 248 to 266 of chain A, was a common

feature in all three mutant variants of *nah*Ac and this particular stretch of amino acid is

very close to the active site residues (Figure 3C). This implies that the "248-266" amino

acid stretch is highly prone to mutation and it may influence on the enzymatic efficiency

and environmental adaptability. Thus, our study provides a new insight, which could be

beneficial for rational approaches of enzyme redesigning.

The biodegradation performance of a bacterial strain is linked with the three

dimensional structure of the catalytic enzyme and ligand binding posture (Singh et al.,

2019). Moreover, alteration of a single amino acid in the catalytic domain of the enzyme

caused different ligand binding postures which eventually lead to a unique

biodegradation pathways (Ferraro et al., 2006). In addition, site directed mutagenesis in the catalytic domain offers superior enzyme activity (Parales et al., 1999; Parales, 2003). The comparative analysis of binding free energy helps to comprehend the superior naphthalene biodegradation performance by the *Pseudomonas* sp. consortium, which is, in fact, the summative activity of all three mutant variants of naphthalene 1, 2-dioxygenase.

The evolutionary trace on the naphthalene 1, 2-dioxygenase among different species were studied through pairwise distance matrix analysis (Figure 4), which suggests a significant intra and inter species difference among *Sphingopyxis sp.*, *Croceicoccus naphthovorans, Burkholderia multivorans, Burkholderia sp. Massilis sp. and Cycloclasticus sp.* Conversely, other *Pseudomonas* sp. particularly, *Pseudomonas benzenivorans, Pseudomonas balearica, Pseudomonas stutzeri, Pseudomonas kunmingensis, Pseudomonas frederikbergensis* have similarities in their version of naphthalene 1, 2-dioxygenase. A few other species such as *Xenophilusa zovorans, Ralsoniam annitolilytica,* and *Paraburkholderia aromaticivorans* does not show significant evolutionary distance in their version of naphthalene 1, 2-dioxygenase (Figure 4). However, a few strains of *Burkholderria multivorans* showed similarities and some other does not (Figure 4). The source of collection of these two *Burkholderria multivorans* species may be the reason for such variation (Li et al., 2007). In a study by (Su et al., 2016), the epigenetic impacts on the metabolic enzymes have been suggested. Moreover, the divergence of evolutionary distances of the same enzyme between different and same bacterial species indicates thrive in a particular microenvironment. However, further studies are needed to identify the underlying cause and mechanisms of such variation intra-species variation.

3.4. Molecular docking

The result depicted from the rigid body molecular docking, suggests *nah*Ac-KD10$_{I250, V256}$ mutant has a little higher binding free energy than that of mutant *nah*Ac encoded by KD9 but little lower than the *nah*Ac mutant variant encoded by the KD6 (Table 1). However, the rigid-flexible molecular docking using MOE algorithms showed *nah*Ac I$_{250}$, G$_{256}$ mutant variant has ΔG_{London} of -7.1 kcal mol^{-1} and $\Delta G_{GBVI/WSA}$ of -1.68 kcal mol^{-1}, which is little higher than other two mutant variants of *nah*Ac encoded by KD6 and KD9 (Table 1). The interacting amino acid residues in the active site were confined to be quite same in three mutant variants, except one variation, *i.e.,* His$_{208}$ which is located about 5.49 Å away from the bicyclic ring of naphthalene (Figure 3D). The altered binding free energy of the *nah*Ac-KD10$_{I250, V256}$ may be due to the fact that the two altered amino acid residues reside in "248-266" and participation of His$_{208}$ as a unique interacting residue (Figure 3C). Furthermore, the non-covalent interactions of the *nah*Ac-KD10$_{I250, V256}$ with naphthalene, phenanthrene and anthracene suggest that Phn 352 was common residue and His 208 is involve in π- staking with a bond angle of 80.58° and 77.19° respectively for naphthalene and anthracene (Table S5).

3.5. Enzyme kinetic assay

Enzyme activity by cell free extract of *Pseudomonas putida* KD10 was studied using Naphthalene as substrate. The cell free extract of strain KD10 degraded Naphthalene, with R^2 of 0.935 indicating the experimental data are well correlated with the model (Table 2). The performance constant (Kcat/ Km) was 0.142×10^3 ml^{-1} mol.$^{-1}$ s^{-1} which further validate the chemical data obtained from chemical analysis (Table 1). In

previous study, the performance constant of *Pseudomonas putida* KD6 was little higher

than *Pseudomonas putida* strain KD10, suggesting the six altered amino acid residues of

the mutant naphthalene 1,2-dioxygenase of the KD6 may be main cause of this

difference (Dutta et al. 2017). The performance constant (Kcat/ Km) is the indicator of

the performance of an enzyme and usually the higher Kcat/Km meaning better

enzymatic performance (Koshland Jr DE, 2002).

3.6. Detection of solvent efflux system

The cellular microenvironment and internal homeostasis are crucial for

maintaining normal cellular functions and cell tends to adopt several strategies in order

to achieve it (Blanco et al., 2016). One such adaptation strategy is the solvent efflux

pump system, that control intracellular toxicity to some extent (Kusumawardhani et al.,

2018). The role of *srp*ABC solvent efflux pump system in the biodegradation of PAHs

are common in literature (Bugg et al., 2000). Besides the other cellular activities,

*srp*ABC also assist in gaining antimicrobial resistance (Schweizer, 2003).

Biodegradation studies on chlorpyrifos indicate that the intermediates formed during

biodegradation, act as an antimicrobial agent to other species, meaning that the

metabolic intermediates may have a role on inter-species competition in a particular

micro-environment to thrive in nutrient-limiting condition (Anwar et al., 2009), (Raes

and Bork, 2008). Besides, the role of *srp*ABC on biodegradation enhancement process

is still poorly understood. The presence of *srp*ABC in *Pseudomonas putida* strain KD10

(Figure S5) and the efficient naphthalene biodegradation property can be interlinked

(Dutta et al., 2018).

3.7. Cell immobilization and viability count

The viscosity of CABs determines its efficiency of the immobilization and its

performance of detoxification of environmental pollutant (Young et al., 2006).

Therefore, sodium alginates of medium viscosity were chosen for this study. The

efficiency of cell immobilization in CABs was evaluated by enumeration of viable cells

(Table S3). Bacteria immobilized in CABs do not significantly lose its viable biomass

after 21 days of incubation at 4°C.

3.8. Naphthalene biodegradation studies

3.8.1 By free Pseudomonas putida strain KD10

Biodegradation kinetics of naphthalene (500 mg L^{-1}) by the *Pseudomonas putida*

strain KD10 was first tested to determine its naphthalene biodegradation potential

(Figure 4). It was found that after 12 days of incubation at 31°C the amount of residual

naphthalene reduced significantly (p<0.05) (Figure 5). Results are also summarised at

Table 3 along with names of the immobilized systems. The first order degradation

kinetic data suggest that at the end of the incubation *Pseudomonas putida* strain KD10

efficiently decompose 95.22 % naphthalene in CSM as a sole source of carbon and

energy. The values of degradation rate constant (k) and half-life ($t_{1/2}$) were 0.2193 and

3.1 days with R^2 of 0.981. In the previous study, 99.1 % of initial naphthalene was

removed within 96 h by strain *Bacillus fusiformis* (BFN). However, the initial

concentration was very low (50 mg L^{-1} of initial naphthalene) (Lin et al., 2010).

After confirmation of naphthalene biodegradation potential of *Pseudomonas*

putida strain KD10, the stain was allowed to grow in association with other two strains

of *Pseudomonas putida*, namely *Pseudomonas putida* strain KD6 and *Pseudomonas*

putida strain KD9 (Dutta et al., 2017; Dutta et al., 2018). These two strains were

selected to develop the blend of *Pseudomonas* sp. consortium, because they were

collected from different isolation points, they encode different variant of *nah*Ac and

they have common optimized growth parameters (temperature 31°C and pH 7.1) (Dutta

et al., 2017 & Dutta et al., 2018). Further, *Pseudomonas putida* strain KD6 encodes a

six point mutant *nah*Ac with the capability to co-degrade high concentration of

naphthalene, phenanthrene (PHN) and pyrene (PYR), (500 mg L^{-1} each). Moreover, the

stain KD9 encodes a four point mutant variant of *nah*Ac with rhamnolipid production

capabilities (Dutta et al. 2018).

In the previous study, $t_{1/2}$ of naphthalene co-biodegradation with phenanthrene

and pyrene by *Pseudomonas putida* strain KD6 was 4.1 days, which was significantly

reduced to 2.2 days when bacterial cells were allowed to grow as *Pseudomonas* sp.

consortium with only naphthalene (Table 3). Further, this value was found as 2.7 days

for *Pseudomonas putida* KD9, suggesting *Pseudomonas putida* strain KD6 might face

some sort of substrate inhibition by the co-presence PHN and PYR in the system (Jiang

et al., 2018) and the cooperative nature of the *Pseudomonas* sp. consortium helps to

enhance the naphthalene bio-utilization. The microbial consortium works several ways,

viz., by the division of labor, cross-feeding, *etc.* (Smid and Lacroix, 2013). Moreover, a

successful consortium could also overcome shortcomings of single bacteria (Bhatia et

al., 2018). However, it is a fact that, bacteria select PAH among mixed PAHs based on

their structural simplicity first (Dutta et al., 2017) and the velocity could be optimized

by reducing the initial concentration of the PAHs gradually according to the structural

simplicity (Jiang et al., 2018).

3.8.2. By immobilized Pseudomonas sp. *consortium*

The biodegradation kinetic of naphthalene by individual *Pseudomonas putida*

strain KD6, KD9, KD10 and *Pseudomonas* sp. consortium immobilized in CABs,

depicted further enhancement of overall naphthalene bio-utilization (Table 3). Cell

immobilization using hydrogel, such as CAB found to be advantageous rather than free

cells (Hameed and Ismail, 2018).

This phenomenon may be attributable due to the increased level of tightness of

the cross-linked polymers of the calcium alginate beads that render bacteria adequate

amount of protection from harsh environment (Chen et al., 2013). However, the free

bacterial cell lacks the capabilities to degrade a high initial concentration of the toxicant

because they followed the conventional growth phases (Marrot et al., 2006). In addition,

exposure of free bacterial cells to the high initial concentration of toxicants may

challenge them to experience shock-concentration (Zhao et al., 2006). Conversely, cells

immobilized in calcium alginate beads can tolerate high concentration of the toxicant

and decrease the lag phase duration (Kao et al., 2014). Moreover, the diffusion

limitation natures of the CABs matrix provide a high local concentration of the cell

population (Bezbaruah et al., 2009). Furthermore, CABs provides remarkable stability

and reusable features that effectively reduces the production cost (Daâssi et al., 2014).

Biodegradation of naphthalene by immobilized *Pseudomonas* sp. consortium

significantly elevated the overall naphthalene bio-utilization efficiency with $t_{1/2}$ and R^2

values of 2 days and 0.998 respectively, suggesting experimental data are well

correlated to the model (Table 3). The individual cell population also displayed

improved biodegradation efficiency. However, the biodegradation efficiency was

maximum by *Pseudomonas putida* strain KD6, which showed a marked reduction of $t_{1/2}$

from 4.1 to 3.0 days (Table 3). It promptly suggests that CABs facilitate KD6 optimized

naphthalene bio-utilization, that might be masked-up by the co-presence of NAP, PHN,

PYR (Jiang et al., 2018). However, further studies are required to investigate

behavioural patterns of a consortium with mixed PAH.

Bacterial cell grown on CSM with naphthalene as a sole source of carbon and

energy showed a delayed growth rate and their total biomass was also low (Figure 1).

Nevertheless, growing cells on CSM prior to immobilization in CABs provide them

essential adaptation to the toxicant and mature them for such stress condition (Table 3).

Immobilized cell system was useful for bioremediation of a toxicant after prior

adaptation to the surrounding environment (Partovinia and Rasekh, 2018). However,

growing cells in Luria-Bertani broth does not adapt cell adequately, and their

naphthalene removal performance was very poor (Figure S3). The inoculum was

normalized ($OD_{600nm} = 0.002$) for each *Pseudomonas putida* strain in order to prepare

the *Pseudomonas* sp. consortium. In aqueous medium, the immobilized bacterial cell

mainly on the surface was exposed to naphthalene and primarily involved in the

naphthalene bio-utilization process. However, due to the micro-porous feature of the

calcium alginate beads, microbial cells immobilized other than surface are also

participates in the bio-utilization and bio-sorption process of naphthalene. Further, with

time due to the mechanical force generated by shaking, a few microbial cells may

release from the calcium alginate beads. However, to fuel up the cell, it is necessary to

uptake naphthalene for bio-utilization through step by step intracellular enzymatic

reactions (Lin et al., 2014).

3.8.3. Effect of initial naphthalene concentration on biodegradation kinetics

The effect of different initial concentration of naphthalene ($150 - 2500$ mg L^{-1}) on degradation kinetics was evaluated with immobilized KD6, KD9, and KD10 as individual and as a consortium (Table S2). The results suggest cell immobilization in CABs facilitates bacteria to cope with a high initial concentration of NAP (Table S2). The substrate inhibition kinetic parameters, *viz.,* the maximum specific degradation rate (q_{max}), and inhibition constant (k_i) were highest for immobilized *Pseudomonas* sp. consortium and these values were 0.707 h^{-1} and 1475 mg L^{-1} respectively (Table S2). However, we did not find any significant change on half saturation constant (k_s), suggesting the reaction does not depend on its initial concentration and the biodegradation process follows pseudo-first order reaction kinetics. In previous study, free *Pseudomonas putida* KD9 in CSM was capable to tolerate relatively low initial concentration of naphthalene with k_i value of 1107 mg L^{-1} and addition of sucrose as SCS provides quite similar potential of naphthalene tolerance (k_i of 1429 mg L^{-1}) with that of immobilized *Pseudomonas* sp. consortium (Dutta et al., 2018). However, sucrose (0.5 gm. %) in the mobile open water system would not be beneficial for bacteria to overcome the high shock concentration of the toxicant, again suggesting cell immobilization and development of effective microbial consortium as systematic optimization of biodegradation process.

3.9. Detection of metabolic end products

The metabolic pathway of naphthalene biodegradation that might be followed by *Pseudomonas putida* strain KD10 was elucidated through GCMS analysis of the

metabolites (Figure S4). In the previous study, the major metabolites ware restricted to

be salicylaldehyde, catechol, D-gluconic acid and pyruvic acid (Dutta et al., 2018).

However, in the present study, we have detected salicylic acid as an additional

metabolite (Table 4). Catechol step into the TCA cycle by two possible pathway one is

ortho-cleavage and another is meta-cleavage pathway. The meta-cleavage pathway is

led by catechol-2, 3-dioxygenase (*nah*H), and presence of *nah*H (Figure S5) suggesting

in *Pseudomonas putida* stain KD10 opt the meta-cleavage pathway. Furthermore, we

assume that D-gluconic acid enters in the bio-conversion of naphthalene, possibly from

glucose as a precursor (Figure 6) and it is finally converted to pyruvate via aldolase. D-

gluconic acid implies impotence between bacteria-plant mutualistic association e.g.,

induction of phosphate solubilization processes (Rodriguez et al., 2004) which can

inhibit fungal growth (Kaur et al., 2006). The metabolic intermediates *viz.,* salicylic

acid, catechol has plant-growth promoting activity (Lee et al., 2010) e.g., the antioxidant

property of the catechol help and promotes seed germination (Schweigert et al., 2001).

The presence of d-gluconic acid as a major metabolite and its subsequent entry in TCA

cycle from glucose precursor, suggest that KD10 may able to sequestered catechol upon

needs to promote the plants' growth (Figure 6).

4. Conclusion

Naphthalene biodegradation by free and immobilized *Pseudomonas putida*

strain KD10 and *Pseudomonas* sp. consortium were studied. HPLC analysis showed

80.1% of initial naphthalene (500 mg L^{-1}) was utilized by immobilized *Pseudomonas*

sp. consortium after 72 h of incubation. Further, initial naphthalene tolerance by immobilized *Pseudomonas* sp. consortium was highest (1475 mg L^{-1}) with maximum specific degradation of 0.707 h^{-1}. The gene sequence analysis of the naphthalene 1, 2-dioxygenase suggests a significant evolutionary distance among different microbial species and in few cases, intra-species variation was observed. A common mutation prone amino acid stretch inside Chain A of all three natural mutant variants of naphthalene 1, 2-dioxygenase were found at close proximity of the active site. Further, the rigid-flexible molecular docking showed better binging free energy of the mutant variant encoded by *Pseudomonas putida* strain KD10 than that of wild-type variant of naphthalene 1,2-dioxygenase. This common mutation prone amino acid stretch could aid the rational approaches of enzyme redesigning. Overall, this study summarises the application of bacterial cell immobilization in calcium alginate beads and development of the microbial consortium together for enhanced naphthalene biodegradation. However, further studies are required for the systematic optimization of naphthalene biodegradation in a real environment.

Authors Contributions

KD and CG conceive the main hypothesis. KD design, performed all experiments, and wrote the manuscript. SS, IK, SB, DJ assisted KD in some experiments. MK, TM, PR, KCG performed the statistical analysis and wrote the manuscript. CG critically proofread and wrote the manuscript. All authors read the manuscript.

Funding

University Grant Commission (UGC) Govt. of India, New Delhi, India (Grant

No. VU/Innovative/Sc/17/2015).

Acknowledgement

University Grant Commission (UGC) Govt. of India, New Delhi, India (Grant

No. VU/Innovative/Sc/17/2015) is acknowledged by C.G. K.D. acknowledges Council

of Scientific and Industrial Research (CSIR), Govt. of India, New Delhi, India for

Senior Research Fellowship (SRF) sanction letter no. 09/599 (0082) 2K19 EMR-Z

dated: 29/03/2019.

Conflict of Interest

The authors declare that they have no conflict of interests.

References

Anwar, S., Liaquat, F., Khan, Q.M., Khalid, Z.M., and Iqbal, S. (2009). Biodegradation
of chlorpyrifos and its hydrolysis product 3, 5, 6-trichloro-2-pyridinol by
Bacillus pumilus strain C2A1. *Journal of Hazardous Materials* 168(1)**,** 400-405.

Bernstein, H.C., Paulson, S.D., and Carlson, R.P. (2012). Synthetic Escherichia coli
consortia engineered for syntrophy demonstrate enhanced biomass productivity.
Journal of biotechnology 157(1)**,** 159-166.

Bezbaruah, A.N., Krajangpan, S., Chisholm, B.J., Khan, E., and Bermudez, J.J.E.
(2009). Entrapment of iron nanoparticles in calcium alginate beads for
groundwater remediation applications. *Journal of Hazardous Materials* 166(2-
3)**,** 1339-1343.

Bhardwaj, T.R., Kanwar, M., Lal, R., and Gupta, A. (2000). Natural gums and modified
natural gums as sustained-release carriers. *Drug development and industrial*
pharmacy 26(10)**,** 1025-1038.

Bhatia, S.K., Bhatia, R.K., Choi, Y.-K., Kan, E., Kim, Y.-G., and Yang, Y.-H. (2018).
Biotechnological potential of microbial consortia and future perspectives.
Critical reviews in biotechnology 38(8)**,** 1209-1229.

Blanco, P., Hernando-Amado, S., Reales-Calderon, J., Corona, F., Lira, F., Alcalde-
Rico, M., et al. (2016). Bacterial multidrug efflux pumps: much more than
antibiotic resistance determinants. *Microorganisms* 4(1)**,** 14.

Bruce, R.M., Haber, L., and McClure, P. (1998). *Toxicological Review of Naphthalene*
(CAS No. 91-20-3): In Support of Summary Information on the Integrated Risk
Information System (IRIS). US Environmental Protection Agency, National
Center for Environmental Assessment.

Bugg, T., Foght, J.M., Pickard, M.A., and Gray, M.R. (2000). Uptake and active efflux
of polycyclic aromatic hydrocarbons by Pseudomonas fluorescens LP6a.
Applied and environmental microbiology 66(12)**,** 5387-5392.

Chen, D.-Z., Fang, J.-Y., Shao, Q., Ye, J.-X., Ouyang, D.-J., and Chen, J.-M. (2013).
Biodegradation of tetrahydrofuran by Pseudomonas oleovorans DT4
immobilized in calcium alginate beads impregnated with activated carbon fiber:
mass transfer effect and continuous treatment. *Bioresource technology* 139**,** 87-
93.

Chen, Q., Li, J., Liu, M., Sun, H., and Bao, M. (2017). Study on the biodegradation of
crude oil by free and immobilized bacterial consortium in marine environment.
PloS one 12(3)**,** e0174445.

Daâssi, D., Rodríguez-Couto, S., Nasri, M., and Mechichi, T. (2014). Biodegradation of
textile dyes by immobilized laccase from Coriolopsis gallica into Ca-alginate
beads. *International Biodeterioration & Biodegradation* 90**,** 71-78.

Dutta, K., Shityakov, S., Das, P.P., and Ghosh, C. (2017). Enhanced biodegradation of
mixed PAHs by mutated naphthalene 1,2-dioxygenase encoded by Pseudomonas
putida strain KD6 isolated from petroleum refinery waste. *3 Biotech* 7(6)**,** 365.
doi: 10.1007/s13205-017-0940-1.

Dutta, K., Shityakov, S., Khalifa, I., Mal, A., Moulik, S.P., Panda, A.K., et al. (2018).
Effects of secondary carbon supplement on biofilm-mediated biodegradation of
naphthalene by mutated naphthalene 1, 2-dioxygenase encoded by Pseudomonas
putida strain KD9. *Journal of Hazardous Materials* 357**,** 187-197. doi:
https://doi.org/10.1016/j.jhazmat.2018.05.024.

Ferraro, D.J., Okerlund, A.L., Mowers, J.C., and Ramaswamy, S. (2006). Structural
basis for regioselectivity and stereoselectivity of product formation by
naphthalene 1, 2-dioxygenase. *Journal of Bacteriology* 188(19)**,** 6986-6994.

Ghazali, F.M., Rahman, R.N.Z.A., Salleh, A.B., and Basri, M. (2004). Biodegradation
of hydrocarbons in soil by microbial consortium. *International Biodeterioration*
& Biodegradation 54(1)**,** 61-67.

Gilbert, E., Walker, A., and Keasling, J. (2003). A constructed microbial consortium for
biodegradation of the organophosphorus insecticide parathion. *Applied*
microbiology and biotechnology 61(1)**,** 77-81.

Hameed, B.B., and Ismail, Z.Z. (2018). Decolorization, biodegradation and
detoxification of reactive red azo dye using non-adapted immobilized mixed
cells. *Biochemical Engineering Journal* 137**,** 71-77.

James, L.C., and Tawfik, D.S. (2003). Conformational diversity and protein evolution–a
60-year-old hypothesis revisited. *Trends in biochemical sciences* 28(7)**,** 361-368.

Jiang, Y., Zhang, Z., and Zhang, X. (2018). Co-biodegradation of pyrene and other
PAHs by the bacterium Acinetobacter johnsonii. *Ecotoxicology and*
environmental safety 163**,** 465-470.

Kao, P.-M., Hsu, B.-M., Huang, K.-H., Tao, C.-W., Chang, C.-M., and Ji, W.-T. (2014).
Biohydrogen production by immobilized co-culture of Clostridium butyricum
and Rhodopseudomonas palustris. *Energy Procedia* 61**,** 834-837.

Kaur, R., Macleod, J., Foley, W., and Nayudu, M. (2006). Gluconic acid: An antifungal
agent produced by Pseudomonas species in biological control of take-all.
Phytochemistry 67(6)**,** 595-604.

Koshland Jr DE. (2002). The application and usefulness of the ratio kcat/KM.
Bioorganic chemistry. 30(3), 211-3. doi: https://doi.org/10.1006/bioo.2002.1246

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: molecular evolutionary
genetics analysis version 7.0 for bigger datasets. *Molecular biology and*
evolution 33(7)**,** 1870-1874.

Kumari, S., Regar, R.K., and Manickam, N. (2018). Improved polycyclic aromatic
hydrocarbon degradation in a crude oil by individual and a consortium of
bacteria. *Bioresource technology* 254**,** 174-179.

Kusumawardhani, H., Hosseini, R., and de Winde, J.H. (2018). Solvent tolerance in
bacteria: fulfilling the promise of the Biotech Era? *Trends in biotechnology*.

Lee, K.-Y., and Heo, T.-R. (2000). Survival of Bifidobacterium longumImmobilized in
calcium alginate beads in simulated gastric juices and bile salt solution. *Applied*
and environmental microbiology 66(2)**,** 869-873.

Lee, S., Kim, S.G., and Park, C.M. (2010). Salicylic acid promotes seed germination
under high salinity by modulating antioxidant activity in Arabidopsis. *New*
Phytologist 188(2)**,** 626-637.

Li, X., He, J., and Li, S. (2007). Isolation of a chlorpyrifos-degrading bacterium,
Sphingomonas sp. strain Dsp-2, and cloning of the mpd gene. *Research in*
Microbiology 158(2)**,** 143-149.

Lin, C., Gan, L., and Chen, Z.-L. (2010). Biodegradation of naphthalene by strain
Bacillus fusiformis (BFN). *Journal of Hazardous Materials* 182(1-3)**,** 771-777.

Lin, C., Gan, L., Chen, Z., Megharaj, M., and Naidu, R. (2014). Biodegradation of
naphthalene using a functional biomaterial based on immobilized Bacillus
fusiformis (BFN). *Biochemical Engineering Journal* 90**,** 1-7.

Marrot, B., Barrios-Martinez, A., Moulin, P., and Roche, N. (2006). Biodegradation of
high phenol concentration by activated sludge in an immersed membrane
bioreactor. *Biochemical Engineering Journal* 30(2)**,** 174-183.

Morris, G.M., Goodsell, D.S., Halliday, R.S., Huey, R., Hart, W.E., Belew, R.K., et al.
(1998). Automated docking using a Lamarckian genetic algorithm and an
empirical binding free energy function. *Journal of computational chemistry*
19(14)**,** 1639-1662.

Mrozik, A., and Piotrowska-Seget, Z. (2010). Bioaugmentation as a strategy for
cleaning up of soils contaminated with aromatic compounds. *Microbiological*
research 165(5)**,** 363-375.

Norris, R.D. (2017). "In-situ bioremediation of soils and ground water contaminated
with petroleum hydrocarbons," in *Handbook of Bioremediation (1993).* CRC
Press), 17-38.

Parales, R.E. (2003). The role of active-site residues in naphthalene dioxygenase.
Journal of Industrial Microbiology and Biotechnology 30(5)**,** 271-278.

Parales, R.E., Parales, J.V., and Gibson, D.T. (1999). Aspartate 205 in the catalytic
domain of naphthalene dioxygenase is essential for activity. *Journal of*
Bacteriology 181(6)**,** 1831-1837.

Partovinia, A., and Rasekh, B. (2018). Review of the immobilized microbial cell
systems for bioremediation of petroleum hydrocarbons polluted environments.
Critical Reviews in Environmental Science and Technology 48(1)**,** 1-38.

Raes, J., and Bork, P. (2008). Molecular eco-systems biology: towards an understanding
of community function. *Nature Reviews Microbiology* 6(9)**,** 693.

Rodriguez, H., Gonzalez, T., Goire, I., and Bashan, Y. (2004). Gluconic acid production
and phosphate solubilization by the plant growth-promoting bacterium
Azospirillum spp. *Naturwissenschaften* 91(11)**,** 552-555.

Schweigert, N., Zehnder, A.J., and Eggen, R.I. (2001). Chemical properties of catechols
and their molecular modes of toxic action in cells, from microorganisms to
mammals: minireview. *Environmental microbiology* 3(2)**,** 81-91.

Schweizer, H.P. (2003). Efflux as a mechanism of resistance to antimicrobials in
Pseudomonas aeruginosa and related bacteria: unanswered questions. *Genet Mol*
Res 2(1)**,** 48-62.

Singh, N.S., Sharma, R., and Singh, D.K. (2019). Identification of enzyme (s) capable
of degrading endosulfan and endosulfan sulfate using in silico techniques.
Enzyme and Microbial Technology 124**,** 32-40.

Smid, E.J., and Lacroix, C. (2013). Microbe–microbe interactions in mixed culture food
fermentations. *Current opinion in biotechnology* 24(2)**,** 148-154.

Su, X., Wellen, K.E., and Rabinowitz, J.D. (2016). Metabolic control of methylation
and acetylation. *Current opinion in chemical biology* 30**,** 52-60.

Surkatti, R., and El-Naas, M. (2018). Competitive interference during the
biodegradation of cresols. *International Journal of Environmental Science and*
Technology 15(2)**,** 301-308.

Team, R.C. (2013). R: A language and environment for statistical computing.

Thomé, A., Reginatto, C., Vanzetto, G., and Braun, A.B. (Year). "Remediation
Technologies Applied in Polluted Soils: New Perspectives in This Field", in:
The International Congress on Environmental Geotechnics: Springer), 186-203.

Tyagi, M., da Fonseca, M.M.R., and de Carvalho, C.C. (2011). Bioaugmentation and
biostimulation strategies to improve the effectiveness of bioremediation
processes. *Biodegradation* 22(2)**,** 231-241.

Usha, M., Sanjay, M., Gaddad, S., and Shivannavar, C. (2010). Degradation of h-acid
by free and immobilized cells of Alcaligenes latus. *Brazilian Journal of*
Microbiology 41(4)**,** 931-945.

Wang, L., Zhang, S., Wang, L., Zhang, W., Shi, X., Lu, X., et al. (2018). Concentration
and risk evaluation of polycyclic aromatic hydrocarbons in urban soil in the
typical semi-arid city of Xi'an in Northwest China. *International journal of*
environmental research and public health 15(4)**,** 607.

Wilson, S.C., and Jones, K.C. (1993). Bioremediation of soil contaminated with
polynuclear aromatic hydrocarbons (PAHs): a review. *Environmental pollution*
81(3)**,** 229-249.

Wongwilaiwalin, S., Rattanachomsri, U., Laothanachareon, T., Eurwilaichitr, L.,
Igarashi, Y., and Champreda, V. (2010). Analysis of a thermophilic
lignocellulose degrading microbial consortium and multi-species
lignocellulolytic enzyme system. *Enzyme and Microbial Technology* 47(6)**,** 283-
290.

Woyke, T., Teeling, H., Ivanova, N.N., Huntemann, M., Richter, M., Gloeckner, F.O.,
et al. (2006). Symbiosis insights through metagenomic analysis of a microbial
consortium. *Nature* 443(7114)**,** 950.

Young, C.C., Rekha, P., Lai, W.A., and Arun, A. (2006). Encapsulation of plant
growth‐promoting bacteria in alginate beads enriched with humic acid.
Biotechnology and bioengineering 95(1)**,** 76-83.

Zhao, X., Wang, Y., Ye, Z., Borthwick, A.G., and Ni, J. (2006). Oil field wastewater
treatment in biological aerated filter by immobilized microorganisms. *Process*
Biochemistry 41(7)**,** 1475-1483.

Figure 1. Growth kinetic model of *Pseudomonas putida* and *Pseudomonas* sp.
consortium during naphthalene biodegradation. **A.** Gompertz's growth kinetic
model fit of the biodegradation of naphthalene by *Pseudomonas putida* strain
KD10 and by the *Pseudomonas* sp. consortium. **B.** Effect of 0.5 gm. % sucrose
supplementation on growth pattern of *Pseudomonas putida* strain KD10 and
Pseudomonas sp. consortium.

Figure 2. Cell immobilization in calcium alginate beads (CABs) with different
cell morphology of the *Pseudomonas putida* strain KD10.
783

Figure 3. Sequence and structure analysis of the naphthalene 1, 2-dioxygenase
(*nah*Ac) encoded by *Pseudomonas putida*. **A**. The amino acid sequence of wild
type and three mutant variants of the naphthalene 1, 2-dioxygenase (*nah*Ac).
Mutated residues are highlighted in blue and the common mutation prone amino
acid stretch is highlighted with red box. **B.** Structural mismatches among mutant
variants and wild type naphthalene 1, 2-dioxygenase. Each mutated residues and
its subsequent neighbour residues of all the mutant variant of naphthalene 1, 2-
dioxygenase is overlaid with cartoon representation. The highlighted red circle
indicates local mismatch among the mutant variant of naphthalene 1, 2-
dioxygenase **C.** Different ligand binding posture of the naphthalene 1, 2-
dioxygenase encoded by *Pseudomonas putida* strain KD10. Naphthalene (red),
phenanthrene (cyan) and anthracene (green) binding postures with major
interacting amino acid residues (left). Two-dimensional presentation of the
interacting residues of *nah*Ac$_{I250, V256}$ (right). **D.** Docking pose of naphthalene 1,
2-dioxygenase (Chain A) encoded by *Pseudomonas putida* strain KD10. The
four major naphthalene interacting residues and their molecular distance are
labelled in black.

Figure 4. Distance matrix and cladogram of the naphthalene 1, 2-dioxygenase.
The evolutionary distance of the naphthalene 1, 2-dioxygenase I_{250}, V_{255} variant
encoded by *Pseudomonas putida* strain KD10 other variants by different
microbial species is scaled by colour index (upper-left side).

807
808

Figure 5. Naphthalene degradation kinetics of *Pseudomonas putida* strain KD10
and *Pseudomonas* sp. consortium free and cell immobilized in calcium alginate
beads.

Figure 6. Proposed pathway of naphthalene biodegradation by *Pseudomonas*
putida strain KD10. Naphthalene (I), salicylic acid (II), salicylaldehyde (III),
catechol (IV), pyruvate (V), glucose (VI), d-gluconic acid (VII). The grey arrow
indicates alternative pathway via d-gluconic acid.

817
818

819   *Table captions*

820   **Table 1.** Summary of rigid-flexible molecular docking for the analysed mutant
821   naphthalene 1, 2-dioxygenase encoded by different *Pseudomonas putida*
822   strains.

824   **Table 2.** Apparent enzyme kinetic parameters of naphthalene 1, 2-dioxygenase
825           $_{I250, V256}$ encoded by *Pseudomonas putida* strain KD10.

827   **Table 3.** Parameters of first-order biodegradation kinetics of naphthalene by
828   different free and immobilized *Pseudomonas putida* strains.

830   **Table 4.** Major metabolite detected by GCMS analysis during naphthalene
831   biodegradation by *Pseudomonas putida* KD10.

Supplementary section

Table S1. Nucleotide sequences used as primer in polymerase chain reactions

Table S2. Parameters of substrate inhibition kinetic model of naphthalene
biodegradation by different *Pseudomonas putida* strains immobilized in calcium
alginate beads.
Table S3. Enumeration of viable cell in calcium alginate beads.

Table S4. Gompertz's growth curve model fit of *Pseudomonas putida* strain
KD10 and *Pseudomonas* sp. consortium.

Table S5. Major amino acid residues of mutant variant of naphthalene 1, 2-
dioxygenase $_{I250,\ V256}$ involve in hydrophobic interaction with different ligands.

Figure S1. Colony morphology and catechol confirmatory test. **A.** Colony
morphology of *Pseudomonas putida* strain KD10 after 24 h of incubation. **B.**
Colony morphology of *Pseudomonas putida* strain KD10 after one week of
incubation.
Figure S2. Molecular phylogenetic analysis of *Pseudomonas putida* strain

KD10.

Figure S3. Effect of cell adaptation on naphthalene biodegradation.

Figure S4. Mass spectra of the major metabolites detected during
biodegradation of naphthalene in CSM as sole source of carbon and energy by
Pseudomonas putida KD10.

Figure S5. Agarose gel electrophoresis of PCR products of catechol 2, 3-
dioxygenase (*nah*H) gene and solvent efflux pump (*srp*ABC) system.
A. Catechol 2, 3-dioxygenase (*nah*H). Lane 1 and 2 *nah*H pcr product of
Pseudomonas putida strain KD9 and *Pseudomonas putida* strain KD10. **B.**
Solvent efflux pump (*srp*ABC) system. Lane 1: *Pseudomonas aeruginoasa*
ATCC 9027, Lane 2: *Pseudomonas putida* strain KD10. Lane M: 100bp DNA
ladder (HiMedia, India), Lane 3: Negative control.

Figure S6. Nucleotide sequence chromatogram of the mutant naphthalene 1, 2-
dioxygenase (*nah*Ac), (chain-A) encoded by *Pseudomonas putida* strain KD10.
870

Table 1. Summary of rigid-flexible molecular docking of naphthalene and different mutant version of naphthalene 1, 2-dioxygenase encoded by different *Pseudomonas putida* strains

ΔG*			LB[4]	IC[5]	IME[6]	VDW[7]	RMSD[c]	RMSD[r]	References
BE[1]	London[2]	GBVI/WSA[3]							
-5.92[a]	-7.18	-4.61	-0.59	46.14	-5.92	-5.91	0.00	77.29	Dutta et al., 2017
-5.41[b]	-7.13	-4.02	-0.54	109.07	-5.41	-5.4	0.00	96.32	Dutta et al., 2019
-5.83[c]	-7.10	-1.68	-0.58	53.13	-5.83	-5.84	0.00	96.08	This study

*PubChem ligand CID 931 = Naphthalene, BE[1] = Binding free energy, [2]London free energy, [3]Generalized born volume integral/weighted surface areas, LB[4] = ligand binding, IC[5] = inhibition constant, IME[6] = Intermolecular energy, VDW[7] = Vdw hb desolv energy, RMSD[c] = Cluster RMSD, RMSD[r] = Reference RMSD. Energy unit = kcal mol[-1]. Inhibition constant unit = µM.at 298.15 K. Naphthalene 1, 2-dioxygenase encoded by (a) *Pseudomonas putida* strain KD6, (b) *Pseudomonas putida* strain KD9 and (c) *Pseudomonas putida* strain KD10.

Table 2. Apparent enzyme kinetic parameters of naphthalene 1, 2-dioxygenase $_{I250, V256}$ encoded by *Pseudomonas putida* strain KD10

Substrate	Kinetic parameters*				
	Km (μmol mL^{-1})	Vmax (μmol min^{-1})	Kcat (s^{-1})	Kcat/Km (mL^{-1} mol^{-1} s^{-1})	R^{2**}
Naphthalene	0.867	2.474	123.72	0.142×10^3	0.938

*The kinetic constants were determined at 30°C and pH 7.5 using 0.5-10μM ml^{-1} substrate concentration (each) by Lineweaver–Burk plot.

** Non-liner regression between initial substrate concentration 1/[S] and degradation rate constant 1/V yielded regression equation and regression coefficient (R^2)

Table 3. Parameters of first-order biodegradation kinetics of naphthalene by different free and immobilized *Pseudomonas putida* strains

| CSM | Cells immobilized in CABs | | | free cells | | | |
	k (Days^{-1})	t ½ (Days^{-1})	R^2	k (Days^{-1})	t ½ (Days^{-1})	R^2	References
A	0.228±0.005	3.0±0.077	0.99	0.167±0.012	4.1±0.307	0.98	Dutta et al., 2017
B	0.312±0.014	2.2±0.102	0.93	0.252±0.019	2.7±0.193	0.99	Dutta et al., 2018
C	0.284±0.017	2.4±0.143	0.97	0.219±0.009	3.1±0.137	0.92	This study
D	0.341±0.294	2.0±0.186	0.99	0.309±0.020	2.2±0.141	0.98	This study

[#]Initial naphthalene concentration = 500 mg L^{-1}
A. *Pseudomonas putida* strain KD6, B. *Pseudomonas putida* strain KD9, C. *Pseudomonas putida* strain KD10. D. *Pseudomonas* sp. consortium.
Each figure of the table represents the mean of three replicates.

Table 4. Major metabolites detected by GCMS analysis during naphthalene biodegradation by *Pseudomonas putida* KD10

Metabolite	R_t (min)	Major ion peaks (m/z)	Suggested structure
I	9.41	64.19, 92.17, 120.21, 138.22, 169.14, 191.18, 211.10	Salicylic acid
II	15.27	65.17, 74.41, 104.73, 121.10, 122.57, 142.21, 174.80, 201.22, 249.31	Salicylaldehyde
III	18.71	64.29, 81.35, 91.61, 110.47, 131.47, 149.71	Catechol
IV	31.89	61.21, 73.44, 76.24, 104.18, 117.41, 133.27, 177.23	D-gluconic acid
V	10.47	43.21, 61.27, 71.37	Pyruvic acid

R_t = Retention time

Figure 1.

978   **Figure 2.**

979 **Figure 3.**

980
981

982
983 **Figure 4.**

996
997
998
999

1000
1001
1002
1003 **Figure 5.**

1017   **Figure 6.**