A bidirectional network for appetite control in zebrafish

*Caroline Lei Wee[1,2, 5], *Erin Yue Song[1], Robert Evan Johnson[1,2], Deepak Ailani[3], Owen

Randlett[1], Jiyoon Kim[1], Maxim Nikitchenko[1], Armin Bahl[1], Misha Ahrens[4], Koichi Kawakami[3],

Florian Engert[1] and **Samuel Kunes[1]**

*Equal contribution first authors

[1]Department of Molecular and Cell Biology, Harvard University, Cambridge, MA, USA

[2]Program in Neuroscience, Harvard University, Boston, MA, USA

[3]Division of Molecular and Developmental Biology, National Institute of Genetics, Department of

Genetics, SOKENDAI (The Graduate University for Advanced Studies), Mishima, Shizuoka,

Japan

[4]Howard Hughes Medical Institute, Janelia Farm Research Campus, Ashburn, Virginia, USA.

[5]Institute of Molecular and Cell Biology, A*STAR, Singapore

ABSTRACT

Medial and lateral hypothalamic loci are known to suppress and enhance appetite, respectively,

but their interactions and dynamics have not yet been explored. Here we report that, in

zebrafish, serotonergic neurons of the ventromedial caudal hypothalamus (cH) becomes

increasingly active during food deprivation, whereas activity in the lateral hypothalamus (LH) is

reduced. Exposure to food sensory and consummatory cues reverses the activity states of

these two nuclei, reflecting an opposing internal hunger state induced by food. An intermediate

activity state is restored as satiety approaches. The overall antagonistic relationship of cH and

LH was confirmed by simultaneous calcium imaging, and a causal relationship was established

by targeted stimulation and ablation of the cH. The collective data allows us to propose a model

in which activities in anti-correlated hypothalamic nuclei direct distinct phases of hunger, and

thus coordinate energy balance via mutually antagonistic control of distinct behavioral outputs.

INTRODUCTION

The regulated intake of food based on caloric needs is a fundamental homeostatically controlled process that is essential for health and survival. The hypothalamus is a highly conserved central convergence point for the regulation of the neural and biochemical pathways underlying these basic mechanisms. Early studies demonstrated by way of electrical stimulation or lesions that specific hypothalamic regions play important roles in the regulation of appetite. For example, while stimulation of ventromedial hypothalamic loci in rodents and cats reduced feeding, activation of more lateral hypothalamic loci increased both hunting behavior and food intake (ANAND and BROBECK, 1951; BROBECK et al., 1956; DELGADO and ANAND, 1953; Krasne, 1962). Conversely, lateral hypothalamic lesions were found to reduce feeding to the point of starvation, whereas medial hypothalamic lesions resulted in overeating(ANAND and BROBECK, 1951; Hoebel, 1965; TEITELBAUM and EPSTEIN, 1962). Thus, the lateral and medial hypothalamic regions came to be regarded as "hunger" and "satiety" centers, respectively.

Recent experiments employing optical and electrophysiological methods have lent support to expectations based on these early studies. For example, GABAergic neurons in the lateral hypothalamus were observed to be activated during feeding and essential for enhanced food intake during hunger (Jennings et al., 2015; Stuber and Wise, 2016). However, these experiments have examined only subsets of hypothalamic neurons; their activity patterns in the context of the entire network remain unknown. This limited view hampers our understanding of the dynamical interactions between the ensemble of brain circuits thought to be important for the initiation, maintenance and termination of food consumption (Sternson and Eiselt, 2017).

Here, we leverage the small and optically accessible larval zebrafish to identify modulatory regions central to the control of appetite and to shed light on their specific roles and dynamical activity patterns in relation to the whole brain and behavior. Using pERK-based brain-wide activity mapping we first identified neuronal populations that display differential neural activity under conditions of hunger and satiety. We show that lateral and medial hypothalamic

regions have anti-correlated activity patterns during hunger, voracious feeding and satiety. Next,

through a combination of calcium imaging, optogenetics and ablation analysis, we show that

serotonergic neurons in the caudal periventricular zone of the medial hypothalamus (cH) are

state-dependent regulators of feeding behavior, likely via their modulation of the lateral

hypothalamus. This allows us to propose a model where mutually antagonistic brain states

regulate energy balance by encoding distinct signals in different facets of appetite control.

RESULTS

Whole brain activity mapping of appetite-regulating regions

Larval zebrafish hunt their prey, paramecia, through a sequence of motor actions that has been

considered a hardwired reflex response to external prey stimuli (Bianco et al., 2011;

Semmelhack et al., 2015; Trivedi and Bollmann, 2013). Only recently has evidence emerged

that this behavior is flexibly modulated by satiation state (Filosa et al., 2016; Jordi et al., 2015,

2018) and that larvae at 7 **d**ays **p**ost-**f**ertilization (dpf) display enhanced hunting and enhanced

food intake after a period of food deprivation. A robust readout of food intake in larval zebrafish

was obtained both by fluorescently-labeled paramecia and behavioral analysis approaches that

have been adapted for this study (Jordi et al., 2015, 2018; Shimada et al., 2012, Johnson et al.,

in preparation, Figure 1a). Given that a 2-hour period of food-deprivation is sufficient to robustly

enhance subsequent food intake, fish at the end of this food-deprivation period are considered

to be in a state of "hunger" and nutrient/caloric deficit. Indeed, up to 15 min after food-

presentation, such food-deprived animals display a strong upregulation of hunting and food

intake relative to fish with continuous access to food (fed fish). As the fish in this state are likely

still in a caloric/nutrient deficit and display enhanced food intake, we will refer to this phase as

"voracious feeding". Finally, as the fish continue to consume food, their rate of food intake

declines to a low level comparable to that of fed fish. As we describe below, this state of

continuous low level feeding in the presence of ample food sources reflects a state of "satiety"

that is the same or similar to that of a continuously fed animal.

As a first step toward understanding the homeostatic control of feeding in this simple

vertebrate system, we employed whole-brain neuronal activity mapping via phosphorylated ERK

visualization in post-fixed animals (MAP-mapping; Randlett et al., 2015). Whole brain confocal

image datasets of phospho-ERK expression were gathered from animals sacrificed after 15

minutes of voracious feeding that followed a 2-hour period of food deprivation. For comparison,

image sets were also gathered from animals that had been fed constantly. These image

volumes were registered to a standardized brain atlas and are displayed as a difference map

(Figure 1b), revealing significant differences in neural activity when comparing voracious

feeding with constant feeding (Figure 1b-d, Video 1, Supplementary Tables 1-2). Since both

experimental groups experienced similar sensory stimuli (i.e. exposure to the same

concentration of paramecia), differences in brain activity should reflect the animal's internal

hunger state, which could also manifest as an enhanced sensitivity to food cues, and/or

enhanced hunting and prey capture. Indeed, multiple sensorimotor loci related to hunting

showed enhanced activity in the food-deprived state. These included stronger activation in

retinal Arborization Fields (AFs; optic tectum and AF7), pretectum, as well as downstream

hindbrain loci, such as reticulospinal and oculomotor neurons that all have been shown to be

engaged during prey capture behavior (Bianco and Engert, 2015; Muto et al., 2017;

Semmelhack et al., 2015). In addition, enhanced activity was observed in the cerebellum,

inferior olive, vagal sensory and motor neurons, area postrema and locus coeruleus, all of which

have been implicated in producing motor programs related to feeding behavior (Ahima and

Antwi, 2008; Ammar et al., 2001; Dockray, 2009; Zhu and Wang, 2008).

We next focused our attention on brain areas likely to be involved in regulating internal

states related to hunger and satiety. These included an area of particularly strong differential

activity in the lateral region of the intermediate hypothalamus (LH; Fig. 1b-d), which has recently

been identified as part of the feeding pathway in larval zebrafish (Muto et al., 2017), and whose

mammalian analog has been strongly implicated in appetite control (Sternson and Eiselt, 2017).

However, the zebrafish LH, unlike its mammalian counterpart, does not express melanin-

concentrating hormone (MCH) or contain orexin (hypocretin)-positive neurons, nor does it

clearly express other major feeding-related peptides (Figure 1- Figure Supplement 1 and 2).

MCH, hypocretin and other appetite-related neuromodulators (AgRP, MSH, CART, NPY) are in

fact expressed in other nearby areas of the hypothalamus (Figure 1 - Figure Supplement 1).

The zebrafish LH region does however contain a variety of glutamatergic and GABAergic cell

types (Figure 1 - Figure Supplement 2) that have been shown to be important for regulation of

feeding in mammals, independent of MCH and orexin (Jennings et al., 2015; Stuber and Wise,

2016). Among areas that showed relatively decreased neural activity upon feeding food-

deprived animals, the most significant was the caudal hypothalamus (cH), which contains

monoaminergic (mainly serotonergic and dopaminergic) neurons (Fig 1c; Kaslin and Panula,

2001; Lillesaar, 2011). Indeed, in all of nine independent MAP-mapping experiments, activity

was reduced in the cH and increased in the LH within 15 min of food presentation (Fig 1c). The

evident inverse relationship between overall LH and cH activity in this context was supported by

independent component analysis (Randlett et al., 2015), which was applied to all feeding-related

MAP-mapping data, and uncovered multiple components where the cH and LH are strongly

anti-correlated (Figure 1e, Figure 1 - Figure Supplement 3). These results led us to hypothesize

that the lateral and caudal hypothalamus may form a functionally interconnected network with

opposing activity patterns.

125
126
Figure 1 with 4 supplements: Whole brain activity mapping reveals anti-correlated hypothalamic
clusters

(a) Top: Schematic of protocols used to evaluate appetite behavior in larval zebrafish. At 7 or 8 dpf,
larvae were either food-deprived for 2 hours, or fed with excess paramecia for this duration. After 2 hrs (2-
4 hours in the case of behavioral imaging), they were subject to a quick wash, followed either by: 1)
addition of excess fluorescent-labeled paramecia (left), 2) high-resolution behavioral imaging (right).
Bottom left: Gut fluorescence measurements of food-deprived (red) or fed (blue) fish as a function of
feeding duration. Groups of fed or food-deprived larvae were fixed at indicated time points after feeding of
labeled paramecia (fed: n=7/18/19/17/17, food-deprived: n= 8/23/20/14/15). Food-deprived fish had
significantly higher gut fluorescence than fed fish overall (p = 7.5859×10^{-10}, Two-way ANOVA, asterisk
indicates corrected p-values<0.05. **Bottom right:** The probability of performing a hunting-related swim
bout across fed and food-deprived fish groups in 3-minute time bins over 45 minutes. Error bars represent
90% confidence intervals. For all bins except those indicated with triangles, the null hypothesis that initial
feeding condition has no effect on hunting-bout probability is rejected (p < 0.00001, Fisher's Exact Test
comparing binomial probability distributions per bin). Fed: n =85655 bouts from 73 fish; Food-deprived n =
75357 bouts from 57 fish. Since the rate of food intake and hunting behavior was highest in the first 15
minutes (voracious feeding phase, gray boxes), we chose this time point for subsequent MAP-mapping
experiments.
(b) Brain-wide activity mapping of food-deprived (Dep.) and fed fish, in response to food. Data from 9
experiments (n = 557 fish total) were combined to generate this map. Activated regions include the
telencephalon (Tel), Arborization field 7 (AF7), cerebellum (CB), hindbrain (HB), Vagal ganglion (VG) and
lateral lobe of the intermediate hypothalamus (LH). Suppression was observed in the caudal
hypothalamus (cH) and some parts of the telencephalon. Scale bar = 100 µm. Also see Video 1.
(c) ROI-specific analysis of LH and cH regions in 9 independent MAP-mapping experiments (20-30 fish
per treatment per experiment). Food-deprived fish constantly had higher LH and lower cH activity in
response to food (p=0.0039 for both cH and LH, Wilcoxon Signed Rank Test).
(d) Z-projection of same MAP-map showing the hypothalamus, where lateral regions (i.e. LH) are strongly
activated and medial regions (e.g. cH) are suppressed. The map is overlaid on an anatomy stack for the
transgenic line *Tg(etVMAT:GFP)* to highlight the location of cH neurons.
(e) Six examples of independent component analysis (ICA) maps. Voxels for each recovered independent
component (IC) are shown as maximum projections, with intensity proportional to the z-score of the
loadings of the ICA signal. These ICs, along with others (16/30) highlight LH and cH regions of opposite
loadings, suggesting they may be part of a network with anti-correlated activity patterns. Positive (+)
loading and Negative (-ve) loadings are reflected in green and magenta respectively.
(f) Higher resolution imaging of dissected brains stained with pERK during phases of feeding. Scale bar:
50 µm. Inset: Higher magnification view of cH neurons. Scale bar: Scale bar = 20 µm. Fish were mounted
ventral side up.
(g) Quantification of cH activity (normalized pERK fluorescence) and LH (medial LH and lateral LH)
activity (normalized pERK fluorescence) (top) and # pERK-positive cells (bottom) in fed and food-
deprived fish (n =13/11/9/9/13/12).
Normalized pERK intensity (cH/mLH/lLH): Fed vs Dep. (p = 0.016/0.17/0.17), Dep. vs Dep + 5 min food
(p=3.1×10^{-4}/9.9×10^{-5}/0.02), Fed vs Dep. + 5 min food (p= 0.0097/0.001/0.08).
Active Cell count (mLH/lLH): Fed vs Dep. (p = 0.001/0.0038), Dep. vs Dep + 5 min food (p= 9.7×10^{-5}
/1.3×10^{-5}), Fed vs Dep. + 5 min food (p= 0.0038/0.048). Asterisks denote p<0.05, one-tail Wilcoxon
Rank Sum Test. Note that mean pERK intensity does not change as significantly as active cell count.

172
173
**Figure 1 - Figure Supplement 1: Anatomical characterization of intermediate hypothalamus
feeding areas**
(a) Expression patterns of a number of feeding-related peptides in the zebrafish hypothalamus, based on
antibody-staining or transgenic labels. HCRT = hypocretin (orexin), CART = cocaine and amphetamine
related transcript MCH = melanin-concentrating hormone, TH = tyrosine hydroxylase (labels
dopaminergic and/or noradrenergic neurons), MSH = alpha-melanocyte stimulating hormone, AgRP =
Agouti-related peptide, NPY = neuropeptide Y, VMAT = vesicular monoamine transporter (labels
dopaminergic (DA) and serotonergic neurons (5-HT)).   Note that MCH and HCRT staining is absent from
the zebrafish LH. Though not apparent from the schematic, HCRT is located more dorsally. The preoptic
area, which contains oxytocinergic as well as other peptidergic neurons, is located more dorsally and not
reflected in this schematic.
(b) Schematic summarizing zebrafish hypothalamic peptide expression. GABA (dark blue) and
glutamatergic (blue) neurons are found in the lateral hypothalamus (see Figure 1- Figure Supplement 2)
and also throughout the medial regions of the hypothalamus. PVO = paraventricular organ, which also
contains DA and 5-HT neurons. A number of peptidergic neurons are located within the anterior and
posterior pituitary/hypophysis (aPit and pPit). Color code corresponds to images in (a).

191
Figure 1- Figure Supplement 2: Characterization of zebrafish LH

(a) Glutamatergic and GABAergic neuron distribution in the hypothalamus. *Tg(VGlut2a:dsRed)* and *Tg(GAD1b:GFP)* transgenic fish were dissected, imaged and registered onto a common reference hypothalamus.

(b) Glutamatergic cells, labeled by *Tg(VGlut2a:dsRed)*, overlap with active (pERK-positive) neurons in both the ILH and outer rim of the mLH. (i) Overview of hypothalamus. (ii) Higher magnification images of LH. (iii-iv) Inset showing overlap of ILH and outer rim of mLH with glutamatergic cells.

(c) GABAergic cells, labeled by *Tg(Gad1b:GFP)*, overlap with active neurons in the inner rim of the mLH but not the ILH. (i) Overview of hypothalamus. (ii) Higher magnification images of LH. (iii-iv) Inset showing overlap of inner rim of mLH with GABAergic cells. White arrows point to examples of overlapping cells. All fish were mounted ventral side up. Scale bar = 50 μm. Inset scale bar = 20 μm.

Figure 1- Figure Supplement 3: All 30 independent components extracted from ICA analysis. This method separates pERK signals into statistically independent components based on their correlated activity, thus identifying putative functional connectivity (both positive or negative relationships) between different brain regions (Randlett et al., 2015; see Methods). To increase the robustness of the analysis, we included fish from other feeding-related treatments that we did not otherwise use in this manuscript (n = 904 fish total).

(a-c) From this analysis, we identified multiple independent component networks (ICs) in which at least part of the LH displayed an inverse activity relationship (i.e. opposite loadings) with the cH (16/30).

(d) 7/30 ICs had correlated LH and cH activity.

(e) The other 7/30 had asymmetrical or noisy activity patterns.

Data compilation (Custom MATLAB scripts)

Figure 1 – Figure supplement 4: Automated quantification of pERK-positive cells

(a) Method by which we quantify pERK-positive ("active") cell count in a high-throughput manner. This method works best with high-resolution images (i.e. dissected brains). Brain stacks are registered onto a reference brain within the same dataset, using the tERK channel, though there is the option of using unregistered images (for which individual ROIs have to be defined for each image). A series of processing steps allows for automated segmentation of pERK-positive cells using the same manually optimized threshold across the entire dataset.

Cellular dissection of hypothalamus neural activity reveals modulation by satiation state

To probe neural activity changes with higher resolution, we performed pERK staining in

dissected brains, and examined the activity of these populations in time course experiments that

spanned the period of food-deprivation and subsequent feeding (Figure 1f-g, Figure 2). We

quantified changes in mean cellular fluorescence as well as changes in the number of active

cells or cell clusters (Figure 1 - Figure Supplement 4). While the high density of labeled cells

and high background fluorescence in the cH made the identification of individual neurons

difficult, we found that in the LH segmentation of individual neurons and classifying activity

based on the thresholded fluorescence levels provided a cleaner and more reliable readout for

overall neuronal activity. Using these respective metrics, we observed that mean fluorescence

in the cH was predictably high in food-deprived fish, while the number of active neurons in the

medial and lateral lobes of the LH (mLH and lLH, respectively) was low when compared to

continuously fed animals. However, within 5 minutes of food presentation, cH activity fell

dramatically to a level significantly below that observed in continuously fed fish (Figure 1f). This

characteristically low cH activity level was accompanied by a large increase in LH mean

fluorescence and neuron activity, which is consistent with our MAP-mapping results. As the

feeding period continued, LH neuronal activity declined and, reciprocally, cH activity increased,

coincident with the decline in voracious hunting and food ingestion (Figure 1f). After two hours,

neural activity in the cH and LH and feeding behavior all converged onto baseline levels similar

to those observed in continuously fed fish (Figure 1f). Thus these two neuronal populations

display anti-correlated activity patterns over time frames that span the progression of hunger

during food-deprivation, voracious feeding and the gradual return to satiety.

Satiation state influences the sensitivity of cH and LH populations to food

The neural activity patterns described above suggest that cH and LH activity may report the

satiation state of the animal. To better align these patterns with the animal's internal state, we

examined these loci over a time course that started with food removal from well-fed animals,

followed by food presentation after a variable food-deprivation period (30 min or 2 hours, Figure

2). We found that food removal resulted in a reduction of the number of active mLH and lLH

neurons within 30 minutes. In contrast, cH activity gradually increased, with a significantly

higher level of activity occurring between 30 min and 2 hours post-food removal. A possible

interpretation of these patterns is that LH activity is directly driven by the presence of food cues

(as noted by Muto et al 2017), whereas the level of activity observed in cH neurons is a

correlate of the animal's nutrient/caloric deficit and resulting hunger state in the absence of food,

possibly even generating the signal necessary to sensitize the LH's responsiveness to such

stimuli.

Food presentation not only rapidly reverses the activity patterns of both loci, but also

does so in a manner correlating with the length of food-deprivation. Indeed, fish that had been

food-deprived for longer periods (2-4 hrs) displayed a relatively enhanced induction of LH

activity upon the introduction of food. Likewise, the reduction in cH activity on food presentation

was significantly more pronounced when it followed a longer period of prior deprivation; both of

these neural responses were strongly correlated with enhanced food consumption (voracious

feeding; Figure 2, Figure 2 – Figure Supplement 1). As prey continues to be consumed, activity

in both loci gradually reverts back to the baseline levels representative for fed animals (Figure

1f-g, 2d).

Figure 2 with 5 supplements: cH and LH activities are modulated by food and satiation state

(a) Representative images showing how cH and LH activity in the presence and absence of food vary with the extent of food-deprivation, from the same dataset quantified below.

(b) Mean pERK fluorescence in the cH increases with food-deprivation, whereas mean LH pERK fluorescence does not change significantly, except during voracious feeding (Dep. 2hr + food). Normalized pERK intensity (cH/mLH/lLH): Fed vs Dep. 2hr (p = 0.0022/0.41/0.59), Fed vs Dep. 2hr + food (0.047/0.0011/0.0011), Dep. 30 min + food vs Dep. 2hr + food (p = 0.041/0.0022/0.0022), Dep. 2hr vs Dep. 2hr + food (p = 0.0022/0.0011/0.0022).

(c) The number of active LH (particularly lLH) cells decline within 30 min of food deprivation, and is significantly enhanced during feeding, particularly after a longer period of food-deprivation. Active cell count (mLH/lLH): Fed vs Dep. 30 min (p = 0.155/5.8x10^{-4}), Fed vs Dep. 2hr (p = 0.047/0.011), Dep. 30 min + food vs Dep. 2hr + food (p = 0.0022/0.0043), Dep. 30 min vs Dep. 30 min + food (p = 0.07/0.013), Dep. 2hr vs Dep. 2hr + food (p = 0.0011/0.0011), Fed vs Dep. 2hr + food (p = 0.0022/0.07), n = 6/7/5/6/6 fish, One-tail Wilcoxon Rank Sum Test.

(d) Schematic of inferred cH and LH activity over phases of feeding. LH active cell count appears to decline more rapidly than the rise in cH activity. More supporting data can be found in the supplements.

**Figure 2 - Figure Supplement 1: Additional data showing modulation of cH and LH by satiation
state, and correlation with food intake**
(a) Gut fluorescence (i.e. food intake) as a function of mean cH pERK fluorescence, mean LH (mLH and
lLH) pERK fluorescence and active cell count.
(b) Top: mLH and lLH mean pERK fluorescence (left) and active cell count (right) as a function of food-
deprivation time (denoted by color intensity). **Bottom:** mLH and lLH mean fluorescence (left) and cell
count (right) as a function of gut fluorescence (i.e. food intake) after 15 min of feeding (denoted by color
intensity).
(c-e) Quantification of gut fluorescence, cH and LH mean pERK fluorescence and LH active cell count
across different food deprivation times (30 min, 2hr and 4hr). Note that in this dataset, these fish brains
have been stained individually, which may have affected cH quantification.
(c) Food intake: 30 min vs 2hr dep. (p = 2.8×10^{-4}), 30 min vs 4hr dep. (p = 4.0×10^{-4}), 2hr vs 4hr dep. (p =
0.56). Asterisk denotes p< 0.05, n = 16/11/14 fish, One-tail Wilcoxon Rank Sum Test.
(d) Mean pERK fluorescence (cH/mLH/lLH): 30 min vs 2hr dep. (p = $0.60/0.001/5.9 \times 10^{-4}$), 30 min vs 4hr
dep. (p=$0.084/8.6 \times 10^{-4}/0.058$), 2hr vs 4hr dep. (p =0.02/0.24/0.54)
(e) Active cell count (mLH/lLH): 30 min vs 2hr dep. (p = 0.0073/0.0094), 30 min vs 4hr dep. (p= $1.6 \times 10^{-4}/0.0017$), 2hr vs 4hr dep. (p = 0.056/0.053).
303

Given the indirect nature of activity mapping in post-fixed animals, as above, we

employed *in vivo* calcium imaging to measure cH and LH neuronal activities during food

deprivation in real time (Figure 2 - Figure Supplement 3-5). Two transgenic Gal4 drivers,

Tg(116A:Gal4) and *Tg(76A:Gal4)*, were used to drive expression of GCaMP6s

(*Tg(UAS:GCaMP6s)*) in large subsets of cH and LH neurons (Figure 2 - Figure Supplement 2-

3). The 116A:Gal4 transgene drives expression mainly in serotonergic neurons in the cH

(88.9±0.8% 5-HT positive) and paraventricular organ (PVO; Figure 2- Figure Supplement 2),

whereas 76A:Gal4 drives expression in a large proportion of LH cells (Figure 2 - Figure

Supplement 3, Muto et al., 2017). Consistent with our pERK results, the initial calcium-mediated

mean fluorescence and firing frequency of a subset of cH neurons scaled with the length of

food-deprivation prior to imaging (Figure 2- Figure Supplement 3d), and increased further as

food-deprivation continued over the 2 hr imaging period (Figure 2 - Figure Supplement 3-5).

The largest rate of increase occurred during the initial hour of food-deprivation (Figure 2 - Figure

Supplement 3-5).

Analysis of LH activity gave more diverse results over the course of the food-deprivation

time course. While some mLH and lLH voxels showed a predicted reduction in baseline

fluorescence and firing rate, many others displayed a significant enhancement of baseline

activity. It is possible that these changes reflect real dynamics in certain cellular or neuropil

subtypes within the LH, or that they are artifacts of head fixation. Despite the significant diversity

in response properties within the LH, we still observe, in line with our expectations, an overall

negative correlation of lLH calcium spikes with the mean value of cH fluorescence (Figure 2 -

Figure Supplement 4 and 5).

326
327

328
329
Figure 2 - Figure Supplement 2: Characterization of the cH and 116A:Gal4 line
(a) pERK-positive cH cells overlaps with anti-5-HT immunostaining and *Tg(116A:Gal4)* cells, and less
with *Tg(TH2:GCaMP5)* (i.e. dopaminergic) cells. Scale bar = 20μm. White arrows point to examples of
overlapping cells.
(b) Z-projection images of whole mount *Tg(116A:Gal4;UAS:GFP)* fish at low (left) and high (right)
intensities. Scale bar = 100 μm.
(c) Overlap of *Tg(116A:Gal4;UAS:GFP)* with anti-5-HT immunostaining is seen in all layers of the caudal
hypothalamus, and also the anterior and posterior paraventricular organ (aPVO and pPVO). Each row
shows a different Z-plane, moving from more dorsal to more ventral. Scale bar = 50 μm.
(d) Higher magnification images of cH, aPVO and pPVO from left side of image in **(c).**
(e) Minimal overlap of *Tg(116A:Gal4;UAS:nfsb-mCherry)* with dopamine neurons labeled by
Tg(TH2:GCaMP5). Note that the *Tg(116A:Gal4;UAS:nfsb-mCherry)* transgenic, which is used in ablation
experiments, shows sparser labeling than with *Tg(UAS:GFP).* Scale bar = 50 μm.
(f) Quantification of 5-HT overlap with *Tg(116A:Gal4;UAS:GFP)* in the cH, aPVO and pPVO.
344

345
346
Figure 2 - Figure Supplement 3: 116A and 76A:Gal4 expression overlap with pERK activity
(a) mLH and lLH activity in voraciously-feeding fish overlaps with *Tg(76A:Gal4;UAS:GCaMP6s)*
expression (dissected brains). All visible pERK-positive neurons were also co-labeled with
GCaMP6s. *Tg(116A:Gal4)* is also epressed. Scale bar = 50 μm.
(b) mLH and lLH activity in voraciously-feeding fish overlaps with *Tg(76A:Gal4;UAS:GCaMP6s)*
expression (whole-mount). All visible pERK-positive neurons were also co-labeled with GFP. Note that
more dorsally and anteriorly other neurons beyond the LH are labeled. Scale bar = 50 μm.
(c) pERK positive cells (food-deprived fish) overlap partially with *Tg(116A:Gal4)* expression. Left:
Tg(116A:Gal4;UAS:GFP) Right: *Tg(116A:Gal4;UAS:nfsb-mCherry).* Scale bar = 20 μm.
(d) Mean cH activity *(Tg(116A:Gal4;UAS:GCaMPHS)* increases as a function of food-deprivation time.
Larvae were food-deprived for 0.5, 2.5, 4.5 or 6.5 hours (n = 12/4/4/8), quickly embedded in agarose and
subsequently imaged for 1.5 hours (every 5 minutes) under a confocal microscope. See Figure 2 - Figure
Supplement 4-5 for simultaneous calcium imaging of cH and LH activity at higher temporal resolution.
360

361
362 **Figure 2 - Figure Supplement 4: Calcium imaging of cH and LH over food-deprivation reveal**
363 **complex dynamics (volumetric imaging)**

(a) Two fish (left and right) are shown. **Top left:** Mean Δf/f across the entire cH, mLH and lLH **Bottom left:** calcium dynamics (integrated (sum) fluorescence and spike frequency, 5 min bins) over the course of a 2 hr long imaging session. Fish were imaged ~20 min after embedding, thus initial food-deprivation time is 20 min. **Right:** Spike-triggered averages based on lLH spikes reveals an accompanying reduction in cH calcium fluorescence, suggesting opposing activity patterns.

(b) Top: K-means clustering (k=4) over all cH and LH voxels reveals diverse clusters of activity. Number of voxels within each cluster is indicated next to the mean Δf/f trace. **Bottom:** Raster plots of clustered neurons sorted from the least active (blue) to most active (red) cluster.

(c) Calcium dynamics (integrated fluorescence and spike frequency) for each cluster over time reveal diverse activity patterns (5 min bins).

(d) Left: Average intensity projection images showing imaged regions. Cell bodies are outlined in white; for the LH they tend to correspond to the edges, whereas neuropil are more concentrated in the center. The cH comprises mainly cell bodies. **Right:** Positions of voxels corresponding to each cluster. Fish 2 is imaged at a more ventral plane than fish 1.

379
Figure 2 - Figure Supplement 5: Calcium imaging of cH and LH over food-deprivation reveal
complex dynamics (single plane)
(a) Two fish (left and right) are shown. **Top left:** Mean Δf/f across the entire cH, mLH and lLH (top left);
Bottom left: calcium dynamics (integrated fluorescence and spike frequency, 5 min bins) over the course
of a 2hr long imaging session. Fish were imaged ~20 min after embedding, thus initial food-deprivation

time is 20 min. **Right:** Spike-triggered averages based on lLH spikes reveals an accompanying reduction
in cH calcium fluorescence. As only a single-plane was imaged, the inverse relationship between the cH
and LH is not as prominent.
(b) Top: K-means clustering (k=4) over all cH and LH voxels reveals diverse clusters of activity. Number
of voxels within each cluster is indicated next to the mean Δf/f trace. **Bottom:** Raster plots of clustered
neurons sorted from the least active (blue) to most active (red) cluster.
(c) Calcium dynamics (integrated fluorescence and spike frequency) for each cluster over time reveal
diverse activity patterns (5 min bins).
(d) Left: Average intensity projection images showing imaged regions. Cell bodies are outlined in white;
for the LH they tend to correspond to the edges, whereas neuropil are more concentrated in the center.
The cH comprises mainly cell bodies. **Right:** Positions of voxels corresponding to each cluster. Since
these fish were not embedded completely symmetrically, the left and right sides of the hypothalamus are
at slightly different dorsal-ventral positions (right sides slightly more dorsal than left sides). Fish 3 is
imaged at a more ventral plane than fish 4.
399
The caudal and lateral hypothalamus respond to food sensory cues and are anti-

correlated over short timescales

We next examined the effects of food sensory cues on cH and LH activity dynamics by

performing calcium imaging on tethered animals during the controlled presentation of food-

related stimuli (Figure 3a). Consistent with the above results of pERK analysis of post-fixed

brains, mLH and lLH neurons were strongly activated and cH neurons suppressed within

seconds of paramecia addition to the water in the vicinity of a food-deprived fish (Figure 3b-c).

Interestingly, neurons in all three hypothalamic loci responded to water flow alone (Figure 3b-c),

with the cH and lLH responding more strongly than the mLH (Figure 3c, bottom panels).

However, these responses were still significantly less than when paramecia were presented

(Figure 3c, bottom panels). Thus, food (and other) sensory cues in the absence of hunting or

food ingestion differentially modulate the activities of neurons in the caudal and lateral

hypothalamic lobes.

We also observed that, across periods in which food cues were either present or absent,

the activities of cH and LH neurons were remarkably anti-correlated; both spontaneous or food-

induced fluctuations in one population were accompanied by a corresponding opposing change

in the other (Figure 3b). This observation was supported by cross-correlation analysis between

cH, mLH and lLH voxels (Figure 3d-f), which revealed high correlation within the same

hypothalamic region (red), and anti-correlation between cH and LH regions (blue) (Figure 3d-e).

Further, lLH voxels showed more spatial heterogeneity than mLH voxels (Figure 3e), though a

small cluster of cells at the most-anterior part of the lLH was found to be consistently anti-

correlated with the cH (Figure 3e, Fish C and D, for example). When ranked according to their

degrees of anti-correlation with voxels from other lobes, the cH and lLH appeared to show the

greatest anti-correlation (Figure 3f). Overall, these results indicate that cH and LH neurons

display generally anti-correlated activity over short timescales in addition to longer epochs

reflecting states of hunger, voracious feeding and satiety.

Figure 3: Anti-correlation on seconds timescale
(a) Top: Schematic showing the calcium imaging setup. Transgenic fish with GCaMP-labeled cH and LH
neurons were paralyzed, tethered in agarose with their eyes and nostrils freed, and exposed to live
paramecia. **Top image:** GCaMP expression in the cH and LH driven by two transgenic lines. **Bottom**
image: Downsampled image stack used for analysis
(b) Left: Mean calcium activity from respective hypothalamic ROIs (shown in (i)) from 4 example fish after
exposure to water (control) or paramecia. Left and right lobes are shown in same color and overlaid.

Paramecia presentation activates the LH and suppresses the cH, which show opposing activity on short
timescales. **Right:** Average Δf/f triggered on lLH calcium spikes shows a mean corresponding reduction
in cH activity (n = 159 lLH spikes extracted from mean Δf/f traces from 14 fish across the entire duration
of the experiment)
(c) Top: Raster plots showing mean calcium activity from the mLH, lLH and cH of 14 fish before and after
presentation of water and food cues. **Bottom:** Quantification of integrated (sum) Δf/f values and spike
frequency (spikes/min) per fish across experimental epochs (300s baseline, 300s after water (W) delivery
or 600 s after food delivery (F). Water delivery was sufficient to significantly modulate both the cH (p
$=6.1\times10^{-5}$ (integrated fluorescence)/0.0497(spike freq.) and the lLH (p=0.029/0.026) but not the mLH (p =
0.48/0.055). Note though from **(b)** that the mLH does transiently respond to water delivery. Food delivery
significantly increased mLH integrated fluorescence (p=1.2×10^{-4}) and spike frequency (p = 1.2×10^{-4})
relative to water delivery. Food delivery also significantly increased lLH integrated fluorescence (p
=0.045) and spike frequency (0.0016) relative to water delivery. Food delivery significantly reduced cH
integrated fluorescence further relative to water delivery (p =3.1×10^{-4}), but not spike frequency (p = 0.52).
W = water, F = food. One-tail Wilcoxon Sign Rank Test.
(d) Cross-correlogram of hypothalamic cell-sized voxels (cells and/or neuropil from downsampled image
stacks, see Figure 2a) from 4 example fish. cH and LH voxels were mostly anti-correlated, whereas
voxels within each cluster had correlated activity.
(e) Correlation coefficients of other hypothalamic voxels relative to a voxel with the cH, mLH or lLH.
(f) Summary of data from 14 fish, showing the probability of the n^{th} most anti-correlated voxel belonging to
each of the other clusters.
455
The activities of cH and LH neurons are differentially modulated by food sensory cues

and ingestion

We next asked whether food sensory cues might have differential and independent effects from

the consummatory cues elicited by the biting and swallowing of prey. Specifically, we tested the

hypothesis that such ingestion cues might be necessary for the more sustained reciprocal

changes in LH and cH activities that accompany voracious feeding. Since consummatory

activity cannot be assessed in head-fixed animals, pERK analysis of activity was performed on

post-fixed animals after free-swimming hunting and feeding behaviors. To distinguish between

sensory and consummatory activities, we compared the neural activity of food-deprived fish

upon exposure to either paramecia or artemia. Artemia are live prey commonly used to feed

adult zebrafish, and are actively hunted by fish at all stages (Figure 4a, Video 2). They are

however too large to be swallowed and consumed by larvae. Thus, the comparison between

these two types of prey dissociates neural activity associated with prey detection and hunting

from the neural consequences of food ingestion.

470   We found that with exposure to artemia it was not possible to detect a change in cH

471 activity, but as observed above with live calcium imaging, exposure to this food cue in the

472 absence of ingestion induced a small increase in lLH neural activity and a larger increase in

473 mLH activity (Figure 4a-b). The artemia-induced hypothalamic activity was, however, less than

474 that observed with consumable prey (Fig 4a-b). These observations suggest that the mLH

475 responds primarily to sensory cues and/or induced hunting behavior whereas the induction of

476 lLH activity largely depends on consumption. These data are furthermore consistent with the

477 strong anti-correlation of cH with lLH activity (compared to mLH activity, Fig 3f), since both

478 respond more strongly to food consumption rather than sensory cues. Thus, in addition to

479 comprising distinct cell types (Figure 1- Figure Supplement 2), the lLH and mLH are also

480 selective for different food cues, raising the possibility that they could be further specialized for

481 distinct behavioral functions (Figure 4c, also see Discussion).

Figure 4: Sensory cues and food consumption differentially regulates cH and LH domains

(a) Representative images of pERK activity induced by paramecia vs artemia larvae. Hatched artemia are sensed and actively hunted by 7-8 dpf larval zebrafish, but too large to consume, allowing for the dissociation of sensory cues/hunting behavior and food consumption. Scale bar = 50 µm. Rightmost panel: Larval zebrafish attempt to hunt live artemia, performing J-bends and pursuits with eyes converged (Bianco et al., 2011). Also see Video 2.

(b) cH activity (normalized pERK intensity) is significantly reduced by paramecia but not by artemia (p = 0.016 (paramecia), 0.648 (artemia)). In contrast, the LH can be activated by artemia alone, though more strongly in the presence of paramecia. The lLH is more weakly activated than the mLH by artemia. Both normalized pERK intensity (mLH: p = 2.06×10^{-5} (paramecia), p = 4.87×10^{-4} (artemia); lLH: p = 2.06×10^{-5} (paramecia), p = 0.033 (artemia)), and active cell count (mLH: p = 1.08×10^{-5} (paramecia), p = 6.02×10^{-5} (artemia); lLH: p = 1.08×10^{-5} (paramecia), p = 5.04×10^{-5} (artemia)) are shown, with n = 8/9/11 fish, One-tail Wilcoxon Rank Sum Test).

(c) Revised schematic showing differential activation of cH and LH domains in response to food sensory vs consummatory cues.

Optogenetic cH activation suppresses ILH neural activity

The anti-correlated activity of the caudal and lateral hypothalamus suggests they might

participate in mutual inhibition. We hypothesized that, during food deprivation, rising cH activity

(along with the absence of food cues) suppresses LH activity, whereas the arrival of food cues

and initial consumption that induces strong LH activity reciprocally shuts down cH activity. The

reduction in cH activity, in turn, would permit higher LH activity, which may underlie voracious

feeding behavior. Conversely, increased cH activity would reduce LH activity and return it to

satiety levels.

In order to test our model that activation of cH neurons is sufficient to suppress LH

activity, we expressed a red-shifted Channelrhodopsin (*Tg(UAS:ReaCHR-RFP)*) in cH neurons

(Dunn et al., 2016; Lin et al., 2013) and visualized LH neuronal activity via calcium imaging

using *Tg(HuC:GCaMP6s)*. Since expression of ReaChR by the *Tg(116A:Gal4)* driver line was

weak, we used a different Gal4 line, *Tg(y333:Gal4),* that labels a smaller fraction of serotonergic

neurons in the cH (57.4±2.1%), but drives robust ReaChR expression (Figure 5 – Figure

Supplement 1). In addition, the *Tg(UAS:GGaMP6s)* transgene was expressed in some animals,

allowing us to monitor optogenetically-induced cH activity. In these animals, ReaChR

stimulation (10-15 seconds, 633 nm laser illumination) and subsequent calcium imaging

confirmed strong activation of the cH. Significantly, optogenetic stimulation of the cH reduced

spontaneous lLH activity (Figure 5b, d), but did not alter mLH activity. Hence it appears that cH

activity is sufficient to inhibit lLH but not mLH activity. This distinction may allow mLH neurons to

remain sensitive and responsive to food cues under food-deprivation conditions, while cH

activity is elevated.

Figure 5 with 1 supplement: Optogenetic cH stimulation is sufficient to reduce lLH activity

(a) Stimulation of cH neurons in *Tg(y333:Gal4;UAS:ReaChR-RFP;UAS:GCaMP6s)* fish with a 633 nm laser induces sustained activation in a fraction of cells. Image shows confocal imaging and stimulation

area, numbers depict individual cells whose activities (Δf/f) are shown on the right. Scale bar = 50 μm.
Bottom: Mean fluorescence across the entire ROI over time. Orange bars = 10 second stimulation period
(no imaging occurs during that period). Gray bars indicate pre- and post-stimulation period over which
activity will be averaged.
(b) Stimulation of cH neurons in a different fish expressing *Tg(y333:Gal4;UAS:ReaChR-RFP;*
HuC:GCaMP6s) reduces spontaneous activity in lLH neurons. The cH was not imaged simultaneously as
ReaChR can be activated by higher-intensity 488 nm light (see methods). Image shows confocal imaging
and stimulation area, numbers depict individual cells or neuropil (NP) whose activities (Δf/f) are shown on
the right. It was not always possible to resolve individual LH cells in other imaged fish. Scale bar = 50 μm.
Bottom: Mean fluorescence across mLH and lLH ROIs over time.
(c-f) Comparison of mean and maximum Δf/f for a 90 s window before and after ReaChR stimulation.
Each data point represents a single stimulation.
(c) cH activity increases after ReaChR stimulation. N = 29 stimulations across 8 fish, p =
$0.0002(max)/0.036(sum)/9.2 \times 10^{-5}(mean)$, One-tail Wilcoxon sign rank test.
(d) mLH activity does not change after ReaChR stimulation (p = 0.74 (max)/0.85 (sum)/0.13 (mean)),
whereas lLH activity is significantly suppressed after ReaChR stimulation (p = $0.0003(max)/1.8 \times 10^{-6}$
$(sum)/0.049(mean)$). N = 108 stimulations across 9 fish. Two-tail Wilcoxon sign rank test.
(e) Stimulation of a control area (i.e. more anterior to cH and unlabeled by ReaChR) tends to increase
activity in the mLH (p = 0.0003(max)/0.039(sum)/0.039(mean) and does not change lLH activity (p =
0.099(max)/0.65(sum)/0.096(mean). N = 37 stimulations from 5 fish. Two-tail Wilcoxon sign rank test.
547

a y333Gal4;UAS:ReaChR:RFP Z-projection

OB

aPVO

pPVO

cH

100 μm

anti-5-HT
y333;Gal4;UAS:ReaChR:RFP

50 μm

aPVO

pPVO

cH

b y333:Gal4;
UAS:ReaChR-RFP   anti-5-HT   Overlay

aPVO

pPVO

cH

More ventral

50 μm

c anti-5-HT   y333;Gal4;UAS:ReaChR:RFP

20 μm   cH(ventral)   cH (dorsal)

aPVO   aPVO   pPVO

d y333:Gal4;UAS:ReaChR-RFP

aPVO

Fish	y333+	y333+/5-HT+	% Overlap
1	12	7	58.3%
2	35	20	57.1%
3	19	10	52.6%
4	16	8	50.0%
MEAN	20.5	11.3	54.5%
ERR	5.04	2.98	1.95%

pPVO

Fish	y333+	y333+/5-HT+	% Overlap
1	13	6	46.2%
2	29	15	51.7%
3	20	9	45.0%
4	19	9	47.4%
MEAN	20.3	9.8	47.6%
ERR	3.30	1.89	1.47%

cH

Fish	y333+	y333+/5-HT+	% Overlap
1	44	26	59.1%
2	87	54	62.1%
3	50	28	56.0%
4	59	31	52.5%
MEAN	60.0	34.8	57.4%
ERR	9.51	6.50	2.05%

548
549
Figure 5- Figure Supplement 1: Characterization of the y333:Gal4 line
(a) We used an alternative cH-labeling Gal4 line, *Tg(y333:Gal4)* (Marquat et al (2015), to drive
Tg(UAS:ReaChR-RFP) expression, as we were unable to detect any ReaChR expression using
Tg(116A:Gal4). **Top:** Whole mount stack of a *Tg(y333:Gal4;UAS:ReaChR-RFP)* (green) shows relatively
specific expression in the caudal hypothalamus, as well as some labeling in the olfactory bulb (white

arrow) and other scattered cells. Scale bar = 100 μm. Bottom: Z-projection image of a dissected fish
brain mounted ventral side up, with anti-5-HT staining shown in magenta. Scale bar = 50 μm.
(b) Overlap of *Tg(y333:Gal4;UAS:ReaChR-RFP)* (green) with anti-5-HT immunostaining (magenta) is
seen in all layers of the caudal hypothalamus, and also the paraventricular organ (PVO), though the
degree of overlap is less for the PVO. Each row shows a different Z-plane, moving from more dorsal to
more ventral. Dissected fish brains mounted ventral side up. Scale bar = 50 μm.
(c) Higher magnification image showing moderate overlap of *Tg(y333:Gal4;UAS:ReaChR-RFP)* with anti-
5-HT staining in the cH and PVO. Arrows indicate cells with overlapping RFP and 5-HT expression. Scale
bar = 20 μm.
(d) Quantification of overlap between 5-HT and *Tg(y333Gal4;UAS:ReaChR-RFP)* expression in the cH
and PVO.
566
Functional dissection of cH serotonergic neurons in feeding behavior

The opposing patterns of cH and LH activities suggest that they might have opposing roles in

the control of hunting and feeding behavior. Given that the cH is composed of neuromodulatory

populations, including serotonergic neurons, and is sufficient to suppress lLH neural activity, we

reasoned that these cH neurons might act as a homeostatic regulator of satiation state-

dependent food intake. In particular, we postulated that: 1) higher cH activation prior to feeding

would encode a state of hunger and enhance the animal's sensitivity to subsequent food cues,

and 2) higher cH activation during feeding would oppose lLH activity to suppress food intake. To

test this hypothesis, we combined ReaChR activation of cH neurons with quantitative

measurements of food intake, again using the y333:Gal4 transgenic line. When satiated

(continuously fed) fish were exposed to whole field orange (630 nm) light for 10 minutes prior to

food presentation, fish in which ReaChR was expressed in the cH consumed significantly more

paramecia than fish that lacked *Tg(y333:Gal4;UAS:ReaChR-RFP)* expression (Figure 6a). This

effect was not observed for food-deprived fish, perhaps because they already display high cH

activity and a high rate of feeding (Figure 6a). These observations are consistent with the

interpretation that optogenetic cH activation simulated a food-deprived state in satiated fish and

thus enhanced their subsequent feeding.

In contrast to the outcome of optogenetic activation prior to feeding, the induction of cH

activity during food presentation reduced feeding, especially in food-deprived fish (Figure 6b).

These observations indicate that optogenetic cH activation can reduce feeding, particularly in

animals in which cH activity is low (Figure 6b, Figure 6 - Figure Supplement 1). Accordingly, we

propose that optogenetic stimulation of cH activity inhibits lLH activity and thereby causes the

feeding rate to decrease.

Finally, we asked what would happen if we directly reduced net cH activity via partial

ablation of the serotonergic population. We hypothesized that this would induce a constitutively

low cH activity, which should enhance food intake regardless of satiation state. Thus, we

performed chemical-genetic ablations of serotonergic cH neurons labeled by *Tg(116A:Gal4;*

UAS:nfsb-Cherry) (Curado et al., 2008), and compared the feeding rates of ablated animals to

that of sibling controls (Figure 6c). Food ingestion was examined in animals in which cH

neurons were partially ablated (see Figure 6 - Figure Supplement 2 and legend for details) and

compared to that of their non-ablated siblings, who lacked *Tg(116A:Gal4;UAS:nfsb-Cherry)*

expression (Figure 6c). We observed a significantly increased food intake for ablated fish in

both food-deprived and fed conditions, indicating that regardless of prior activity patterns, low

cH activity in the presence of food is what ultimately controls food consumption.

Figure 6 with 3 supplements: Role of the cH in behavioral control

(a) Optogenetic activation (orange bar in schematic) of the cH 10 min prior to feeding increases food intake in fed fish, but not food-deprived fish, during subsequent food presentation. Fed: n = 27/26 (ReaChR-/ReaChR+), p = 0.005. Food-deprived: n = 25/29 (ReaChR-/ReaChR+), p = 0.36, One-tail Wilcoxon Rank Sum Test. Since ReaChR expression via 116A:Gal4 was negligible, we used another Gal4 (*Tg(y333:Gal4)*) line that is also specific to the cH when ReaChR is expressed. Fed and food-deprived fish were assayed simultaneously, thus all results was normalized to fed controls. ReaChR-controls do not have visible *Tg(y333:Gal4;UAS:ReaChR-RFP)* expression, and thus are a mixture of siblings expressing *Tg(y333:Gal4) only*, *Tg(UAS:ReaChR-RFP)* or neither of these transgenes, each with ⅓ probability.

(b) Left: Optogenetic activation of the cH (orange bar in schematic) during feeding in fed fish does not

significantly reduce food intake. n = 19/16 (ReaChR-/ReaChR+), p = 0.44 (N.S.), Right**:** Optogenetic
activation of the cH during feeding in food-deprived fish reduces food intake. n = 53/44 (ReaChR-
/ReaChR+), p = 0.042. Since fed and food-deprived fish were assayed in different experiments, gut
fluorescence normalized to their respective controls, One-tail Wilcoxon Rank Sum Test.
(c) Nitroreductase-mediated ablation of the cH in (*Tg(116A:Gal4;UAS-nfsb:mCherry)* or negative fish
treated with metronidazole from 5-7 dpf significantly enhances food intake in 8 dpf fish. p =
0.004/0.04/1.4x10^{-5} (fed control vs ablated, dep. control vs ablated, fed vs dep.). Controls do not have
visible *Tg(116A:Gal4;UAS:nfsb-mcherry)* expression, and thus are a mixture of siblings expressing
Tg(116A:Gal4) only, *Tg(UAS:nfsb-mcherry)* or neither of these transgenes, each with ⅓ probability.
(d) Schematic summarizing our results. We propose distinct roles of the cH during hunger, depending on
the presence or absence of food. See Supplementary Document 1 – Conceptual Circuit Model for
elaboration.
625

a Whole-field optogenetic illumination

626
627
Figure 6 – Figure Supplement 1: ReaChR activation by whole-field optogenetic illumination
(a) *Tg(y333:Gal4;UAS:ReaChR-RFP)* stimulation during feeding is sufficient to induce pERK activity in
many transgene-positive neurons. White arrows point to a few examples where *Tg(y333:Gal4;ReaChR-*
RFP) expression corresponds to more intense pERK staining. Yellow arrows point to examples in which
ReaChR expression is visibly absent, which appears to correspond to weaker pERK staining. ReaChR
expression looks hazy as it was photobleached by the end of the 10 min stimulation period. Scale bar =
20 µm.
635

a

Control (no MTZ)

116A:Gal4; UAS:nfsb-mCherry
anti-5-HT

aPVO
(i)   (ii)

pPVO   (iii)

cH   50 μm

20 μm   20 μm

aPVO (i)

cH (ventral)

aPVO (ii)

pPVO (iii)   cH (dorsal)

Ablated (+ MTZ)

(i)   (ii)

(iii)

cH

aPVO (i)

cH (ventral)

aPVO (ii)

pPVO (iii)   cH (dorsal)

b 116A:Gal4; UAS:nfsb-Cherry   anti-5-HT

More dorsal   50 μm More ventral aPVO (i)

(ii)   (i)
aPVO   (iii) pPVO   cH

cH   aPVO (ii)

cH (dorsal)   cH (ventral)   pPVO (iii)

20 μm   20 μm

c Transgene control

Figure 6 - Figure Supplement 2: Nitroreductase-mediated ablation of cH neurons

(a) Effective ablation of *Tg(116A:Gal4;UAS:nfsb-mCherry)*-labeled neurons using MTZ. Note that due to sparser expression, ablation of the cH/PVO populations are likely to be partial (<50%). Representative projection images are shown of non-ablated (left) and ablated fish brains (right). Scale bar = 50 μm. Insets roughly labeled by white boxes show higher-magnification single-plane images of cH, aPVO and pPVO labeling by this transgene, and overlap with 5-HT expression. Overall, since the labeling of cH neurons with nitroreductase-mCherry is relatively weak (~ 6-8 cells in the aPVO, ~2-4 cells in pPVO, and ~30-40 cells in the cH), our ablations are only partial and may include a few PVO neurons. Scale bar = 20 μm.

(b) Similar to Tg (116A:Gal4;UAS:GFP, there is high overlap of *Tg(116A:Gal4;UAS:nfsb-mCherry)* with anti-5-HT immunostaining. Scale bar = 50 μm. Insets roughly labeled by white boxes show higher-magnification single-plane images of cH, aPVO and pPVO labeling by this transgene, and overlap with 5-HT expression. Scale bar = 20 μm.

(c) The *Tg(116A:Gal4;UAS:nfsb-mCherry)* transgene does not affect feeding in the absence of MTZ, relative to non-expressing siblings. Fed: $p = 0.64$, $n = 11$(negative)/10(positive); Dep.: $p = 0.91$, $n = 11$(negative)/10(positive), Fed vs Dep.: $p = 0.035$ (negative)/7.7×10^{-4}(positive).

Figure 6 - Figure Supplement 3: A conceptual circuit model of the cH/LH hypothalamic network
HEC = Hunger Encoding Circuit, SEC = Satiety Encoding Circuit, which should have anti-correlated activities and report the animal's energy/caloric status. The cH represents both hunger and satiety state and sensitizes (or primes) the LH during hunger. It may drive other behaviors such as exploration or aversive behavior, but also suppresses feeding. Other HEC components may also be involved in LH sensitization. We propose mutual inhibition between the cH and LH, though we have only demonstrated unidirectional inhibition (cH on lLH) thus far. The mLH, normally responsive to food cues, may promote hunting, though not necessarily coupled with ingestion, whereas the lLH, which is more responsive to ingestive cues, should enhance further ingestion (i.e. eating). The LH "gate" is a conceptual representation of how its sensitivity to food cues could be modulated by other signals (i.e. reduced by the SEC and enhanced by cH-mediated sensitization). It does not necessarily represent a physical neuronal population. More elaboration can be found in Supplementary Document 1 – Conceptual Circuit Model.

DISCUSSION

Decades-old studies on appetite regulation in mammals have suggested modular hypothalamic

units that work to suppress or enhance food intake respectively. Here, we show that the larval

zebrafish hypothalamic network can similarly be functionally divided into its medial and lateral

units. These units show anti-correlated activity patterns during various states that relate to

ingestive behavior, such as hunger and voracious eating, and energy homeostasis is reflected

by a restoration of balance between these areas (Figure 6d). Furthermore, we show that within

these broad neural response classes lies a diversity of neurons that encode specific stimuli and

perform distinct functions depending on the timing of activation.

Mutually opposing hypothalamic networks control zebrafish appetite

We show that the medial hypothalamic zone, especially the caudal hypothalamus (cH) in the

zebrafish, is strongly activated by food-deprivation, and strongly inhibited during voracious

feeding, and that this happens on a timescale of seconds to minutes. Here, we focused mainly

on the serotonergic cH neurons, although many medially localized neurons may show similar

activity patterns. In contrast, the lateral hypothalamus (LH), which contains GABAergic and

glutamatergic neurons, is inhibited in the absence of food and most strongly activated during

voracious eating. Interestingly, satiated fish exhibit intermediate activity levels in both

hypothalamic regions. Thus, hunger in the presence and absence of food is represented by two

distinct states of activity in opposing brain regions, with restoration of energy homeostasis

paralleled by a balance of the network.

While generally anti-correlated, the cH and LH also appear to be differentially modulated

by both internal (i.e. hunger cues) and external factors (i.e. food). In the absence of food, LH

cellular activity decreases rapidly, suggesting a requirement of food/other external cues to drive

LH activity, though some modest rate of spontaneous activity is still observed. On the other

hand, the slower timescales of cH activation appears to reflect the animal's rising caloric deficit.

Notably, many of the cH neurons are cerebrospinal fluid-contacting and thus have access to

circulatory information (Lillesaar, 2011; Pérez et al., 2013).

Further, despite clear reductions in pERK cellular activity, calcium imaging has revealed

more complex dynamics of the LH over food-deprivation. These changes could be induced by

the generally aversive and potentially unnatural internal state of a head-fixed preparation, or,

more intriguingly, could reflect an increased sensitivity of LH neurons over the course of hunger.

These hypotheses can potentially be distinguished in future work by performing calcium imaging

of hunting and feeding behavior in a free-swimming setting (Kim et al., 2017).

Once food becomes available (but when caloric deficit is still high), a state change

occurs, and LH activity is strongly enhanced whereas cH activity is strongly suppressed.

Importantly, the degree of cH suppression and LH activation are correlated with the extent of

prior food-deprivation, suggesting a role for these nuclei in regulating food intake based on

caloric needs. This striking anti-correlation between the cH and LH suggests a mutual inhibition,

and that an acute reduction in cH activity is what allows for the enhanced LH release.

We have partially confirmed the hypothesis of such mutual inhibition using optogenetic

stimulation of the cH and simultaneous calcium imaging of the LH. We show that activation of

the cH is sufficient to drive down lLH, but not mLH activity. Consistent with these results, the cH

appears to be more strongly anti-correlated with the lLH than the mLH.

However, the mechanisms by which cH might influence LH activity, and vice versa, are

still unknown. It is possible that the cH may act via inhibitory GABAergic neurons, and/or exert

their effects through direct secretion of monoamines into the ventricles or perineuronal space.

The effect of cH optogenetic activation on lLH activity appears to persist for minutes, allowing

for the possibility of direct neuromodulatory action. At the same time, there appears to be a fast

(seconds) anti-correlation between cH and LH calcium activity, suggesting faster inhibitory

connections. The LH, which was previously characterized in Muto et. al (2017), similarly does

not appear to send direct projections to the cH, but could potentially interact via intermediary

neurons in the medial/periventricular regions of the hypothalamus.

Food cues differentially regulate cH and LH domains

Ingestive behavior has been proposed to comprise a number of temporal stages: 1) the

initiation phase, triggered by energy deficit, in which the animal begins to forage; 2) the

procurement phases, triggered by the presence of food sensory cues, in which the animal seeks

and pursues food; and 3) the consummatory phase, which involves a more stereotyped motor

program (Berthoud, 2002; Watts, 2000). An animal's energy status is sensed internally and may

influence the initiation, procurement and consummatory stages of ingestive behavior. Thus, a

hungry animal will be more alert to food cues, seek food more persistently and also eat it more

voraciously.

In mammals, LH neurons are responsive to both external food sensory cues and

consummatory cues (Jennings et al., 2015). Here, we show that the LH lobes in zebrafish also

respond differentially to food cues in an anatomically segregated manner. In this "sensory"

stage, the mLH is already activated, which may reflect an enhanced sensitivity to food cues

during hunger. However, mLH activation during this sensory stage is not as strong as post-food

consumption. In contrast, the lLH is only weakly activated by food cues, and cH activity

transiently falls but remains overall high. Thus, taken together with our optogenetic and calcium

imaging data, the mutually inhibitory circuit model is most consistent between the cH and lLH

(though the mLH is still generally anti-correlation with cH activity, especially in the presence of

food).

Since lLH and cH activity are modulated within minutes of food consumption they are

unlikely to reflect satiety signals, and rather might play a role in further driving voracious food

consumption, at least until the activity of both populations returns to baseline. In contrast, the

mLH may pay a role in enhancing the sensitivity of the animal to external food sensory cues,

even prior to initial food consumption.

It is unclear which consummatory cues modulate lLH and cH activity. Based on live

imaging results from Muto et al (2017), the greatest enhancement of LH activity was observed

almost immediately (milliseconds to seconds) after paramecia consumption. Thus, the cue is

likely a fast pregastric signal (taste/tactile/swallowing), rather than postgastric absorption or

hormone secretion.

Functional roles of the cH and LH in and beyond appetite control

Finally, we test the hypothesis that the cH and LH form mutually antagonistic functional units

that dominate different phases of hunger and drive appropriate behavioral responses during

each phase. In particular, we show that the activation state of the cH is a crucial regulator of

satiation-state dependent food intake. Artificial cH activation in satiated fish *prior* to feeding is

sufficient to drive subsequent voracious feeding. Based on observed cH dynamics, we propose

that the degree on cH inhibition during voracious feeding is proportional to the degree of cH

activation prior to feeding. This could be mediated by the release of serotonin/other

neuromodulators over the course of food-deprivation, which may be capable of sensitizing the

LH even in the absence of food cues. An intriguing, though untested hypothesis is that the rise

in LH calcium fluorescence during food-deprivation, that tends to parallel that of cH activity, may

reflect such sensitization. In this way, zebrafish are able to retain a "memory" of their hunger

state, which is released once food is presented, and up-regulate their feeding behavior

accordingly. This motif might help ensure that the animal eventually returns to a stable

equilibrium, that is, satiety.

We furthermore show that the acute effect of cH activation *during* feeding is suppression

of food intake, whereas cH ablation enhances food intake, which is again consistent with

mammalian studies of medial hypothalamic areas.   At first glance, the observation that the cH

acutely suppresses food intake is inconsistent with the idea that it is most active during hunger.

However, our optogenetic experiments show that the context of cH activation needs to be taken

into consideration, and can have opposing results on feeding. In the presence of food, activation

of the cH may simply drive down LH activity, hence reducing food intake. This is assuming that

any sensitizing effect of cH activation is weaker than the acute inhibitory effect of cH activation

on ILH activity, a conclusion that appears to be validated by our behavioral results.

The seemingly paradoxical roles of the cH during hunger may also make sense when

considering that, in the absence of food, consummatory behavior would in fact be

counterproductive. Thus, during food-deprivation, the cH may play complementary roles such as

the sensitization of the LH and/or other feeding-related circuits (as discussed above), or drive

alternative behavioral programs, like foraging or energy-conserving measures during this stage

of hunger (see Supplementary Document 1 - Conceptual Circuit Model for a more in-depth

discussion). Given that cH neurons appear also to be activated by aversive stimuli (Randlett et

al., 2015), it may also more generally encode a negative valence state in the absence of food.

Similar features of hunger-related (i.e. AgRP) neurons have also been described in mammals

(Betley et al., 2015; Chen et al., 2015; Dietrich et al., 2015; Mandelblat-Cerf et al., 2015).

Although the cH does not have an exact mammalian homolog, its functions have been

proposed to be adopted by other modulatory populations, such as the serotonergic raphe

nucleus in mammals (Gaspar and Lillesaar, 2012; Lillesaar, 2011). While known to be a potent

appetite suppressant, serotonin is also released during food deprivation, and has been shown to

enhance food-seeking behavior (Elipot et al., 2013; Kantak et al., 1978; Pollock and Rowland,

1981; Voigt and Fink, 2015). Thus, our results showing opposing cH activity patterns during

hunger could reflect similarly complex roles of serotonin in zebrafish, potentially explaining

some of its paradoxical functions. The cH and PVO also express dopaminergic (intermingled

with 5-HT) and histaminergic neurons (in the surrounding cell-layer of the cH), which appear to

be densely interconnected (Kaslin and Panula, 2001). We note that our data, while confirming a

role of serotonergic neurons, does not rule out an involvement of these other neuromodulators

in appetite control.

Further, our results do not rule out the involvement of other circuits in appetite control; in

fact, they suggest that there are numerous players involved. For example, the PVO appears to

be modulated by food cues and food-deprivation, is anti-correlated with LH activity, and labeled

by our transgenic lines (albeit more sparsely), suggesting it may complement the role of the cH.

Our conclusions are also limited by available tools and methodologies -- since different

transgenic lines were utilized for stimulation and ablation, we cannot be certain that we are

manipulating the same population of neurons, though both share mutual overlap with

serotonergic cells. Also, due to the lack of complete transgene specificity, there is a possibility

that our manipulations may affect non-specific targets such as the olfactory bulb.

Similarly, while the strong LH activation after food-deprivation suggests that it might

promote voracious feeding, we were unable to assay the effect of LH activation in larval

zebrafish, due to broad and unspecific expression within the LH-labeling transgenic line.

However, Muto et al (2017) recently demonstrated that inhibition of the LH impairs prey capture

behavior, though they did not implicate the LH in the regulation of food intake based on hunger

state. Furthermore, electrical stimulation of the homologous region (lateral recess nuclei) in

adult cichlids and bluegills (Demski, 1973; Demski and Knigge, 1971) can elicit feeding

behavior, which is consistent with our hypothesis. Interestingly, while stimulating some of these

regions induced food intake, other induced behaviors, such as the "snapping of gravel", were

reminiscent of food search or procurement. In mammals, electrical or optogenetic stimulation of

LH neurons triggers voracious eating, again consistent with our findings that the LH is highly

activated during the voracious eating phase in hungry fish (DELGADO and ANAND, 1953). In

particular, GABAergic neurons that do not co-express MCH or Orexin have been shown to be

responsive to food cues and are sufficient to stimulate food intake in mammals (Jennings et al.,

2015). Whether these GABAergic and glutamatergic neurons of the zebrafish LH co-express

other neuromodulators, as has been recently discovered in mammals (Mickelsen et al., 2019),

remains to be explored. Overall, these data suggest that the zebrafish LH may similarly play an

important role in driving food intake during hunger, despite some differences in peptidergic

expression from mammalian LH. Certainly, since cues such as water flow and optogenetic

stimulation light are sufficient to modulate cH and/or LH neurons, these hypothalamic loci are

likely also involved in other sensorimotor behaviors beyond appetite regulation.

In conclusion, we have shown here how anatomically segregated hypothalamic networks

might interact to control energy homeostasis. We argue that the medial-lateral logic of

hypothalamic function may be conserved even in non-mammalian vertebrates, though their

activity patterns may possibly be more complex than originally believed. Our data suggests

diverse roles of neuromodulators such as serotonin in regulating behavioral responses during

hunger, which may complement mammalian observations. Finally, we propose that investigating

large-scale network dynamics may reveal an additional layer of insights into the principles

underlying homeostatic behavior, which might be overlooked when studies are restricted to the

observation and perturbation of a small subpopulation.

837
838   **SUPPLEMENTARY FIGURE LEGENDS**
839
840   **Supplementary Table 1:** Z-brain anatomical regions that are more activated in voraciously
841   feeding (food-deprived + food) fish as compared to fed fish.
842
843   **Supplementary Table 2:** Z-brain anatomical regions that are more activated in fed fish as
844   compared to voraciously feeding (food-deprived + food) fish.
845
846   **Video 1:** Z-stack (dorsal to ventral) of brain activity map shown in Figure 1b.
847
848   **Video 2:** Video of larval zebrafish hunting artemia larvae. Prey-capture behavior, such as J-
849   bends and pursuits, but no capture swims, were observed in response to artemia larvae.
850   Recording rate: 30 fps. Playback rate: Real time.
851
852   **Supplementary File 1: Conceptual Circuit Model**
853   A comprehensive overview of our circuit model and current understanding, including a circuit
854   diagram, detailed elaboration and testable predictions.
855
856
857
858

859   **MATERIALS AND METHODS**
860
861   **Key Resource Table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
genetic reagent (danio rerio)	Tg(pGal4FF:116A)	Characterized in this manuscript		Dr. Koichi Kawakami (NIG, Japan)
genetic reagent (danio rerio)	Tg(pGal4FF:76A)	PMID: 28425439		Dr. Koichi Kawakami (NIG, Japan)
genetic reagent (danio rerio)	Tg(y333:Gal4)	PMID: 26635538		Dr. Harold Burgess (NIH)
genetic reagent (danio rerio)	Tg(HuC:GCaMP6s)	PMID: 28892088		Dr. Florian Engert (Harvard)
genetic reagent (danio rerio)	Tg(UAS:GCaMP6s)	PMID: 28425439		Dr. Koichi Kawakami (NIG, Japan)
genetic reagent (danio rerio)	Tg(UAS:GCaMPHS)	PMID: 22046464		Dr. Koichi Kawakami (NIG, Japan)
genetic reagent (danio rerio)	Tg(UAS:ReaChR-RFP)	Characterized in this manuscript		Dr. Misha Ahrens (Janelia Research Campus)
genetic reagent (danio rerio)	Tg(UAS-E1b:NTR-mCherry)	PMID: 17335798		Available from ZIRC
genetic reagent (danio rerio)	Tg(Vglut2a:dsRed)	PMID: 19369545		
genetic reagent (danio rerio)	Tg(Gad1b:loxP-dsRed-loxP-GFP)	PMID: 23946442		
genetic reagent (danio rerio)	Tg(Gad1b:GFP)	PMID: 23946442		
genetic reagent (danio rerio)	Tg(TH2:GCamP5)	PMID: 26774784		Dr. Adam Douglass (University of Utah)

genetic reagent (danio rerio)	Tg(ETvmat2:GFP)	PMID:18164283		
genetic reagent (danio rerio)	Tg(HCRT:RFP)	PMID: 25725064		
antibody	Rabbit polyclonal anti-pERK	Cell Signaling	4370 RRID:AB_2315112	IHC(1:500)
antibody	mouse monoclonal anti-ERK	Cell Signaling	4696 RRID:AB_390780	IHC (1:500)
antibody	rabbit polyclonal anti-5-HT	Sigma-Aldrich	S5545 RRID:AB_477522	IHC (1:500)
antibody	goat polyclonal anti-5-HT	AbCam	ab66047 RRID:AB_1142794	IHC(1:500), 2% BSA in PBS blocking solution)
antibody	goat polyclonal anti-MSH	EMD Millipore	AB5087 RRID:AB_91683	IHC(1:500), 2% BSA in PBS blocking solution)
antibody	rabbit polyclonal anti-AGRP	Phoenix Pharmaceuticals	H-003-53 RRID:AB_2313908	IHC (1:500)
antibody	rabbit polyclonal anti-MCH	Phoenix Pharmaceuticals	H-070-47 RRID:AB_10013632	IHC (1:500)
antibody	rabbit polyclonal anti-CART	Phoenix Pharmaceuticals	55-102 RRID:AB_2313614	IHC (1:500)
antibody	rabbit polyclonal anti-NPY	Immunostar	22940 RRID:AB_2307354	IHC (1:500)
antibody	mouse monoclonal anti-TH	Immunostar	22941 RRID:AB_1624244	IHC (1:500)
chemical compound, drug	DiD' solid (lipid dye)	Thermo Fisher Scientific	D-7757	Stock solution (10mg/ml), working solution (2.5mg/ml), in ethanol

Fish husbandry and transgenic lines

Larvae and adults were raised in facility water and maintained on a 14:10 hr light:dark cycle at

28°C. All protocols and procedures involving zebrafish were approved by the Harvard
University/Faculty of Arts & Sciences Standing Committee on the Use of Animals in Research
and Teaching (IACUC). WIK wildtype larvae and mit1fa-/- (nacre) larvae in the AB background,
raised at a density of ~40 fish per 10 cm petri dish, were used for behavioral and MAP-
mapping experiments.
871
Transgenic lines *Tg(UAS-E1b:NTR-mCherry)*(Davison et al., 2007) (referred to as UAS:nfsb-
mCherry), *Tg(UAS:GCaMPHS)* and *Tg(UAS:GCaMP6s)*(Muto and Kawakami, 2011; Muto et al.,
2017)*, Tg(HuC:GCaMP6s)*(Kim et al., 2017)*, Tg(Vglut2a:dsRed)* (Miyasaka et al., 2009),
Tg(Gad1b:loxP-dsRed-loxP-GFP and *Tg(Gad1b:GFP)* (Satou et al., 2013), *Tg(TH2:GCamP5)*
(McPherson et al., 2016)*, Tg(ETvmat2:GFP)* (referred to as VMAT:GFP) (Wen et al., 2008)*,
Tg(HCRT:RFP)* (Liu et al., 2015) have all been previously described and characterized.
Tg(pGal4FF:116A) (referred to as 116A:Gal4) was isolated from a gene trap screen by the
Kawakami group (Kawakami et al., 2010); *Tg(pGal4FF:76A)* was recently published by the
same group(Muto et al., 2017). *Tg(y333:Gal4)* from a different enhancer trap screen was used
to drive expression in the cH in cases where 116A:Gal4-driven expression was sparse
(Marquart et al., 2015). *Tg(UAS:ReaChR-RFP)* was a gift from Misha Ahrens and Chao-Tsung
Yang, made using Tol2 transgenesis in the Ahrens lab at Janelia Research Campus. The same
construct was previously validated in Dunn et al, 2016.
885
886   *MAP-mapping of appetite regions*
887
More details on the MAP-mapping procedure can be found in Randlett et al (2015). 5-6 dpf,
mit1fa-/- (nacre) larvae in the AB background larvae were fed an excess of paramecia once
daily. On the day of the experiment (at 7dpf), the larvae were distributed randomly into two
treatment groups: 1) FOOD-DEPRIVED, where larvae were transferred into a clean petri dish of
facility water, taking care to rinse out all remaining paramecia or 2) FED, where after washing
and transferring they were fed again with an excess of paramecia. After two hours, larvae in
both groups were fed with paramecia. After 15 minutes, larvae were quickly funneled through a
fine-mesh sieve, and the sieve was then immediately dropped into ice-cold 4%
paraformaldehyde (PFA) in PBS (PH 7.2-7.4). Fish were then immunostained with procedures
as reported below (see Immunostaining methods). The rabbit anti-pERK antibody (Cell
Signaling, #4370) and mouse anti-ERK (p44/42 MAPK (Erk1/2) (L34F12) (Cell Signaling,
#4696) were used at a 1:500 dilution. Secondary antibodies conjugated with alexa-fluorophores
(Life Technologies) were diluted 1:500. For imaging, fish were mounted dorsal-up in 2% (w/v)
low melting agarose in PBS (Invitrogen) and imaged at ~0.8/0.8/2 μm voxel size (x/y/z) using an
upright confocal microscope (Olympus FV1000), using a 20x 1.0NA water dipping objective. All
fish to be analyzed in a MAP-Mapping experiment were mounted together on a single imaging
dish, and imaged in a single run, alternating between treatment groups.
905
906   *Whole-mount Immunostaining*
907
24 hours after fixation (4% paraformaldehyde (PFA) in PBS), fish were washed in PBS + 0.25%
Triton (PBT), incubated in 150mM Tris-HCl at pH 9 for 15 min at 70°C (antigen retrieval),

washed in PBT, permeabilized in 0.05% Trypsin-EDTA for 45 min on ice, washed in PBT,
blocked in blocking solution (10% Goat Serum, 0.3% Triton in BSS or 2% BSA in PBS, 0.3%
Triton) for at least an hour and then incubated in primary and secondary antibodies for up to 3
days at 4°C diluted in blocking solution. In-between primary and secondary antibodies, fish were
washed in PBT and blocked for an hour. If necessary, pigmented embryos were bleached for 5
min after fixation with a 5%KOH/3%H2O2 solution.

The protocol was similar for dissected brains, except that the brains were dissected in PBS after
24 hours of fixation, and the Tris-HCL antigen retrieval/permeabilization step in Trypsin-EDTA
was omitted. Dissected brains were mounted ventral up on slides in 70% glycerol prior to
imaging. Confocal images of dissected brains were obtained using either a Zeiss LSM 700 or
Olympus FV1000.

Quantification of 5-HT overlap with transgenic lines

The same individual manually quantified overlap of all transgenic lines with whole-mount or
dissected 5-HT staining, to maintain standardization.

Quantification of food intake

Paramecia cultures (~1-2 500 ml bottles) were harvested, spun down gently (<3000 rpm) and
concentrated, and subsequently incubated with lipid dye (DiD' solid, D-7757, Thermo Fisher
Scientific, dissolved in ethanol) for > 2 hrs (5 µl of 2.5mg/ml working solution per 1 ml of
concentrated paramecia) on a rotator with mild agitation. They were then spun down gently
(<3000 rpm), rinsed and reconstituted in deionized water. An equal amount (100µl, ~500
paramecia) was pipetted into each 10 cm dish of larvae. This method was adapted from
Shimada et al., 2012. After the experiment, larvae were fixed and mounted on their sides on
glass slides or placed in wells of a 96 well plate. They were then imaged using the AxioZoom
V16 (Zeiss) and analyzed using custom Fiji software (Schindelin et al., 2012). In cases where
identity of larvae needed to be maintained, for example, to correlate food intake with brain
activity, larvae were imaged and subsequently stained individually in 96 well plates. This led to
more variable staining which precludes analysis of mean fluorescence.

Larvae were always distributed randomly into experimental groups.

Quantification of LH and cH activity in dissected brains

Brains within each dataset were usually registered onto a selected reference image from the
same dataset using the same CMTK registration software used in MAP-mapping. Further
analysis was then performed using custom Fiji and MATLAB software.

For quantification of cH, mLH and pERK fluorescence intensity, ROIs were manually defined
using the reference image, and pERK intensity was quantified over all registered images and
averaged across the entire lobe (multiple z-planes) as well as across both lobes. Analysis of cH

pERK fluorescence was restricted to the most ventral planes, as more dorsal cH neurons show
weaker correlation with feeding states (as seen in Figure 2 – Figure Supplement 4).
956
957 For quantification of mLH and lLH active cell count, automated analysis of cell count was again
958 performed using custom Fiji software, namely: 1) Image processing to reduce background and
959 enhance contrast 2) Adaptive thresholding to isolate strongly-stained cells 3) Applying the
960 "Analyze Particles" function to quantify the number of cells within each manually-defined ROI.
961
962 Aggregation and visualization of results was performed using custom MATLAB software.
963
Note that, in experiments in which the data was collected without the tERK channel (e.g. from
Figure 2), thus prohibiting image registration, ROIs were drawn manually over each region
across all z-planes and averaged to obtain mean fluorescence values.
For Figure 2 - Figure Supplement 1, where individual fish were stained, all measurements,
including cell count, were made manually. In addition, background fluorescence was measured
for each sample and subtracted from measured values.
970
971 *Calcium imaging*
972
For confocal calcium imaging of the cH and LH simultaneously in the presence of food,
Tg(76A:Gal4;116A:Gal4; UAS:GCaMP6s) triple transgenic fish were embedded in 1.8%
agarose, with their eyes/nostrils were released. GCaMP activity from a single z-plane (where
the cH and LH neurons could be seen) was imaged using a confocal microscope (Olympus
FV1000) at 1 fps. After a 5 min habituation period and a 10 min baseline period, a dense drop of
paramecia was pipetted into the dish. Due to paramecia phototaxis, most of the paramecia
moved into close vicinity of the fish's head under the laser, allowing for strong visual/olfactory
exposure to paramecia. After image registration (TurboReg Fiji Plugin, Thevenaz et al., 1998),
and downsampling (Fiji/MATLAB), manually-segmented ROIs were selected and total
fluorescence within the ROI was calculated. Cross-correlation and other analyses were
performed using custom MATLAB software.
984
For confocal calcium imaging of the caudal hypothalamus (Figure 2- Figure Supplement 3), 4 to
6 food-deprived (2-6 hrs) or fed larvae expressing *Tg(116A:Gal4; UAS:GCaMPHS)*, were
embedded in 1.5% agarose on a large petri dish, and a z-stack covering the entire caudal
hypothalamus imaged using multi-area time lapse imaging every 5 minutes for 2 hrs. Maximum
projection images from the timelapse series were aligned to the first image of the series and
total fluorescence of both caudal hypothalamic nuclei was subsequently measured using
manually-drawn ROIs in ImageJ, to obtain the average calcium activity for each fish at each
time point.
993
For 2P imaging of the cH and LH simultaneously in the absence of food (Figure 2 - Figure
Supplement 4-5), *Tg(76A:Gal4;116A:Gal4; UAS:GCaMP6s)* triple transgenic fish were
embedded in 1.8% agarose. GCaMP activity from either multiple slices (3 z-planes spanning a
~20 µm volume of the intermediate hypothalamus using an electrically-tunable liquid lens

(Edmund Optics, 83-922), 237 ms per z-plane) or a single z-plane where the cH and LH
neurons (1.5 fps) could be seen was imaged using custom 2P microscopes. After image
registration and downsampling to cell-sized voxels (Fiji/MATLAB), manually segmented ROIs
were selected and total fluorescence within the ROI was calculated. Clustering, spike detection
and other analyses were again performed using custom MATLAB software.
1003
1004 *Optogenetic stimulation and simultaneous calcium imaging*
1005
Optogenetic stimulation and calcium imaging was performed on a confocal microscope (Zeiss
LSM 880) using a 633 nm laser for ReaChR activation, and a 488 nm laser for calcium imaging.
Tg(y333:Gal4;UASReaChR-RFP; HuCGCaMP6s) triple-transgenic fish were used to record LH
activity after ReaChR activation. As *Tg(HuC:GCaMP6)* does not label the cH, in some cases we
used fish that also had *Tg(UAS:GCaMP6s)* co-expressed in cH, allowing for monitoring of cH
activity directly.
1012
The ReaChR activation spectrum is wide and 488 nm laser power at sufficiently high intensities
is sufficient to activate ReaChR. Since *Tg(y333:Gal4;UASGCaMP6s)* is expressed strongly in
the cH, weak 488 nm laser power can be used to monitor cH activity after ReaChR activation of
cH. On the other hand, *Tg(HuC:GCaMP6s)* expression in the LH is considerably weaker than
Tg(UAS:GCaMP6s) expression driven by *Tg(y333:Gal4)*, and recording LH activity requires high
laser power. Thus, during LH recording trials, we could not simultaneously image the cH.
1019
1020 Fed fish were embedded in 1.8%-2% agarose, with tails, mouth and eyes freed, 15-20 minutes
1021 before imaging in the absence of food. For baseline recording, spontaneous activities in cH or
1022 LH were recorded. ReachR activation was then induced in one side of cH periodically for 10-15
1023 s, and ensuing activity in one or both sides of LH or cH was recorded continuously during
1024 intervals (of 120-180s) between stimuli.
1025
1026 *Nitroreductase-mediated ablations*
1027
Larvae expressing *Tg(116A:Gal4;UAS:nfsb-mCherry)*, or their non-transgenic siblings were
incubated in 2.5mM Metronidazole (Sigma-Aldrich, M3761) from 4-6 dpf/5-7 dpf. MTZ was
subsequently washed out, and food intake was measured at 7 or 8 dpf. For these experiments,
the MTZ-treated siblings were used as the control group. Each control or ablated group was
food-deprived or fed for 2 hrs, and labeled food was added to quantify food intake. In the case
of fed fish, unlabeled food was very gently washed out 15 mins before the experiment and the
food-deprived fish were also agitated slightly to simulate a short washout.
1035
1036 *Optogenetic stimulation with behavior*
1037
1038 Optogenetic stimulation was done by placing a square LED panel (630 nm, 0.12mW/mm^2 driven
1039 at full current, Soda Vision, Singapore) directly on top of petri dishes containing ReaChR
1040 positive or negative fish, for 10 minutes continuously before or during feeding. We had
1041 attempted other methods of stimulating the fish (e.g. pulsed LED stimulation) but found that it

1042 was disruptive to behavior.
1043
1044 *Artemia Hunting Video*
1045
1046 7 dpf larval fish were food-deprived for 2 hours, acclimatized in 24 well plates for 30 minutes,
1047 and then fed either an excess of hatched artemia or paramecia. Raw videos of hunting behavior
1048 were then recorded for 10 min at 30 fps using a high-resolution monochrome camera (Basler
1049 acA4924) and custom Python-based acquisition software.
1050
1051 *High-resolution behavioral tracking*
1052
We developed a system (Johnson et al., manuscript in preparation) in which a high-speed
infrared camera moves on motorized rails to automatically track a zebrafish larvae in a large
pool (300 x 300 x 4mm). A single fish is recruited to the arena center with motion cues delivered
from a projector to initiate each trial. Paramecia are dispersed throughout the middle of the pool
For analysis 60 Hz image frames are centered and aligned. In every frame, the tail was
skeletonized and the gaze angle of each eye is calculated. The eyes can each move from
around zero degrees (parallel to body-axis) to 40 degrees (converged for hunting). Each bout
was then represented as a point in 220-dimensional posture space by accumulating 22 posture
measurements (20 tail tangent angles to encode tail shape, and 2 eye gaze angles) across 10
image frames (~167 ms) from the beginning of each bout. All bouts were then mapped to a 2-D
space with t-distributed stochastic neighbor embedding (t-SNE), Four major hunting bout types
can be identified from this embedding. Hunts begin with the "j-turn", and fish follow and advance
toward prey objects with "pursuit" bouts. Hunts end with an "abort" or a "strike". When the fish is
not actively involved in a hunt, it explores the arena with "exploratory" bouts. Fractions of
hunting bouts were then compared between fed and food-deprived fish in 3-minute time bins
over 45 min.
1069
1070 *Statistics*
1071
All error bars show mean ± SEM over fish. Significance was reported as follows: *p<0.05.
Significance was determined using the non-parametric Wilcoxon Sign Rank test for paired data
and the Wilcoxon rank Sum test for independent samples. One-tail tests were performed in
cases where there was a prior prediction regarding the direction of change. A one-or two-way
ANOVA (Tukey-Kramer correction, MATLAB statistical toolbox) was used in cases where
multiple comparisons were involved.
1078
1079 **ACKNOWLEDGEMENTS**
1080

We thank Harold Burgess kindly provided us with the y333:Gal4 transgenic line. We further
thank Thomas Panier who assisted Robert Johnson in construction of the rig used for high
resolution behavioral imaging. Support from Steve Turney and the CBS imaging facility, and the
Harvard Center for Biological Imaging were essential for the successful completion of many
experiments. Finally, we would like to thank Jessica Miller, Steve Zimmerman, Karen Hurley

and Brittany Hughes at Harvard for providing invaluable fish care.
1087
1088 **COMPETING INTERESTS**
1089 The authors declare no competing interests.
1090
1091
1092 **REFERENCES**
1093

Ahima, R.S., and Antwi, D.A. (2008). Brain regulation of appetite and satiety. Endocrinol. Metab.
Clin. North Am. *37*, 811–823.
Ammar, A.A., Södersten, P., and Johnson, A.E. (2001). Locus coeruleus noradrenergic lesions
attenuate intraoral intake. Neuroreport *12*, 3095–3099.
ANAND, B.K., and BROBECK, J.R. (1951). Hypothalamic control of food intake in rats and cats.
Yale J. Biol. Med. *24*, 123–140.
Berthoud, H.-R. (2002). Multiple neural systems controlling food intake and body weight.
Neurosci. Biobehav. Rev. *26*, 393–428.
Betley, J.N., Xu, S., Cao, Z.F.H., Gong, R., Magnus, C.J., Yu, Y., and Sternson, S.M. (2015).
Neurons for hunger and thirst transmit a negative-valence teaching signal. Nature *521*, 180–
185.
Bianco, I.H., and Engert, F. (2015). Visuomotor Transformations Underlying Hunting Behavior in
Zebrafish. Curr. Biol. *25*, 831–846.
Bianco, I.H., Kampff, A.R., and Engert, F. (2011). Prey capture behavior evoked by simple
visual stimuli in larval zebrafish. Front. Syst. Neurosci. *5*, 101.
BROBECK, J.R., LARSSON, S., and REYES, E. (1956). A study of the electrical activity of the
hypothalamic feeding mechanism. J. Physiol. *132*, 358–364.
Chen, Y., Lin, Y.-C., Kuo, T.-W., and Knight, Z.A. (2015). Sensory Detection of Food Rapidly
Modulates Arcuate Feeding Circuits. Cell *160*, 829–841.
Davison, J.M., Akitake, C.M., Goll, M.G., Rhee, J.M., Gosse, N., Baier, H., Halpern, M.E.,
Leach, S.D., and Parsons, M.J. (2007). Transactivation from Gal4-VP16 transgenic insertions
for tissue-specific cell labeling and ablation in zebrafish. Dev. Biol. *304*, 811–824.
DELGADO, J.M.R., and ANAND, B.K. (1953). Increase of food intake induced by electrical
stimulation of the lateral hypothalamus. Am. J. Physiol. *172*, 162–168.
Demski, L.S. (1973). Feeding and aggressive behavior evoked by hypothalamic stimulation in a
cichlid fish. Comp. Biochem. Physiol. Part A Physiol. *44*, 685–692.
Demski, L.S., and Knigge, K.M. (1971). The telencephalon and hypothalamus of the bluegill
(Lepomis macrochirus): evoked feeding, aggressive and reproductive behavior with
representative frontal sections. J. Comp. Neurol. *143*, 1–16.
Dietrich, M.O., Zimmer, M.R., Bober, J., and Horvath, T.L. (2015). Hypothalamic Agrp neurons
drive stereotypic behaviors beyond feeding. Cell *160*, 1222–1232.
Dockray, G.J. (2009). The versatility of the vagus. Physiol. Behav. *97*, 531–536.
Dunn, T.W., Mu, Y., Narayan, S., Randlett, O., Naumann, E.A., Yang, C.-T., Schier, A.F.,
Freeman, J., Engert, F., and Ahrens, M.B. (2016). Brain-wide mapping of neural activity
controlling zebrafish exploratory locomotion. Elife *5*.
Elipot, Y., Hinaux, H., Callebert, J., and Rétaux, S. (2013). Evolutionary shift from fighting to
foraging in blind cavefish through changes in the serotonin network. Curr. Biol. *23*, 1–10.
Filosa, A., Barker, A.J., Dal Maschio, M., and Baier, H. (2016). Feeding State Modulates
Behavioral Choice and Processing of Prey Stimuli in the Zebrafish Tectum. Neuron *90*, 596–
608.
Gaspar, P., and Lillesaar, C. (2012). Probing the diversity of serotonin neurons. Philos. Trans.
R. Soc. Lond. B. Biol. Sci. *367*, 2382–2394.

Hoebel, B.G. (1965). Hypothalamic Lesions by Electrocauterization: Disinhibition of Feeding and
Self-Stimulation. Science *149*, 452–453.
Jennings, J.H., Ung, R.L., Resendez, S.L., Stamatakis, A.M., Taylor, J.G., Huang, J., Veleta, K.,
Kantak, P.A., Aita, M., Shilling-Scrivo, K., et al. (2015). Visualizing Hypothalamic Network
Dynamics for Appetitive and Consummatory Behaviors. Cell *160*, 516–527.
Johnson, R., Linderman, S., Panier, T., Wee, C.L., Song, E.Y., Herrera, K., Miller, A., and
Engert, F. Point Process Models of Larval Zebrafish Behavior: Structure on Many Scales.
(Manuscript in Preparation)
Jordi, J., Guggiana-Nilo, D., Soucy, E., Song, E.Y., Wee, C.L., and Engert, F. (2015). A high-
throughput assay for quantifying appetite and digestive dynamics. Am. J. Physiol. Regul. Integr.
Comp. Physiol. ajpregu.00225.2015.
Jordi, J., Guggiana-Nilo, D., Bolton, A.D., Prabha, S., Ballotti, K., Herrera, K., Rennekamp, A.J.,
Peterson, R.T., Lutz, T.A., and Engert, F. (2018). High-throughput screening for selective
appetite modulators: A multibehavioral and translational drug discovery strategy. Sci. Adv. *4*,
eaav1966.
Kantak, K.M., Wayner, M.J., and Stein, J.M. (1978). Effects of various periods of food
deprivation on serotonin turnover in the lateral hypothalamus. Pharmacol. Biochem. Behav. *9*,
529–534.
Kaslin, J., and Panula, P. (2001). Comparative anatomy of the histaminergic and other
aminergic systems in zebrafish (Danio rerio). J. Comp. Neurol. *440*, 342–377.
Kawakami, K., Abe, G., Asada, T., Asakawa, K., Fukuda, R., Ito, A., Lal, P., Mouri, N., Muto, A.,
Suster, M.L., et al. (2010). zTrap: zebrafish gene trap and enhancer trap database. BMC Dev.
Biol. *10*, 105.
Kim, D.H., Kim, J., Marques, J.C., Grama, A., Hildebrand, D.G.C., Gu, W., Li, J.M., and Robson,
D.N. (2017). Pan-neuronal calcium imaging with cellular resolution in freely swimming zebrafish.
Nat. Methods.
Krasne, F.B. (1962). General Disruption Resulting from Electrical Stimulus of Ventromedial
Hypothalamus. Science *138*, 822–823.
Lillesaar, C. (2011). The serotonergic system in fish. J. Chem. Neuroanat. *41*, 294–308.
Lin, J.Y., Knutsen, P.M., Muller, A., Kleinfeld, D., and Tsien, R.Y. (2013). ReaChR: a red-shifted
variant of channelrhodopsin enables deep transcranial optogenetic excitation. Nat. Neurosci. *16*,
1499–1508.
Liu, J., Merkle, F.T., Gandhi, A.V, Gagnon, J.A., Woods, I.G., Chiu, C.N., Shimogori, T., Schier,
A.F., and Prober, D.A. (2015). Evolutionarily conserved regulation of hypocretin neuron
specification by Lhx9. Development *142*, 1113–1124.
Mandelblat-Cerf, Y., Ramesh, R.N., Burgess, C.R., Patella, P., Yang, Z., Lowell, B.B., and
Andermann, M.L. (2015). Arcuate hypothalamic AgRP and putative POMC neurons show
opposite changes in spiking across multiple timescales. Elife *4*, e07122.
Marquart, G.D., Tabor, K.M., Brown, M., Strykowski, J.L., Varshney, G.K., LaFave, M.C.,
Mueller, T., Burgess, S.M., Higashijima, S., and Burgess, H.A. (2015). A 3D Searchable
Database of Transgenic Zebrafish Gal4 and Cre Lines for Functional Neuroanatomy Studies.
Front. Neural Circuits *9*, 78.
McPherson, A.D., Barrios, J.P., Luks-Morgan, S.J., Manfredi, J.P., Bonkowsky, J.L., Douglass,
A.D., and Dorsky, R.I. (2016). Motor Behavior Mediated by Continuously Generated
Dopaminergic Neurons in the Zebrafish Hypothalamus Recovers after Cell Ablation. Curr. Biol.
26, 263–269.
Mickelsen, L.E., Bolisetty, M., Chimileski, B.R., Fujita, A., Beltrami, E.J., Costanzo, J.T.,
Naparstek, J.R., Robson, P., and Jackson, A.C. (2019). Single-cell transcriptomic analysis of
the lateral hypothalamic area reveals molecularly distinct populations of inhibitory and excitatory
neurons. Nat. Neurosci. *22*, 642–656.
Miyasaka, N., Morimoto, K., Tsubokawa, T., Higashijima, S., Okamoto, H., and Yoshihara, Y.

(2009). From the olfactory bulb to higher brain centers: genetic visualization of secondary
olfactory pathways in zebrafish. J. Neurosci. *29*, 4756–4767.
Muto, A., and Kawakami, K. (2011). Imaging functional neural circuits in zebrafish with a new
GCaMP and the Gal4FF-UAS system. Commun. Integr. Biol. *4*, 566–568.
Muto, A., Lal, P., Ailani, D., Abe, G., Itoh, M., and Kawakami, K. (2017). Activation of the
hypothalamic feeding centre upon visual prey detection. Nat. Commun. *8*, 15029.
Pérez, M.R., Pellegrini, E., Cano-Nicolau, J., Gueguen, M.-M., Menouer-Le Guillou, D., Merot,
Y., Vaillant, C., Somoza, G.M., and Kah, O. (2013). Relationships between radial glial
progenitors and 5-HT neurons in the paraventricular organ of adult zebrafish - potential effects
of serotonin on adult neurogenesis. Eur. J. Neurosci. *38*, 3292–3301.
Pollock, J.D., and Rowland, N. (1981). Peripherally administered serotonin decreases food
intake in rats. Pharmacol. Biochem. Behav. *15*, 179–183.
Randlett, O., Wee, C.L., Naumann, E.A., Nnaemeka, O., Schoppik, D., Fitzgerald, J.E.,
Portugues, R., Lacoste, A.M.B., Riegler, C., Engert, F., et al. (2015). Whole-brain activity
mapping onto a zebrafish brain atlas. Nat. Methods *12*, 1039–1046.
Satou, C., Kimura, Y., Hirata, H., Suster, M.L., Kawakami, K., and Higashijima, S. (2013).
Transgenic tools to characterize neuronal properties of discrete populations of zebrafish
neurons. Development *140*, 3927–3931.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for
biological-image analysis. Nat. Methods *9*, 676–682.
Semmelhack, J.L., Donovan, J.C., Thiele, T.R., Kuehn, E., Laurell, E., and Baier, H. (2015). A
dedicated visual pathway for prey detection in larval zebrafish. Elife *4*.
Shimada, Y., Hirano, M., Nishimura, Y., and Tanaka, T. (2012). A high-throughput fluorescence-
based assay system for appetite-regulating gene and drug screening. PLoS One *7*, e52549.
Sternson, S.M., and Eiselt, A.-K. (2017). Three Pillars for the Neural Control of Appetite. Annu.
Rev. Physiol. *79*, 401–423.
Stuber, G.D., and Wise, R.A. (2016). Lateral hypothalamic circuits for feeding and reward. Nat.
Neurosci. *19*, 198–205.
TEITELBAUM, P., and EPSTEIN, A.N. (1962). The lateral hypothalamic syndrome: recovery of
feeding and drinking after lateral hypothalamic lesions. Psychol. Rev. *69*, 74–90.
Thevenaz, P., Ruttimann, U.E., and Unser, M. (1998). A pyramid approach to subpixel
registration based on intensity. IEEE Trans. Image Process. *7*, 27–41.
Trivedi, C.A., and Bollmann, J.H. (2013). Visually driven chaining of elementary swim patterns
into a goal-directed motor sequence: a virtual reality study of zebrafish prey capture. Front.
Neural Circuits *7*, 86.
Voigt, J.-P., and Fink, H. (2015). Serotonin controlling feeding and satiety. Behav. Brain Res.
277, 14–31.
Watts, A.G. (2000). Understanding the neural control of ingestive behaviors: helping to separate
cause from effect with dehydration-associated anorexia. Horm. Behav. *37*, 261–283.
Wen, L., Wei, W., Gu, W., Huang, P., Ren, X., Zhang, Z., Zhu, Z., Lin, S., and Zhang, B. (2008).
Visualization of monoaminergic neurons and neurotoxicity of MPTP in live transgenic zebrafish.
Dev. Biol. *314*, 84–92.
Zhu, J.-N., and Wang, J.-J. (2008). The cerebellum in feeding control: possible function and
mechanism. Cell. Mol. Neurobiol. *28*, 469–478.
1232
1233
1234