Temporally coherent perturbation of neural dynamics during retention alters human multi-item working memory

Jiaqi Li[1-3, 7], Qiaoli Huang[1-3, 7], Qiming Han[1-4], Yuanyuan Mi[5, 6, *], Huan Luo[1-3, *]

[1] School of Psychological and Cognitive Sciences, Peking University

[2] PKU-IDG/McGovern Institute for Brain Research, Peking University

[3] Beijing Key Laboratory of Behavior and Mental Health, Peking University

[4] Peking-Tsinghua Center for Life Sciences, Peking University, Beijing 100871, China

[5] Center for Neurointelligence, Chongqing University, Chongqing 400044, China;

[6] AI Research Center, Peng Cheng Laboratory, Shenzhen 518005, China

[7] These authors contributed equally to the work

Running title:

Dynamic perturbation manipulates human working memory

Address for Correspondence:

* Yuanyuan Mi

Center for Neurointelligence, Chongqing University, Chongqing 400044, China;

miyuanyuan0102@cqu.edu.cn

* Huan Luo

School of Psychological and Cognitive Sciences, Peking University

PKU-IDG/McGovern Institute for Brain Science, Peking University

Room 1703, 52 Haidian Road, Beijing, 100087, China

huan.luo@pku.edu.cn

Summary

Temporarily storing a list of items in working memory (WM), a fundamental ability in cognition, has been posited to rely on the temporal dynamics of multi-item neural representations during retention. Here, we develop a "dynamic perturbation" approach to manipulate the relative memory strength of a list of WM items, by interfering with their neural dynamics during the delay period in a temporally correlated way. Six experiments on human subjects confirm the effectiveness of this WM manipulation method. A computational model combining continuous attractor neural network (CANN) and short-term synaptic plasticity (STP) principles further reproduces all the empirical findings. The model shows that the "dynamic perturbation" modifies the synaptic efficacies of WM items through STP principles, eventually leading to changes in their relative memory strengths. Our results support the causal role of temporal dynamics of neural network in mediating multi-item WM and offer a promising, non-invasive approach to manipulate WM.

Keywords:

working memory, recency effect, short-term plasticity (STP), hidden state, memory manipulation

Introduction

Temporarily holding a sequence of items in working memory (WM) for subsequence planning or action is an essential function in many cognitive processes, such as language, movement control, episodic memory, and decision making (Dell et al., 1997; Burgess and Hitch, 1999; Cowan, 2001; Doyon et al., 2003; Giraud and Poeppel, 2012). Psychological evidence and computational models suggest that this process relies on attentional refresh (Awh et al., 1998; Baddeley, 2003; Camos et al., 2018; Oberauer, 2019) or neural reactivations during retention (Lisman and Idiart, 1995; Horn and Opher, 1996; Compte et al., 2000; Raffone and Wolters, 2001; Siegel et al., 2009; Heusser et al., 2016; Lundqvist et al., 2016; Michelmann et al., 2016; Huang et al., 2018; Liu et al., 2019; Schuck and Niv, 2019). Interestingly, recent studies and computational modellings advocates a hidden-state WM network view that does not demand persistent firing during maintenance (Stokes, 2015; Fiebig and Lansner, 2017; Trübutschek et al., 2017; Wolff et al., 2017). Instead, items could be temporarily held in synaptic weights via short-term plasticity (STP) principles (Fusi, 2008; Mongillo et al., 2008; Barak and Tsodyks, 2014; Mi et al., 2017). In fact, the two seemingly contradictory proposals – activated state and hidden state – could be well combined into a unified framework whereby the reactivation and STP mutually facilitate each other and together subserve WM storage (Masse et al., 2018, 2020; Miller et al., 2018). In this WM network, multiple items, by sharing the same connection to an inhibitory pool, compete with each other in time and take turns to elicit brief bouts of reactivation to strengthen their individual synaptic weights (Fino and Yuste, 2011; Mi et al., 2017). Therefore, the dynamic rivalry between item reactivations and the accompanying slowly decaying synaptic weight changes that accumulate over the retention interval would essentially determine the memory strength of each item in the WM network.

Previous studies have examined the neural coding of items and the top-down modulation of WM representations (Sauseng et al., 2009; Rose et al., 2016; Mallett and Lewis-Peacock, 2018; Van Ede et al., 2018), yet the causal evidence for the role

of retention-stage neural network dynamics in mediating multi-item WM is largely

absent. In fact, unlike optogenetics tools widely used in animal study, there is yet no

time-resolved approach to manipulate the dynamics of WM network in the human

brain. Here we developed a novel "dynamic perturbation" approach aiming to modify

the relative memory strength of a list of items maintained in WM. Specifically,

previous studies show that a task-irrelevant color feature will be automatically bound

to the to-be-memorized feature (e.g., orientation) in WM when they belong to the

same object (Luck et al., 1997; Johnson et al., 2008; Hyun et al., 2009; Huang et al.,

2018). Therefore, by manipulating the ongoing temporal relationship between the

continuously flickering color probes that are bound to each WM item respectively, we

could presumably apply temporally correlated perturbations to these item-specific

neural assemblies in the WM network. Consequently, this temporal perturbation

would impact the dynamical WM network in certain way and modulate the ongoing

competition between items, leading to the changes in their relative STP-based

memory strength and finally the subsequent recalling performance.

Six experiments consistently show that the "dynamic perturbation" approach

successfully alters the multi-item WM recalling performance on a trial-by-trial basis.

We focused on the recency effect (i.e., better memory performance for recently than

early presented item), a typical WM behavioral index to characterize the relative

memory strength of a sequence of items (Burgess and Hitch, 1999; Gorgoraptis et al.,

2011; Baddeley, 2012; Jones and Oberauer, 2013). First, when applying synchronized

continuous luminance inputs to the color probes ("Synchronization" manipulation)

during retention, the recency effect is significantly disrupted. In contrast, the recency

effect keeps intact when luminance inputs are temporally uncorrelated ("Baseline"

condition). Second, when the luminance sequences of the color probes are temporally

shifted with each other in an order that is either the same or reversed as the stimulus

sequence ("Order reversal" manipulation), they lead to distinct and even reversed

recency effect. Finally, we established a theoretical continuous attractor neural

network (CANN) with STP effects (Brody et al., 2003; Romani and Tsodyks, 2015;

Trübutschek et al., 2017; Seeholzer et al., 2019) to mimic the "dynamic perturbation"

course. The simulation successfully replicated all our experiment findings, further

corroborating the neural mechanism for our manipulation approach. Taken together,

our results constitute new causal evidence for the essential role of STP-based neural

dynamics in multi-item WM and provides an efficient approach to manipulate WM in

humans.

Results

"Dynamic perturbation" approach: temporal association of memory-related but task-irrelevant color probes during retention

Subjects performed a WM task in all the experiments. As shown in Figure 1A, participants viewed two serially presented bar stimuli and needed to memorize the orientations of the two bars ("encoding period"). After 5 sec memory retention, participants rotated a horizontal bar to the memorized 1^{st} or 2^{nd} orientation based on the instruction ("recalling period"). Importantly, the two bar stimuli, in addition to having distinct to-be-memorized orientations, also had different colors. During the "maintaining period", subjects performed a central fixation task, while two discs that had color feature of either the 1^{st} or the 2^{nd} memorized bars (1^{st} memory-related probe, WM1; 2^{nd} memory-related probe, WM2) were displayed at left or right side of the fixation. The color (red, green, or blue) and the spatial locations (left or right) of the two probes (WM1 and WM2) were balanced across trials.

Crucially, throughout the 5 sec retention period, the luminance of the two color probes (WM1 and WM2) was continuously modulated according to two 5 sec temporal sequences respectively (i.e., Seq1 for WM1, Seq2 for WM2), the relationship of which varied in certain way. Under Baseline condition, the two luminance sequences (Seq1 and Seq2) were two independently generated random time series, and thus no temporal association between items were introduced (Figure 1B, left). Importantly, we also developed two types of manipulation that aim to apply temporally correlated perturbation – Synchronization (Figure 1B) and Order reversal (Figure 3A) – to the two color probes. For "synchronization" condition, the flickering probes were modulated by the same luminance sequence, either the same random temporal sequence (F-Sync; Figure 1B, right), or the same alpha-band filtered random time series (A-Sync; Figure 1B, middle). Note that including A-Sync condition is motivated by previous findings revealing prominent alpha-band (8-11Hz) activations in WM (Klimesch, 1999; Jensen, 2002). We then examined whether and how the dynamic manipulation impacts the subsequent memory performance, particularly the

recency effect, a behavioral index reflecting the relative memory strength of WM

items (Burgess and Hitch, 1999; Baddeley, 2012). For Baseline condition when no

temporal associations are introduced, we would expect to see intact recency effect, as

shown previously (Gorgoraptis et al., 2011; Huang et al., 2018). In contrast,

conditions that bring temporally correlated perturbation such as Synchronization or

Order reversal would presumably interfere with neural dynamics and impact the

recency effect.

It is noteworthy that all the luminance sequences were generated anew in each

trial. To quantify the memory performance for the 1st and the 2nd orientation, we used

Paul Bays' model (Bays et al., 2009), a largely used approach that could be

implemented through an open-source toolbox, to calculated the target probability (see

the precision results in Supplemental Figure 1). We then converted the target

probability results using an arcsine square root transformation (Bartlett et al., 1936) to

make it normally distributed.

Figure 1 about here

"Synchronization" manipulation disrupts recency effect (Experiment 1–2)

Twenty-seven subjects participated in Experiment 1 (Figure 2A). First, consistent with

our hypothesis, the Baseline condition showed significant recency effect ($2^{nd} > 1^{st}$

item), as typically observed when subjects load a list of items into WM (Gorgoraptis

et al., 2011; Huang et al., 2018), suggesting that temporally uncorrelated perturbation

would not affect the relative memory strength of WM items. In fact, a control

experiment that employed the same paradigm but without flickering color probes

presented during retention showed the same recency effect as the Baseline condition

(see Supplementary Figure 2). In contrast, when Seq1 and Seq2 are temporally

synchronized with each other (A-Sync and F-Sync) by sharing the same luminance

sequence, the recency effect was largely disrupted, suggesting their relative memory

strengths were successfully modified. Two-way repeated ANOVA (Position * Sync)

revealed significant interaction effect ($F_{(2,52)} = 3.337$, p = 0.043, $\eta_p^2 = 0.114$), and

nonsignificant main effect for either position ($F_{(1,26)} = 0.372$, p = 0.547, $\eta_p^2 = 0.014$) or

condition ($F_{(2,52)} = 0.268$, p = 0.766, $\eta_p^2 = 0.010$), indicating that this dynamic

perturbation specifically disrupted recency effect without affecting the overall

memory performance. Post-hoc comparisons further confirmed the disruption of

recency effect for both A-Sync and F-Sync compared to Baseline (2-way repeated

ANOVA; A-Sync and Baseline: interaction effect, $F_{(1,26)} = 5.274$, p = 0.030, $\eta_p^2 =$

0.169; F-Sync and Baseline: interaction effect, $F_{(1,26)} = 3.503$, p = 0.073, $\eta_p^2 = 0.119$).

Moreover, the two synchronization conditions did not show significant difference,

indicating their similar disruption effects (2-way repeated ANOVA; A-Sync and F-

Sync: interaction effect, $F_{(1,26)} = 0.494$, p = 0.488, $\eta_p^2 = 0.019$).

To further confirm the findings, we ran a separate group of subjects (N = 24) for

only the A-Sync and Baseline conditions (Experiment 2). As shown in Figure 2B, the

results were well consistent with Experiment 1, such that the Baseline condition

showed recency effect (paired t-test, $t_{(23)} = -2.721$, p = 0.012, Cohen's d = -0.555),

whereas the A-Sync condition did not (paired t-test, $t_{(23)} = 1.300$, p = 0.207, Cohen's d

= 0.265). Two-way repeated ANOVA (Positions * Sync) revealed significant

interaction effect ($F_{(1,23)} = 7.493$, p = 0.012, $\eta_p^2 = 0.246$) but nonsignificant main

effect for either position ($F_{(1,23)} = {}_{0.6}59$, p = 0.425, $\eta_p^2 = 0.028$) or sync condition

($F_{(1,23)} = 0.283$, p = 0.600, $\eta_p^2 = 0.012$), further supporting the synchronization-

induced disruption effect.

Moreover, to examine whether the disruption effect was specifically introduced

by the memory-related probes, we ran another experiment by using memory-

irrelevant color probes (Experiment 2-IC, N = 25). As shown in Figure 2B (right), the

Baseline and Synchronization conditions now did not show significant difference in

recency effect (two-way repeated ANOVA, interaction effect, $F_{(1,24)} = 0.006$, p =

0.938, $\eta_p^2 < 0.001$). Two-sample two-way ANOVA (across Experiment 2 and

Experiment 2-IC) showed marginally significant interaction effect

(Experiment*Position*Sync, $F_{(1,47)} = 3.391$, p = 0.072, $\eta_p^2 = 0.067$), further supporting

that the manipulation is effective for memory-related probes but not for the non-

memory-related ones.

As summarized in Figure 2C, temporal synchronization of the memory-related

probes during retention essentially disturbed the recency effect (see similar but

nonsignificant trend when spatial locations were synchronized, Experiment 7,

Supplementary Figure 3). It is noteworthy that the color probes were completely task-

irrelevant in terms of both memory task (memorizing orientations) and attentional

task (central fixation task). Moreover, this disruption effect was not accompanied by

any overall memory performance changes (i.e., no main effect), indicating that the

manipulation does not interrupt memory retention at a general level but indeed

specifically modulates the relative memory strength of WM items.

188   Figure 2 about here

"Order reversal" manipulation reverses recency effect (Experiment 3-5)

After establishing the efficacy of the synchronization manipulation, we next examined

the "order reversal" manipulation. Specifically, the same experimental paradigm as

Experiment 1 was used (see Figure 1A), except that the luminance sequences of the

two color discs were modulated in a different way. As shown in Figure 3A, Seq1 was

a temporally shifted version of Seq2, either leading (Same, upper panel) or lagging

Seq2 (Reversed, lower panel) by 200 ms. The selection of 200 ms was motivated by

the time constant used in STP neural model (Mongillo et al., 2008; Mi et al., 2017;

Trübutschek et al., 2017) and previous neurophysiological studies revealing a central

role of theta-band rhythm in organizing memory replay (Lisman and Idiart, 1995;

Gyorgy, 2002; Bahramisharif et al., 2018; Huang et al., 2018; Herweg et al., 2020).

Two separate groups of subjects participated in Experiment 3 (N = 24; Same,

Reversed, Baseline) and Experiment 4 (N=27; Same, Reversed, F-Sync), respectively.

First, as shown in Figure 3B, consistent with previous results (Experiment 1-2),

recency effect persisted for Baseline condition (paired t-test, $t_{(23)} = -1.835$, p = 0.0790,

Cohen's d = -0.375; left panel) and vanished for Synchronization condition (paired t-

test, $t_{(26)}$ = -0.366, p = 0.718, Cohen's d = -0.0704; right panel). Most importantly, the

recency effect significantly decreased for Reversed compared to Same condition

(paired t-test, Expt. 3: $t_{(23)}$ = 2.184, p = 0.039, Cohen's d = 0.446; Expt. 4: $t_{(26)}$ =

1.854, p = 0.075, Cohen's d = 0.357) in both experiments (Figure 3B). Two-sample

two-way repeated ANOVA (2*2*2, Experiment*Position*Order) revealed significant

Position*Order interaction effect ($F_{(1,49)}$ = 7.543, p = 0.008, η_p^2 = 0.133), supporting

that the "order reversal" manipulation successfully altered the WM behavior. We

further combined the Same and Reversed conditions across Experiment 3 and 4 since

the two conditions were exactly the same and derived from two independent subject

groups. As shown in Figure 3C (left panel), the pooled results (N = 51) showed clear

decrease and even reversal in recency effect for the Reversed condition compared to

the Same condition (paired t-test, $t_{(50)}$ = 2.787, p = 0.007, Cohen's d = 0.390). It is

noteworthy that the only difference between the Same and Reversed conditions was

the relative order of the two luminance sequences (i.e., Seq1 leads Seq2, Seq2 leads

Seq1), yet they led to distinct recency effect. This strongly advocates that the dynamic

perturbation in terms of different temporal direction interferes with the WM neural

network in opposite ways.

Finally, we assessed whether introducing a temporal lag outside of the STP time

window would also influence WM behavior. Instead of 200 ms, Experiment 5 (N =

27) employed a 500 ms temporal lag between Seq1 and Seq2. As shown in the right

panel of Figure 3C, the Same and Reversed conditions now did not show significant

recency difference (paired t-test, $t_{(26)}$ = 0.331, p = 0.743, d = 0.064). Thus, the

dynamic perturbation should be applied within the STP-related temporal window

(e.g., short-term depression (STD) time constant) to successfully influence WM.

231   Figure 3 about here

Meta-analysis

Finally, to increase the detecting sensitivity and examine the overall effects, we

performed a meta-analysis (Comprehensive Meta-Analysis Software, CMA) across all

8 experiments, including Expt. 1 -5 and Expt. 2-IC in main texts as well as Expt. 6

(probe-absent experiment) and Expt. 7 (spatial location experiment) in Supplementary

Materials. As shown in Figure 4, the Baseline condition (Experiment 1, 2, 3, 5, 6, 7, 2-

IC) showed significant recency effect ($p = 3e^{-8}$), whereas the Synchronization

conditions (Experiment 1, 2, 4, 7) did not ($p = 0.424$). Furthermore, Baseline and

Synchronization conditions (Experiment 1, 2, 7) revealed significant difference ($p =$

$1e^{-4}$). The meta-analysis on precision measurements showed the same results

(Supplementary Figure 4).

246   Figure 4 about here

247
248

CANN-STP WM computational model

After confirming the effectiveness of the dynamic perturbation approach on WM

manipulation, we further developed a computational model that combines CANN and

STP principles to unveil the underlying neural mechanism. CANN has been widely

used to model orientation tuning (Ben-Yishai et al., 1995; Blumenfeld et al., 2006)

and WM-related network (Brody et al., 2003; Parthasarathy et al., 2019; Seeholzer et

al., 2019) in neural systems. STP, referring to a phenomenon that synaptic efficacy

changes over time with neuronal activity, is also commonly used to implement WM

functions (Mongillo et al., 2008; Barak and Tsodyks, 2014; Mi et al., 2017; Miller et

al., 2018). Specifically, as shown in Fig. 5A, in the CANN, excitatory neurons are

aligned according to their preferred orientations, connected in a translation-invariant

manner. Furthermore, all excitatory neurons are reciprocally connected to a common

inhibitory neuron pool (Kim et al., 2017), so that only a single bump-shaped neural

response could be generated at any moment. The STP effect is modeled by two

dynamical variables, u and x, representing the release probability of neural

transmitters (leading to short-term facilitation, STF) and the fraction of resources

available after neurotransmitter depletion (leading to STD), respectively (Markram et al., 1998; Mongillo et al., 2008). The instantaneous synaptic efficacy is given by Jux, with J as the maximum synaptic efficacy. The dynamics of the synaptic efficacy variation is determined by the time constants of u and x, denoted as τ_f and τ_d, respectively. Since STF is dominated in PFC，i.e., $\tau_f \gg \tau_d$, we can effectively approximate the STD variable x to be a constant, and only compare the simplified synaptic efficacies Ju (see Supplementary Materials and Methods for more details).

We then used the CANN-STP WM model to mimic the intrinsic dynamics of neural activities and synaptic efficacies during the delay period as well as the memory recalling performance, for each experimental condition (200 trials in each run, 20 runs for each condition), respectively. Taking one condition as example (Figure 5B), two orientation features are first loaded into the CANN sequentially and evoke two strong transient response (i.e., population spikes). Notably, during retention when stimuli are not present anymore, the synaptic efficacies of the two memory-related neuronal populations would not fall to baseline, and instead, they decay slowly and remain at high values as a result of STP principles. Finally, during recall, a noisy signal containing partial information of a to-be-recalled item is presented to the network, and correct retrieval is achieved if the corresponding neuron group is successfully elicited.

The critical manipulation – flickering color probes – are simulated as continuous and weak inputs, which are endowed with associated temporal characteristics as used in each experiment condition, to the corresponding neural groups during retention (Figure 6, left column, I_{ext}; see more details in Supplementary Materials). These input sequences will perturb the neuronal dynamic through the CANN structure and STP principles and in turn modulate the relative size of the synaptic efficacies of neuron groups (Figure 6, middle column), ultimately determining the memory strength of WM items and the recalling performance (Figure 6, right column).

Figure 5 about here

Model simulations

First, we examined whether our model could reproduce the typical recency effect shown in previous studies (Burgess and Hitch, 1999; Gorgoraptis et al., 2011; Baddeley, 2012; Jones and Oberauer, 2013) as well as our own (Probe-absent experiment, Supplementary Figure 2), when there is no probe stimulation during retention. As shown in Fig. 5B, the loaded items evoke PSs, and the network activity decays to silence rapidly afterwards. Meanwhile, during retention when inputs are removed, the synaptic efficacies of the two neuron groups persist at high values as a result of STP (Figure 5B, left column). Since the two WM items are loaded at different moments, the averaged synaptic efficacy of the early presented item (1st WM item) would be generally smaller than that of the recently presented item (2nd WM item) (Figure 5B, middle column). Consequently, given their difference in synaptic efficacy developed during retention, the 2nd WM item would have larger probability to be successfully retrieved than the 1st WM item (Figure 5B, right column), resulting in the recency effect (paired t-test, $t_{(19)}$ = -7.389, p = $5e^{-7}$, Cohen's d = -1.652).

We next examined the Baseline condition, by applying two independent random inputs to the two neuron groups during retention. The probes occasionally trigger PSs in the network (Figure 6A, left column), but they do not essentially disturb the relative size of the synaptic efficacies of the two neuron groups, i.e., $\langle Ju_2 \rangle > \langle Ju_1 \rangle$ still holds, and the mean firing rate for the 2nd item is larger than the 1st item, ($\langle r_2 \rangle > \langle r_1 \rangle$) (Figure 6A, middle column). As a result, the Baseline manipulation still leads to recency effect in recalling performance (Figure 6A, right column; paired t-test, $t_{(19)}$ = -3.381, p = 0.003, Cohen's d = -0.756), consistent with the empirical findings (Figure 2).

For Synchronization conditions, instead of two independent random inputs, the same input series (white noise for F-Sync and alpha-band series for A-Sync) were applied to the two neuron groups (Figure 6BC, left column). Crucially, the synchronized inputs largely disturb the network dynamics and change the relative size of synaptic efficacies, i.e., fluctuating pattern in the synaptic efficacies (Figure 6BC,

middle column) as well as the neural activities. Consequently, during recalling (Figure 6BC, right column), the recency effect is largely disrupted (F-Sync: paired t-test, $t_{(19)}$ = 0.971, p = 0.344, Cohen's d = 0.217; A-Sync: paired t-test, $t_{(19)}$ = 1.905, p = 0.072, Cohen's d = 0.426), also in line with the experimental observations (Figure 2).

For the "Order reversal" simulation, the input to the neural group that encodes the 1^{st} orientation would either lead (same-order condition) or lag (reversed-order condition) the input to the neural group encoding the 2^{nd} orientation (Figure 6DE, left column). Specifically, the same-order manipulation enhances the relative sizes of the synaptic efficacies of two neuron groups, i.e., enlarged $\langle Ju_2 \rangle - \langle Ju_1 \rangle$ difference and $\langle r_2 \rangle > \langle r_1 \rangle$ (Figure 6D, middle column), finally leading to the recency effect (Figure 6D, right column; paired t-test, $t_{(19)}$ = -6.652, p = $2e^{-6}$, Cohen's d = -1.487). In contrast, the "Order reversal" manipulation keeps disturbing the relative size of synaptic efficacies and flips the relationship between them, i.e., $\langle Ju_1 \rangle > \langle Ju_2 \rangle$ and $\langle r_1 \rangle > \langle r_2 \rangle$ (Figure 6E, middle column). This process causes the reversal of the recency effect to primary effect during recalling (Figure 6E, right column; paired t-test, $t_{(19)}$ = 4.363, p = $3e^{-4}$, Cohen's d = 0.976), again consistent with the experimental results (Figure 3).

Taken together, the CANN-STP WM model successfully reproduces all the empirical results (Figure 5-6). The model demonstrates that the changes on the ratio of synaptic efficacies of neuron groups, which are caused by the dynamic perturbation during the delay period, result in the manipulation of relative memory strength of WM items.

Figure 6 about here

Model predictions

Our experiments only tested "Order reversal" manipulation using 200 ms and 500 ms time differences (Figure 3). We further used the model to predict how systematic changes in the correlation time window between the two probe sequences, in the

same- or reversed-order conditions, would affect the recency effect (Figure 7). First, when the time lag is close to zero, both the same- and reversed-order conditions show disrupted recency effect and tend to shift to a nonsignificant trend of primary effect, similar to "Synchronization" conditions. Second, when the time lag is too large, both conditions show the recency effect, similar to "Baseline" condition. Most interestingly, when the time lag is within [110 ms, 210 ms], a range close to the temporal constant of STD, i.e., $\left(0.5 \sim 1\right)*\tau_d$, the recalling performance for the same- or reversed-order conditions begin to display divergence, i.e., recency and primacy effects for the same-order and reversed-order conditions, respectively. Therefore, the temporal window for efficient WM manipulation is determined by the STD time constant.

Figure 7 about here

Discussion

We developed a "dynamic perturbation" approach to manipulate the relative memory strength of a list of items held in human WM, by interfering with their neural dynamics in a temporally correlated or uncorrelated way. Six experiments confirm the effectiveness of this method in multi-item WM manipulation. A computational model incorporating CANN and STP principles further reproduces the empirical findings. Importantly, the model demonstrates that this "dynamic perturbation" induces changes in the relative synaptic efficacies of WM items and eventually causes the modulation of their relative memory strength. Taken together, the temporal dynamics of WM neural representation, determined by both active-state and STP-based hidden-state, plays a fundamental role in mediating the storage of multiple items. Our results also provide a promising non-invasive behavioral approach to manipulate human WM.

It has been long posited that temporary maintenance of information relies on a refreshing process (Awh et al., 1998; Baddeley, 2003; Camos et al., 2018; Oberauer, 2019). Animal recordings and human neuroimaging studies show memory-related neural reactivations during retention (Skaggs and Mcnaughton, 1996; Compte, 2000; Wang, 2001; Foster and Wilson, 2006; Siegel et al., 2009; Michelmann et al., 2016; Huang et al., 2018; Liu et al., 2019; Schuck and Niv, 2019), supporting an active-state WM view. Meanwhile, neural response are also known to induce changes in the synaptic efficacy via STP mechanisms (Mongillo et al., 2008; Barak and Tsodyks, 2014; Mi et al., 2017; Trübutschek et al., 2017). Specifically, the synaptic efficacy, given its slow temporal properties, could remain for a while to preserve memory information without relying on sustained neural firing, namely a hidden-state WM view (Stokes, 2015; Wolff et al., 2017; Miller et al., 2018). In fact, the two theories could be well integrated into a framework where information is maintained in the synaptic weights between reactivations through STP principles, and the dynamic interplay between the active-state and hidden-state over the delay period ultimately leads to memory maintenance. Interestingly, sustained firing and hidden-state have recently been posited to be involved in manipulation and passive storage of WM information, respectively (Trübutschek et al., 2017; Masse et al., 2018, 2020). Hence,

manipulation of stored information might require reinstatement of the hidden-state

representation into activated state (Stokes, 2015; Trübutschek et al., 2019; Masse et

al., 2020). Here, by applying certain temporal perturbation patterns to interfere with

the active-state of the WM network, we modulate the hidden-state via STP principles

and alter the memory strength.

Holding a list of items in WM elicits an item-by-item sequential reactivation

pattern (Siegel et al., 2009; Heusser et al., 2016; Michelmann et al., 2016;

Bahramisharif et al., 2018; Huang et al., 2018; Liu et al., 2019; Schuck and Niv,

2019), which arises from an ongoing competition between items through lateral

inhibition and shared inhibitory inputs (Fino and Yuste, 2011; Kim et al., 2017; Mi et

al., 2017). During this rivalry process, each item takes turns to reactivate and

accordingly strengthen its respective synaptic efficacy which slowly builds up and

decays over the retention interval. Therefore, the dynamic competition between items

and the resulted memory trace held in the synaptic weights would eventually

determine their relative memory strength. Here, the two WM items are perturbed in a

temporally correlated manner so that their dynamic competition is modulated,

initiating changes in their relative memory strength. In contrast, temporally

uncorrelated perturbation, i.e., baseline condition, given its lack of stable influence on

the competition between WM items, shows no modulation effects. Moreover, it is

worth noting that the luminance sequences used in our experiment were generated

anew in each trial, and the manipulation thus does not depend on specific sequences

but on the temporal associations between them.

We build a computational model that combines CANN and STP to explore the

dynamics of the WM neural network– both neural firing rate (active state) and

synaptic efficacy (hidden state) – and how the dynamics leads to memory strength

changes. In this model, STF and STD, as two forms of STP, play different functions,

i.e., STF serves to hold memory information and STD controls the time window for

effective manipulation. The dynamic sequences brought by the flickering probes

continuously perturb the network state and lead to changes in synaptic efficacies and

memory performance. When items are independently disturbed in time (Baseline), their competition are not affected and so the recency effect keeps intact, whereas synchronous perturbance would instead change the competition, which in turn modulates their relative synaptic weights and disrupts recency effect (Synchronization). In the case of same-order condition, the perturbation strengthens the relative synaptic efficacies for the two consecutively loaded WM items, resulting in the recency effect. In contrast, under the reversed-order condition, the flipped perturbation pattern indeed reverses the relative synaptic efficacies, i.e.., $\langle Ju_2 \rangle > \langle Ju_1 \rangle$ becomes $\langle Ju_2 \rangle < \langle Ju_1 \rangle$, leading to a shift from recency to primacy effect. Notably, in both cases, the temporal correlation of the two probe sequences needs to fall into the STD temporal range for the manipulation to be effective. How to understand this phenomenon? Previous work (Mi et al., 2017), using STP-based mechanism to explain WM capacity, demonstrates that the time separation between consecutive WM is close to the STD time constant. Therefore, when two temporal sequences are applied to perturb the order of memorized items, their correlation time needs to fall into the STD time constant, so that the changes they induce to synaptic efficacy can effectively disturb the retrieval performances of two memory items. Together, our model sheds light on a challenging WM issue and provides promising ways to efficiently manipulate WM from brain's perspective.

Previous studies, by using behavioral or non-invasive stimulation approaches in the encoding or delay period, have also successfully modulated the recalling performance (Sauseng et al., 2009; Rose et al., 2016; Clouter et al., 2017; Hanslmayr et al., 2019; Martorell et al., 2019). Here, benefiting from using the time-resolved luminance sequences to disturb the WM network dynamics, our study largely differs from previous interference findings in two aspects. First, previous studies are mainly focused on the modulation of general memory performance while our study aims at particularly manipulating the relative memory strength of WM items (i.e., recency). Second, by using temporally correlated luminance sequences, our approach could precisely modulate the ongoing temporal relationship between items (e.g.,

synchronization, 200 ms or 500 ms order reversal). Generally speaking, our

"synchronization manipulation" condition shares similar motivations as associative

memory paradigms, during which pairing two stimuli would lead to changes in their

association in memory (Shohamy and Wagner, 2008; Duncan et al., 2012; Gershman

et al., 2017). Meanwhile, instead of long-term memory, our manipulation modulates

working memory on a trial-by-trial basis. In addition, we adopted a bottom-up

approach by modulating the luminance of task-irrelevant color probes, thus largely

different from previous interference methods.

Finally, recent studies advocate a close connection between WM and attention,

and WM has been proposed to be an internal attentional process that shares similar

properties and operation principles as attention (Chun et al., 2011; D'Esposito and

Postle, 2015; Oberauer, 2019). For instance, the recency effect could be readily

explained in terms of attentional gain, such that the lately presented item enters the

focus of attention (FOA) and obtains more attention (Oztekin et al., 2010; Oberauer

and Lin, 2017). In general, WM and attention are likely to be mediated via largely

overlapping neural networks and similar mechanism, making them hard to be fully

disentangled. In fact, the memory-related reactivation might also reflect attentional

modulation of neural response that would affect subsequent attentional performance

(Soto et al., 2005; Olivers et al., 2006; Mallett and Lewis-Peacock, 2018). Similarly,

attention-induced response might leave memory traces to affect following memory

behavior (Cowan et al., 2005; Rose et al., 2016; Camos et al., 2018; deBettencourt et

al., 2019). Moreover, attention has been shown to sample different objects

sequentially and rhythmically, also resembling the dynamic reactivation pattern in

WM (Jensen et al., 2014; Jia et al., 2017; Fiebelkorn and Kastner, 2019). Therefore,

the "dynamic perturbation" might impact the attentional process as well and thus

constitutes a promising way to also manipulate attention, planning and decision

making.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

We thank Dr. David Poeppel, Dr. Fang Fang, and Dr. Nai Ding for their helpful

comments, as well as anonymous reviews to our previous submission. This work was

supported by the National Natural Science Foundation of China (31930052 to H.L.,

31771146, 11734004 to Y.Y. Mi), and Beijing Municipal Science & Technology

Commission (Z181100001518002), Beijing Nova Program (Z181100006218118, Y. Y.

Mi), Guangdong Province with Grant No. 2018B030338001.

Methods

Participants

One hundred and ninety-six participants (90 males, age ranging from 18 to 26 years) took part in eight experiments (Expt. 1–5 and Expt. 2-IC in main texts, Expt. 6 (probe-absent experiment), Expt. 7 (spatial location experiment) in Supplementary Materials). One subject in Expt. 1, two in Expt. 2, two in Expt. 2-IC, three in Expt. 3, four in Expt. 4, three in Expt. 5, one in Expt. 6 and five subjects in Expt. 7 were excluded due to their overall low memory performance, or self-reported strategies, or not finishing the whole experiment, resulting in 27, 24, 24, 27, 27, 25, 16, 26 subjects for Expt. 1 to Expt. 5, Expt. 2-IC, and Expt. 6-7 respectively. All the participants were naïve to the purpose of the experiments and had normal or corrected-to-normal vision with no history of neurological disorders, and have provided written informed consent before the experiments. All experiments were carried out in accordance with the Declaration of Helsinki and have been approved by the Research Ethics Committee at Peking University.

Stimuli and tasks

Experiment paradigm

Participants sat in a dark room, 58 cm in front of a CRT monitor (Display++ monitor in Experiment 4) with 100 Hz refresh rate and a resolution of 1024 * 768. Subjects performed a sequence memory task, with their head stabilized on a chin rest. Each trial consisted of three periods: encoding, maintaining, and recalling (Fig. 1A). In the "encoding period", subjects were serially presented with two bars (Experiment 1, 2, 2-IC, 3, 4, 5, 6, 7) displayed at the center. Each bar stimulus lasted for 1 s with a 0.5 s inter-stimulus interval and had different orientation and color features. Subjects were instructed to memorize the orientations of the bar stimuli. In the "maintaining period", participants performed a central fixation task by monitoring an abrupt luminance change (within 25% of total trials) and reported if they detected changes after the maintenance period. All trials were included into further analysis. Finally, in the "recalling period", an instruction cue was presented for 1.5 s to indicate which

orientation should be recalled (1^{st} or 2^{nd} orientation), and participants needed to rotate a horizontal white bar by pressing corresponding keys to the memorized orientation as precise as possible, without time limit.

Experiment 1–2 ("temporal synchronization" manipulation)

In each trial, after a 0.5 s fixation period, participants performed a two-item sequence memory task. The 1^{st} orientation was chosen randomly between 0° and 180°, and the 2^{nd} orientation was generated based on the 1^{st} orientation with the tiled angles of ±17°, ±24°, ±38°, ±52°, ±66°, ±80° (Liu and Becker, 2013). The color of the 1^{st} and 2^{nd} bar was different (randomly selected from red, green, or blue in each trial). During the maintenance period, after a 0.6-1 s blank interval, two task-irrelevant discs (3° in radius) that had the color feature of either the 1^{st} or 2^{nd} memorized bar, referred as the 1^{st} memory-related probe (WM1) and 2^{nd} memory-related probe (WM2) respectively, were displayed at the left or right side of the fixation (7° visual angle) for 5 secs. The color (red, green, or blue) and the spatial locations (left or right) of the two probes (WM1 and WM2) were carefully balanced across trials. Importantly, throughout the 5 s maintenance period, the luminance of the two color discs (WM1 and WM2) was continuously modulated according to two 5 s temporal sequences, respectively (i.e., Seq1 for WM1, Seq2 for WM2), ranging from dark (0 cd/m^2) to bright (15 cd/m^2). The Seq1 and Seq2 were generated anew in each trial.

In Experiment 1, the relationship between Seq1 and Seq2 during the maintenance period varied in three ways: Baseline, alpha-band synchronization (A-Sync), or full-spectrum synchronization (F-Sync). For Baseline condition, Seq1 and Seq2 were two independently generated random time series (Seq1 ≠ Seq2). For F-Sync condition, Seq1 and Seq2 were the same random temporal series (Seq1 = Seq2). For A-Sync condition, Seq1 and Seq2 were the same alpha-band (8-11 Hz) time series (Seq1 = Seq2). Experiment 1 had 216 trials in total (72 trials for each of the three synchronization conditions), divided into three blocks. The three synchronization conditions were randomly mixed within each experimental block.

Experiment 2 only examined the Baseline and A-Sync conditions. Experiment 2 had 144 trials in total (72 trials for each of the two synchronization conditions), divided into three blocks. The two synchronization conditions were randomly mixed within each experimental block.

Experiment 2-IC employed the same design as Experiment 2 (Baseline condition and A-Sync condition), except that the color probes during the delay period had memory-irrelevant colors. Specifically, when the to-be-remembered two bars were blue and red in color, the color probes presented in the delay period would be set to yellow and green, and vice versa.

Experiment 3–5 ("order reversal" manipulation)

Experiment 3–5 employed the same paradigm as Experiment 1–2, except that the temporal relationship between Seq1 and Seq2 during the maintenance period were manipulated in a different manner. Specifically, Seq1 was a randomly generated temporal sequence, and Seq2 was a temporal shifted version of Seq1. Specifically, Seq1 either leads or lags Seq2 by 200 ms, corresponding to the "same" and "reversed" conditions, respectively, compared to the stimulus sequence. Furthermore, in order to replicate findings of Experiment 1-2, Baseline and F-Sync conditions were added to Experiment 3 and 4, respectively. Experiment 3 included Same, Reversed, and Baseline conditions, and Experiment 4 included Same, Reversed, and F-Sync conditions. We then assessed the behavioral results using a two-sample (Experiment 3 and 4) two-way (Position*Sync) repeated ANOVA analysis to examine the overall manipulation effect. Each condition had 72 trials, resulting in 216 trials for Experiment 3 and 4, respectively.

Experiment 5 only examined Same and Reversed conditions. Moreover, rather than 200 ms lag in Experiment 3-4, the Seq1-Seq2 temporal lag was set to 500 ms in Experiment 5. Each condition had 72 trials, resulting in 144 trials for Experiment 5.

Data analysis

A mixture-modelling analysis (Bays et al., 2009) was used to quantify the multi-item

WM behavioral performance for each of the item, by quantifying the Von Mises

distribution variability of the reported error. The three-parameter mixture model

calculates the precision of the reports, as well as the respective proportions of

target, distractor and guess responses. In the present study, we used the target

probability (see the precision estimation results in Supplementary Materials) that has

been widely used to estimate memory performance (Bays et al., 2009; Gorgoraptis et

al., 2011; Van Ede et al., 2018). Moreover, the target probability was further

normalized by using an arcsine square root transformation (Bartlett et al., 1936). The

normalized target probabilities were further used to examine the effect of temporal

manipulation on multi-item WM.

The Meta-analysis across experiments were performed by using Comprehensive

Meta-Analysis Software (CMA) across all 8 experiments (Expt. 1-5 and Expt. 2-IC in

main text, Expt. 6-7 in Supplementary Materials). We calculated the recency effect for

Baseline condition across Experiment 1, 2, 3, 5, 6, 7, 2-IC, the recency effect for

Synchronization condition across Experiment 1, 2, 4, 6, 7, and the interaction effect

between Baseline and Synchronization conditions across Experiment 1, 2, 7.

Computational modelling

Continuous attractor neural network model (CANN)

In this model, excitatory neurons that represent specific orientation (θ , $(-\pi/2,+\pi/2]$)

are aligned in the network according to its preferred orientation. They are connected

to each other ($J(\theta,\theta')$ as the synaptic strength) and are further reciprocally connected

to a global inhibitory neural pool (J_{EI} as the synaptic strength). Specifically, for each

excitatory neuron at θ , its dynamics of synaptic inputs $h_E(\theta,t)$ is calculated as:

$$\tau \frac{\partial h_E(\theta,t)}{\partial t} = -h_E(\theta,t) + \int_{-\pi/2}^{+\pi/2} J(\theta,\theta')u(\theta',t)x(\theta',t)r_E(\theta',t)d\theta'$$

608
$$- J_{EI}r_I + I_{ext}(\theta,t) + I_0 + \sigma_1\zeta_1(\theta,t)$$

In other words, the dynamics of synaptic current of excitatory neuron at θ

($h_E(\theta,t)$) would be decided by its own dynamics, its stimulus input ($I_{ext}(\theta,t)$), inputs

from other excitatory neurons ($r_E(\theta',t)$), background activities ($I_0 + \sigma_1\varsigma_1(\theta,t)$), and

inhibitory inputs (r_I). For each neuron, the relationship between firing rate and

synaptic current is characterized by $r(h) = \alpha Ln(1 + \exp(h/\alpha))$.

The dynamics of inhibitory neuron is given by $\tau\dfrac{dh_I}{dt} = -h_I + J_{IE}\displaystyle\int_{-\pi/2}^{+\pi/2} r_E(\theta,t)$, where

r_I and h_I denote the firing rate and synaptic current of the inhibitory neural pool,

and $r_I(h_I) = \alpha\ln\left(1 + \exp(h_I/\alpha)\right)$.

Short-term synaptic plasticity (STP)

To model STP effect, $u(\theta,t)$ and $x(\theta,t)$ are used to quantify the short-term

facilitation (STF) and short-term depression (STD) effects, respectively. In fact,

$u(\theta,t)$ represents the release probability of neural transmitters, and $x(\theta,t)$ denotes

the fraction of resources available after neurotransmitter depletion, and their dynamics

are given by:

$$\frac{\partial u(\theta,t)}{\partial t} = \frac{-u(\theta,t)}{\tau_f} + U\left(1 - u(\theta,t)\right)r_E(\theta,t),$$

$$\frac{\partial x(\theta,t)}{\partial t} = \frac{1 - x(\theta,t)}{\tau_d} - u(\theta,t)x(\theta,t)r_E(\theta,t).$$

625   where τ_f and τ_d are the time constants of STF and STD, respectively, and U

626   determines the increment of u due to neuronal firing. We followed the settings in

627   previous STF-dominating WM models, i.e., $\tau_f \gg \tau_d$ and a large value of U

628   ($\tau_d = 0.214s$, $\tau_f = 4.9s$, U $= 0.45$).

630   *Model simulation*

We mimicked the whole experimental procedure to do the model simulation. First, during the encoding period, two color bars are loaded into the network sequentially, with each lasting 1 s and the time interval between them being 0.5 s. This is simulated as applying inputs $I_{ext}(\theta_1, t)$ and $I_{ext}(\theta_2, t)$ to the CANN, corresponding to the 1st and 2nd orientation features, sequentially. Next, after 0.6-1.4 s, during the delay period, given that the color probes are presumably bound to its associated orientation feature, the 5 sec flickering color probes are simulated by applying a weak, continuous inputs to the CANN respectively. Afterwards, we presented a recalling signal as a loaded item for 1.5 s to retrieve the corresponding memory. Crucially, we mimicked different dynamic perturbation conditions by applying temporal sequences with the corresponding characteristics (e.g., Baseline, Synchronization, Order reversal, etc.) as used in behavioral experiments.

In the recall phase, we used the population vector method to read out the orientation value retrieved by the network, and then followed the same measurement used in psychophysical experiments to calculate the recalling performance. We defined neurons with preferred orientations in the range of $|\theta - \theta_1| < 7^o$ as the neuronal group 1, and neurons with preferred orientations in the range of $|\theta - \theta_2| < 7^o$ as the neuronal group 2. We also calculated the mean firing rates and the means of synaptic efficacies of two neuronal groups.

The simulation was performed for each experimental condition, respectively. Specifically, 6 pairs of orientation angles with difference (17°, 24°, 38°, 52°, 66°, and 80°) were used for the simulation. For each condition (probe-absent, baseline, synchronization manipulation, same-order, reversed-order, etc.), there were 20 simulation runs and each run contained 200 simulation trials. The results of each run, similar as that of one subject, were then calculated by averaging across the 200 trials that belonged to this run.

For Baseline condition, two independent probe sequences were presented to the 1st and 2nd neuron group during the delay period (Figure 6A). For Synchronization conditions, the same temporal sequences were presented, i.e., seq1 = seq2, during the

delay period, either as a white-noise temporal sequence (F-sync condition; Figure 6B)

or an alpha-band filtered temporal sequence (A-sync condition; Figure 6C). For the

order reversal manipulation, seq1 and seq 2 were a 200 ms temporally shifted version

of each other, either seq1 lead seq2 (Same-order condition; Figure 6D) or seq1 lags

seq2 (Reversed-order condition; Figure 6E).

For model predictions, the same-order and reversed-order conditions were

simulated using varying time lags (10, 50, 90, 110, 130, 150, 170, 190, 210, 230, 270,

310, and 350 ms).

See more model details in Supplementary Materials.

References

Awh E, Jonides J, Reuter-lorenz PA (1998) Rehearsal in Spatial Working Memory. J Exp Psychol Hum Percept Perform 24:780–790.

Baddeley A (2003) Working memory: Looking back and looking forward. Nat Rev Neurosci.

Baddeley A (2012) Working Memory: Theories, Models, and Controversies. Annu Rev Psychol 63:1–29.

Bahramisharif A, Jensen O, Jacobs J, Lisman J (2018) Serial representation of items during working memory maintenance at letter-selective cortical sites. PLOS Biol 16:e2003805 Available at: http://dx.plos.org/10.1371/journal.pbio.2003805.

Barak O, Tsodyks M (2014) Working models of working memory. Curr Opin Neurobiol 25:20–24 Available at: http://dx.doi.org/10.1016/j.conb.2013.10.008.

Bartlett AMS, Supplement S, Society S (1936) The square root transformation in analysis of variance. Suppl to J R Stat Soc 3:68–78.

Bays PM, Catalao RFG, Husain M (2009) The precision of visual working memory is set by allocation of a shared resource. J Vis 9:7–7 Available at: http://jov.arvojournals.org/Article.aspx?doi=10.1167/9.10.7.

Ben-Yishai R, Lev Bar-Or R, Sompolinsky H (1995) Theory of orientation tuning in visual. Pnas 92:3844–3848.

Blumenfeld B, Bibitchkov D, Tsodyks M (2006) Neural network model of the primary visual cortex: From functional architecture to lateral connectivity and back. J Comput Neurosci 20:219–241.

Brody CD, Romo R, Kepecs A (2003) Basic mechanisms for graded persistent activity: Discrete attractors, continuous attractors, and dynamic representations. Curr Opin Neurobiol 13:204–211.

Burgess N, Hitch GJ (1999) Memory for serial order: A network model of the phonological loop and its timing. Psychol Rev 106:551–581.

Camos V, Johnson M, Loaiza V, Portrat S, Souza A, Vergauwe E (2018) What is attentional refreshing in working memory? Ann N Y Acad Sci 1424:19–32.

Chun MM, Golomb JD, Turk-Browne NB (2011) A Taxonomy of External and Internal Attention. Annu Rev Psychol 62:73–101.

Clouter A, Shapiro KL, Hanslmayr S (2017) Theta Phase Synchronization Is the Glue that Binds Human Associative Memory. Curr Biol 27:3143-3148.e6 Available at: https://doi.org/10.1016/j.cub.2017.09.001.

Compte A (2000) Synaptic Mechanisms and Network Dynamics Underlying Spatial Working Memory in a Cortical Network Model. Cereb Cortex 10:910–923.

Compte A, Brunel N, Goldman-rakic PS, Wang X (2000) Dynamics Underlying Spatial Working Memory in a Cortical Network Model e. :910–923.

Cowan N (2001) The magical number 4 in short-term memory: A reconsideration of mental storage capacity. Behav Brain Sci.

Cowan N, Elliott EM, Saults SJ, Morey CC, Mattox S, Hismjatullina A, Conway ARA (2005) On the capacity of attention: Its estimation and its role in working memory and cognitive aptitudes. Cogn Psychol 51:42–100.

D'Esposito M, Postle BR (2015) The Cognitive Neuroscience of Working Memory.

Ssrn.

deBettencourt MT, Keene PA, Awh E, Vogel EK (2019) Real-time triggering reveals
concurrent lapses of attention and working memory. Nat Hum Behav 3 Available
at: http://dx.doi.org/10.1038/s41562-019-0606-6.

Dell GS, Burger LK, Svec WR (1997) Language production and serial order: A
functional analysis and a model. Psychol Rev 104:123–147.

Doyon J, Penhune V, Ungerleider LG (2003) Distinct contribution of the cortico-
striatal and cortico-cerebellar systems to motor skill learning. Neuropsychologia
41:252–262.

Duncan K, Sadanand A, Davachi L (2012) Memory's Penumbra: Episodic memory
decisions induce lingering mnemonic biases. Science (80-) 337:485–487.

Fiebelkorn IC, Kastner S (2019) A Rhythmic Theory of Attention. Trends Cogn Sci
23:87–101 Available at: https://doi.org/10.1016/j.tics.2018.11.009.

Fiebig XF, Lansner XA (2017) A spiking working memory model based on Hebbian
short-term potentiation. J Neurosci 37:83–96.

Fino E, Yuste R (2011) Dense inhibitory connectivity in neocortex. Neuron 69:1188–
1203 Available at: http://dx.doi.org/10.1016/j.neuron.2011.02.025.

Foster DJ, Wilson MA (2006) Reverse replay of behavioural sequences in
hippocampal place cells during the awake state. Nature 440:680–683.

Fusi S (2008) A quiescent working memory. Science (80-) 319:1495–1496.

Gershman SJ, Monfils M-H, Norman KA, Niv Y (2017) The computational nature of
memory modification. Elife 6:1–41.

Giraud AL, Poeppel D (2012) Cortical oscillations and speech processing: Emerging
computational principles and operations. Nat Neurosci 15:511–517 Available at:
http://dx.doi.org/10.1038/nn.3063.

Gorgoraptis N, Catalao RFG, Bays PM, Husain M (2011) Dynamic updating of
working memory resources for visual objects. J Neurosci 31:8502–8511
Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.0208-
11.2011.

Gyorgy B (2002) Theta oscillations in the hippocampus. Neuron 33:325–340
Available at: papers3://publication/uuid/90B140FB-AC03-4F7C-9968-
EF84D84A3009.

Hanslmayr S, Axmacher N, Inman CS (2019) Modulating Human Memory via
Entrainment of Brain Oscillations. Trends Neurosci 42:485–499 Available at:
https://doi.org/10.1016/j.tins.2019.04.004.

Herweg NA, Solomon EA, Kahana MJ (2020) Theta Oscillations in Human Memory.
Trends Cogn Sci 24:208–227 Available at:
https://doi.org/10.1016/j.tics.2019.12.006.

Heusser AC, Poeppel D, Ezzyat Y, Davachi L (2016) Episodic sequence memory is
supported by a theta-gamma phase code. Nat Neurosci 19:1374–1380.

Horn D, Opher I (1996) Temporal segmentation in a neural dynamic system. Neural
Comput 8:373–389.

Huang Q, Jia J, Han Q, Luo H (2018) Fast-backward replay of sequentially
memorized items in humans. Elife 7:1–21.

Hyun J seok, Woodman GF, Vogel EK, Hollingworth A, Luck SJ (2009) The comparison of visual working memory representations with perceptual inputs. J Exp Psychol Hum Percept Perform 35:1140–1160.

Jensen O (2002) Oscillations in the alpha band (9-12 Hz) increase with memory load during retention in a short-term memory task. Cereb Cortex 12:877–882.

Jensen O, Gips B, Bergmann TO, Bonnefond M (2014) Temporal coding organized by coupled alpha and gamma oscillations prioritize visual processing. Trends Neurosci 37:357–369 Available at: http://dx.doi.org/10.1016/j.tins.2014.04.001.

Jia J, Liu L, Fang F, Luo H (2017) Sequential sampling of visual objects during sustained attention. PLoS Biol 15:1–20 Available at: http://dx.doi.org/10.1371/journal.pbio.2001903.

Johnson JS, Hollingworth A, Luck SJ (2008) The role of attention in the maintenance of feature bindings in visual short-term memory. J Exp Psychol Hum Percept Perform 34:41–55.

Jones T, Oberauer K (2013) Serial-position effects for items and relations in short-term memory. Memory 21:347–365.

Kim SS, Rouault H, Druckmann S, Jayaraman V (2017) Ring attractor dynamics in the Drosophila central brain. Science (80-) 356:849–853.

Klimesch W (1999) EEG alpha and theta oscillations reflect cognitive and memory performance: a rKlimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. Brain Research Reviews, 29(2-3), 169–195. doi:10.1016/S016. Brain Res Rev 29:169–195 Available at: http://www.sciencedirect.com/science/article/pii/S0165017398000563.

Lisman JE, Idiart MAP (1995) Storage of 7 ± 2 short-term memories in oscillatory subcycles. Science (80-) 267:1512–1515.

Liu T, Becker MW (2013) Serial consolidation of orientation information into visual short-term memory. Psychol Sci 24:1044–1050.

Liu Y, Dolan RJ, Kurth-Nelson Z, Behrens TEJ (2019) Human replay spontaneously reorganizes experience. Cell 178:640–652.

Luck S, Vogel J, Edward K (1997) The capacity of visual working memory for features and conjuctions. Nature 390:279–281.

Lundqvist M, Rose J, Herman P, Brincat SLL, Buschman TJJ, Miller EKK (2016) Gamma and beta bursts underlie working memory. Neuron 90:152–164.

Mallett R, Lewis-Peacock JA (2018) Behavioral decoding of working memory items inside and outside the focus of attention. Ann N Y Acad Sci 1424:256–267.

Markram H, Wang Y, Tsodyks M (1998) Differential signaling via the same axon of neocortical pyramidal neurons. Proc Natl Acad Sci U S A.

Martorell AJ, Paulson AL, Suk HJ, Abdurrob F, Drummond GT, Guan W, Young JZ, Kim DNW, Kritskiy O, Barker SJ, Mangena V, Prince SM, Brown EN, Chung K, Boyden ES, Singer AC, Tsai LH (2019) Multi-sensory Gamma Stimulation Ameliorates Alzheimer's-Associated Pathology and Improves Cognition. Cell 177:256-271.e22 Available at: https://doi.org/10.1016/j.cell.2019.02.014.

Masse NY, Rosen MC, Freedman DJ (2020) Reevaluating the Role of Persistent

Neural Activity in Short-Term Memory. Trends Cogn Sci 24:242–258 Available at: https://doi.org/10.1016/j.tics.2019.12.014.

Masse NY, Yang GR, Song HF, Wang X-J, Freedman DJ (2018) Circuit mechanisms for the maintenance and manipulation of information in working memory. bioRxiv:305714 Available at: https://www.biorxiv.org/content/early/2018/05/21/305714.

Mi Y, Katkov M, Tsodyks M (2017) Synaptic correlates of working memory capacity. Neuron 93:323–330 Available at: http://dx.doi.org/10.1016/j.neuron.2016.12.004.

Michelmann S, Bowman H, Hanslmayr S (2016) The Temporal Signature of Memories: Identification of a General Mechanism for Dynamic Memory Replay in Humans. PLoS Biol 14:1–27.

Miller EK, Lundqvist M, Bastos AM (2018) Working Memory 2.0. Neuron.

Mongillo G, Barak O, Tsodyks M (2008) Synaptic theory of working memory. Science (80-) 319:1543–1546.

Oberauer K (2019) Working Memory and Attention – A Conceptual Analysis and Review. J Cogn 2.

Oberauer K, Lin HY (2017) An interference model of visual working memory. Psychol Rev 124:21–59.

Olivers CNL, Meijer F, Theeuwes J (2006) Feature-based memory-driven attentional capture: Visual working memory content affects visual attention. J Exp Psychol Hum Percept Perform 32:1243–1265.

Oztekin I, Davachi L, Mcelree B (2010) Are representations in working memory distinct from representations in long-term memory? Neural evidence in support of a single store. Psychol Sci 21:1123–1133.

Parthasarathy A, Tang C, Herikstad R, Cheong LF, Yen SC, Libedinsky C (2019) Time-invariant working memory representations in the presence of code-morphing in the lateral prefrontal cortex. Nat Commun 10:1–11 Available at: http://dx.doi.org/10.1038/s41467-019-12841-y.

Raffone A, Wolters G (2001) A cortical mechanism for binding in visual working memory. J Cogn Neurosci.

Romani S, Tsodyks M (2015) Short-term plasticity based network model of place cells dynamics. Hippocampus 25:94–105.

Rose NS, LaRocque JJ, Riggall AC, Gosseries O, Starrett MJ, Meyering EE, Postle BR (2016) Reactivation of latent working memories with transcranial magnetic stimulation. Science (80-) 354:1136–1139.

Sauseng P, Klimesch W, Heise KF, Gruber WR, Holz E, Karim AA, Glennon M, Gerloff C, Birbaumer N, Hummel FC (2009) Brain oscillatory substrates of visual short-term memory capacity. Curr Biol 19:1846–1852 Available at: http://dx.doi.org/10.1016/j.cub.2009.08.062.

Schuck NW, Niv Y (2019) Sequential replay of nonspatial task states in the human hippocampus. Science (80-) 364.

Seeholzer A, Deger M, Gerstner W (2019) Stability of working memory in continuous attractor networks under the control of shortterm plasticity. Available at:

http://dx.doi.org/10.1371/journal.pcbi.1006928.

Shohamy D, Wagner AD (2008) Integrating memories in the human brain:
Hippocampal-midbrain encoding of overlapping events. Neuron 60:378–389
Available at: http://dx.doi.org/10.1016/j.neuron.2008.09.023.

Siegel M, Warden MR, Miller EK (2009) Phase-dependent neuronal coding of objects
in short-term memory. Proc Natl Acad Sci 106:21341–21346 Available at:
http://www.pnas.org/lookup/doi/10.1073/pnas.0908193106.

Skaggs WE, Mcnaughton BL (1996) Replay of neuronal firing sequences in rat
hippocampus during sleep following spatial experience. Science (80-)
271:1870–1873.

Soto D, Heinke D, Humphreys GW, Blanco MJ (2005) Early, involuntary top-down
guidance of attention from working memory. J Exp Psychol Hum Percept
Perform 31:248–261.

Stokes MG (2015) "Activity-silent" working memory in prefrontal cortex: A dynamic
coding framework. Trends Cogn Sci 19:394–405 Available at:
http://dx.doi.org/10.1016/j.tics.2015.05.004.

Trübutschek D, Marti S, Ojeda A, King JR, Mi Y, Tsodyks M, Dehaene S (2017) A
theory of working memory without consciousness or sustained activity. Elife
6:1–29.

Trübutschek D, Marti S, Ueberschär H, Dehaene S (2019) Probing the limits of
activity-silent non-conscious working memory. Proc Natl Acad Sci U S A
116:14358–14367.

Van Ede F, Chekroud SR, Stokes MG, Nobre AC (2018) Decoding the influence of
anticipatory states on visual perception in the presence of temporal distractors.
Nat Commun 9 Available at: http://dx.doi.org/10.1038/s41467-018-03960-z.

Wang X-J (2001) Synaptic reverberations underlying mnemonic persistent activity.
Trends Neurosci 24:455–463.

Wolff MJ, Jochim J, Akyürek EG, Stokes MG (2017) Dynamic hidden states
underlying working-memory-guided behavior. Nat Neurosci 20:864–871.

Figures

Figure 1

Figure 1.

"Dynamic perturbation" approach: temporal association during memory

retention

(A) Experimental paradigm. In each trial, in the "encoding period", two bar stimuli

(item 1, item 2) were presented serially at the center. Subjects needed to memorize the

orientations of the two bars (1st and 2nd orientation). In the "recalling period",

participants rotated a horizontal bar to the memorized orientation based on the

instruction (recalling the 1st or the 2nd orientation). During the 5 s "maintaining

period", subjects performed a central fixation task, while two discs that had color

feature of either the 1st or the 2nd memorized bars (1st memory-related probe, WM1;

2nd memory-related probe, WM2) were displayed at left or right side of the fixation. The color (red, blue, or green) and the spatial locations (left or right) of the two probes (WM1 and WM2) were balanced across trials. Throughout the 5 s retention period, the luminance of the two color discs (WM1 and WM2) was continuously modulated according to two 5 sec temporal sequences respectively (Seq1 for WM1, Seq2 for WM2). Note that the luminance sequences were generated anew in each trial. (B) Temporal synchronization of WM1 and WM2 during maintenance. Upper: Illustration of 500 ms segment of the 5 s Seq1 (denoted in dark line) and Seq2 (denoted grey line). Lower: Spectrum of the 5 s Seq1 (dark) and Seq2 (grey). The temporal relationship between Seq1 (black) and Seq2 (grey) varied in one of the three ways (left: Baseline; Middle: A-Sync; Right: F-Sync). In Baseline condition (Left), Seq1 and Seq2 were two independently generated random time series. In A-Sync condition (Middle), Seq1 and Seq2 shared the same alpha-band (8-11 Hz) time series. In F-Sync condition (Right), Seq1 and Seq2 were the same random time series. Note that Seq1 and Seq2 were generated anew in each trial.

Figure 2

Figure 2

"Temporal synchronization" manipulation disrupts recency effect

A probabilistic mixture model was used to quantify the memory performance for the

1st and the 2nd item. The target probability results were further converted using an

arcsine square root transformation to make it normally distributed. (A) Experiment 1

results (N = 27). Grand averaged (mean±SEM) normalized target probability for the

1st (light grey) and 2nd (dark grey) orientation for Baseline, A-Sync, and F-Sync

conditions. (B) Experiment 2 (N = 24) and Experiment 2-IC (irrelevant color probes)

(N = 25) results. Grand averaged (mean±SEM) normalized target probability for the

1st (light grey) and 2nd (dark grey) orientation for Baseline and A-Sync conditions, in

Experiment 2 (left) and Experiment 2-IC (right). (C) Grand averaged (mean±SEM)

recency effect (2nd minus 1st) for Baseline (white box), A-Sync (rightward checked

box), and F-Sync (leftward checked box) conditions in Experiment 1, Experiment 2,

and Experiment 2-IC. Noting the significant synchronization-induced recency

disruption in Experiment 1 and 2, but not in Experiment 2-IC. (**: p<=0.05, *:

p<=0.1).

Figure 3

Figure 3

"Order reversal" manipulation disrupts recency effect

(A) Experiment 3-5 employed the same experimental paradigm as Experiment 1 (see

Figure 1A), except that the luminance sequences (Seq1 and Seq2) were modulated

differently. Specifically, Seq1 was a temporally shifted version of Seq2, either leading

("Same-order", upper panel) or lagging Seq2 ("Reversed-order", lower panel) by 200

ms. Right: the Seq1-Seq2 cross-correlation coefficients as a function of temporal lag

(-1 to 1 s). Note that the luminance sequences were generated anew in each trial. (B)

Two separate subject groups participated in Experiment 3 (N=24; Baseline, Same-

order, Reversed-order) and Experiment 4 (N=27; F-Sync, Same-order, Reversed-

order) respectively. Grand averaged (mean$\pm$SEM) recency effect (2^{nd} minus 1^{st}) for

all conditions in Experiment 3 and Experiment 4 (positive and negative valued

represents recency and primacy effect, respectively). Noting the significant decrease

in recency effect for Reversed than Same conditions in both experiments. A two-

sample two-way repeated ANOVA was performed across Experiment 3 and 4. (C)

Left: pooled recency effects (N=51; mean$\pm$SEM) for Same-order and Reversed-order

conditions across Experiment 3 and 4. Right: Grand average recency effects (N= 27;

mean$\pm$SEM) for Same-order and Reversed-order conditions in Experiment 5 (500 ms

temporal shift). (***: $p<=0.01$, **: $p<=0.05$, *: $p<=0.1$.)

Figure 4

946
Figure 4

Meta-analysis across eight experiments

Meta-analysis on recency effect across all 8 experiments (Expt. 1-5 and Expt. 2-IC in

main text, Expt. 6-7 in Supplementary Mate rials). (A) Significant recency effect for

Baseline condition (across Experiment 1, 2, 2-IC, 3, 5, 6, 7). (B) Non-significant

recency effect for Synchronization condition (across Experiment 1, 2, 4, 6, 7). (C)

Significant difference in recency effect between Baseline and Synchronization

conditions (across Experiment 1, 2, 7).

Figure 5

A   STP-based WM network model

B   Network performance in Probe-absent Condition

Figure 5

CANN-STP WM network model and simulation of probe-absent condition

(A) CANN and STP. Left: Excitatory neurons in the CANN are aligned according to

their preferred orientations $\theta \in \left(-\pi/2, +\pi/2\right]$, and they are connected in a translation-

invariant manner. All excitatory neurons are reciprocally connected to a common

inhibitory neuron pool (blue solid circle). In response to an external input $I_{ext}\left(\theta, t\right)$,

the network generates a bump response (red curve) at the corresponding location

according to stimulus strength. **Right**: Illustration of STP principle. In response to a

transient external input, the neuron encoding orientation at θ generates a transient

response (firing rate, $r(\theta, t)$), which induces changes in both the neurotransmitter

release probability $u(\theta, t)$ (red) and the fraction of available neurotransmitter

$x(\theta, t)$ (green), representing STF and STD effect, respectively. The neuronal

synaptic efficacy Jux, with J as the synaptic weight, will remain at high values for a

while, due to the larger time constant for STF than STD (i.e., $\tau_d = 0.214s$,

$\tau_f = 4.9s$).

(B) Simulation results for probe-absent condition. Left: Individual trial simulation

example. Top: Two orientations (1^{st} and 2^{nd}) at $\theta_1 = -\pi/4$ and $\theta_2 = -\pi/4 + 66^o$ (as

examples), respectively, are loaded into the network sequentially and trigger

population spikes in the corresponding neuron groups (1^{st} and 2^{nd} neuron group).

During retention, the two neuron groups still show response, due to STP, but fall to

baseline gradually. In the recalling period, a noisy signal containing partial

information of the 2^{nd} memory item (i.e., recalling the 2^{nd} orientation as an example

here) is presented and elicits the firing of the 2^{nd} neuron group. Middle: the temporal

course of synaptic efficacies of the 1^{st} (red) and 2^{nd} (blue) neuron group throughout

the trial. Bottom: the temporal course of external inputs to the 1^{st} (red) and 2^{nd} (blue)

neuron group throughout the trial. **Middle:** Grand averaged (mean$\pm$SEM, across 20

simulation runs with each having 200 simulation trials) synaptic efficacy difference

$\langle Ju_1 \rangle - \langle Ju_2 \rangle$ (top) and firing rate difference $\langle r_1 \rangle - \langle r_2 \rangle$ (bottom) between the 1^{st} and

2^{nd} neuron groups over retention. **Right**: Grand averaged (mean$\pm$SEM, across 20

simulation runs) recalling performance characterized by the normalize target

probability for the 1^{st} (red) and 2^{nd} (blue) orientation. (***: p<=0.01.)

Figure 6

Figure 6

Model simulations for different manipulation conditions

Simulation results for (A) Baseline condition, (B) F-Synchronization condition, (C)

A-Synchronization condition, (D) Same-order condition and (E) Reversed-order

condition. **Left**: Individual trial example for $\theta_1 = -\pi/4$ and $\theta_2 = -\pi/4 + 66^o$. **Middle**: Grand averaged (mean$\pm$SEM, across 20 simulation runs with each having 200 simulation trials) synaptic efficacy difference $\langle Ju_1 \rangle - \langle Ju_2 \rangle$ (top) and firing rate difference $\langle r_1 \rangle - \langle r_2 \rangle$ (bottom) between the 1st and 2nd neuron groups over retention. **Right**: Grand averaged (mean$\pm$SEM, across 20 simulation runs) recalling performance characterized by the normalize target probability for the 1st (red) and 2nd (blue) orientation. (***: p<=0.01, **: p<=0.05, *: p<=0.1.). Parameters and mathematical details of probe stimulations are given in Supplementary Materials.

1005 **Figure 7**

1006

Figure 7

Model predictions

Grand averaged (mean±SEM, across 20 simulation runs with each having 200

simulation trials) normalized target probability difference between the 1st and 2nd

orientation (2nd minus 1st) at varying time lags (10-350 ms) for (A) same-order and

(B) reversed-order conditions. τ_d (STD time constant) and $0.5\tau_d$ are marked by

dark blue arrows (214 ms) and light blue arrows (107 ms), respectively. Note that

when the time lag is close to zero (similar to Synchronization), both the same- or

reversed-order conditions show disruption in recency effect, whereas when the time

lag is large (similar to Baseline), both conditions show recency effect. Only when the

correlation time is in the range of [110 ms, 210 ms], the two conditions show

divergence, i.e., recency effect for same-order condition and primacy effect for

reversed-order condition. Parameters are given in Supplementary Materials.

1020