Regeneration Rosetta: An interactive web application to explore regeneration-associated gene expression and chromatin accessibility

Andrea Rau[1,2], Sumona P. Dhara[3], Ava J. Udvadia[3], Paul L. Auer[2]

[1] GABI, INRA, AgroParisTech, Universite Paris-Saclay, 78350, Jouy-en-Josas, France
[2] Joseph J. Zilber School of Public Health, University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA
[3] Department of Biological Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

Abstract Time-course high-throughput assays of gene expression and enhancer usage in zebrafish provide a valuable characterization of the dynamic mechanisms governing gene regulatory programs during CNS axon regeneration. To facilitate the exploration and functional interpretation of a set of fully-processed data on regeneration-associated temporal transcription networks, we have created an interactive web application called *Regeneration Rosetta*. Using either built-in or user-provided lists of genes in one of dozens of supported organisms, our web application facilitates the (1) visualization of clustered temporal expression trends; (2) identification of proximal and distal regions of accessible chromatin to expedite downstream motif analysis; and (3) description of enriched functional gene ontology categories. By enabling a straightforward interrogation of these rich data without extensive bioinformatic expertise, *Regeneration Rosetta* is broadly useful for both a deep investigation of time-dependent regulation during regeneration in zebrafish and hypothesis generation in other organisms.

Keywords CNS axon regeneration; gene expression; chromatin accessibility; functional enrichment; zebrafish; R/Shiny

Introduction

Axon degeneration accompanying central nervous system (CNS) injury or disease leads to a permanent loss of function in human patients. This is largely due to an inability of mammals to reinitiate axon growth in adult CNS neurons[1]. In contrast to mammals, adult teleost fish can fully regenerate CNS axons that reinnervate appropriate targets, enabling functional recovery from CNS injury [2]. Interestingly, fish and mammals share common mechanisms for wiring the nervous system during development, and both are known to downregulate developmental growth and

guidance signaling pathways during nervous system maturation[3-4]. Thus, what appears to set fish

apart is the ability to re-initiate a sustained program of axon growth in response to CNS injury.

Transcriptional changes have long been correlated with the intrinsic capacity for regenerative

axon growth[5,6]. In order to understand the precise mechanisms governing gene regulatory

programs during CNS axon regeneration, Dhara et al. (2019)[7] recently identified the dynamic

changes in gene expression and enhancer usage in zebrafish over the full time-course of axon

regeneration in CNS neurons that are capable of successful regeneration. Adult zebrafish were

subjected to optic nerve crush injury, and regenerating retinas were dissected at various time-

points post injury in order to identify the interactions among expressed genes, open chromatin,

and transcription factor expression during CNS axon regeneration.

These data on regeneration-associated temporal transcription networks in zebrafish represent a

rich source of information with wide potential use and insight for the broader regeneration

community. To this end, we provide fully processed data from Dhara et al. (2019) in an interactive

web application, *Regeneration Rosetta*, as a means to explore, visualize, and functionally

interpret regeneration-associated gene expression and chromatin accessibility. Using either built-

in lists of differentially expressed (DE) genes from Dhara et al. (2019) or user-provided gene lists

in one of 69 supported organisms from Ensembl (Table 1), our web application facilitates (i)

customized visualization of clustered temporal expression trends during optic nerve regeneration;

(ii) identification of proximal and distal regions of open chromatin relative to the gene list to

expedite downstream motif analysis via the MEME suite[8]; and (iii) gene ontology (GO) functional

enrichment analysis. Similarly, using either built-in lists of differentially accessible chromatin from

Dhara et al. (2019) or user-provided genomic coordinates of accessible chromatin, the application

identifies proximal and distal genes relative to their position and their corresponding enriched GO

categories.

Species	Genome version	Species	Genome version
Danio rerio	`GRCz10`	*Meleagris gallopavo*	`UMD2`
Homo sapiens	`GRCh38.p5`	*Microcebus murinus*	`micMur1`
Mus musculus	`GRCm38.p4`	*Monodelphis domestica*	`monDom5`
Rattus norvegicus	`Rnor_6.0`	*Mustela putorius furo*	`MusPutFur1.0`
Ailuropoda melanoleuca	`ailMel1`	*Myotis lucifugus*	`myoLuc2`
Anas platyrhynchos	`BGI_duck_1.0`	*Nomascus leucogenys*	`Nleu1.0`
Anolis carolinensis	`AnoCar2.0`	*Ochotona princeps*	`OchPri2.0`
Astyanax mexicanus	`AstMex102`	*Oreochromis niloticus*	`Orenil1.0`
Bos taurus	`UMD3.1`	*Ornithorhynchus anatinus*	`OANA5`
Caenorhabditis elegans	`WBcel235`	*Oryctolagus cuniculus*	`OryCun2.0`
Callithrix jacchus	`C_jacchus3.2.1`	*Oryzias latipes*	`HdrR`
Canis familiaris	`CanFam3.1`	*Otolemur garnettii*	`OtoGar3`
Cavia porcellus	`cavPor3`	*Ovis aries*	`Oar_v3.1`
Chlorocebus sabaeus	`ChlSab1.1`	*Pan troglodytes*	`CHIMP2.1.4`
Choloepus hoffmanni	`choHof1`	*Papio anubis*	`PapAnu2.0`
Ciona intestinalis	`KH`	*Pelodiscus sinensis*	`PelSin_1.0`
Ciona savignyi	`CSAV2.0`	*Petromyzon marinus*	`Pmarinus_7.0`
Dasypus novemcinctus	`Dasnov3.0`	*Poecilia formosa*	`PoeFor_5.1.2`
Dipodomys ordii	`dipOrd1`	*Pongo abelii*	`PPYG2`
Drosophila melanogaster	`BDGP6`	*Procavia capensis*	`proCap1`
Echinops telfairi	`TENREC`	*Pteropus vampyrus*	`pteVam1`
Equus caballus	`EquCab2`	*Saccharomyces cerevisiae*	`R64-1-1`
Erinaceus europaeus	`eriEur1`		
Felis catus	`Felis_catus_6.2`	*Sarcophilus harrisii*	`DEVIL7.0`
Ficedula albicollis	`FicAlb_1.4`	*Sorex araneus*	`sorAra1`
Gadus morhua	`gadMor1`	*Sus scrofa*	`Sscrofa10.2`
Gallus gallus	`Galgal4`	*Taeniopygia guttata*	`taeGut3.2.4`
Gasterosteus aculeatus	`BROADS1`	*Takifugu rubripes*	`FUGU4.0`
Gorilla gorilla	`gorGor3.1`	*Tarsius syrichta*	`tarSyr1`
Ictidomys tridecemlineatus	`spetri2`	*Tetraodon nigroviridis*	`TETRAODON8.0`
Latimeria chalumnae	`LatCha1`	*Tupaia belangeri*	`tupBel1`
Lepisosteus oculatus	`LepOcu1`	*Tursiops truncatus*	`turTru1`
Loxodonta africana	`loxAfr3`	*Vicugna pacos*	`vicPac1`
Macaca mulatta	`MMUL_1`	*Xenopus tropicalis*	`JGI4.2`
Macropus eugenii	`Meug_1.0`	*Xiphophorus maculatus*	`Xipmac4.4.2`

Table 1. List of supported organisms and their associated genome version for user-provided gene set queries in the *Regeneration Rosetta* app.

The *Regeneration Rosetta* app represents a new paradigm to facilitate data sharing and re-use

in the field of regeneration. This type of data sharing directly promotes the National Institutes of

Health guidelines for ensuring rigor and reproducibility in pre-clinical research

(https://www.nih.gov/research-training/rigor-reproducibility/principles-guidelines-reporting-

preclinical-researchas). As large-scale genomic data become more common, tools that allow

rapid querying and exploration of fully processed data (without the need for additional coding or

pre-processing steps) will be critical in accelerating advances in the regeneration field.

Results

Regeneration Rosetta yields insight into cholesterol and lipid biosynthesis regulation during regeneration

Cholesterol biosynthesis pathways were found to be enriched during regeneration in Dhara et al. (2019) and have been previously shown to be important in axon regeneration in mouse[17]. To more deeply investigate their behavior during optic nerve regeneration in zebrafish and demonstrate some of the capabilities of the *Regeneration Rosetta*, we input a list of 125 genes with cholesterol-related GO terms obtained from the Mouse Genome Informatics (MGI) database[18] (Table S2), corresponding to 236 expressed zebrafish transcripts, into the *Regeneration Rosetta* app (Fig. 1).

Figure 1: Workflow of *Regeneration Rosetta* app. Workflow for investigating temporal patterning of regeneration-associated genes classified within specific biological processes and/or comparative evolutionary analysis of the conserved mechanism among regenerative species, using the *Regeneration Rosetta* app.

Of these 236 transcripts, 64 were DE in Dhara et al. (2019); focusing on this subset of transcripts, the *Regeneration Rosetta* produces a clustered heatmap of expression Z-scores across distinct stages of optic nerve regeneration (Figure 2A). Using on the associated GO terms from MGI, we found that the twenty transcripts with peak expression early in regeneration during the phase of growth towards the midline (2-4 dpi) were predominantly enriched in cholesterol metabolic genes, while the majority of those peaking during the midline crossing and target selection phases (4-7 dpi) were enriched in cholesterol biosynthetic pathways (Figure 2B). Interestingly, transcripts differentially down-regulated during regeneration were enriched in negative regulation of cholesterol biosynthetic processes.

Among the cholesterol biosynthetic genes, we observed upregulation during mid-regeneration (4-7 dpi) of SREBF2, a known cholesterol master-regulatory transcription factor[18,19,20]. After downloading the FASTA files of the sequences for peaklets proximal and distal to the 64 DE cholesterol metabolic genes from the *Regeneration Rosetta*, AME motif analysis [21] revealed that mostly proximal open chromatin were enriched in the SREBF2 motif resulting in 23% sequence enrichment. We found a number of genes with temporal expression profiles similar to SREBF2 that are proximal to these accessible binding sites, including *dhcr24*, *hmgcs1*, *hsd17b7*, *insig1*, *sqlea*, and *idi1* (Figure 2A). Interestingly, the proximal and distal peaks of *srebf1*, which exhibited peak expression during brain reinnervation (7-12 dpi), were also enriched for SREBF2 motifs; SREBF1 is a transcription factor related to SREBF2 that is known to regulate genes involved in fatty acid synthesis and lipid homeostasis [21,22,23], both of which have been shown to be important for axon growth and myelination during neurogenesis [23–25]. This suggests that SREBF2 could potentially regulate the transcriptional activity of SREBF1, which, in turn, promotes the expression of genes associated with later regenerative processes.

Figure 2: *Regeneration Rosetta* **app identifies process-specific analysis after optic nerve injury**. (A)
Temporal transcript profiles of genes in the cholesterol metabolic pathway. Relative transcript counts from
retinas dissected 2-, 4-, 7- and 12-days post injury (dpi) were compared with those from uninjured animals
(0 dpi). Transcript expression is presented as TPM Z-scores; putative SREBF2 target genes are indicated
to the right of the heatmap (biosynthetic enzymes in black; transcription factors are in red). (B) Specific
enrichment of cholesterol metabolic and biosynthetic genes early in regeneration. Fisher's exact test of
over-representation was used to identify cholesterol-related GO-terms correlated with specific stages of
regeneration.

Regeneration Rosetta provides insight into evolutionary conserved regenerative

mechanisms

Nervous system function is dependent on the development of highly specific connections between

neurons and their direct targets. The molecular mechanisms regulating this network of

connections are highly conserved across evolution[26, 27]. Unlike mammals, vertebrates such as fish

and amphibians exhibit regenerative abilities of complex tissues and structures. Therefore,

comparing regenerative capabilities across species will enable researchers to identify genes and

transcriptional networks that are critical to the regenerative program.

To facilitate a cross-species comparison, the *Regeneration Rosetta* app enables queries of

patterns of gene expression across injury models with different regenerative capacities. To

illustrate, we compared CNS regeneration in lamprey and zebrafish. Following [27,28], we used the

regenerating lamprey transcriptional profiles from cell bodies located in the brain and spinal cord

following spinal cord injury over a course of 12 weeks. This study identified 3,664 and 3,999

differentially expressed regeneration-associated genes at one or more post-injury time points in

lamprey brain and spinal cord, respectively. After removing duplicates, we filtered these lists to

2,325 (brain) and 2,520 (spinal cord) differentially expressed genes. We looked for an overlap of

genes that were differentially regulated after injury to the zebrafish optic nerve and the lamprey

spinal cord and brain. We found 3,298 transcripts (corresponding to 1,971 genes) and 3,722

transcripts (corresponding to 2,151 genes) in the zebrafish optic regeneration data corresponding

to the lamprey spinal cord and brain, respectively. After subsetting to those that were differential

in the zebrafish (FDR < 5%), we identified 674 (spinal cord) and 728 (brain) differentially

expressed genes (nearly 28% of genes identified as differentially expressed in any one study or

more) common to the zebrafish optic regeneration, lamprey brain, and lamprey spinal cord (Fig.

3), suggesting a considerable overlap between the two regenerative models and three different

tissue types.

To identify the core transcription factors that could potentially regulate stage-specific

regeneration-associated gene transcription in both injury models, we cross referenced the

lamprey list of DE transcripts to a recently compiled list of human transcription factors[29]. We

identified 105 brain- and 72 spinal- transcription factor encoding genes that were differentially

expressed at one or more post-injury time points (Table S3 and S4). Assessing the combined list

of DE transcription factors, the *Regeneration Rosetta* revealed that 17 transcripts corresponding

to 8 transcription factor encoding genes are common among neurons in the lamprey spinal cord

and brain, and zebrafish retina (Fig. 3). Thus, the *Regeneration Rosetta* highlights potential

regulatory factors driving regeneration-associated gene expression among regenerative

organisms.

Figure 3. *Regeneration Rosetta* **app identifies conserved core regulators of CNS axon regeneration.**
Venn diagram of axon growth-associated genes from regenerating CNS neurons from zebrafish and
lamprey. Approximately 10-15% of regeneration-associated genes are shared between neurons
regenerating axons in brain, spinal cord and optic nerve, including a core set of 8 regeneration-associated
transcription factors.

Discussion

The *Regeneration Rosetta* interactive web app represents a rich resource of fully processed,

analyzed, and queryable data from a unique study of regeneration-associated gene expression

and chromatin accessibility during optic nerve regeneration in Dhara et al. (2109). The app was

a crucial component for generating and interpreting results in Dhara et al. (2019), as it facilitated

a deep interrogation of the data that would have otherwise only been possible with extensive

bioinformatic expertise. In addition, we have illustrated the broad utility of the *Regeneration*

Rosetta app through examples focusing on time-dependent regulation during regeneration for

specific biological processes of interest and regenerative mechanisms that are evolutionarily

conserved across species and tissue types. The *Regeneration Rosetta* app will be widely useful,

both for further investigation and interpretation of the data from Dhara et al. (2019) and for

hypothesis generation in other organisms. We expect these use cases of the app to inform the

design of future functional studies that are crucial for translating basic biological insights into new

therapeutics for optic nerve injury.

Methods

Experimental design, data generation, and bioinformatic analyses

Comprehensive experimental details may be found in Dhara et al. (2019). Briefly, whole retinas

were dissected from 7-9 month old adult zebrafish at 0, 2, 4, 7, or 12 days post injury (dpi),

following an optic nerve crush lesion (*n*=3 at each time point). After extracting total RNA from

these samples, RNA-seq libraries were prepared and sequenced as previously described. Retinal

ganglion cells (RGCs) were collected from retinas at each time point using fluorescent activated

cell sorting (FACS) and chromatin was subsequently isolated. ATAC-seq libraries were prepared

and sequenced as previously described[7]. One ATAC-seq library was dropped due to poor

sequence quality.

For the RNA-seq data, after merging technical replicates and validating sequence quality we

quantified transcript abundance[9] and tested differential expression with respect to the initial time

point (0dpi)[10], controlling the false discovery rate at 5%. For the ATAC-seq data, sequences were

aligned [11] to the zebrafish reference and open chromatin regions were called[12]. For each region

with a p-value < 10e-10, a 500bp "peaklet" was defined by anchoring on the mode of the peak

signal[13]. Chromatin accessibility was quantified by counting the number of overlapping reads for

each retained peaklet, and differential accessibility was calculated with respect to the initial time

point (0dpi)[14], controlling the false discovery rate at 5%. Software versions and parameters are

provided in Dhara et al. (2019). All genomic coordinates and annotations are reported with respect

to the GRCz10 Danio rerio genome assembly and Ensembl 90 gene annotation for the zebrafish.

Integration of RNA-seq and ATAC-seq data

To link regions of accessible chromatin with gene expression, we calculated peaklet-to-gene

distance based on the coordinates of the peaklet mode and the gene's transcription start site

(TSS). A proximal peaklet was then defined as one that overlaps the TSS and/or is within ±1kb of

the TSS, while a distal peaklet was defined as one within ±100kb of the TSS but not proximal.

Users can optionally remove exonic peaklets from these lists, defined as those within 50bp of

exonic regions but not overlapping a TSS. To identify genes that are proximal or distal to a given

set of accessible chromatin (whether user-provided or through the built-in lists of differentially

accessible chromatin provided in the app), users may choose to include all genes or only a subset

of those identified to be DE at a particular time point.

For peaklets identified as proximal or distal to the query set of genes, a FASTA file of sequences

and BED file of genomic coordinates may be downloaded by the user for further analysis; in

addition, a CSV file providing the potentially many-to-many correspondences of proximal and

distal peaklets to genes may also be downloaded.

Gene expression visualization and queries for alternative species

Several built-in gene lists, based on the results described in Dhara et al. (2019), are directly

available within the *Regeneration Rosetta* app. These include the lists of DE genes (based on an

FDR adjusted p-value < 0.01) compared to 0dpi, as well as pre-identified clusters with expression

patterns roughly corresponding to established events in the regeneration process (down-

regulation during early-, mid-, or late-regeneration, growth toward the midline, midline crossing,

target selection, and brain innervation). Users may also employ the *Regeneration Rosetta* app to

explore gene sets for one of 68 species (Table S1) in addition to zebrafish by providing the

relevant Ensembl gene IDs[15]. These converted gene lists can be further narrowed to include only

those genes found to be DE post-injury.

For a given set of genes, expression heatmaps using log fold-changes, log transcripts per million

(TPM), or Z-scores of these measures are produced using *ComplexHeatmap*[16], where transcript

clusters are identified for a given number of clusters using the K-means algorithm, and rows are

ordered within each cluster according to hierarchical clustering (Euclidean distance, complete

linkage). Samples may also be hierarchically clustered. A high-resolution heatmap may be resized

and downloaded by the user.

Functional enrichment analysis

The *Regeneration Rosetta* performs on-the-fly functional enrichment analyses of GO terms for

Biological Processes (BP), Cellular Components (CC), and Molecular Function (MF) for a given

gene set using *topGO* (`weight01` algorithm, Fisher test statistic, and gene universe defined as

the set of expressed transcripts from Dhara et al. (2019). P-values are not adjusted for multiple

correction, and only GO terms with raw p-values < 0.05 are reported; tables of enriched GO terms

are displayed in an HTML table in the app and may be optionally downloaded as a CSV file, Excel

spreadsheet, or PDF file.

Technical details of the Regeneration Rosetta

The *Regeneration Rosetta* interactive web app was built in R using the *Shiny* and *flexdashboard*

packages. In addition to the other software packages already cited above, it makes use of the

data.table and *RSQLite* R packages for fast data manipulation, *DT* for rendering HTML tables

using JavaScript, *readxl* for parsing data from Excel spreadsheets, *dplyr* for data manipulation,

and *tokenizers* to convert user-provided gene IDs into tokens.

Web resources:

- The *Regeneration Rosetta* R/Shiny application is available at http://ls-shiny-prod.uwm.edu/rosetta/. A FAQ page is available directly on the app website.

- Source code for the *Regeneration Rosetta* app is available from GitHub: https://github.com/andreamrau/rosetta. The processed data used within the app are directly located in https://github.com/andreamrau/rosetta/tree/master/data; scripts used to process the raw data from Dhara et al. (2019) may be found at https://github.com/andreamrau/OpticRegen_2019.

- Archived source code at the time of publication can be found at https://doi.org/10.5281/zenodo.2658771.

- Software licence (GPL-3)

Acknowledgments

We thank the UWM High Performance Computing (HPC) Service for computing resources used in this work. We are grateful to Maria Replogle for comments on the manuscript. We wish to thank the University of Wisconsin Biotechnology Center DNA Sequencing Facility, Madison for providing RNA and ATAC sample sequencing facilities. We gratefully acknowledge support from the Research Growth Initiative Grant -RGI (to P.L.A. and A.J.U.), University of Wisconsin-Milwaukee. A.R. was supported by the AgreenSkills+ fellowship program, which received funding from the EU's Seventh Framework Program under grant agreement number FP7-609398 (AgreenSkills+ contract). S.P.D. was supported by the Clifford Mortimer Distinguished Scholar Award, University of Wisconsin-Milwaukee.

Literature Cited

1.   Crair, M. C. & Mason, C. A. Reconnecting Eye to Brain. *J. Neurosci.* **36**, 10707–10722

(2016).

2.   Diekmann, H., Kalbhen, P. & Fischer, D. Characterization of optic nerve regeneration using

transgenic zebrafish. *Front. Cell. Neurosci.* **9**, 118 (2015).

3.   Erskine, L. & Herrera, E. Connecting the retina to the brain. *ASN Neuro* **6**, (2014).

4.   Skene, J. H. Axonal growth-associated proteins. *Annu. Rev. Neurosci.* **12**, 127–156 (1989).

5.   Moore, D. L. & Goldberg, J. L. Multiple transcription factor families regulate axon growth

and regeneration. *Dev. Neurobiol.* **71**, 1186–1211 (2011).

6.   Smith, D. S. & Skene, J. H. A transcription-dependent switch controls competence of adult

neurons for distinct modes of axon growth. *J. Neurosci.* 17, 646–658 (1997).

7.   Dhara, S. *et al*. Regulation of CNS regeneration-associated genes is driven by a temporally

changing cast of transcription factors. *bioRxiv*, doi: https://doi.org/10.1101/638734.

8.   Bailey, T. L. *et al.* MEME SUITE: tools for motif discovery and searching. *Nucleic Acids*

Research **37**, W202–W208 (2009).

9.   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

10. Pimentel, H., Bray, N. L., Puente, S., Melsted, P. & Pachter, L. Differential analysis of RNA-

seq incorporating quantification uncertainty. *Nat. Methods* **14**, 687–690 (2017).

11. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754–1760 (2009).

12. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

13. Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS Comput.*

Biol. **9**, e1003118 (2013).

14. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

15. Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the integration

of genomic datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* **4**, 1184–1191

(2009).

16. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

17. Wang, Z. *et al.* KLF6 and STAT3 co-occupy regulatory DNA and functionally synergize to

promote axon growth in CNS neurons. *Sci. Rep.* **8**, 12565 (2018).

18. Smith, C. L. *et al.* Mouse Genome Database (MGD)-2018: knowledgebase for the

laboratory mouse. *Nucleic Acids Res.* **46**, D836–D842 (2018).

19. Roux, P. & Fisher, R. M. Chest injuries in children: an analysis of 100 cases of blunt chest

trauma from motor vehicle accidents. *J. Pediatr. Surg.* **27**, 551–555 (1992).

20. Madison, B. B. Srebp2: A master regulator of sterol and fatty acid synthesis. *J. Lipid Res.*

57, 333–335 (2016).

21. McLeay, R. C. & Bailey, T. L. Motif Enrichment Analysis: a unified framework and an

evaluation on ChIP data. *BMC Bioinformatics* **11**, 165 (2010).

22. Ye, J. & DeBose-Boyd, R. A. Regulation of cholesterol and fatty acid synthesis. *Cold Spring*

Harb. Perspect. Biol. **3**, (2011).

23. Ferré, P. & Foufelle, F. SREBP-1c transcription factor and lipid homeostasis: clinical

perspective. *Horm. Res.* **68**, 72–82 (2007).

24. Salles, J. *et al.* Acetyl-CoA carboxylase and SREBP expression during peripheral nervous

system myelination. *Biochim. Biophys. Acta* **1631**, 229–238 (2003).

25. Dietschy, J. M. & Turley, S. D. Thematic review series: brain Lipids. Cholesterol metabolism

in the central nervous system during early development and in the mature animal. *J. Lipid*

Res. **45**, 1375–1397 (2004).

26. Jiang, X. & Nardelli, J. Cellular and molecular introduction to brain development. *Neurobiol.*

Dis. **92**, 3–17 (2016).

27. Kaprielian, Z., Imondi, R. & Runko, E. Axon guidance at the midline of the developing CNS.

Anat. Rec. **261**, 176–197 (2000).

28. Herman, P. E. *et al.* Highly conserved molecular pathways, including Wnt signaling,

promote functional recovery from spinal cord injury in lampreys. *Sci. Rep.* **8**, 742 (2018).

29. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

324