Regeneration Rosetta: An interactive web application to explore regeneration-associated gene expression and chromatin accessibility

Andrea Rau[1,2], Sumona P. Dhara[3], Ava J. Udvadia[3], Paul L. Auer[2]

[1] GABI, INRA, AgroParisTech, Universite Paris-Saclay, 78350, Jouy-en-Josas, France
[2] Joseph J. Zilber School of Public Health, University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA
[3] Department of Biological Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

Abstract Time-course high-throughput assays of gene expression and enhancer usage in zebrafish provide a valuable characterization of the dynamic mechanisms governing gene regulatory programs during CNS axon regeneration. To facilitate the exploration and functional interpretation of a set of fully-processed data on regeneration-associated temporal transcription networks, we have created an interactive web application called *Regeneration Rosetta*. Using either built-in or user-provided lists of genes in one of dozens of supported organisms, our web application facilitates the (1) visualization of clustered temporal expression trends; (2) identification of proximal and distal regions of accessible chromatin to expedite downstream motif analysis; and (3) description of enriched functional gene ontology categories. By enabling a straightforward interrogation of these rich data without extensive bioinformatic expertise, *Regeneration Rosetta* is broadly useful for both a deep investigation of time-dependent regulation during regeneration in zebrafish and hypothesis generation in other organisms.

Keywords CNS axon regeneration; gene expression; chromatin accessibility; functional enrichment; zebrafish; R/Shiny

Introduction

Axon degeneration accompanying central nervous system (CNS) injury or disease leads to a permanent loss of function in human patients. This is largely due to an inability of mammals to reinitiate axon growth in adult CNS neurons (Crair and Mason 2016). In contrast to mammals, adult teleost fish can fully regenerate CNS axons that reinnervate appropriate targets, enabling functional recovery from CNS injury (Diekmann *et al.* 2015). Interestingly, fish and mammals

share common mechanisms for wiring the nervous system during development, and both are

known to downregulate developmental growth and guidance signaling pathways during nervous

system maturation (Skene 1989; Erskine and Herrera 2014). Thus, what appears to set fish apart

is the ability to re-initiate a sustained program of axon growth in response to CNS injury.

Transcriptional changes have long been correlated with the intrinsic capacity for regenerative

axon growth (Smith and Skene 1997; Moore and Goldberg 2011). In order to understand the

precise mechanisms governing gene regulatory programs during CNS axon regeneration, Dhara

et al. (2019) recently identified the dynamic changes in gene expression and enhancer usage in

zebrafish over the full time-course of axon regeneration in CNS neurons that are capable of

successful regeneration. Adult zebrafish were subjected to optic nerve crush injury, and

regenerating retinas were dissected at various time-points post injury in order to identify the

interactions among expressed genes, open chromatin, and transcription factor expression during

CNS axon regeneration.

These data on regeneration-associated temporal transcription networks in zebrafish represent a

rich source of information with wide potential use and insight for the broader regeneration

community. To this end, we provide fully processed data from Dhara et al. (2019) in an interactive

web application, *Regeneration Rosetta*, as a means to explore, visualize, and functionally

interpret regeneration-associated gene expression and chromatin accessibility. Using either built-

in lists of differentially expressed (DE) genes from Dhara et al. (2019) or user-provided gene lists

in one of 69 supported organisms from Ensembl (Table 1), our web application facilitates (i)

customized visualization of clustered temporal expression trends during optic nerve regeneration;

(ii) identification of proximal and distal regions of open chromatin relative to the gene list to

expedite downstream motif analysis via the MEME suite (Bailey *et al.* 2009); and (iii) gene

ontology (GO) functional enrichment analysis. Similarly, using either built-in lists of differentially

accessible chromatin from Dhara et al. (2019) or user-provided genomic coordinates of accessible

chromatin, the application identifies proximal and distal genes relative to their position and their

corresponding enriched GO categories.

Species	Genome version	Species	Genome version
Danio rerio	`GRCz10`	*Meleagris gallopavo*	`UMD2`
Homo sapiens	`GRCh38.p5`	*Microcebus murinus*	`micMur1`
Mus musculus	`GRCm38.p4`	*Monodelphis domestica*	`monDom5`
Rattus norvegicus	`Rnor_6.0`	*Mustela putorius furo*	`MusPutFur1.0`
Ailuropoda melanoleuca	`ailMel1`	*Myotis lucifugus*	`myoLuc2`
Anas platyrhynchos	`BGI_duck_1.0`	*Nomascus leucogenys*	`Nleu1.0`
Anolis carolinensis	`AnoCar2.0`	*Ochotona princeps*	`OchPri2.0`
Astyanax mexicanus	`AstMex102`	*Oreochromis niloticus*	`Orenil1.0`
Bos taurus	`UMD3.1`	*Ornithorhynchus anatinus*	`OANA5`
Caenorhabditis elegans	`WBcel235`	*Oryctolagus cuniculus*	`OryCun2.0`
Callithrix jacchus	`C_jacchus3.2.1`	*Oryzias latipes*	`HdrR`
Canis familiaris	`CanFam3.1`	*Otolemur garnettii*	`OtoGar3`
Cavia porcellus	`cavPor3`	*Ovis aries*	`Oar_v3.1`
Chlorocebus sabaeus	`ChlSab1.1`	*Pan troglodytes*	`CHIMP2.1.4`
Choloepus hoffmanni	`choHof1`	*Papio anubis*	`PapAnu2.0`
Ciona intestinalis	`KH`	*Pelodiscus sinensis*	`PelSin_1.0`
Ciona savignyi	`CSAV2.0`	*Petromyzon marinus*	`Pmarinus_7.0`
Dasypus novemcinctus	`Dasnov3.0`	*Poecilia formosa*	`PoeFor_5.1.2`
Dipodomys ordii	`dipOrd1`	*Pongo abelii*	`PPYG2`
Drosophila melanogaster	`BDGP6`	*Procavia capensis*	`proCap1`
Echinops telfairi	`TENREC`	*Pteropus vampyrus*	`pteVam1`
Equus caballus	`EquCab2`	*Saccharomyces cerevisiae*	`R64-1-1`
Erinaceus europaeus	`eriEur1`		
Felis catus	`Felis_catus_6.2`	*Sarcophilus harrisii*	`DEVIL7.0`
Ficedula albicollis	`FicAlb_1.4`	*Sorex araneus*	`sorAra1`
Gadus morhua	`gadMor1`	*Sus scrofa*	`Sscrofa10.2`
Gallus gallus	`Galgal4`	*Taeniopygia guttata*	`taeGut3.2.4`
Gasterosteus aculeatus	`BROADS1`	*Takifugu rubripes*	`FUGU4.0`
Gorilla gorilla	`gorGor3.1`	*Tarsius syrichta*	`tarSyr1`
Ictidomys tridecemlineatus	`spetri2`	*Tetraodon nigroviridis*	`TETRAODON8.0`
Latimeria chalumnae	`LatCha1`	*Tupaia belangeri*	`tupBel1`
Lepisosteus oculatus	`LepOcu1`	*Tursiops truncatus*	`turTru1`
Loxodonta africana	`loxAfr3`	*Vicugna pacos*	`vicPac1`
Macaca mulatta	`MMUL_1`	*Xenopus tropicalis*	`JGI4.2`
Macropus eugenii	`Meug_1.0`	*Xiphophorus maculatus*	`Xipmac4.4.2`

Table 1. List of supported organisms and their associated genome version for user-provided gene set queries in the *Regeneration Rosetta* app.

The *Regeneration Rosetta* app represents a new paradigm to facilitate data sharing and re-use

in the field of regeneration. This type of data sharing directly promotes the National Institutes of

Health   guidelines   for   ensuring   rigor   and   reproducibility   in   pre-clinical   research

(https://www.nih.gov/research-training/rigor-reproducibility/principles-guidelines-reporting-

preclinical-researchas). As large-scale genomic data become more common, tools that allow

rapid querying and exploration of fully processed data (without the need for additional coding or

pre-processing steps) will be critical in accelerating advances in the regeneration field.

Materials and Methods

Experimental design, data generation, and bioinformatic analyses

Comprehensive experimental details may be found in Dhara et al. (2019). Briefly, whole retinas

were dissected from 7-9 month old adult zebrafish at 0, 2, 4, 7, or 12 days post injury (dpi),

following an optic nerve crush lesion (*n*=3 at each time point). For the RNA-seq data, we tested

differential expression with respect to the initial time point (0dpi), controlling the false discovery

rate at 5%. For the ATAC-seq data, sequences were aligned (Li and Durbin 2009) to the zebrafish

reference and open chromatin regions were called (Zhang *et al.* 2008). For each region with a p-

value < 10e-10, a 500bp "peaklet" was defined by anchoring on the mode of the peak signal

(Lawrence *et al.* 2013). Chromatin accessibility was quantified by counting the number of

overlapping reads for each retained peaklet, and differential accessibility was calculated with

respect to the initial time point (0dpi) (Love *et al.* 2014), controlling the false discovery rate at 5%.

Software versions and parameters are provided in Dhara et al. (2019). All genomic coordinates

and annotations are reported with respect to the GRCz10 Danio rerio genome assembly and

Ensembl 90 gene annotation for the zebrafish.

Integration of RNA-seq and ATAC-seq data

To link regions of accessible chromatin with gene expression, we calculated peaklet-to-gene

distance based on the coordinates of the peaklet mode and the gene's transcription start site

(TSS). A proximal peaklet was then defined as one that overlaps the TSS and/or is within ±1kb of

the TSS, while a distal peaklet was defined as one within ±100kb of the TSS but not proximal.

Users can optionally remove exonic peaklets from these lists, defined as those within 50bp of

exonic regions but not overlapping a TSS. To identify genes that are proximal or distal to a given

set of accessible chromatin (whether user-provided or through the built-in lists of differentially

accessible chromatin provided in the app), users may choose to include all genes or only a subset

of those identified to be DE at a particular time point.

For peaklets identified as proximal or distal to the query set of genes, a FASTA file of sequences

and BED file of genomic coordinates may be downloaded by the user for further analysis; in

addition, a CSV file providing the potentially many-to-many correspondences of proximal and

distal peaklets to genes may also be downloaded.

Gene expression visualization and queries for alternative species

Several built-in gene lists, based on the results described in Dhara et al. (2019), are directly

available within the *Regeneration Rosetta* app. These include the lists of DE genes (based on an

FDR adjusted p-value < 0.01) compared to 0dpi, as well as pre-identified clusters with expression

patterns roughly corresponding to established events in the regeneration process (down-

regulation during early-, mid-, or late-regeneration, growth toward the midline, midline crossing,

target selection, and brain innervation). Users may also employ the *Regeneration Rosetta* app to

explore gene sets for one of 68 species (Table 1) in addition to zebrafish by providing the relevant

Ensembl gene IDs (Durinck *et al.* 2009). These converted gene lists can be further narrowed to

include only those genes found to be DE post-injury.

For a given set of genes, expression heatmaps using log fold-changes, log transcripts per million

(TPM), or Z-scores of these measures are produced using *ComplexHeatmap* (Gu *et al.* 2016),

where transcript clusters are identified for a given number of clusters using the K-means

algorithm, and rows are ordered within each cluster according to hierarchical clustering (Euclidean

distance, complete linkage). Samples may also be hierarchically clustered. A high-resolution

heatmap may be resized and downloaded by the user.

Functional enrichment analysis

The *Regeneration Rosetta* performs on-the-fly functional enrichment analyses of GO terms for

Biological Processes (BP), Cellular Components (CC), and Molecular Function (MF) for a given

gene set using *topGO* (`weight01` algorithm, Fisher test statistic, and gene universe defined as

the set of expressed transcripts from Dhara et al. (2019). P-values are not adjusted for multiple

correction, and only GO terms with raw p-values < 0.05 are reported; tables of enriched GO terms

are displayed in an HTML table in the app and may be optionally downloaded as a CSV file, Excel

spreadsheet, or PDF file. We remark that a list of enriched GO terms must be interpreted with

respect to the context of the user-provided gene list. Most users of the *Regeneration Rosetta*

would typically make use of lists of regeneration-related genes from one of the supported

organisms to understand their behavior in the gene expression and chromatin accessibility data

of Dhara et al. (2019); however, if provided with genes unrelated to regeneration, the app will

typically return lists of enriched GO terms also unrelated to regeneration. We note that a variety

of other ontology databases and enrichment methods exist in the literature, and the interested

user could easily export a gene list obtained from the app to other tools, if desired.

Technical details of the Regeneration Rosetta

The *Regeneration Rosetta* interactive web app was built in R using the *Shiny* and *flexdashboard*

packages. In addition to the other software packages already cited above, it makes use of the

data.table and *RSQLite* R packages for fast data manipulation, *DT* for rendering HTML tables

using JavaScript, *readxl* for parsing data from Excel spreadsheets, *dplyr* for data manipulation,

and *tokenizers* to convert user-provided gene IDs into tokens. The *Regeneration Rosetta* R/Shiny

app is available at http://ls-shiny-prod.uwm.edu/rosetta/. A FAQ page is available directly on the

app website. Source code for the *Regeneration Rosetta* is available from GitHub:

https://github.com/andreamrau/rosetta. The processed data used within the app are directly

located in https://github.com/andreamrau/rosetta/tree/master/data; scripts used to process the

raw   data   from   Dhara   et   al.   (2019)   may   be   found   at

https://github.com/andreamrau/OpticRegen_2019.   Archived   source   code   at   the   time   of

publication can be found at https://doi.org/10.5281/zenodo.2658771. The *Regeneration Rosetta*

app was developed under a GPL-3 software license.

Results

Regeneration Rosetta yields insight into cholesterol and lipid biosynthesis regulation

during regeneration

Cholesterol biosynthesis pathways were found to be enriched during regeneration in Dhara et al.

(2019) and have been previously shown to be important in axon regeneration in mouse (Wang *et*

al. 2018). To more deeply investigate their behavior during optic nerve regeneration in zebrafish

and demonstrate some of the capabilities of the *Regeneration Rosetta*, we input a list of 125

genes with cholesterol-related GO terms obtained from the Mouse Genome Informatics (MGI)

database (Smith *et al.* 2018) (Table S1), corresponding to 232 expressed zebrafish transcripts,

into the *Regeneration Rosetta* app (Fig. 1).

Figure 1: Workflow of *Regeneration Rosetta* app. Workflow for investigating temporal patterning of regeneration-associated genes classified within specific biological processes and/or comparative evolutionary analysis of the conserved mechanism among regenerative species, using the *Regeneration Rosetta* app.

Of these 232 transcripts, 62 were differentially expressed (DE) in Dhara et al. (2019); focusing on this subset of transcripts, the *Regeneration Rosetta* produces a clustered heatmap of expression Z-scores across distinct stages of optic nerve regeneration (Table S2; Figure 2A). Using on the associated GO terms from MGI, we found that the twenty transcripts with peak expression early in regeneration during the phase of growth towards the midline (2-4 dpi) were predominantly enriched in cholesterol metabolic genes, while the majority of those peaking during the midline crossing and target selection phases (4-7 dpi) were enriched in cholesterol biosynthetic pathways

(Figure 2B). Interestingly, transcripts differentially down-regulated during regeneration were

enriched in negative regulation of cholesterol biosynthetic processes.

Among the cholesterol biosynthetic genes, we observed upregulation during mid-regeneration (4-

7 dpi) of SREBF2, a known cholesterol master-regulatory transcription factor (Madison 2016;

Smith *et al.* 2018). After downloading the FASTA files of the sequences for peaklets proximal and

distal to the 62 DE cholesterol metabolic genes from the *Regeneration Rosetta*, AME motif

analysis (McLeay and Bailey 2010) revealed that mostly proximal open chromatin were enriched

in the SREBF2 motif resulting in 23% sequence enrichment. We found a number of genes with

temporal expression profiles similar to SREBF2 that are proximal to these accessible binding

sites, including *dhcr24*, *hmgcs1*, *hsd17b7*, *insig1*, *sqlea*, and *idi1* (Figure 2A). Interestingly, the

proximal and distal peaks of *srebf1*, which exhibited peak expression during brain reinnervation

(7-12 dpi), were also enriched for SREBF2 motifs; SREBF1 is a transcription factor related to

SREBF2 that is known to regulate genes involved in fatty acid synthesis and lipid homeostasis

(Ferré and Foufelle 2007; McLeay and Bailey 2010; Ye and DeBose-Boyd 2011), both of which

have been shown to be important for axon growth and myelination during neurogenesis (Salles

et al. 2003; Dietschy and Turley 2004; Ferré and Foufelle 2007). This suggests that SREBF2

could potentially regulate the transcriptional activity of SREBF1, which, in turn, promotes the

expression of genes associated with later regenerative processes.

Figure 2: *Regeneration Rosetta* app identifies process-specific analysis after optic nerve injury. (A)
Temporal transcript profiles of genes in the cholesterol metabolic pathway. Relative transcript counts from
retinas dissected 2-, 4-, 7- and 12-days post injury (dpi) were compared with those from uninjured animals
(0 dpi). Transcript expression is presented as TPM Z-scores; putative SREBF2 target genes are indicated
to the right of the heatmap (biosynthetic enzymes in black; transcription factors are in red). (B) Specific
enrichment of cholesterol metabolic and biosynthetic genes early in regeneration. Fisher's exact test of
over-representation was used to identify cholesterol-related GO-terms correlated with specific stages of
regeneration.

Regeneration Rosetta provides insight into evolutionary conserved regenerative

mechanisms

Nervous system function is dependent on the development of highly specific connections between

neurons and their direct targets. The molecular mechanisms regulating this network of

connections are highly conserved across evolution (Kaprielian *et al.* 2000; Jiang and Nardelli

2016). Unlike mammals, vertebrates such as fish and amphibians exhibit regenerative abilities of

complex tissues and structures. Therefore, comparing regenerative capabilities across species

will enable researchers to identify genes and transcriptional networks that are critical to the

regenerative program.

To facilitate a cross-species comparison, the *Regeneration Rosetta* app enables queries of

patterns of gene expression across injury models with different regenerative capacities. To

illustrate, we compared CNS regeneration in lamprey and zebrafish. Following (Kaprielian *et al.*

2000; Herman *et al.* 2018), we used the regenerating lamprey transcriptional profiles from cell

bodies located in the brain and spinal cord following spinal cord injury over a course of 12 weeks.

This study identified 3,664 and 3,998 differentially expressed regeneration-associated genes at

one or more post-injury time points in lamprey brain and spinal cord, respectively. After removing

duplicates, we filtered these lists to 2,325 (brain) and 2,519 (spinal cord) differentially expressed

genes. We looked for an overlap of genes that were differentially regulated after injury to the

zebrafish optic nerve and the lamprey brain and spinal cord. We found 3,712 transcripts

(corresponding to 2,150 genes) and 4,076 transcripts (corresponding to 2,404 genes) in the

zebrafish optic regeneration data corresponding to the lamprey brain and spinal cord,

respectively. After subsetting to those that were differential in the zebrafish (FDR <5%, Wald tests

vs 0dpi), we identified 734 (brain) and 846 (spinal cord) differentially expressed genes, 483 were

shared between the three data sets (Fig. 3). This suggests a considerable overlap between

responses to nerve injury in two different species and three different tissue types.

To identify the core transcription factors that could potentially regulate stage-specific

regeneration-associated gene transcription in both injury models, we cross referenced the

lamprey list of DE transcripts to a recently compiled list of human transcription factors (Lambert

et al. 2018). We identified 105 brain- and 131 spinal- transcription factor encoding genes that

were differentially expressed at one or more post-injury time points (Table S3 and S4). The

Regeneration Rosetta revealed 32 (brain) and 53 (spinal cord) DE transcription factor encoding

genes that were differential in the zebrafish optic nerve regeneration data after subjecting to FDR

5%, Wald tests vs 0dpi. Assessing the combined list of DE transcription factors, we found that 25

transcripts corresponding to 21 transcription factor encoding genes are shared among CNS

neuron responses to axonal injury in the lamprey spinal cord and brain, and zebrafish optic nerve

(Fig. 3). Thus, the *Regeneration Rosetta* highlights potential regulatory factors driving

regeneration-associated gene expression among regeneration capable organisms.

Figure 3. *Regeneration Rosetta* app identifies conserved core regulators of CNS axon regeneration.
Venn diagram of axon growth-associated genes from regenerating CNS neurons after zebrafish optic nerve
injury (ONI; retina response) and lamprey spinal cord injury (SCI; spinal cord (SC) and brain responses).
Approximately 10-20% of regeneration-associated genes are shared between neurons regenerating axons
in brain, spinal cord and optic nerve, including a core set of 21 regeneration-associated transcription factor
encoding genes that are homologous to human genes (listed in the middle with HGNC gene symbol)

Discussion

The *Regeneration Rosetta* interactive web app represents a rich resource of fully processed,

analyzed, and queryable data from a unique study of regeneration-associated gene expression

and chromatin accessibility during optic nerve regeneration in Dhara et al. (2019). The app was

a crucial component for generating and interpreting results in Dhara et al. (2019), as it facilitated

a deep interrogation of the data that would have otherwise only been possible with extensive

bioinformatic expertise. In addition, we have illustrated the broad utility of the *Regeneration*

Rosetta app through examples focusing on time-dependent regulation during regeneration for

specific biological processes of interest and regenerative mechanisms that are evolutionarily

conserved across species and tissue types. All of the source code for running our analyses and

implementing the app is posted on GitHub; in this way researchers can modify the code for their

own applications or run a local version of the app if desired. As such, the *Regeneration Rosetta*

app (and its open-source code) provide a useful a framework for sharing results and data. The

Regeneration Rosetta app will be widely useful, both for further investigation and interpretation of

the data from Dhara et al. (2019) and for hypothesis generation in other organisms. Indeed, we

have provided example user-provided gene lists from a variety of published regeneration studies

into the app. The lists include regeneration-associated genes from 10 different cell/tissue types

from five different species. Our applications allows users to explore how the expression of these

genes may change over the course of optic nerve regeneration. We expect these use cases of

the app to inform the design of future functional studies that are crucial for translating basic

biological insights into new therapeutics for optic nerve injury.

Acknowledgments

We thank the <u>UWM High Performance Computing (HPC) Service</u> for computing resources used

in this work. We are grateful to Maria Replogle for comments on the manuscript. We wish to thank

the University of Wisconsin Biotechnology Center DNA Sequencing Facility, Madison for providing

RNA and ATAC sample sequencing facilities. We gratefully acknowledge support from the

Research Growth Initiative Grant -RGI (to P.L.A. and A.J.U.), University of Wisconsin-Milwaukee.

A.R. was supported by the AgreenSkills+ fellowship program, which received funding from the

EU's Seventh Framework Program under grant agreement number FP7-609398 (AgreenSkills+

contract). S.P.D. was supported by the Clifford Mortimer Distinguished Scholar Award, University

of Wisconsin-Milwaukee.

Reference:

Bailey, T. L., M. Boden, F. A. Buske, M. Frith, C. E. Grant *et al.*, 2009 MEME Suite: Tools for

motif discovery and searching. Nucleic Acids Res. 37: 202–208.

Bray, N. L., H. Pimentel, P. Melsted, and L. Pachter, 2016 Near-optimal probabilistic RNA-seq

quantification. Nat. Biotechnol. 34: 525–527.

Crair, M. C., and C. A. Mason, 2016 Reconnecting eye to brain. J. Neurosci. 36: 10707–10722.

Diekmann, H., P. Kalbhen, and D. Fischer, 2015 Characterization of optic nerve regeneration

using transgenic zebrafish. Front. Cell. Neurosci. 9: 1–11.

Dietschy, J. M., and S. D. Turley, 2004 Cholesterol metabolism in the central nervous system

during early development and in the mature animal. J. Lipid Res. 45: 1375–1397.

Dhara, S. P., Rau, A., Flister, M. J., Recka, N. M., Laiosa M. D., Auer, P.L., and Udvadia, A. J.

2019. Cellular reprogramming for successful CNS axon regeneration is driven by a

temporally changing cast of transcription factors. Scientific Reports *(in press)*

Durinck, S., Spellman, P. T., Birney, E. and Huber, W. 2009 Mapping identifiers for the

integration of genomic datasets with the R/Bioconductor package biomaRt. Nat. Protoc. 4,

1184–1191

Erskine, L. & Herrera, E., 2014 Connecting the retina to the brain. ASN Neuro 6 :

1759091414562107.

Ferré, P., and F. Foufelle, 2007 SREBP-1c transcription factor and lipid homeostasis: Clinical

perspective. Horm. Res. 68: 72–82.

Gu, Z., Eils, R. & Schlesner, M., 2016 Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics 32, 2847–2849

Herman, P. E., A. Papatheodorou, S. A. Bryant, C. K. M. Waterbury, J. R. Herdy *et al.*, 2018

Highly conserved molecular pathways, including Wnt signaling, promote functional

recovery from spinal cord injury in lampreys. Sci. Rep. 8: 1–15.

Jiang, X. & Nardelli, J., 2016 Cellular and molecular introduction to brain development.

Neurobiol. Dis. 92, 3–17.

Kaprielian, Z., R. Imondi, and E. Runko, 2000 Axon guidance at the midline of the developing

CNS. Anat. Rec. 261: 176–197.

Lambert, S. A., A. Jolma, L. F. Campitelli, P. K. Das, Y. Yin *et al.*, 2018 The Human

Transcription Factors. Cell 172: 650–665.

Lawrence, M., W. Huber, H. Pagès, P. Aboyoun, M. Carlson *et al.*, 2013 Software for

Computing and Annotating Genomic Ranges. PLoS Comput. Biol. 9: 1–10.

Li, H., and R. Durbin, 2009 Making the Leap: Maq to BWA. Mass Genomics 25: 1754–1760.

Love, M. I., W. Huber, and S. Anders, 2014 Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol. 15: 1–21.

Madison, B. B., 2016 Srebp2: A master regulator of sterol and fatty acid synthesis1. J. Lipid

Res. 57: 333–335.

McLeay, R. C., and T. L. Bailey, 2010 Motif Enrichment Analysis: A unified framework and an

evaluation on ChIP data. BMC Bioinformatics 11:.

Moore, D. L., and J. L. Goldberg, 2011 Multiple transcription factor families regulate axon

growth and regeneration. Dev. Neurobiol. 71: 1186–1211.

Pimentel, H., N. L. Bray, S. Puente, P. Melsted, and L. Pachter, 2017 Differential analysis of

RNA-seq incorporating quantification uncertainty. Nat. Methods 14: 687–690.

Salles, J., F. Sargueil, A. Knoll-Gellida, L. A. Witters, C. Cassagne *et al.*, 2003 Acetyl-CoA

carboxylase and SREBP expression during peripheral nervous system myelination.

Biochim. Biophys. Acta - Mol. Cell Biol. Lipids 1631: 229–238.

Skene, J. H., 1989 Axonal growth-associated proteins. Annu. Rev. Neurosci. 12, 127–156.

doi: 10.1146/annurev.ne.12.030189.001015

Smith, C. L., J. A. Blake, J. A. Kadin, J. E. Richardson, and C. J. Bult, 2018 Mouse Genome

Database (MGD)-2018: Knowledgebase for the laboratory mouse. Nucleic Acids Res. 46:

D836–D842.

Smith, D. S., and J. H. Skene, 1997 A transcription-dependent switch controls competence of

adult neurons for distinct modes of axon growth. J. Neurosci. 17: 646–58.

Wang, Z., V. Mehra, M. T. Simpson, B. Maunze, A. Chakraborty *et al.*, 2018 KLF6 and STAT3

co-occupy regulatory DNA and functionally synergize to promote axon growth in CNS

neurons. Sci. Rep. 8: 1–16.

Ye, J., and R. A. DeBose-Boyd, 2011 Regulation of cholesterol and fatty acid synthesis. Cold

Spring Harb. Perspect. Biol. 3: 1–13.

Zhang, Y., T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson *et al.*, 2008 Model-based analysis

of ChIP-Seq (MACS). Genome Biol. 9:.