A new clinical tool to predict outcome in early-stage melanoma patients

Filippo Mancuso[1,±], Sergio Lage[1,±], Javier Rasero[2,±], José Luis Díaz-Ramón[2,3], Aintzane Apraiz[1], Gorka Pérez-Yarza[1,2], Pilar Ezkurra[1,2], Cristina Penas[1], Ana Sánchez-Diez[2,4], María Dolores García-Vazquez[2], Jesús Gardeazabal[2,3], Rosa Izu[2,4], Karmele Mujika[5,6], Jesús Cortés[2], Aintzane Asumendi[1,2] and María Dolores Boyano[1,2]

[1]Department of Cell Biology and Histology, Faculty of Medicine and Nursing, UPV/EHU, Leioa, Spain.

[2]Biocruces Bizkaia Health Research Institute, Barakaldo, Spain.

[3]Department of Dermatology, Cruces University Hospital, Barakaldo, Spain.

[4]Department of Dermatology, Basurto University Hospital, Bilbao, Spain.

[5]Department of Medical Oncology, Onkologikoa Hospital, Donostia, Spain

[6]Biodonostia Institute, Donostia, Spain

[±] These authors contributed equally to this work.

Correspondence: María Dolores Boyano, Department of Cell Biology and Histology, Faculty of Medicine and Nursing, University of the Basque Country UPV/EHU, Barrio Sarriena s/n, 48940-Leioa, Spain. E-mail: lola.boyano@ehu.eus

Short title: Prognostic serum signature in early-stage melanoma

Key words: melanoma, serum biomarkers, prognosis, Interleukin, Dermcidin

Abbreviations: Interleukin (IL), Interferon (IFN), Transforming Growth Factor (TGF), Granulocyte-Macrophage Colony-stimulating Factor (GM-CSF), Dermcidin (DCD), T helper (Th)

NOVELTY AND IMPACT

We have developed a prognostic equation that considers the serum IL-4 and GM-CSF values, along with the Breslow thickness to accurately classify melanoma outcome in patients. In this sense, a rigorous follow-up is recommended for early-stage melanoma patients with a high Breslow thickness, high serum IL-4 levels and low GM-CSF at the time of diagnosis, given the elevated risk that these patients will develop metastasis during follow-up.

ABSTRACT

Around 25% of early-stage melanoma patients eventually develop metastasis. Thus, we set out to define serological biomarkers that could be used along with clinical and histopathological features of the disease to predict these events. We previously demonstrated that in stage II melanoma patients, serum levels of Dermcidin (DCD) are associated with metastatic progression. Since DCD was recently associated with a local and systemic immune response against cancer cells, the serum levels of DCD were studied using ELISA and Luminex, along with those of IL-4, IL-6, IL-10, IL-17A, IFNγ, TGFβ and GM-CSF, in order to define potential prognostic factors in a group of 448 melanoma patients, 323 of whom were diagnosed as stages I-II according AJCC. The data were analyzed employing machine learning and Kaplan-Meier techniques in order to define an algorithm that was capable of accurately classifying early stage melanoma patients with a high and low risk of developing metastasis within 2 years, the period of highest recurrence in such patients. The results show that in early stage melanoma patients and in addition to the Breslow thickness, the serum levels of the cytokines IL-4 and GM-CSF are those that best predict of melanoma metastasis. As a result, these parameters can be incorporated into a tool to discriminate subjects with a better and worse melanoma prognosis.

INTRODUCTION

Early and accurate classification of patients is the cornerstone of precision medicine, intimately linked to the optimal management of cancer. This is especially relevant for melanoma, the most deadly type of skin cancer due to its high metastatic capacity and the limited therapeutic tools available to combat the spread of the disease. In fact, distant metastasis are associated with median survival rates ranging from 6 to 15 months[1]. Despite all the efforts to devise prevention and detection strategies, the incidence of melanoma is expected to increase in the forthcoming years[2], further supporting the benefits to be gained by investing in the development of predictive tools.

The prognosis of melanoma is currently assigned almost entirely on the basis on a limited set of histopathological markers[3,4]. In this context, tumor thickness is the most important histopathological characteristic included in the AJCC staging system and it is officially considered as a prognostic factor for melanoma progression in clinical practice[5,6]. However, due to the clinical and biological heterogeneity of primary melanoma, survival can vary widely even among individuals considered to be within the same stage[7,8], highlighting the need for new prognostic tools to improve the management of primary melanoma patients[9]. Precision medicine focuses on classifying early stage melanoma patients on the basis of genetic and other biochemical features in order to identify profiles that are most likely to develop into more advanced disease stages, and to define more effective treatments for the metastatic disease[10].

Serum is a highly accessible and valuable source of biomarkers, containing tumor and host-related factors that are correlated with tumor behavior and patient prognosis[11]. Cytokines are key mediators of the immune system with either pro-inflammatory or anti-inflammatory activity, and they are serum factors with potential value as biomarkers. In fact, cytokine profiling is providing valuable data regarding patient classification in a wide range of diseases, including cancer[12,13,14]. In terms of tumor activity, elevated Th2 cytokines (IL-4 -Interleukin-4, IL-5, IL-10 and IL-13) and decreased Th1 cytokines (IL-2, TGFβ and IFNɣ - Transforming growth factor β and Interferon-ɣ) suppress effective spontaneous anti-tumor immunity[15,16,17]. In addition, the IL-17A pro-inflammatory cytokine has been associated with poor prognosis in some tumors[18]. GM-CSF (Granulocyte-Macrophage Colony-Stimulating Factor) is a hematopoietic growth factor that fulfils a fundamental role in macrophage and granulocyte differentiation. While classically linked to anti-tumor activities[19] there is growing evidence that GM-CSF can also promote tumor progression[20,21,22], supporting its inclusion in biomarker studies.

In a previous study carried out on a large group of melanoma patients, and based on serum proteomic analysis and immunoassays, we established prognostic value of serum Dermcidin (DCD) in stage II melanoma patients[23]. DCD is considered to play an important role in the cutaneous microenvironment due to its antimicrobial activity[24]. Nevertheless, DCD is not just an antimicrobial peptide as it can stimulate keratinocytes to produce cytokines though G-protein and mitogen-activated protein kinase activation[25]. These data suggest a possible relationship between *in situ* and systemic immune responses. Accordingly, the purpose of this study was to determine whether selected cytokine and DCD levels in serum could be used to develop a tool with clinical applications to improve the prognostic prediction of patients diagnosed at early stages of melanoma. To achieve this, we adopted a machine learning approach that incorporated the serum measurements of GM-CSF, IFN-γ, TGFβ1, IL-4, IL-6, IL-10, IL-17A and DCD, in conjunction with clinical-pathological features of such melanoma patients to determine the prognostic value of these parameters.

MATERIALS AND METHODS

Patients

A total of 448 patients were recruited at the Dermatology Units at the Basurto and Cruces University Hospitals between 2008 and 2016, of whom 261 (58%) were women and 187 (42%) men. To be eligible for inclusion in the study, patients had to fulfill the following criteria: 1) a histologically confirmed diagnosis of malignant melanoma; 2) no treatment except primary surgery; 3) no infection as judged by clinical evaluation and the absence of increased infectious parameters in the blood.

After surgery of the primary tumor, clinical check-ups were scheduled every three months for the first two years of the follow-up, and every six months thereafter, until a five-year follow-up had been completed. Annual revisions were then scheduled up to the tenth year post-surgery. The patients who developed metastasis during the follow-up period were again examined every three months for two years after metastasis had been diagnosed. The presence or absence of metastasis was assessed in all patients by physical examination, as well as through laboratory and radiological testing (X-rays and/or computed tomography –CT- scanning). Metastases were detected in 119 of the 448 melanoma patients (27%) during this study, including those in whom the disease had spread at the moment of diagnosis, and 79 patients died due to distant metastasis (40 women and 39 men).

Disease stages were classified according to the AJCC[7]. The clinical and diagnostic data for each patient were collected retrospectively from centralized electronic and/or paper medical records. For the statistical prediction analysis, only melanoma patients at early disease stages (I and II) were included, and the inclusion of the "disease-free" group required a minimum tracking of 2 years.

Ethics statement

This study was approved by the Euskadi Ethics Committee (reference 16-99) and written informed consents were obtained from all the subjects. The serum samples collected were stored at -80 ºC at the Basque Biobank until use.

Serum samples

Venous blood samples were drawn at the time of diagnosis and these samples were used to obtain serum following the protocol established at the Basque BioBank for Research. Accordingly, blood samples were allowed to clot at room temperature for at least 30 minutes and they were then centrifuged at 1000 g for 10 minutes. The serum was removed and was it subsequently divided into 500 µL aliquots, which were stored at -80 ºC until use.

Quantification of Granulocyte-Macrophage Colony-Stimulating Factor (GM-CSF), Interferon-γ (IFN-γ), Transforming Growth Factor beta (TGF-β1) and Interleukins (IL) 4, 6, 10 and 17A in serum

Upon reception, the serum samples were divided into 25 µL aliquots to avoid multiple freeze/thaw cycles, and the GM-CSF, IFN-γ, IL-4, 6, 10, 17A, and TGF-β1 in the samples was measured using magnetic bead-based multiple immunoassays (MILLIPLEX® MAP kit, Human High Sensitivity T Cell Magnetic Bead Panel: EMD Millipore Corporation, Germany). Each assay included two calibration curves for each of the proteins to be measured (calibration ranges: GM-CSF 1.22-5,000 pg/mL; IFN-γ 0.61-2,500 pg/mL; IL-4 1.83-7,500 pg/mL; IL-6 0.18-750 pg/mL; IL-10 1.46-6,000 pg/mL; IL-17A 0.73-3,000 pg/mL; and TGF-β1 9.8-10,000 ng/mL), with 8 calibration points in each curve. Two low and two high quality controls were also included in the assays. In the case of TGF-β1, serum samples were treated with 1N HCl, diluting the samples 1:4 and then adding 2 µL of 1.0 N HCl before incubating the mixture for 1 h at room temperature. The samples were then further diluted 1:6 in assay buffer to achieve a final dilution of 1:30. In the assays we followed the

protocol established by the manufacturer to quantify aforementioned proteins. Briefly, 25 µL of each standard or quality control were added to the wells of a 96-well plate and 25 µL of assay buffer was then added. Subsequently, 25 µL of the solution containing the magnetic beads was added to the wells, and the plate was sealed and incubated overnight in the dark at 4º C. The contents of each well were then removed, and the plate was washed three times before adding 25 µL of the detection antibodies and incubated at room temperature for 1 h. A streptavidin-phycoerythrin solution was then added (25 µL) to the wells and the plate was left at room temperature for 30 min. After washing 3 times, sheath fluid (150 µL) was added to each well (or 100 µL in the case of TGFβ1), before running the plate on a Luminex 100™ apparatus (Luminex Corporation, The Netherlands): 50 events per bead; 150 µL of sample (or 100 µL in the case of TGFβ1); gate settings from 8,000 to 15,000; reported gain as default and time out 100 s. The serum concentration of each of the proteins was calculated through a 5-parameter logistic curve-fitting method using the xPONENT® software (Luminex Corporation).

DCD Quantification in serum

Serum DCD was measured with an ELISA kit (Cusabio Biotech Co., Ltd, Houston, US) according to manufactures´ instructions, using two calibration curves for the protein to be measured in each assay (calibration range 0.625-40 ng/mL), with 8 calibration points for each curve. Briefly, 100 µL of the standards and samples (previously diluted 1/500) were added to the appropriate wells of a 96-well plate. The plate was sealed and incubated at 37 ºC for 2 h, the contents of the well were removed, and 100 µL of a biotin-antibody solution was added to the wells and incubated for 1 h. Subsequently, the well contents were removed and the plate was washed 3 times before 100 µL of a HRP-avidin solution was added to each well and incubated for 1 h at 37 ºC. The wells were then washed 5 times as above, 90 µL of a TMB substrate solution was then added and incubated at 37 ºC for 20 min, before 50 µL of the stop solution was added to each well. The optical density was determined on a microplate reader (Synergy HT, Biotek Instruments, Inc., Vermont, USA) set to 450 and 540 nm. Readings at 540 nm were subtracted from those obtained at 450 nm to correct for optical imperfections in the plate, and the serum DCD levels were calculated using the Gen5 software (2005, Biotek Instruments, Inc., Vermont, USA) with a 4-parameter logistic curve-fitting.

Statistical analysis

Variables were expressed as the median along with the 95% confidence interval (CI), computed by bootstrap resampling in which 10,000 samples were extracted with replacement for each variable from the original data and calculating the 95% percentile interval. Inter-group comparisons were carried out using the Kruskal-Wallis test when more than two groups were involved and a two-sided Mann–Whitney U test when only two groups were compared. In the latter case, in addition to the p-values, the effect sizes were also reported, measured through the absolute Cliff's Delta value that can be calculated from the U statistic as follows[26]:

$$\delta = 1.0 - \frac{2U}{N_1 \cdot N_2}$$

where N_1 and N_2 are the sample sizes of both groups. The effect size lies between -1 and 1, and it estimates the difference between the probability that a value from one of the groups is higher than that value from the other group, and vice versa. The P-values were corrected for multiple comparisons by controlling the False Discovery Rate (FDR) and those whose significance level was below the threshold of 0.05 were considered significant. Likewise, further inspection of statistical significance was addressed by means of the shift function[27], where the differences between the deciles of both groups are calculated in deciles. For independent groups this function is implemented in *shifthd_pbci* and it can be found in the *rogme* R package[28], where quantiles are compared using a Harrell-Davis estimator. Levels of confidence were computed by a bootstrap estimation of the standard error of the quantiles and the type I error was controlled to remain around 0.05 across all the quantile comparisons.

Machine learning analysis

A machine learning analysis was performed in order to assess the power of the data to correctly classify the prognosis of melanoma patients. We focused on patients diagnosed at stage I and II due to the clinical relevance of early metastasis prediction, of which there were 328 subjects in our cohort. Among these patients 244 remained disease free and 84 developed metastasis during the follow-up period. The predictive power of different biomarkers was inspected in three different variable domains: the Histological domain, represented by the Breslow thickness; the Serum domain, which involved all the serum variables indicated above; and a Multi-modal domain, a conjunction of the variables from the two previous domains. Missing information was imputed by removing instances containing unknown components, which reduced the input data to 211 disease-free and 56 metastatic samples, respectively. Subsequently, a nested cross-validation was employed

to assess both the optimization and generalization of the model. In the outer loop, a 5-times stratified 10-fold repeated cross-validation repeated 5 times with different randomization in each repetition was performed to estimate the generalization error of the model. In the inner loop, a stratified 10-fold cross-validation was implemented for model optimization, which involved the tuning of a pipeline assembled by robust scaling of the data, random over sampling of the class minority, L1-based feature selection and classifier hyperparameter fitting. A scheme of the workflow is shown in Figure S1 (Supplementary Material).

Classification scores were computed using a battery of five classification algorithms: Logistic Regression (LR) with a L2 regularization term; Support Vector Machine (SVM) with a radial basis kernel; a Decision Tree (DT); Gaussian Naive Bayes (NB) classifier; and the K-Nearest Neighbors vote (KNN) algorithm. All the different hyperparameters of the mentioned classifiers, the level of shrinkage and the number of features to select were tuned by an exhaustive grid search within the inner loop. For each classifier, the following scores were calculated:

$$balanced\,accuracy = 0.5 \cdot \left(\frac{TP}{P} + \frac{TN}{N} \right)$$

$$Precision = \frac{TP}{TP+FP}$$

$$Recall = \frac{TP}{TP+FN}$$

$$F_1^{weighted} = 2 \cdot \frac{Precision \cdot Recall}{Precision + Recall}$$

where P and N are the number of positive and negative cases, TP and TN the number of True Positive and Negative cases, FN and FP the number of False Negative and Positive cases. In addition, a Receiver Operating Characteristic (ROC) Curve was also computed, which displays the True Positive Rate (TPR: Sensitivity) against the False Positive Rate (FPR: 1-Specificity) as the threshold of the classifier varied. The area under this curve (AUC) provides a measure to evaluate the classifier quality, 1.0 being optimal since the rate of true positives is maximal and the rate of false negatives is minimal. The classifier with the highest ROC area was finally considered the most efficient one.

All the machine learning analysis was performed using *scikit-learn,* a library for machine learning written in python[29] and *Imbalanced-learn,* a Python toolbox to "tackle the curse of imbalanced datasets in Machine Learning"[30].

Survival analysis

Once the best algorithm and subset of biomarkers to reflect the evolution of metastasis had been found, we used this combination to fit the entire stage I/II sub-population, defining the ROC curve. Subsequently, the optimal cut-off point on this curve was determined using the Index of Union method, which corresponds to computing the value where the sensitivity and specificity are the closest to the AUC, and the absolute difference between the specificity and sensitivity is minimal[31]. This cut-off point allows us to define a class partition criterion, which separates subjects with a high probability of developing metastasis from those with a low probability, as witnessed through a Kaplan-Meier estimator implemented in a *lifelines* library in Python[32].

RESULTS

Patient characteristics

As mentioned above, 448 patients were included in this study (187 male, 261 female), with a median age at diagnosis of 56 years (95% CI 54.0-60.0: Table 1). Melanoma was most often diagnosed in patients' trunks (158 patients), followed by the lower limb (121 patients). Head and neck, upper limb and acral locations for melanomas were identified in 69, 49 and 34 patients, respectively. A large proportion of the lesions had a Breslow thickness less than 1 mm (231 melanomas) and between 1.01-4.00 mm (131 melanomas). Staging was based on the AJCC staging system[7] and most patients were diagnosed as stage I or II (224 and 99 patients, respectively), while only 38 were considered to be at a stage related with metastasis, stage III (30) or IV (8). However, 119 (27%) of the 448 patients recruited developed metastasis, including those with spread disease at the moment of diagnosis and those who suffered from disease recurrence during the follow-up. Distant and ganglion metastases were the main subtypes detected. Considering patients at AJCC stages I and II as early-stage melanoma patients (323 patients), 84 of these patients (25.6%) developed metastasis during the first years of the follow-up. In fact, the median interval from the removal of the primary tumor until the diagnosis of a metastasis was 1.9 years. By contrast, 239 patients remained disease-free (without recurrence or metastasis) and the median follow-up of these patients was 4.5 years.

Analysis of serum GM-CSF, IL-4, IL-6, IL-10, IL-17A, IFN-γ, TGFβ and DCD

The amount of GM-CSF, IL4, IL-6, IL-10, IL-17A and TGFβ detected in serum was independent of the age of the melanoma patients and it did not vary between the sexes. However, there were significant differences between the sexes in the levels of IFNγ and DCD ($|\delta|= 0.2$, $p_{FDR}<0.01$ and $|\delta| = 0.2$, $p_{FDR}<0.01$, respectively: data not shown). Of the proteins studied, the median serum level in the melanoma patients at the time of diagnosis was considered according to the stage of the tumor (Table 2). There appeared to be no significant differences in the median serum levels in patients of different AJCC stages, nor were any differences found between the distinct histological subtypes of melanoma (data not shown).

To analyze the prognostic value of these proteins, we assessed the serum cytokine and DCD levels in melanoma patients diagnosed at stages I/II and that remained disease-free at the end of the follow-up period, comparing these with those in patients who developed metastasis. There were 84 of 323 (25.6%) stage I/II melanoma patients who developed metastasis during the follow-up and there appeared to be a significant difference in the serum IL-4 and IL-6 levels between these two groups of patients, and associated with a moderate effect size ($|\delta|= 0.30$ $p_{FDR} <0.01$ and $|\delta|= 0.20$ $p_{FDR} = 0.04$: Table 3). At the time of diagnosis, the serum IL-4 levels of patients who developed metastasis were double those observed in patients who remained disease-free (62.27 pg/mL *versus* 31.96 pg/mL, respectively, p<0.01). In addition, there was a significant difference in the serum IL-6 levels between the two groups of patients (4.71 pg/mL *versus* 3.29 pg/mL respectively, p<0.04). No significant differences in serum IL-10, IL-17A, IFN-γ, TGFβ and DCD were observed.

Prognostic power of the melanoma markers

The performance of the different classifiers was assessed for the subpopulation of subjects at AJCC stages I and II (Table S1). In all the three domains, a LR classifier exhibited the best performance through the ROC area, and the most generalizable results reflected by the smallest gap between the training and test scores. The Breslow thickness represents a biomarker of melanoma metastasis that correctly classified 73% of the patients and generating 83% of the ROC area (Figure 1A). Although the serum levels had a poorer prognostic value, when combined with the Breslow thickness they significantly improved the overall performance of this biomarker (Wilcoxon signed-rank test, p<0.01), exceeding a balanced accuracy of 80% and a ROC area close to the 90%. Furthermore, these data clearly pointed to the cytokines IL-4 and GM-CSF as the biomarkers that best predicted melanoma metastasis in conjunction with the Breslow thickness. Indeed, these two parameters were

selected in almost all the partitions within the inner cross-validation (see panel B of Figure 1). This subset of variables was followed at a distance by the IL-10, IL-6 and DCD proteins.

Similarly, the decile distribution for these potential biomarkers exhibited a clear tendency to separate between subjects who developed metastasis and those who remained disease-free (panel A in Figure 2). Both the distributions of the serum IL-4 levels and the Breslow thickness were higher in the metastatic subpopulation, whereas this tendency switched towards lower levels of GM-CSF. Moreover, when the differences in the distribution of these variables was addressed by means of the shift function, their predictive power was clearly evident, especially that of the Breslow thickness where the separation of the melanoma outcome was significant across its whole spectrum. This was followed by that of IL-4, which began to display a significant separation around the median, whereas GM-CSF started to discriminate these subpopulations above its 8^{th} decile (panel B in Figure 2).

Importantly, these findings can be easily incorporated into clinical protocols by providing a general optimum cut-off from the data from which a prediction of metastasis can be performed. Using the same subpopulation of subjects (I/II melanoma patients), we fitted the entire data using the best classifier and the subset of features found previously (i.e.: a LR classifier with the features Breslow thickness, GM-CSF and IL-4), and we computed the optimal point on the ROC curve that corresponded in this case to FPR (1-Specificity) = 0.15 and TPR (Sensitivity) = 0.80 (see panel A, Figure 3). This point defines a critical threshold that allows us to separate subjects in terms of their prognosis, which for our classifier can be easily translated into a constraint as follows:

$$\left(\frac{Breslow}{1.41\,mm}\right)-\left(\frac{GM-CSF}{571.28\,pg/mL}\right)+\left(\frac{IL-4}{168.1\,pg/mL}\right)=1.46$$

This equation therefore defines a plane in our 3D feature space such that any subject lying above it is classified as metastatic and those below it are considered disease-free. Furthermore, subjects stratified with respect to this critical threshold could be differentiated by their probability of eventually developing metastasis during the follow-up period (Kaplan-Meier, p = 2.82 x10^{-12}, as shown in panel B of Figure 3).

Remarkably, we found a *prognostic plane* involving the serum levels of IL-4 and GM-CSF in conjunction with the Breslow thickness that could accurately classify subjects according to their melanoma outcome. This equation could be easily translated to a clinical setting and inspecting the

signs of the coefficients in this equation, we can clearly see that an increase in IL-4 and the Breslow thickness tend to shift subjects above this plane, indicating a worse prognosis, whereas GM-CSF levels act in the opposite direction.

DISCUSSION

An accurate diagnosis is an essential first step in cancer management. Most melanoma cases are detected at early disease stages and when possible, excision biopsy can be a successful procedure to treat suspicious melanocytic lesions. According to the AJCC classification, Breslow thickness, together with the ulceration and mitotic index, are important variables that should be considered in tumor staging (https://cancerstaging.org/). In addition, several histological biomarkers (e.g. Melan-A, Pmel) are routinely employed for diagnostic porpoises[33]. In this regard, important efforts are being made in order to achieve less invasive techniques or to improve the accuracy of diagnostic markers[33]. Indeed, early and precise prognostic markers are urgently needed for melanoma due to its strong metastatic capacity, particularly given the low survival rate of metastatic patients and the poor 5-year relapse free rate of just 56%, even when melanoma is detected at early stages (stage II)[34,1]. The highest recurrence rate is observed between 2 and 3 years after surgical treatment, even though the probability of recurrence is less than 5% in stage I-III melanoma patients that receive treatment, as was the 5 year disease-free follow-up[35]. According to these figures, intense medical monitoring should be implemented in the first 2-3 years after treatment, even for stage II cases, representing an important medical and economic burden. Therefore, it would be useful if early patients could be rapidly classified into high or low recurrence risk groups when contemplating efficient and sustainable personalized follow-up programs. Moreover, to assuage the unpredictable clinical behavior of melanoma much research has focused on the discovery of prognostic factors to improve the prognostic accuracy for this type of skin cancer[3]. In this regard, our study focused on the discovery of prognostic biomarkers capable of evaluating the metastatic risk of patients identified at early stages of the disease (stage I-II). Moreover, we defined a clinically applicable mathematical tool to accurately classify such melanoma patients.

Currently, predicting patient outcome mainly relies on staging based on the histopathological parameters described previously, while treatment options are often based on the BRAF mutation[36]. Nevertheless, patient monitoring, especially upon surgical removal of the primary tumor, requires

other variables to be analyzed. As a systemic system for information transfer, serum represents a complex but accessible sensor. To date, LDH has been one and perhaps the only clinical serological biomarker for melanoma, with increasing values interpreted as reflecting disease progression. However, an increase in serum LDH levels may also occur in other settings, which means employing some caution before reaching any conclusions[36].

Our previous attempt to identify novel serological prognostic markers identified a threshold for serological DCD that was associated with a poor prognosis value for melanoma patients diagnosed specifically at AJCC stage II[23]. Consistent with this finding, DCD, a major human antimicrobial peptide in human skin[24,25] was also recently proposed as a serological marker for the diagnosis and staging of hepatocellular carcinoma[37]. The current study of 448 new patients revealed DCD to be a marker of metastatic progression, although other serological parameters appear to have greater predictive potential than DCD, such as IL-4, GM-CSF or IL-6. These differences may be due to the patient stratification, as both stage I and stage II melanoma patients were previously included in separate groups, or the length of the follow-up period (10 *vs* 4.5 years for disease-free patients). DCD is expressed in melanoma cell lines, and it could be involved in autophagy and apoptotic cell death, yet the underlying molecular mechanisms and the true clinical-pathological relevance of this is not fully understood[38].

Serological cytokines reflect the general immunological state of the body, offering information regarding the cytokines released by tumors and those that accumulate in the tumor microenvironment[39]. The melanoma microenvironment contains stromal cells and immune cells like T or B lymphocytes, NK cells or tumor-associated macrophages[39]. Most of these cells secrete cytokines that may play a key role in inhibiting or promoting tumor progression[40]. The pro-inflammatory cytokines IL-4 and IL-6, produced either by host immune cells or tumor cells themselves, are associated with tumor malignancy in patients and animal cancer models[41,42,43]. At the cutaneous level, keratinocytes secrete IL-6 in order to enhance T cell-mediated antitumor activity and therefore, high IL-6 levels are considered a marker for immune system upregulation[42,43]. IL-4 is the most important Th2 cytokine, and it is mainly produced by activated T cells, mast cells, basophils and eosinophils in order to regulate lymphocyte proliferation and survival[42]. Interestingly, elevated serum IL-6 was correlated with a poor prognosis in melanoma, while IL-4 is thought to promote the proliferation and survival of several cancer cells[44,45,46]. In line with previous findings, here early stage (I or II) melanoma patients that developed metastasis had significantly higher levels of serum IL-4 and IL-6 than patients who did not develop metastasis during the follow-up.

The Breslow thickness is a crucial prognostic factor, with substantial evidence confirming a direct relationship between Breslow thickness and survival[6]. Accordingly, we show that Breslow thickness is an important risk factor for the malignant progression of melanoma. Nonetheless, a significant increase in the predictive power of Breslow thickness was achieved by combining it with data regarding serum IL-4 and GM-CSF, resulting in the development of an algorithm to identify early stage melanoma patients with a high risk of developing metastasis during the follow-up. According to this algorithm, a high Breslow thickness and serum IL-4 levels in early stage melanoma patients are associated with a poor prognosis, whereas GM-CSF levels decrease in patients in whom the disease outcome is poor. These results are consistent with other studies describing an antitumor effect of GM-CSF[47]. Our data also revealed the importance of IL-10 and IL-6 in predicting metastatic progression, although they did not appear to add additional information to the predictive equation involving the Breslow thickness, and serum IL-4 and GM-CSF.

In summary, the use of machine learning techniques has helped to define an algorithm capable of accurately classifying early stage melanoma patients with a high and low risk of developing metastasis during the period of highest recurrence, 2 years. The equation generated took into account the serum IL-4 and GM-CSF levels, and the Breslow index, and it could stratify melanoma patients to be triaged at the time of diagnosis and initial surgery, or it could also be used clinically to determine whether stage in I or II melanoma patients should receive adjuvant therapy to prevent metastatic progression.

ACKNOWLEDGEMENTS

We are grateful to the Basque Biobank for providing the serum samples. We are also most grateful to Drs Arantza Arrieta and Natalia Maruri (Cruces University Hospital) for their technical support with the serum marker detection. This work was supported by grants from the Basque Government (KK2016-036 and KK2017-041 to MDB) and UPV/EHU (GIU17/066 to MDB).

REFERENCES

1. Tracey EH, Vij A. Updates in Melanoma. Dermatol Clin. 2019; 37(1):73-82.

2. Whiteman DC, Green AC, Olsen CM. The Growing Burden of Invasive Melanoma: Projections of Incidence Rates and Numbers of New Cases in Six Susceptible Populations through 2031. J Invest Dermatol. 2016; 136(6):1161-1171.

3. Kashani-Sabet M, Nosrati M, Miller JR 3rd, Sagebiel RW, Leong SPL, Lesniak A, Tong S, Lee SJ, Kirkwood JM. Prospective Validation of Molecular Prognostic Markers in Cutaneous Melanoma: A Correlative Analysis of E1690. Clin Cancer Res. 2017; 23(22):6888-6892.

4. Kashani-Sabet M. Molecular markers in melanoma. Br J Dermatol. 2014; 170(1):31-5.

5. Foth M, Wouters J, de Chaumont C, Dynoodt P, Gallagher WM. Prognostic and predictive biomarkers in melanoma: an update. Expert Rev Mol Diagn. 2016; 16(2):223-37.

6. Stiegel E, Xiong D, Ya J, Funchain P, Isakov R, Gastman B, Vij A. Prognostic value of sentinel lymph node biopsy according to Breslow thickness for cutaneous melanoma. J Am Acad Dermatol. 2018; 78(5):942-948.

7. Balch CM, Gershenwald JE, Soong SJ, Thompson JF, Atkins MB, Byrd DR, Buzaid AC, Cochran AJ, Coit DG, Ding S, Eggermont AM, Flaherty KT, Gimotty PA, Kirkwood JM, McMasters KM, Mihm MC Jr, Morton DL, Ross MI, Sober AJ, Sondak VK. Final version of 2009 AJCC melanoma staging and classification. J Clin Oncol. 2009; 27(36):6199-206.

8. Elmore JG, Elder DE, Barnhill RL, Knezevich SR, Longton GM, Titus LJ, Weinstock MA, Pepe MS, Nelson HD, Reisch LM, Radick AC, Piepkorn MW. Concordance and Reproducibility of Melanoma Staging According to the 7th vs 8th Edition of the AJCC Cancer Staging Manual. JAMA Netw Open. 2018; 1(1).

9. Weiss SA, Hanniford D, Hernando E, Osman I. Revisiting determinants of prognosis in cutaneous melanoma. Cancer. 2015; 121(23):4108-23.

10. Gogas H, Eggermont AMM, Hauschild A, Hersey P, Mohr P, Schadendorf D, Spatz A, Dummer R. Biomarkers in melanoma. Ann Oncol. 2009; 20 Suppl 6:vi8-13.

11. Palmer SR, Erickson LA, Ichetovkin I, Knauer DJ, Markovic SN. Circulating serologic and molecular biomarkers in malignant melanoma. Mayo Clin Proc. 2011; 86(10):981-90.

12. Johdi NA, Mazlan L, Sagap I, Jamal R. Profiling of cytokines, chemokines and other soluble proteins as a potential biomarker in colorectal cancer and polyps. Cytokine. 2017; 99:35-42.

13. Obraztsov IV, Shirokikh KE, Obraztsova OI, Shapina MV, Wang MH, Khalif IL. Multiple Cytokine Profiling: A New Model to Predict Response to Tumor Necrosis Factor Antagonists in Ulcerative Colitis Patients. Inflamm Bowel Dis. 2019; 25(3):524-531.

14. D'Angelo C, Reale M, Costantini E, Di Nicola M, Porfilio I, de Andrés C, Fernández-Paredes L, Sánchez-Ramón S, Pasquali L. Profiling of Canonical and Non-Traditional Cytokine Levels in Interferon-β-Treated Relapsing-Remitting-Multiple Sclerosis Patients. Front Immunol. 2018; 9:1240.

15. Nevala WK, Vachon CM, Leontovich AA, Scott CG, Thompson MA, Markovic SN; Melanoma Study Group of the Mayo Clinic Cancer Center. Evidence of systemic Th2-driven chronic inflammation in patients with metastatic melanoma. Clin Cancer Res. 2009; 15(6):1931-9.

16. Boyano MD, Garcia-Vázquez MD, López-Michelena T, Gardeazabal J, Bilbao J, Cañavate ML, García de Galdeano A, Izu R, Díaz-Ramón L, Raton JA, Díaz-Pérez JL. Soluble interleukin-2 receptor, intercellular adhesion molecule-1 and interleukin-10 serum levels in patients with melanoma. Br J Cancer. 2000; 83(7):847-52.

17. Boyano MD, García-Vázquez MD, Gardeazabal J, García de Galdeano A, Smith-Zubiaga I, Cañavate ML, Raton JA, Bilbao I, Díaz-Pérez JL. Serum-soluble IL-2 receptor and IL-6 levels in patients with melanoma. Oncology. 1997; 54(5):400-6.

18. Ma YF, Chen C, Li D, Liu M, Lv ZW, Ji Y, Xu J. Targeting of interleukin (IL)-17A inhibits PDL1 expression in tumor cells and induces anticancer immunity in an estrogen receptor-negative murine model of breast cancer. Oncotarget. 2017; 8(5):7614-7624.

19. Bhattacharya P, Thiruppathi M, Elshabrawy HA, Alharshawi K, Kumar P, Prabhakar BS. GM-CSF: An immune modulatory cytokine that can suppress autoimmunity. Cytokine. 2015; 75(2):261-71.

20. Singel KL, Segal BH. Neutrophils in the tumor microenvironment: trying to heal the wound that cannot heal. Immunol Rev. 2016; 273(1):329-43.

21. Reggiani F, Labanca V, Mancuso P, Rabascio C, Talarico G, Orecchioni S, Manconi A, Bertolini F. Adipose Progenitor Cell Secretion of GM-CSF and MMP9 Promotes a Stromal and Immunological Microenvironment That Supports Breast Cancer Progression. Cancer Res. 2017; 77(18):5169-5182.

22. Wang TT, Zhao YL, Peng LS, Chen N, Chen W, Lv YP, Mao FY, Zhang JY, Cheng P, Teng YS, Fu XL, Yu PW, Guo G, Luo P, Zhuang Y, Zou QM. Tumour-activated neutrophils in gastric cancer foster immune suppression and disease progression through GM-CSF-PD-L1 pathway. Gut. 2017; 66(11):1900-1911.

23. Ortega-Martínez I, Gardeazabal J, Erramuzpe A, Sanchez-Diez A, Cortés J, García-Vázquez MD, Pérez-Yarza G, Izu R, Luís Díaz-Ramón J, de la Fuente IM, Asumendi A, Boyano MD. Vitronectin and dermcidin serum levels predict the metastatic progression of AJCC I-II early-stage melanoma. Int J Cancer. 2016; 139(7):1598-607.

24. Zeth K, Sancho-Vaello E. The Human Antimicrobial Peptides Dermcidin and LL-37 Show Novel Distinct Pathways in Membrane Interactions. Front Chem. 2017; 5: 86.

25. Paulmann M, Arnold T, Linke D, Özdirekcan S, Kopp A, Gutsmann T, Kalbacher H, Wanke I, Schuenemann VJ, Habeck M, Bürck J, Ulrich AS, Schittek B. Structure-Activity Analysis of the

Dermcidin-derived Peptide DCD-1L, an Anionic Antimicrobial Peptide Present in Human Sweat. J Biol Chem. 2012; 287(11): 8434-43.

26. Cliff, N. Dominance statistics: Ordinal analyses to answer ordinal questions. Psychol Bull 1993; 114(3):494-509.

27. Rand R. Wilcox. Introduction to Robust Estimation and Hypothesis Testing, 3rd ed., California, Elsevier, 2012. 608p

28. Rousselet GA, Pernet CR, Wilcox RR. Beyond differences in means: robust graphical methods to compare two groups in neuroscience. Eur J Neurosci 2017; 46(2):1738-1748.

29. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, et al. Scikit-learn: Machine Learning in Python. J Mach Learn Res 2011; 12:2825–30.

30. Lemaître G, Nogueira F, Aridas CK. Imbalanced-learn: A Python Toolbox to Tackle the Curse of Imbalanced Datasets in Machine Learning. J Mach Learn Res 2017; 18:1–5.

31. Ilker Unal. Defining an Optimal Cut-Point Value in ROC Analysis: An Alternative Approach. Comput Math Methods Med 2017; 2017:14.

32. Davidson-Pilon C, Kalderstam J, Kuhn B, Fiore-Gartland A, Moneda L, Zivich P, Parij A, Stark K, Anton S, Besson, Jona, Harsh Gadgil, et al. Software Open Access CamDavidsonPilon/lifelines: v0.14.3. 2018.

33. Mohammadpour A, Derakhshan M, Darabi H, Hedayat P, Momeni M. Melanoma: Where we are and where we go. J Cell Physiol. 2019; 234(4):3307-3320.

34. Svedman FC, Pillas D, Taylor A, Kaur M, Linder R, Hansson J. Stage-specific survival and recurrence in patients with cutaneous malignant melanoma in Europe - a systematic review of the literature. Clin Epidemiol. 2016; 8:109-22.

35. Rutkowski P, Lugowska I. Follow-up in melanoma patients. Memo. 2014; 7(2):83-86.

36. Karagiannis P, Fittall M, Karagiannis SN. Evaluating biomarkers in melanoma. Front Oncol. 2015; 4:383.

37. Qiu F, Qiu F, Liu L, Liu J, Xu J, Huang X. The Role of Dermcidin in the Diagnosis and Staging of Hepatocellular Carcinoma. Genet Test Mol Biomarkers. 2018; 22(4):218-223.

38. Trzoss L, Fukuda T, Costa-Lotufo LV, Jimenez P, La Clair JJ, Fenical W. Seriniquinone, a selective anticancer agent, induces cell death by autophagocytosis, targeting the cáncer-protective protein dermcidin. Proc Natl Acad Sci U S A. 2014; 111(41):14687-92.

39. Wang M, Zhao J, Zhang L, Wei F, Lian Y, Wu Y, Gong Z, Zhang S, Zhou J, Cao K, Li X, Xiong W, Li G, Zeng Z, Guo C. Role of tumor microenvironment in tumorigenesis. J Cancer. 2017; 8(5):761-773.

40. Jiang X, Wang J, Deng X, Xiong F, Ge J, Xiang B, Wu X, Ma J, Zhou M, Li X, Li Y, Li G, Xiong W, Guo C, Zeng Z. Role of the tumor microenvironment in PD-L1/PD-1-mediated tumor immune escape. Mol Cancer. 2019; 18(1):10.

41. Ito SE, Shirota H, Kasahara Y, Saijo K, Ishioka C. IL-4 blockade alters the tumor microenvironment and augments the response to cancer immunotherapy in a mouse model. Cancer Immunol Immunother. 2017; 66(11):1485-1496.

42. Setrerrahmane S and Xu H. Tumor-related interleukins: old validated targets for new anti-cancer drug development. Mol Cancer. 2017; 16(1):153.

43. Surcel M, Constantin C, Caruntu C, Zurac S, Neagu M. Inflammatory Cytokine Pattern Is Sex-Dependent in Mouse Cutaneous Melanoma Experimental Model. J Immunol Res. 2017; 2017:9212134.

44. Jobe NP, Živicová V, Mifková A, Rösel D, Dvořánková B, Kodet O, Strnad H, Kolář M, Šedo A, Smetana K Jr, Strnadová K, Brábek J, Lacina L. Fibroblasts potentiate melanoma cells in vitro invasiveness induced by UV-irradiated keratinocytes. Histochem Cell Biol. 2018; 149(5):503-516.

45. Yu SH, Maynard JP, Vaghasia AM, De Marzo AM, Drake CG, Sfanos KS. A role for paracrine interleukin-6 signaling in the tumor microenvironment in prostate tumor growth. Prostate. 2019; 79(2):215-222.

46. Gocheva V, Wang HW, Gadea BB, Shree T, Hunter KE, Garfall AL, Berman T, Joyce JA. IL-4 induces cathepsin protease activity in tumor-associated macrophages to promote cancer growth and invasion. Genes Dev. 2010; 24(3):241-55.

47. Hong IS. Stimulatory versus suppressive effects of GM-CSF on tumor progression in multiple cancer types. Exp Mol Med. 2016; 48(7):e242.

Figure Legends

Figure 1. Logistic Regression analysis. A) Classification of the three variable domains considered: Breslow thickness, Cytokines and DCD serum variables. B) In the scenario combining histological and serum variables, their participation across the folds provided by the feature selection step in the inner cross-validation loop.

Figure 2. A) Decile distribution of metastatic and disease-free subjects for the best groups of biomarkers. B) Shift function displaying the difference between the deciles for both data distributions. Positive values of the shift function are in blue, corresponding to larger decile values in the disease-free group than in the metastatic group, while red values illustrate the opposite scenario.

Figure 3. A) ROC curve from the whole stage I/II dataset. The optimal cut-off point on this curve defines a plane that maximally separates metastatic and disease-free progression. The best subset of biomarkers corresponds to Breslow thickness, IL-4 and GM-CSF. B) Kaplan-Meier analysis. The cut-off plane provides a condition to significantly separate subjects with a worse prognosis from those with a better prognosis.

Figure S1. Workflow of the machine learning analysis.

Table 1. Patient characteristics

Characteristic	Number (m/f)
Melanomas	448 (187/261)
Disease-free	329 (136/193)
Metastasis	119 (51/68)
Tumor location	
Head/neck	69 (33/36)
Trunk	158 (98/60)
Upper limb	49 (16/33)
Lower limb	121 (22/99)
Hand/foot	34 (10/24)
Others	7 (1/6)
Unknown	10 (7/3)
Stage[1]	
0 (*in situ*)	78 (29/49)
I	224 (94/130)
II	99 (40/59)
III	30 (16/14)
IV	8 (4/4)
Unknown	9 (4/5)
Breslow Thickness (mm)	
≤ 1.00	231 (98/133)
1.01-4.00	131 (43/88)
> 4.01	41 (25/16)
Unknown	45 (21/24)

[1]The American Joint Committee of Cancer (AJCC) staging system for melanomas was used. Abbreviations: m/f, males/females; SSM, Superficial Spreading Melanoma; NM, Nodular Melanoma; ALM, Acral Lentiginous Melanoma; LMM, Lentigo Malignant Melanoma.

Table 2. Cytokines and DCD serum levels in melanoma patients.

	Melanomas (n = 448)	AJCC stage[1]					p_{FDR}
		In situ (n = 78)	I (n =224)	II (n =99)	III (n = 30)	IV (n = 8)	
GM-CSF	120.32 [103.60-131.55]	108.49 [78.96-163.75]	122.56 [107.17-140.39]	137.86 [104.20-170.74]	84.68 [42.66-116.94]	63.48 [18.09-280.48]	0.49
IL-4	35.53 [30.20-39.06]	27.76 [20.07-37.21]	35.95 [28.88-41.5]	37.81 [28.42-56.31]	52.82 [27.86-63.00]	33.33 [8.49-136.30]	0.41
IL-6	3.32 [3.06 - 3.87]	3.11 [2.02-3.84]	3.30 [2.91-3.98]	4.27 [3.23-5.46]	2.62 [1.80-5.42]	4.97 [1.11-6.65]	0.41
IL-10	10.12 [8.29-13.11]	8.29 [5.16-13.15]	10.23 [7.95 -13.76]	11.34 [7.38-16.35]	10.83 [6.67-18.51]	10.71 [5.51-30.81]	0.61
IL-17A	17.46 [15.55-19.37]	12.88 [10.87-17.60]	19.35 [17.12-22.24]	16.51 [12.85-22.07]	17.84 [10.0-22.92]	16.43 [10.97-26.67]	0.41
IFN-γ	17.91 [15.61-19.34]	15.08 [10.67-18.88]	17.95 [15.92-20.04]	19.07 [13.94-24.57]	12.60 [10.09-24.98]	21.26 [9.66 -50.75]	0.41
TGF-β	48.58 [45.93-5281]	49.36 [47.34-57.40]	49.18 [45.3-54.21]	46.67 [40.13-55.30]	43.54 [39.46-66.04]	63.04 [34.91-85.40]	0.61
DCD	4.66 [4.42-4.86]	4.82 [4.40-5.65]	4.81 [4.31-5.01]	4.43 [3.98-4.88]	4.18 [3.08-4.88]	4.10 [1.63-6.4]	0.49

Serum levels of GM-CSF, IL-4, IL-6, IL-10, IL-17A and IFN-γ are expressed in pg/mL. TGFβ levels are expressed in ng/mL and DCD in μg/mL. p_{FDR} means False Discovery Rate. Square brackets reflect the Lower and Upper confidence intervals of the median. [1]The American Joint Committee of Cancer (AJCC) staging system for melanomas was used.

Table 3. Comparison of serum cytokine and DCD levels between AJCC stage I and II patients who were disease-free or developed metastasis during follow-up.

	Disease progression		$(|\delta|, p_{FDR})$
	Disease-free	**Metastasis**	
GM-CSF	121.39	131.55	(0.03, 0.72)
	[103.17-138.63]	[101.01-153.24]	
IL-4	31.96	62.27	**(0.30, <0.01)**
	[27.76-38.06]	[39.06-92.21]	
IL-6	3.29	4.71	**(0.20, 0.04)**
	[2.82-3.87]	[3.3- 5.9]	
IL-10	11.23	10.03	(0.05, 0.67)
	[7.89-14.83]	[7.78 - 15.31]	
IL-17A	17.88	20.09	(0.04, 0.67)
	[16.33-20.06]	[14.89-24.33]	
IFN-γ	17.23	22.40	(0.12, 0.37)
	[14.84-19.34]	[16.99-26.91]	
TGF-β	49.71	48.18	(0.04, 0.67)
	[45.32-53.68]	[40.13-60.44]	
DCD	4.78	4.38	(0.07, 0.67)
	[4.39-5.01]	[3.92-4.84]	

Serum levels of GM-CSF, IL-4, IL-6, IL-10, IL-17A and IFN-γ are expressed in pg/mL. TGFβ levels are expressed in ng/mL and DCD in μg/mL. $|\delta|$= Cliff's Delta and p_{FDR} means False Discovery Rate. Square brackets reflect the Lower and Upper confidence intervals for the median.

Table S1. Train and test prediction scores in the three domains of variables for the battery of algorithms used.

	Train Balanced accuracy	Test Balanced accuracy	Train Recall	Test Recall	Train Precision	Test Precision	Train Area ROC	Test Area ROC
BRESLOW THICKNESS								
LR	74.92	73.02	72.65	69.50	52.98	51.69	83.45	83.72
SVM	74.03	73.57	65.65	64.94	56.37	58.4	82.28	81.51
DT	77.93	72.56	76.81	69.06	57.07	53.5	84.07	78.72
NB	72.66	72.56	54.76	54.67	66.65	69.17	83.08	83.28
KNN	78.69	73.26	80.32	72.94	54.99	49.67	87.65	79.17
CYTOKINES + DCD								
LR	64.16	57.53	60.75	49.13	33.41	27.32	70.36	64.02
SVM	69.54	58.67	56.34	39.00	46.85	32.14	75.39	63.02
DT	74.63	55.55	74.68	45.87	47.05	28.34	79.66	57.18
NB	60.87	55.16	49.14	39.07	40.04	27.00	64.89	58.98
KNN	72.96	53.81	76.97	47.87	41.04	24.23	80.51	56.16
BRESLOW THICKNESS + CYTOKINES + DCD								
LR	82.60	80.39	79.41	76.13	59.98	59.74	90.87	89.50
SVM	79.87	76.70	73.38	68.20	59.09	57.53	87.51	84.19
DT	86.25	76.70	87.71	68.27	62.12	50.22	92.57	83.41
NB	79.90	77.33	70.36	65.47	64.07	61.74	87.41	84.94
KNN	85.25	75.60	86.62	71.20	59.15	50.13	92.38	82.93

Figure 1

Figure 2

Figure 3

Figure S1