The landscape of myeloid and astrocyte phenotypes in acute multiple sclerosis lesions

Calvin Park[1], Gerald Ponath[1], Maya Levine-Ritterman[1], Edward Bull[1], Eric Swanson[2], Benjamin M. Segal[3], David
Pitt[1]

[1] Department of Neurology, Yale School of Medicine, New Haven, CT, USA
[2] Fluidigm Corporation, Markham, ON, Canada
[3] Department of Neurology, University of Michigan, Ann Arbor, MI, USA

Author contact information
Calvin Park, Email: calvin.park@yale.edu
Gerald Ponath, Email: gerald.ponath@yale.edu
Maya Levine-Ritterman, Email: maya.levine-ritterman@yale.edu
Edward Bull, Email: embull@bowdoin.edu
Eric Swanson, Email: Eric.Swanson@fluidigm.com
Benjamin M. Segal, Email: bmsegal@med.umich.edu

Corresponding author
David Pitt, MD
Department of Neurology
Yale School of Medicine
300 George Street, Suite 353I
New Haven, CT 06511
david.pitt@yale.edu
Phone: 203-737-7951
Fax: 203-737-2704

Abstract

Activated myeloid cells and astrocytes are the predominant cell types in acute multiple sclerosis (MS) lesions. The

polarization states of these cells play a critical role in lesion formation and resolution. To identify myeloid and

astrocyte subsets present during lesion development, we profiled the cellular composition of an acutely demyelinating

and an acute post-demyelinating MS lesion with thirteen immune markers, using imaging mass cytometry (IMC). In

the early demyelinating lesion, distinct myeloid and astrocyte phenotypes populated separate lesion zones and engaged

in specific interactions. In the late active lesion, phenotypes were less diverse and more uniformly distributed

throughout the lesion. Our study demonstrates that multiplexed tissue imaging, combined with new computational

tools, is a powerful approach to study cellular heterogeneity and spatial organization in MS lesions. Identifying glial

phenotypes and their key interactions at different lesion stages may elucidate novel therapeutic targets for inhibiting

acute demyelination and low-grade neuro-inflammation.

Introduction

Multiple sclerosis (MS) is a common neurological disease, characterized by formation of inflammatory demyelinating lesions in the central nervous system (CNS) [21]. Inflammation is driven by infiltrating lymphocytes and monocytes, in concert with resident activated microglia and astrocytes. Macrophages and reactive astrocytes are the most abundant cell types in acute lesions [14,24]. These cells are highly plastic and can adopt pro-inflammatory, anti-inflammatory, neurotoxic, neuroprotective, and tissue-regenerating functions [3,5,6,18,23,34].

Evaluating these cell phenotypes *in situ* is challenging. Previous studies have identified macrophage phenotypes in MS lesions based on the expression of single classical (M1) and alternative (M2) activation markers; however, those studies have produced limited and sometimes conflicting results [5,34]. Moreover, a recent study of traumatic brain injury in mice demonstrated random expression of M1/M2 polarization markers in infiltrating macrophages in seemingly incompatible combinations [12]. This suggests that the M1/M2 polarization paradigm, which originated as an *in vitro* concept, is of limited value for characterizing myeloid cells in the inflamed CNS [28]. Instead, macrophages within MS lesions exhibit complex phenotypes that cannot be captured by one, or several, M1/M2 markers, but instead require high parameter evaluation.

Powerful new technologies, such as mass cytometry (CyTOF) and single cell RNA-sequencing, make it possible to explore the heterogeneity of individual cell types. However, these methods do not capture the spatial information contained in tissue, such as distribution and interactions of cellular subpopulations. This problem can now be overcome with a novel imaging technology, imaging mass cytometry (IMC), which allows for quantitative profiling with up to 37 antibodies on a single tissue section. Like CyTOF, IMC relies on metal isotope-labeled antibodies, and combines immunohistochemistry with high-resolution laser ablation followed by time-of-flight mass spectrometry [35]. In parallel, novel computational tools have become available to extract single-cell information from highly multiplexed histological data.

We applied IMC and single-cell analytics to two active MS lesions – one acutely demyelinating and one active but post-demyelinating – to examine the cellular heterogeneity of myeloid cells and astrocytes, based on thirteen immune function markers. We demonstrate here that multiplexed tissue imaging, in combination with the appropriate computational tools, can extract previously unattainable information on single cells from histological sections. This includes cellular subpopulations and their distribution within the lesion environment, specific cell-cell interactions,

phenotypic transitions, and the impact of the lesion environment on marker expression. High-dimensional

characterization of MS lesions will ultimately provide a complete picture of polarization states, and of the nodal points

of cellular communication that are critical for lesion formation, chronic low-grade inflammation, and lesion resolution.

Results

Histology and cell clustering overview

We analyzed one actively demyelinating and one active post-demyelinating MS lesion, according to a classification

by Kuhlmann et al. [13], referred to throughout this report as the "early" and "late" lesion, respectively. Both lesions

were located in the brain stem and characterized by complete loss of myelin, hypercellularity with the highest cellular

density at the lesion rim, and diffuse infiltration with foamy macrophages. The laser-scanned area of the early lesion

involved predominantly white matter (WM) but also interspersed gray matter (G/WM), while the scanned area of the

late lesion consisted of only WM (Figure 1).

Consistent with demyelinating activity, macrophages at the rim of the early lesion contained myelin basic protein

(MBP)-positive myelin debris, which was absent in macrophages from the late lesion (Figure 1d, j) [13]. Moreover,

foamy macrophages were numerous and large in size in the rim of the early lesion, whereas macrophages in the

rim of the late lesion were smaller and less concentrated (Figure 1e, k).

Perivascular infiltrates in the early lesion contained mostly lymphocytes and only a few undifferentiated monocytes,

while the perivascular cuffs in the late lesion consisted predominantly of lipid-laden macrophages, as described

previously (Figure 1i, l) [17,32]. We defined single cells by segmenting cell bodies outlined by the markers CD68

(macrophages/microglia), S100B (astrocytes), and CD3 (T cells) using CellProfiler (Figure 2) [11]. This allowed for

a clear delineation of cell types in the vast majority of cells. In a small fraction of cells, we found overlap between

cellular markers, especially between myelin fibers and juxtaposed microglial processes in normal-appearing white

matter (NAWM), and between astrocyte processes and closely neighboring macrophages (Figure 2c, d). We did not

exclude these cells in order to avoid bias in our spatial analysis. However, we excluded perivascular lymphocytes

from the early lesion, as they were too densely packed to be identifiable as individual cells.

Segmentation masks were overlaid with corresponding images from thirteen antibody channels (Figure 3—figure

supplement 1; Figure 5—figure supplement 1) in histoCAT [29], which clustered myeloid cells and astrocytes into

phenotypic subpopulations based on their marker expression profiles. The optimal number of phenotype clusters was

determined based on separation of the principal cell types (i.e. myeloid cells, astrocytes and T cells) and distinctive

expression profiles of phenotypes, using expression heatmaps and t-SNE plots. With twelve phenotypes per lesion,

we demonstrated that the main cell types and their subpopulations separated into distinct clusters (Figures 3, 5).

Phenotype heterogeneity and distribution in early demyelination

In the early lesion, we analyzed a total of 4,397 cells, of which 66.3% were myeloid cells and 32.5% were astrocytes

(Figure 3—figure supplement 2a). This ratio was higher at the rim than the lesion center. The cells clustered into six

myeloid and five astrocyte subtypes (Figure 3). The myeloid phenotypes were spatially sequestered in four lesional

regions, i.e. NAWM, outer lesion rim, inner lesion rim, and center. Within the outer lesion rim, we observed one

myeloid cell phenotype at the G/WM interface (m1) and two phenotypes located in the WM rim (m4, m5). The other

lesion areas were all occupied by one myeloid phenotype (Figure 4a). Most myeloid cell activation markers were

highly expressed in the outer lesion rim and decreased in intensity towards the lesion center (Figure 3a, b). The m1

phenotype at the G/WM interface showed the highest activation profile. m5 cells in the outer WM rim were the largest

in size, whereas m2 cells in the lesion center were the smallest (Figure 3—figure supplement 2b); this may reflect the

amount of myelin fragments phagocytosed by foamy macrophages at the advancing lesion edge, and the degradation

of myelin in phagocytes within the lesion center [25].

Astrocyte phenotypes were each defined by one distinct, highly expressed marker (Figure 3a, b). Analogous to the

myeloid phenotypes, the five astrocyte phenotypes were also spatially stratified into the NAWM, lesion rim, and lesion

center (Figure 4b). Furthermore, a5 astrocytes located in the outer rim within WM were larger than all other

phenotypes (Figure 3—figure supplement 2b). Unlike in myeloid cells, marker expression in astrocytes did not follow

a gradient from the rim to the lesion center, but was uniform throughout the lesion.

T cells made up the smallest population of all immune infiltrates, and were concentrated within the lesion center

(Figure 4c; Figure 3—figure supplement 2a). These cells uniformly expressed CD45 and HLA-DR (Figure 3a, b), and

did not separate into different clusters.

To determine possible transitions between phenotypes, we applied Potential of Heat-diffusion Affinity-based

Transition Embedding (PHATE) mapping and Pseudotime to the myeloid cell and astrocyte populations. PHATE

mapping improves on t-SNE by better visualizing phenotype transitions, where smooth continua from one phenotype

to another indicate a transition trajectory [19]. Rather than a rim-to-center continuum that aligned with the sequential

organization of phenotypes in the lesion, our analysis showed only a partial continuum between the G/WM outer rim

phenotype (m1) and the lesion center phenotype (m2) (Figure 4d), without including the inner rim phenotype (m3).

Similarly, analysis with Pseudotime, an alternative tool for visualizing phenotype transitions [33], showed that

myeloid phenotypes did not follow a linear transition trajectory but branched into several independent fates (Figure

4e). PHATE mapping and Pseudotime analysis of astrocyte phenotypes also suggested independent phenotypic fates

rather than linear phenotype trajectories (Figure 4—figure supplement 1, Pseudotime analysis not shown).

Low phenotype heterogeneity and random phenotype distributions in the late demyelinating lesion

In the late, post-demyelinating lesion we analyzed 6,698 cells, with myeloid cells far outnumbering astrocytes (91.1%

myeloid cells; Figure 5—figure supplement 2a), particularly at the lesion rim. The same clustering criteria used for

the early lesion resulted in nine myeloid phenotypes and two astrocyte phenotypes in this late lesion (Figure 5).

Myeloid phenotypes separated into lesion rim (m3), perivascular space (m4), and NAWM (m7) zones (Figure 6a). In

contrast to the early lesion, the other six myeloid phenotypes were intermixed throughout the lesion center. These

phenotypes showed a low degree of separation on the t-SNE plot, indicating similar marker expression profiles (Figure

5c). The phenotypes in the lesion rim and the perivascular space (m3, m4) were characterized by high expression of

the majority of markers, and shared a similar expression profile with the G/WM rim phenotype in the early lesion

(m1) (Figure 5a, b). Other myeloid subpopulations did not overlap between the two lesions. Phenotypes in the lesion

rim (m3) and perivascular space (m4) were significantly larger than the lesion center phenotypes (Figure 5—figure

supplement 2b). Overall, myeloid cells were significantly smaller in the late lesion than in the early lesion (data not

shown).

Astrocytes were clustered into two phenotypes, with one phenotype localizing primarily to the lesion rim, and the

other to the lesion center (Figure 6b). The rim phenotype (a2) exhibited a marker expression profile similar to the rim

phenotype in the early lesion (a5), while the lesion center phenotype did not overlap with any astrocyte phenotypes in

the early lesion (Figure 5a, b). Finally, T cells were few (Figure 6c; Figure 5—figure supplement 2a), and expressed

the activation markers CD45 and HLA-DR (Figure 5a, b), as observed in the early lesion.

PHATE mapping demonstrated a partial continuum between two lesion center myeloid phenotypes (m2, m6) and the

perivascular space phenotype (m4) (Figure 6d), which was confirmed by Pseudotime (Figure 6e). These analyses

support a lesion center-to-perivascular transition, but not a continuum where all phenotypes align along a rim-to-center

transition axis. The same analyses for astrocytes showed substantial overlap between both phenotypes but not a linear

transition (Figure 6—figure supplement 1, Pseudotime analysis not shown).

In the early lesion, we found no correlation between the expression intensities of different markers at a single-cell

level (Figure 7—figure supplement 1a, b). This was true for an analysis of all cells and cells of each phenotype (data

not shown). In the late lesion, we found strong correlations between the M2 markers MerTK and Mac2, and MerTK

and LAMP1 in both myeloid cells and astrocytes (Figure 7—figure supplement 1c, d). This resulted from high and

continuous dynamic ranges of marker expression (Figure 7). Correlations were weak between the other marker pairs.

Phenotypes in the early and late acute lesions engaged in specific cell-cell interactions

We next investigated the spatial relationships between different phenotypes with a computational tool integrated into

histoCAT that performs an unbiased, systematic analysis of pairwise phenotype interactions and avoidances [29].

After excluding interactions between cells of the same or spatially adjacent phenotypes, our analysis demonstrated

distinct interaction signatures for both lesions, (Figure 8a, b). At a significance cut-off of $p < 0.01$, these included

interactions between inner rim myeloid phenotype m3 (MerTK and CD45 high) and astrocyte phenotype a1 (Mac2

high), as well as interaction of highly activated myeloid rim phenotype m1 and the rim/center phenotype m4 (HLA-

DR and ferritin high) with astrocyte phenotype a2 (HLA-DR high) in the early lesion. In the late lesion, highly

activated perivascular macrophages (m4) interacted with most myeloid cell phenotypes and both astrocyte phenotypes.

There were also significant interactions between myeloid and astrocyte phenotypes m6 and a1, and among lesion

center myeloid phenotypes (m6 with m7 and m8). At a significance cut-off of $p < 0.05$, we found that T cells in the

late lesion interacted with myeloid phenotypes in the perivascular space (m4), which expressed HLA-DR, and in the

lesion center (m8).

Influence of lesion environment on marker expression

Finally, we used spatial variance component analysis (SVCA) to model the effects of extrinsic factors (neighboring

cells and non-cellular environmental stimuli) and cell-intrinsic factors on variations in cell marker expression [2]. In

the rim of both lesions, the expression of several markers was highly influenced by neighboring cells, including CD86,

PLP, and Mac2 in the early lesion, and CD86, PLP and LAMP1 in the late lesion. Other markers, such as ferritin,

vimentin, and CD3 (early lesion), and ferritin, vimentin, and TIM-3 (late lesion), were driven predominantly by non-

cellular environmental stimuli (Figure 8c, d). The relative influence of these factors changed toward the inner lesion

rim and center, leading to an overall increased impact of cell-intrinsic factors and a decrease in the influence of external

factors. In the lesion center, the primary agents influencing marker expression were cell-intrinsic factors and, to a

lower degree, environmental stimuli (Figure 8c, d).

Discussion

Our study examines the landscape of myeloid and astrocyte phenotypes in early and late acute MS brain lesions using

imaging mass cytometry. To our knowledge, this is the first application of highly multiplexed imaging to MS lesions.

We applied thirteen glia-related markers to MS lesions, which clustered myeloid cells and astrocytes into eleven

phenotypes. The phenotypes localized to different lesional zones, suggesting functional specificity. We further

characterized specific spatial interactions between phenotypes, examined phenotypic transitions within lesions and

determined the effect of cell-extrinsic and -intrinsic factors on marker expression in individual cells. Thus, our study

demonstrates that highly multiplexed imaging, combined with computational tools, provides a wealth of data on

cellular spatial organization that is not accessible with standard histology.

Our findings contrast with recent attempts to cluster myeloid cells in experimental autoimmune encephalomyelitis

(EAE), a mouse model of MS, with single-cell cytometry [20], and in white matter MS lesions with RNA-sequencing

data [10], which resulted in only two myeloid cell subpopulations, respectively. The higher number of clusters in our

study might be explained by the targeted set of markers, which are relevant to glial biology. In contrast, RNA-

sequencing emphasizes highly expressed genes, and might not detect low-expression genes or subtle differences in

gene expression that may be important for glial phenotype clustering.

We found that the myeloid cell phenotypes separated into different layers within the lesion, including the NAWM,

outer and inner lesion rims located in the WM or mixed G/WM matter, and the lesion center. This alignment of

phenotypes with lesional zones confirms the validity of our clustering approach. Consistent with the different stages

of myelin phagocytosis and degradation, the phenotypes in the rim and lesion center displayed different cellular sizes.

Only one myeloid phenotype was shared by both lesions, located in the lesion rim (m1 in the early lesion; m3 in the

late lesion), while all other phenotypes were unique to their respective lesions. Phenotypes were distinctly stratified

in the early demyelinating lesion, while their distribution in the post-demyelinated lesion appeared less ordered,

especially in the lesion center. Moreover, expression of all activation markers, with the exception of CD68, was high

in myeloid cells at the lesion rim and diminished substantially toward the lesion center. While the rim-to-center

gradient also existed in the late lesion, it was less pronounced. Thus, whereas the high activation profile of myeloid

cells at the lesion rim remained consistent between the two lesions, the center phenotypes lost their location-specific

polarization in the late lesion and became functionally less distinct. An additional feature of the late lesion was highly

activated macrophages that populated perivascular spaces in large numbers throughout the lesion. Since these

macrophages are believed to exit the CNS [17], this indicates that macrophages leave the CNS in a highly activated

state. Marker expression in astrocytes did not follow a rim-to-center gradient, but remained comparable throughout

the lesion area in both lesions.

Our findings argue that macrophages/microglia in MS lesions do not transition from a pro- to anti-inflammatory state,

as previously suggested [5], but convert from a highly activated to a less activated state as they move from the active

edge to the lesion center. This is consistent with a study by Vogel and colleagues, who demonstrated that pro- and

anti-inflammatory markers were simultaneously expressed by macrophages/microglia in MS lesions [34]. Our results

therefore provide further evidence that activated macrophages in inflamed tissue do not follow a M1/M2 polarization

dichotomy [28]. However, clarification of this question requires functional assays and/or a more detailed evaluation

of pro-inflammatory and immunosuppressive pathways in these cells. Two recent reports, including our own, suggest

a dichotomy in MS and EAE between myeloid cells expressing inducible nitric oxide synthase (iNOS) or arginase,

two enzymes that both use the substrate L-arginine to generate either reactive nitric oxide species or substrates such

as urea and L-ornithine that contribute to tissue repair [7,16].

Using PHATE mapping and Pseudotime to determine transitions between phenotypes, we did not find that myeloid

cell or astrocyte phenotypes developed along a linear transition continuum, i.e. from the outer to the inner lesion rim

and lesion center. Instead, the different phenotypes branched into multiple independent trajectories, suggesting that

they emerge within the lesion environment independently from each other. We cannot exclude that the parametric

depth of our study is too low to detect transition states, especially since Pseudotime has been developed to delineate

phenotypic trajectories from whole transcriptome sequencing data [33]. Nevertheless, since clustering with our set of

markers resulted in lesion zone-specific phenotypes, it is conceivable that the same markers can be used to trace

transitions between the lesion phenotypes.

Neighborhood analysis demonstrated distinct cellular interaction signatures for both lesions, e.g. between phagocytic

macrophages in the inner rim and center astrocytes in the early lesion, and between T cells and myeloid phenotypes

in the late lesion. Thus, interactions in the hypercellular lesion environment are not random, but occur between specific

subpopulations and cell types such as lymphocytes. The low parametric depth of our study does not allow us to identify

the functional implications of these interactions; however, they may represent nodal points of cellular communication

that are key for lesion formation and maintenance of low-grade lesional inflammation.

Finally, spatial variance component analysis (SVCA) suggests that cell-extrinsic factors drive marker expression to a

higher degree at the rim than in the lesion center. Conversely, cell-intrinsic factors have a more prominent influence

on marker expression in the lesion center than at the rim. This indicates that glia cells in the lesion rim are more

responsive to cues from the microenvironment, such as cytokines or receptor-ligand interactions, while glia responses

in the lesion center are more affected by cell-intrinsic programs such as cellular changes that are set in motion by

myelin phagocytosis.

In summary, we found that phenotypic clustering, determined through differential expression of thirteen markers,

produced multiple myeloid cell and astrocyte phenotypes that occupied specific lesion zones in the early acute lesion,

and were more randomly distributed in the late lesion. Myeloid cells were activated along a rim-to-center axis, and

specific myeloid cell-astrocyte-lymphocyte interactions were present in both lesions. Thus, our study highlights the

potential of imaging mass cytometry and other highly multiplexed imaging techniques [4,8] to provide insight into

lesion-forming phenotypes and their spatial formation. In addition, our study provides novel insight into the dynamics

of phenotypes and their spatial organization in MS lesions over time.

The study is limited by the small sample size and the small number of markers, which does not allow for deep

functional characterization of phenotypes. In ongoing studies, we are combining clustering based on single nuclear

RNA-sequencing data of glial cells, derived from MS lesion tissue, with highly multiplexed imaging of the same

lesion, to simultaneously obtain maximal parametric depth and high spatial resolution at single-cell level. This will

eventually define the phenotypes and key interaction networks that drive active demyelination and chronic, low-grade

inflammation in established lesions and may provide targets for therapeutic intervention in relapsing and progressive

MS.

Materials and methods

MS lesions.

Human CNS tissue was obtained at autopsy from two patients with relapsing-remitting MS according to Institutional

Review Board-approved protocols. After autopsy, brain tissue was fixed in 10% formalin, and lesions were cut based

on MRI. Lesion tissue was subsequently embedded in paraffin and sectioned at 5 μm thickness.

A highly inflamed, active lesion was chosen for analysis from each patient: the actively demyelinating lesion was

selected from a 42-year-old male with 5 years disease duration (5.5 hour *post mortem* interval), while the post-

demyelinating lesion was chosen from a 32-year-old female with 6 years disease duration (8 hour *post mortem*

interval). Lesions from both patients have been characterized in previous studies [7,22].

Brightfield histology.

For basic characterization, lesions were stained against CD68, myelin basic protein (MBP) and neuronal marker

MAP2, and examined with brightfield microscopy. De-paraffinized and rehydrated sections were subjected to antigen

retrieval in pH 6, 10 mM citrate buffer at 96°C for 20 minutes, cooled, quenched in 0.3% peroxide, and blocked with

FC receptor binding inhibitor and normal serum before incubation with primary antibody (Table 1) overnight at 4°C.

Sections were subsequently incubated with appropriate biotinylated secondary antibodies, processed with an

avidin/biotin staining kit with 3,3-diaminobenzidene (DAB) as the chromogen (Vector ABC Elite Kit and DAB Kit,

Vector Laboratories, Burlingame, CA, USA), then counterstained with hematoxylin [22]. Adequate controls using

isotype control antibodies were performed for each primary antibody. Sections were rinsed with distilled water,

dehydrated, and cover-slipped with Permount (Vector Laboratories). Images were acquired using a Leica DM5000 B

microscope with a Leica color camera DFC310 Fx and Leica Application Suite (version 4.9.0) imaging software

(Leica Microsystems, Buffalo Grove, IL, USA). Images were processed with Panoramic Viewer (3DHISTECH,

Budapest, Hungary) and Photoshop (Adobe, San Jose, CA, USA) software.

Antibody validation and conjugation to metal isotopes for IMC.

Several lanthanide-conjugated antibodies were purchased from Fluidigm (Markham, ON, Canada). Antibodies not

available in metal-conjugated form were purchased in carrier-free solution and validated by brightfield

immunohistochemistry using the appropriate isotype control antibodies. Subsequently, antibodies were conjugated to

lanthanide metal isotopes following the Maxpar® Antibody Labeling Kit protocol (Fluidigm). Metal-conjugated

antibodies were stored at 0.5 mg/mL in PBS-based Antibody Stabilizer (Candor Bioscience, Wangen, Germany) with

0.05% sodium azide at 4°C. Working concentrations for all metal-conjugated antibodies were optimized by IMC

(Table 2) on MS lesion tissue.

Imaging mass cytometry.

For IMC histology, tissue sections were de-paraffinized and rehydrated, and antigen retrieval was performed in pH 8,

1 mM EDTA buffer at 96°C for 20 minutes. Sections were cooled at room temperature and rinsed in tap water and

TBS (20 mM Tris with 500 mM NaCl, pH 7.5). Tissue was blocked for 3 hours at room temperature with 0.3% BSA,

25% FBS and 0.1 mg/mL FC receptor binding inhibitor in TBS-T (TBS + 0.05% Tween-20). All antibodies (Table 2)

were diluted in 0.3% BSA in TBS-T and applied to the tissue for overnight incubation at 4°C. Sections were then

rinsed in TBS-T and TBS, and counterstained with 125 nM Maxpar® Intercalator-Ir (Fluidigm) in PBS for 1 hour at

room temperature. Sections were rinsed in TBS-T, TBS and two washes of distilled water before air-drying at room

temperature. Antibody-labeled tissue areas (1000x1000 µm) were raster-ablated using a Hyperion™ Laser Scanning

Module (Fluidigm) with a 1 µm diameter spot size at 200 Hz. This process was coupled to a Helios™ Mass Cytometer

(Fluidigm) for lanthanide metal detection [35]. Images for each antibody channel were acquired on CyTOF Software

(Fluidigm, version 6.7). MCD Viewer (Fluidigm, version 1.0) was used to export raw 16-bit tiff images for

computational analyses on histoCAT (version 1.75) [29]. For visualization purposes, images were processed in MCD

Viewer and ImageJ [31].

Computational analyses.

Single-cell segmentation. Merged CD68 (macrophages/microglia), S100B (astrocytes), and CD3 (T cells) antibody

channel images were used to segment single cell objects on CellProfiler (version 3.0.0) [11]. The resulting

segmentation mask images, which outline cell borders, were loaded into histoCAT with corresponding antibody

channel images.

Identification of cellular phenotypes. Mean single-cell marker intensity values were extracted from segmented

antibody channel images on histoCAT and Z-score normalized. Based on the expression intensities of thirteen markers

(Table 2), cell clusters were defined using the PhenoGraph algorithm [15] integrated into histoCAT. Default

parameters with 75 nearest neighbors for the early lesion and 50 nearest neighbors for the late lesion were used. These

nearest neighbor values were chosen such that over- and under-clustering of phenotypes were avoided. Additional

normalization steps were performed internally, as previously described [29].

Analysis of cellular phenotypes. To visualize clusters, the Barnes-Hut t-SNE algorithm implemented in histoCAT was

executed with the same image and marker inputs used in PhenoGraph, as well as default parameters (initial

dimensions, 110; perplexity, 30; theta, 0.5) and internal normalization [1,29]. t-SNE plots were colored to highlight

clusters or show relative marker expression intensity. Images of cell phenotypes visualized in the tissue, as well as

segmentation masks overlaid with histology images, were generated in histoCAT. For the remaining analyses, ".csv"

files containing single-cell parameters were exported from histoCAT and appropriately processed for their application.

To produce an expression heatmap for clusters, normalized marker intensity values were processed using the R

ComplexHeatmap package, which hierarchically clusters single cells within clusters using Ward's method [30]. Violin

plots showing single-cell marker expression variability for each cluster were generated using the R *ggplot2* package

[9]. To study phenotype transitions, Potential of Heat-diffusion Affinity-based Transition Embedding (PHATE)

mapping and Monocle Pseudotime analyses were performed [19,26,27,33]. PHATE mapping was performed using

normalized marker intensity values, with an adjusted number of nearest neighbors (k = 100) and other parameters left

as their default specifications. Pseudotime analysis was performed with negative binomial normalization of marker

intensity values and q < 1 to incorporate all markers, based on a test for differential expression among phenotypes.

Figures showing phenotype cell size and relative quantity of cells per cluster, as well as correlation matrices and

corresponding marker scatter plots with linear regression lines were produced in Prism (version 7). FlowJo software

(version 10.5.3) was used to visualize single-cell marker data on flow cytometry plots. Images and figures were

recolored when necessary in Photoshop.

Analysis of cell spatial relationships. To study the spatial relationships of cell clusters, neighborhood analysis was

performed on histoCAT using the PhenoGraph-generated clusters pertaining to each lesion. Significant pairwise

phenotype interactions and avoidances were determined by an unbiased permutation test which compared the

frequency of one cell type neighboring another to the frequency in 999 random permutations of cell cluster labels.

Neighbors were identified within 4 μm from each cell, and neighboring frequency was normalized to the number of

interacting cells [29]. Analyses were performed for different degrees of significance ($p < 0.05$ and $p < 0.01$), and the

results were reported on a heatmap. To identify sources of single-cell marker variation, spatial variance component

analysis (SVCA) [2] was performed on Python for different lesion zones, using standardized marker intensity values.

SVCA plots were generated in R and Prism.

Statistical analyses. In plots showing phenotype cell size, data represent mean cell areas + standard deviation.

Comparisons of phenotype cell sizes were analyzed by one-way ANOVA followed by the Tukey-Kramer multiple

comparison test. Comparisons of two samples were carried out by unpaired Student's t-tests. For correlation analyses,

Pearson correlation coefficients were computed. *$p < 0.0001$.

Figure supplement list

Figure 3—figure supplement 1: Imaging mass cytometry histology images of the early lesion
Figure 3—figure supplement 2: Early lesion cell phenotype frequency and size
Figure 4—figure supplement 1: PHATE mapping of astrocyte phenotypes in the early lesion
Figure 5—figure supplement 1: Imaging mass cytometry histology images of the late lesion
Figure 5—figure supplement 2: Late lesion cell phenotype frequency and size
Figure 6—figure supplement 1: PHATE mapping of astrocyte phenotypes in the late lesion
Figure 7—figure supplement 1: Pearson correlation matrices for myeloid cells and astrocytes

Data availability

Source data generated and/or analyzed during the current study as well as R source code are available in a GitHub

repository: https://github.com/PittLab/IMC_Park_et_al_2019.

Acknowledgments

The authors thank Fluidigm Corporation (Markham, ON, Canada) and the Yale University CyTOF core facility for

providing IMC tissue imaging service and metal-conjugated antibodies, as well as Denis Schapiro, Damien Arnol and

Kevin Moon for expert assistance with computational analyses.

References

1. Amir E-aD, Davis KL, Tadmor MD, Simonds EF, Levine JH, Bendall SC, Shenfeld DK, Krishnaswamy S, Nolan
GP, Pe'er D (2013) viSNE enables visualization of high dimensional single-cell data and reveals phenotypic
heterogeneity of leukemia. Nature Biotechnology 31:545. doi:10.1038/nbt.2594
2. Arnol D, Schapiro D, Bodenmiller B, Saez-Rodriguez J, Stegle O (2018) Modelling cell-cell interactions from
spatial molecular data with spatial variance component analysis. bioRxiv:265256. doi:10.1101/265256
3. Bartnik BL, Juurlink BH, Devon RM (2000) Macrophages: their myelinotrophic or neurotoxic actions depend upon
tissue oxidative stress. Multiple sclerosis (Houndmills, Basingstoke, England) 6:37-42.
doi:10.1177/135245850000600108
4. Bolognesi MM, Manzoni M, Scalia CR, Zannella S, Bosisio FM, Faretta M, Cattoretti G (2017) Multiplex Staining
by Sequential Immunostaining and Antibody Removal on Routine Tissue Sections. The journal of
histochemistry and cytochemistry : official journal of the Histochemistry Society 65:431-444.
doi:10.1369/0022155417719419
5. Boven LA, Van Meurs M, Van Zwam M, Wierenga-Wolf A, Hintzen RQ, Boot RG, Aerts JM, Amor S,
Nieuwenhuis EE, Laman JD (2006) Myelin-laden macrophages are anti-inflammatory, consistent with foam
cells in multiple sclerosis. Brain : a journal of neurology 129:517-526. doi:10.1093/brain/awh707
6. Farina C, Aloisi F, Meinl E (2007) Astrocytes are active players in cerebral innate immunity. Trends in immunology
28:138-145. doi:10.1016/j.it.2007.01.005
7. Giles DA, Washnock-Schmid JM, Duncker PC, Dahlawi S, Ponath G, Pitt D, Segal BM (2018) Myeloid cell
plasticity in the evolution of central nervous system autoimmunity. Annals of neurology 83:131-141.
doi:10.1002/ana.25128
8. Gut G, Herrmann MD, Pelkmans L (2018) Multiplexed protein maps link subcellular organization to cellular states.
Science 361:eaar7042. doi:10.1126/science.aar7042
9. Hadley W (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York,
10. Jakel S, Agirre E, Mendanha Falcao A, van Bruggen D, Lee KW, Knuesel I, Malhotra D, Ffrench-Constant C,
Williams A, Castelo-Branco G (2019) Altered human oligodendrocyte heterogeneity in multiple sclerosis.
Nature 566:543-547. doi:10.1038/s41586-019-0903-2
11. Kamentsky L, Jones TR, Fraser A, Bray MA, Logan DJ, Madden KL, Ljosa V, Rueden C, Eliceiri KW, Carpenter
AE (2011) Improved structure, function and compatibility for CellProfiler: modular high-throughput image
analysis software. Bioinformatics (Oxford, England) 27:1179-1180. doi:10.1093/bioinformatics/btr095
12. Kim CC, Nakamura MC, Hsieh CL (2016) Brain trauma elicits non-canonical macrophage activation states. J
Neuroinflammation 13:117. doi:10.1186/s12974-016-0581-z
13. Kuhlmann T, Ludwin S, Prat A, Antel J, Brück W, Lassmann H (2017) An updated histological classification
system for multiple sclerosis lesions. Acta Neuropathologica 133:13-24. doi:10.1007/s00401-016-1653-y
14. Lassmann H (2018) Multiple Sclerosis Pathology. Cold Spring Harbor perspectives in medicine 8.
doi:10.1101/cshperspect.a028936
15. Levine JH, Simonds EF, Bendall SC, Davis KL, Amir el AD, Tadmor MD, Litvin O, Fienberg HG, Jager A,
Zunder ER, Finck R, Gedman AL, Radtke I, Downing JR, Pe'er D, Nolan GP (2015) Data-Driven Phenotypic
Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis. Cell 162:184-197.
doi:10.1016/j.cell.2015.05.047
16. Locatelli G, Theodorou D, Kendirli A, Jordao MJC, Staszewski O, Phulphagar K, Cantuti-Castelvetri L, Dagkalis
A, Bessis A, Simons M, Meissner F, Prinz M, Kerschensteiner M (2018) Mononuclear phagocytes locally
specify and adapt their phenotype in a multiple sclerosis model. Nature neuroscience 21:1196-1208.
doi:10.1038/s41593-018-0212-3
17. Love S, Louis D, Ellison DW, Sobel RA, Moore GRW (2008) Chapter 20: Demyelinating diseases. In: Greenfield's
Neuropathology. Boca Raton, FL: CRC Press; p. 1526.
18. Mayo L, Trauger SA, Blain M, Nadeau M, Patel B, Alvarez JI, Mascanfroni ID, Yeste A, Kivisakk P, Kallas K,
Ellezam B, Bakshi R, Prat A, Antel JP, Weiner HL, Quintana FJ (2014) Regulation of astrocyte activation
by glycolipids drives chronic CNS inflammation. Nature medicine 20:1147-1156. doi:10.1038/nm.3681
19. Moon KR, van Dijk D, Wang Z, Gigante S, Burkhardt D, Chen W, van den Elzen A, Hirn MJ, Coifman RR,
Ivanova NB, Wolf G, Krishnaswamy S (2018) Visualizing Transitions and Structure for Biological Data
Exploration. bioRxiv:120378. doi:10.1101/120378
20. Mrdjen D, Pavlovic A, Hartmann FJ, Schreiner B, Utz SG, Leung BP, Lelios I, Heppner FL, Kipnis J, Merkler D,
Greter M, Becher B (2018) High-Dimensional Single-Cell Mapping of Central Nervous System Immune

Cells Reveals Distinct Myeloid Subsets in Health, Aging, and Disease. Immunity 48:380-395.e386.
doi:10.1016/j.immuni.2018.01.011
21. Nylander A, Hafler DA (2012) Multiple sclerosis. The Journal of clinical investigation 122:1180-1188.
doi:10.1172/jci58649
22. Pitt D, Nagelmeier IE, Wilson HC, Raine CS (2003) Glutamate uptake by oligodendrocytes: Implications for
excitotoxicity in multiple sclerosis. Neurology 61:1113-1120
23. Ponath G, Lincoln MR, Levine-Ritterman M, Park C, Dahlawi S, Mubarak M, Sumida T, Airas L, Zhang S, Isitan
C, Nguyen TD, Raine CS, Hafler DA, Pitt D (2018) Enhanced astrocyte responses are driven by a genetic
risk allele associated with multiple sclerosis. Nature Communications 9:5337. doi:10.1038/s41467-018-
07785-8
24. Ponath G, Park C, Pitt D (2018) The Role of Astrocytes in Multiple Sclerosis. Frontiers in immunology 9:217-
217. doi:10.3389/fimmu.2018.00217
25. Prineas J (1975) Pathology of the early lesion in multiple sclerosis. Human pathology 6:531-554
26. Qiu X, Hill A, Packer J, Lin D, Ma YA, Trapnell C (2017) Single-cell mRNA quantification and differential
analysis with Census. Nature methods 14:309-315. doi:10.1038/nmeth.4150
27. Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, Trapnell C (2017) Reversed graph embedding resolves
complex single-cell trajectories. Nature methods 14:979-982. doi:10.1038/nmeth.4402
28. Ransohoff RM (2016) A polarizing question: do M1 and M2 microglia exist? Nature neuroscience 19:987.
doi:10.1038/nn.4338
29. Schapiro D, Jackson HW, Raghuraman S, Fischer JR, Zanotelli VRT, Schulz D, Giesen C, Catena R, Varga Z,
Bodenmiller B (2017) histoCAT: analysis of cell phenotypes and interactions in multiplex image cytometry
data. Nature methods 14:873-876. doi:10.1038/nmeth.4391
30. Schlesner M, Gu Z, Eils R (2016) Complex heatmaps reveal patterns and correlations in multidimensional genomic
data. Bioinformatics (Oxford, England) 32:2847-2849. doi:10.1093/bioinformatics/btw313
31. Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nature
methods 9:671-675
32. Tanaka R, Iwasaki Y, Koprowski H (1975) Ultrastructural studies of perivascular cuffing cells in multiple sclerosis
brain. The American journal of pathology 81:467-478
33. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ, Mikkelsen TS, Rinn JL
(2014) The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single
cells. Nat Biotechnol 32:381-386. doi:10.1038/nbt.2859
34. Vogel DYS, Vereyken EJF, Glim JE, Heijnen PDAM, Moeton M, van der Valk P, Amor S, Teunissen CE, van
Horssen J, Dijkstra CD (2013) Macrophages in inflammatory multiple sclerosis lesions have an intermediate
activation status. Journal of neuroinflammation 10:35-35. doi:10.1186/1742-2094-10-35
35. Wang HA, Grolimund D, Giesen C, Borca CN, Shaw-Stewart JR, Bodenmiller B, Gunther D (2013) Fast chemical
imaging at high spatial resolution by laser ablation inductively coupled plasma mass spectrometry. Analytical
chemistry 85:10107-10116. doi:10.1021/ac400996x
445

446
Figure 1 Brightfield images of the MS lesions examined with IMC. **(a-f, i)** Staining of the early, actively demyelinating lesion for **(a)** myelin basic protein (MBP), **(b)** CD68 (macrophages/microglia) and **(c)** the neuronal marker MAP2. The black rectangle in **(a)** shows the region that was laser-ablated for IMC, capturing the NAWM, hypercellular rim on both sides and lesion center. Red boxes in **a-c** correspond to magnified rim areas in **d-f,** demonstrating in **(d)** condensed myelin fragments, indicative of ongoing demyelination, **(e)** dense infiltration with foamy macrophages and **(f)** dispersed neuronal cell bodies in the right lesion rim.
(g, h, j-l) Staining against MBP and CD68 in the late, post-demyelinating lesion. The black rectangle in **(g)** shows the region laser-ablated for IMC. **(j, k)** Details from **(g)** and **(h)** showing absence of myelin fragments and reduced myeloid cell density at the lesion rim compared to **(e)**. CD68$^+$ cells in the perivascular spaces of **(i)** the early lesion and **(l)** the late lesion. Images show hematoxylin counterstaining. Scale bars **a-c, g, h** = 500 μm. Scale bars **d-f, i, j-l** = 25 μm.

Figure 2 Single-cell segmentation. **(a-c)** Segmentation of myeloid cells (CD68, magenta), astrocytes (S100B, green)
and T cells (CD3, cyan) performed on CellProfiler. **(b)** shows cells in **(a)** at higher magnification. **(c)** shows
examples of S100B[+] astrocyte processes (white arrows) that closely neighbor CD68[+] cells, which resulted in the co-
segmentation of these markers in a small fraction of cells. **(d)** Gating of CD68[+] and S100B[+] populations on a flow
cytometry plot as a quality control for segmentation. The overlap population consists of cells with co-segmented
CD68 and S100B like those in **(c)**. Scale bar in **(a)** = 30 μm; **(b)** = 15 μm; **(c)** = 30 μm, inset = 10 μm. Z-score
normalized expression intensities are shown **(d)**. CellProfiler segmentation files and corresponding IMC histology
images are available in a GitHub repository (see data availability statement).

Figure 3 Early lesion cell phenotype profiles. **(a)** Marker expression heatmap for the myeloid, astrocyte and T cell
phenotypes, identified by PhenoGraph clustering on histoCAT using segmented cells (n = 4397). The heatmap
displays relative expression levels based on Z-score normalized marker intensity values, and single cells are
hierarchically clustered within each phenotype group. Labels at the bottom of the heatmap indicate the area of the
lesion to which each phenotype localizes. **(b)** Violin plot representation of the data in **(a)**. **(c)** t-SNE plot showing the
distinct phenotype clusters. **(d)** t-SNE plot colored by marker intensity, confirming the separation of CD68[+], S100B[+]
and CD3[+] cell types. G/WMoR = gray and white matter outer rim; WMoR = white matter outer rim; iR = inner rim;
WM R/C = white matter rim/center; C = center; NAWM = normal-appearing white matter. Source single-cell and
histoCAT session data are available in a GitHub repository (see data availability statement).

476
Figure 3—figure supplement 1 Imaging mass cytometry histology images of the early lesion. The thirteen markers
used for cell clustering as well as DNA counterstaining are shown. Scale bar = 200 μm.

479
480 **Figure 3—figure supplement 2** Early lesion cell phenotype frequency and size. **(a)** Relative quantities of each cell
481 phenotype. **(b)** Cell phenotype size. Data represent means + standard deviation. *p < 0.0001 by one-way ANOVA
482 followed by the Tukey-Kramer multiple comparison test. Source data is available in a GitHub repository (see data
483 availability statement).

Figure 4 Early lesion phenotype spatial distributions and transition analyses. **(a, b)** Spatial separation of **(a)** myeloid cell and **(b)** astrocyte phenotypes into NAWM, rim and center lesion zones. **(c)** T cells are primarily located in the lesion center. **(d)** PHATE mapping of myeloid cells, indicating that the G/WM outer rim (m1) and lesion center phenotypes (m2) are on a transition continuum (white arrow). **(e)** Pseudotime analysis of myeloid cells shows that phenotypes transition along independent trajectories. Phenotype color schemes on the PHATE and Pseudotime plots reflect the color palettes specific to each analysis. Scale bar for **a-c** = 200 μm. G/WMoR = gray and white matter outer rim; WMoR = white matter outer rim; iR = inner rim; WM R/C = white matter rim/center; C = center; NAWM = normal-appearing white matter. Source data is available in a GitHub repository (see data availability statement).

a1: Center 1, a2: Inner Rim, a3: Center 2, a4: NAWM, a5: WM Outer Rim

493
Figure 4—figure supplement 1 PHATE mapping of astrocyte phenotypes in the early lesion. WMoR = white
matter outer rim; iR = inner rim; C = center; NAWM = normal-appearing white matter. Source data is available in a
GitHub repository (see data availability statement).

Figure 5 Late lesion cell phenotype profiles. **(a)** Marker expression heatmap for the myeloid, astrocyte and T cell
phenotypes, identified by PhenoGraph clustering on histoCAT using segmented cells (n = 6698). The heatmap
displays relative expression levels based on Z-score normalized marker intensity values, and single cells are
hierarchically clustered within each phenotype group. Labels at the bottom of the heatmap indicate the area of the
lesion to which each phenotype localizes. **(b)** Violin plot representation of the data in **(a)**. **(c)** t-SNE plot showing the
phenotype clusters. Compared to the early lesion, myeloid cell phenotypes show a low degree of separation. **(d)** t-
SNE plot colored by marker intensity, confirming the separation of CD68[+], S100B[+] and CD3[+] cell types. R = rim; C
= center; PVS = perivascular space; NAWM = normal-appearing white matter. Source single-cell and histoCAT
session data are available in a GitHub repository (see data availability statement).

507
Figure 5—figure supplement 1 Imaging mass cytometry histology images of the late lesion. The thirteen markers
used for cell clustering as well as DNA counterstaining are shown. Scale bar = 200 μm.

510
511 **Figure 5—figure supplement 2** Late lesion cell phenotype frequency and size. **(a)** Relative quantities of each cell
512 phenotype. **(b)** Cell phenotype size. Data represent means + standard deviation. *$p < 0.0001$ by one-way ANOVA
513 followed by the Tukey-Kramer multiple comparison test. Source data is available in a GitHub repository (see data
514 availability statement).

515
Figure 6 Late lesion phenotype spatial distributions and transition analyses. **(a)** Spatial organization of myeloid cell
phenotypes in the lesion. Phenotypes in the rim (m3), perivascular space (m4) and NAWM (m7) separate into distinct
zones, while lesion center phenotypes (m1, m2, m5, m6, m8, m9) are uniformly distributed. **(b)** Spatial distribution of
astrocyte phenotypes. One phenotype (a1) predominantly occupies the lesion center and the other (a2) occupies the
rim. **(c)** T cells are primarily distributed in the lesion center. **(d)** PHATE mapping of myeloid cells, showing that two
lesion center phenotypes (m2, m6) are on a continuum with perivascular space cells (m4, white arrows). **(e)**
Pseudotime analysis of myeloid cells shows a similar trajectory to PHATE mapping. Phenotype color schemes on the
PHATE and Pseudotime plots reflect the color palettes specific to each analysis. Scale bar for **a-c** = 200 μm. R = rim;
C = center; PVS = perivascular space; NAWM = normal-appearing white matter. Source data is available in a GitHub
repository (see data availability statement).

a1: Center, a2: Rim

526
527 **Figure 6—figure supplement 1** PHATE mapping of astrocyte phenotypes in the late lesion. R = rim; C = center.
528 Source data is available in a GitHub repository (see data availability statement).

529
Figure 7 Single-cell marker correlations in the late lesion. **(a, b)** Co-expression of Mac2 and MerTK, and LAMP1
and MerTK in **(a)** myeloid cells (n = 6100), and **(b)** astrocytes (n = 528). Co-expression plots are shown with linear
expression values and regression lines with Pearson correlation coefficients, and below in flow cytometry contour plot
form with log10-transformed expression values. Source data is available in a GitHub repository (see data availability
statement).

Figure 7—figure supplement 1 Pearson correlation matrices for myeloid cells and astrocytes. Matrices for early lesion **(a)** myeloid cells (n = 2913) and **(b)** astrocytes (n = 1431), and late lesion **(c)** myeloid cells (n = 6100) and **(d)** astrocytes (n = 528) are shown. Source data is available in a GitHub repository (see data availability statement).

539
Figure 8 Neighborhood and spatial variance component analyses for the early and late lesions. **(a, b)** Neighborhood
analysis heatmaps of all significant pairwise phenotype interactions (red) and avoidances (blue) in the **(a)** early and
(b) late lesions. White represents no significant spatial relationship. Dark boxes are highly significant spatial
relationships ($p < 0.01$). Lightly shaded boxes are less significant relationships ($p < 0.05$) and interactions between
cells of the same or spatially adjacent phenotypes. Rows visualize the significance of a phenotype surrounded by other
phenotypes, and columns visualize the significance of a phenotype surrounding other phenotypes. **(c, d)** Spatial
variance component analysis (SVCA) for the **(c)** early and **(d)** late lesions, showing the proportion of marker
expression variance attributable to neighboring cell influences, environmental stimuli, cell-intrinsic factors and
residual noise in different lesion zones. Additional plots show the average proportion of marker variance attributable
to each factor in different lesion zones. G/WM = gray and white matter; WM = white matter; PVS = perivascular
space; NAWM = normal-appearing white matter. Source histoCAT session and SVCA data are available in a GitHub
repository (see data availability statement).

Table 1 Antibodies used for brightfield histology

Primary Antibody	Host	Clone	Company	Target/Significance	Primary Antibody Dilution	Secondary Antibody	Secondary Antibody Dilution
MBP	Rat	12	Millipore-Sigma	Myelin Basic Protein	1:500	Invitrogen, 629540	1:500
CD68	Rabbit	D4B9C	Cell Signaling	Activated macrophage/microglia transmembrane glycoprotein	1:500	Vector Laboratories, BA-1000	1:500
MAP2	Mouse	5H11	Novus	Neuron structural protein	1:200	Vector Laboratories, BA-9200	1:500

553 **Table 2** Metal-conjugated antibodies used for IMC

Antibody	Clone	Company	Target/Significance	Conjugated Metal Isotope	Conjugation Source	Dilution
LAMP1/CD107a	H4A3	Fluidigm	Lysosomes	151Eu	Fluidigm	1:100
CD3	Polyclonal	Fluidigm	T cells	170Er	Fluidigm	1:75
CD45	2B11	Fluidigm	Leukocyte activation	152Sm	Fluidigm	1:100
CD68	KP1	Fluidigm	Activated macrophages/microglia	159Tb	Fluidigm	1:300
CD86	Polyclonal	R&D	"M1" co-stimulatory molecule	171Yb	Our Lab	1:100
Ferritin heavy chain	1-2.3.1.2	Millipore-Sigma	Iron storage	158Gd	Our Lab	1:400
HLA-DR	YE2/36 HLK	Fluidigm	MHC Class II	174Yb	Fluidigm	1:200
Mac2	M3/38	Fluidigm	"M2" activation	153Eu	Fluidigm	1:600
MerTK	y323	abcam	"M2" phagocytosis	148Nd	Our Lab	1:50
PLP	plpc1	Bio-Rad	Myelin	142Nd	Our Lab	1:100
S100B	S100B/1706R	Novus	Astrocytes	167Er	Our Lab	1:400
TIM-3	D5D5R	Fluidigm	Co-inhibitory molecule	154Sm	Fluidigm	1:100
Vimentin	RV202	Fluidigm	Hypertrophic astrocytes	143Nd	Fluidigm	1:100