A comparative assessment of adult mosquito trapping methods to

estimate spatial patterns of abundance and community composition in

southern Africa

Gorsich, E.E*[1,2,3]., Beechler, B[4]., Bodegom, P.M. van[1], Govender, D.[5], Guarido, M.M.[6],

Venter, M.[6], Schrama, M.[1]

1. Institute of Environmental Sciences, Leiden University, Leiden, NL.

2. School of Life Sciences, University of Warwick, Coventry, UK.

3. The Zeeman Institute for Systems Biology & Infectious Disease Epidemiology Research,

University of Warwick, Coventry, UK.

4. College of Veterinary Medicine, Oregon State University, Corvallis, OR, United States

5. SANPARKS, Scientific Services, Skukuza, South Africa.

6. Zoonotic Arbo- and Respiratory Virus Program, Centre for Viral Zoonoses, Faculty of

Health Sciences, University of Pretoria, Pretoria, South Africa

*Corresponding author address: erin.gorsich@warwick.ac.uk

Abstract

Background. Assessing adult mosquito populations is an important component of disease

surveillance programs and ecosystem health assessments. Inference from adult trapping

datasets involves comparing populations across space and time, but comparisons based on

different trapping methods may be biased if traps have different efficiencies or sample

different subsets of the mosquito community.

Methods. We compared four widely-used trapping methods for adult mosquito data

collection in Kruger National Park (KNP), South Africa: Centers for Disease Control

miniature light trap (CDC), Biogents Sentinel trap (BG), Biogents gravid *Aedes* trap (GAT),

and a CO_2-baited net trap. We quantified how trap choice and sampling effort influence

inferences on the regional distribution of mosquito abundance, richness, and community

composition.

Results. The CDC and net traps together collected 96% (47% and 49% individually) of the

955 female mosquitoes sampled and 100% (85% and 78% individually) of the 40 species

or species complexes identified. The CDC and net trap also identified similar regional

patterns of community composition. However, inference on the regional patterns of

abundance differed between these traps because mosquito abundance in the net trap was

influenced by variation in weather conditions. The BG and GAT traps collected significantly

fewer mosquitoes, limiting regional comparisons of abundance and community

composition.

Conclusion: This study represents the first systematic assessment of trapping methods in

in natural savanna ecosystems in southern Africa. We recommend the CDC trap or the

combined use of the net and CDC trap for future monitoring and surveillance programs.

Keywords

Arboviruses, community composition; Kruger National Park; Mosquito; South Africa; Trap

Bias; Vector

Background

Adult mosquito sampling is a key component of mosquito surveillance [1–4], but

trapping success may vary across studies due to differences in trapping methods. Different

traps vary in their ability to catch certain species [5–8]. For example, the dominant species

attracted with light-baited traps may differ from those attracted to traps baited with

carbon dioxide or live hosts [9]. Additionally, sampling conditions such as the number of

nights over which sampling occurs and weather conditions may also influence trapping

success [10], with some species and traps potentially more affected than others. This

variation in trapping outcome may not limit inference on the presence or absence of

common species (e.g. the information used in global risk maps [11,12]). However, it does

limit inference based on comparing patterns of diversity or abundance across space or time

[7]. Given that these comparisons are required to evaluate the ecological or anthropogenic

drivers of mosquito populations and disease risk, choice of trapping methods presents

challenges and opportunities for optimizing sampling efforts.

Previous studies have evaluated trapping methods in Europe, North America, and

South America [5–8,13] while studies in southern Africa remain relatively limited. The

mosquito fauna (Diptera: Cluicidae) of southern Africa consist of over 216 species, many of

which are endemic to the region [14]. Additional studies evaluating trapping methods are

needed to evaluate if traps designed for other locations and species perform equally well in

the rich communities in southern Africa. For example, a recent comparison of four traps in

Germany found that the Biogents Sentinel trap (BG trap) collected the highest number of

individuals in urban and snowmelt forest environments where *Culex* species predominate.

In contrast, the Centers for Disease Control miniature light trap (CDC trap) collected the

most mosquitoes in floodplain environments where *Aedes vexans* predominate, suggesting

that the preferred trapping device may vary by habitat and community composition [7].

Qualitative evaluations in southern Africa suggest different traps likely collect different

subsets of the mosquito community [15]. However, the two studies systematically

evaluating trapping methods in southern Africa focus only on house-based trapping in

residential areas [16,17].

An evaluation of trapping methods is also needed to influence the design of future

mosquito surveillance efforts. Such efforts could provide baseline information for public

health interventions by identifying hotspots with a high abundance of vector species

[18,19]. There were more than 26,000 estimated cases of malaria in southern Africa in

2017 (Botswana, Namibia, South Africa, Lesoto, Swaziland) [20]; key malaria vectors

include members of the *Anopheles gambiae* complex *(An. arabiensis, An. gambiae, An.*

merus) [21–23] and the *An. funestus* complex [24]. Mosquito-borne livestock infections are

also a concern, as outbreaks of West Nile virus, Rift Valley fever, Sindbis, and Wesselsbron

occur [25,26]. Key vectors for these viral infections include, *Aedes caballus, Ae.*

circumluteolus, Ae. dentatus, Ae. juppi, Ae. mcintoshi, Ae. ochraceus, Culex pipiens, Cx.

quinquefasciatus, Cx. univittatus, and *Cx. theileri* (based on suspected or known prime

vectors in Africa, reviewed elsewhere [26]). Previous work has characterized the

distribution [27], ecological drivers [28], and consequences for malaria risk [29] of key

Anopheline species. Understanding the distribution and drivers of non-Anophelinae

mosquito species in southern Africa (but see [15,30]) could facilitate informed

management of a broader range of mosquito-borne disease or the development of vector

control programs that target multiple infections.

In this study, we assessed four commonly used trapping methods for adult mosquito

data collection in natural savanna ecosystems. We compared estimates of abundance and

community composition from four traps: Centers for Disease Control miniature light trap

(CDC), Biogents Sentinel trap (BG), Biogents gravid *Aedes* trap (GAT), and a CO_2-baited net

trap. We also evaluated the importance of trapping method for inferring the regional

distribution of mosquito abundance, mosquito community composition, and the

distribution of key disease vectors.

Methods

Study location

We sampled twenty sites within five regions of Kruger National Park (KNP), South

Africa (Fig 1). Sampling sites within KNP were chosen to cover the geographic extent of the

park and to capture the variability in rainfall, geology and vegetation types within KNP

(between 22°31' and 25°31 S and 30°45 and 32°00 E) [31]. This region experiences

summer rainfall between November and April, and we selected our sampling to occur from

March to April 2017, when mosquito populations are generally high. We sampled 5 out of

the 22 management regions within the park: Malelane, Skukuza, Satara, Shingwedzi, and

Punda Maria. Within each region, we sampled at four sites, which were selected based on

multiple criteria (Fig 1). The primary selection criterion was to sample from water bodies

that represented diverse types of wetlands (temporary ponds, permanent ponds, rivers).

Additional criteria stipulated that the water bodies were at least 2 km away from one

another to avoid sampling mosquitoes from adjacent water bodies and within 25 km from

one another for sampling logistics. These distances are justified based on mean mosquito

dispersal rates, which range from 35m to 1.4 km depending on the species and habitat [32–

35]. Although dispersal farther than 2 km is possible, it is uncommon [36] and we do not

expect it to influence our results because only one pair of sites in Shingwedzi was located

within 1km of each other due limited surface water availability in the area.

Trapping and identification

We trapped at each site within a region for four consecutive nights and moved

sequentially between regions each week in random order (Skukuza: 20-March to 23-

March-2017; Malelane 27-March to 30-March-2017; Satara 3-April to 6-April-2017;

Shingwedzi 17-April to 20-April; Punda Maria 24-April to 27-April-2017). The four traps

we used were the Centers for Disease Control miniature light trap with an incandescent

light (Bioquip Inc, USA), a Biogents Sentinel trap (Biogents AG, Germany), the Biogents

gravid *Aedes* trap (Biogents AG, Germany), and the net trap. Although the net trap is not

commercially available, it is easily and inexpensively made from netting and poles (see

[15,30]). For consistency in sampling and to minimize any bias due to the traps' location,

we set up the traps at a similar distance away from the water body (15-25m) and

consistently rotated their positions around the water body at each site. The four positions

were defined based on their cardinal direction around the water body. Position 1 was set

directly east of the water body and positions 2-4 were set approximately 30-50m away

from each other moving northward around the water body. For the first two nights of

trapping, the BG trap was set in position 1, the GAT trap in position 2, the net trap in

position three, and the CDC trap in position 4. For the second two nights of trapping, the

traps were rotated one position, such that the CDC trap was moved to position 1.

Our trap-use protocol was based on preliminary trapping work and expertise from

mosquito surveillance in southern Africa [15,30]. Specifically, we equipped the CDC trap,

BG trap, and net trap with a closed container containing 200-400g of dry ice. To ensure the

dry ice would last overnight, we reduced sublimation by wrapping the dry ice in multiple

layers of newspaper and ensuring the container was closed except for 2 small holes. We

placed the dry ice containers inside the BG traps, at the center of the net trap, and hanging

with the CDC trap according to the manufacturer's instructions. In addition to the spatial

locations defined above, we hung the CDC light trap on a branch so it was approximately

1m from the ground, and we placed the BG and GAT traps on the ground. We ensured the

traps were placed close to the vegetation, but not directly under it, which has been shown

to improve trapping success [10]. If available, we placed the net trap in slight shade and

pulled the netting down to leave an approximately 10cm opening from the ground. Our

trapping schedule required traps to be set up in the field before dusk (4-6pm) and emptied

again at dawn (6-7am). The timing of this sample collection was used because the net trap

requires clearing at dawn, as mosquitoes may leave the trap after sunrise. Due to rainfall,

we excluded one trapping night at all sites in Satara and Punda Maria from all analyses.

During the study, five sampling events (nights at one site) with the BG trap, two

sampling events with the CDC trap, and one sampling event with the net trap were lost due

to animal interference that damaged the trap. An additional 11 nights of BG trap sampling

were lost due to other logistical reasons, primarily permanent damage by hyenas, which we

hypothesize is due to the Octenol bait. Our statistical analyses account for trap loss by

excluding data or comparisons from damaged traps.

Directly after emptying the traps, we stored the mosquitoes on dry ice in the field

and at -20°C until identification. For identification of Anophelinae mosquitoes, we used

identification literature from Gilles & Coutzee [37]. All Anophelinae were identified to

species except for members of *An. funestus* complex and the *An. gambiae* complex (referred

to here as *An. gambiae s.l*)*.,* which require molecular methods for identification [38]. For

identification of Culicinae, we used the key by Peter Jupp [14]. Species were identified

independently in duplicate by coauthors. Because species identification in the *Aedes vexans*

complex (*Ae. vexans, Ae. hirsutus, Ae. fowleri, Ae. durbanensis, Ae. natronius*), and the *Aedes*

dentatus complex (*Ae. dentatus, Ae. subdentatus Ae. pachyurus, Ae. bevisi, Ae. cumminsii*)

were inconsistent at the species level, we aggregated them to the species complex level.

Abiotic measurements

Temperature and rainfall within KNP follow a north-south gradient, with the highest

values in the south-west [39]. Temperature, relative humidity, and wind speed may

influence trapping success; they were monitored using a Kestrel 3000 handheld weather

meter (NK Inc., Boothwyn, PA, USA). We calculated the median temperature, relative

humidity, and wind speed across sites within a region for each morning at the time of

collection (Table S1). All environmental variables were standardized for analysis, by

centering and dividing by one standard deviation.

Assessing how the number of traps and trapping nights influences mosquito richness

We assessed how observed species richness (the total number of unique species)

saturates with sampling effort. We aggregated the data successively over 1, 2, 3, or 4 nights

and calculated the cumulative proportion of species identified with an increasing

proportion of nights. We also evaluated each trap's ability to estimate richness by

comparing richness estimated in pairs of trap types at each site aggregated across all

sampling nights using a Wilcoxon signed-rank test and a Bonferroni correction for multiple

comparisons. This analysis accounts for trap losses by excluding comparisons from

damaged traps.

Assessing whether trap type influences estimates of mosquito abundance and inference on the

regional patterns of abundance

To test for differences in abundance, we compared the number of mosquitoes

collected per night between pairs of traps. First, we quantified the relationship between

trap type and abundance with a Wilcoxon signed-rank test and a Bonferroni correction for

comparisons among the 4 traps. Then, we quantified the influence of trap type for

inferences on the regional patterns of abundance using linear regressions with Poisson

errors and a log-link function. The regression analyses assessed the relationship between a

trap's nightly mosquito counts with region of the park (*region*) and weather conditions

(wind speed, temperature, relative humidity (*RH*)). These assumptions result in the

following full model,

$\log(\mu_i) = \propto_{j[i]} + \beta_{1,k} region_k + \beta_2 windspeed_i + \beta_3 temperature_i + \beta_4 RH_i + \varepsilon_i$ (1)

where μ_i is the expected number of mosquitoes captured at evening *i* in the trap, ε_i is the

Poisson distributed random error, and $\propto_{j[i]}$ is the site-specific intercept.

We fit the regression model separately to data from the CDC trap, the net trap, the

BG trap, and data from aggregating abundance across all traps at a site. Because no

individuals were collected in the GAT trap on most nights, we did not fit the model to data

from the GAT trap alone. For each dataset, we conducted model selection using backward

selection based on AICc and select the model with the lowest AICc value. From the selected

model, we evaluate $\beta_{1,k}$ to determine how the average counts of mosquitoes in region k

differed from counts in Malelane, and we evaluate $\beta_2, \beta_3, \beta_4$ to determine how the average

counts of mosquitoes vary with wind speed, temperature, and relative humidity,

respectively. If the relative values of $\beta_{1,k}$ are consistent among traps (e.g. $\beta_{1,1} > \beta_{1,2} > \beta_{1,3}$)

for models fit to data from different traps, then we conclude that trap choice does not

influence spatial comparisons among regions. We fit all regression models in R [40] using

the lme4 package [41].

Assessing whether trap type influences estimates of mosquito community composition and

inference on the regional patterns of mosquito communities

To assess if different traps provide different estimates of community composition,

we first evaluated if certain species were particularly attracted to one trap over the other.

We assessed species-specific trap bias by calculating the difference in the number of

individuals for each species sampled between each pair of traps collected at a site over all

nights of trapping. Because traps were paired at each site, we tested for differences

between the traps using a Wilcoxon signed-rank test and a Bonferroni correction for

multiple comparisons. We only compared the species-specific trap bias of common species,

defined as being observed in the dataset more than three times. Because 23 common

species were compared (k = 23), significant differences between traps occur when p-values

are less than p = 0.05/23. We assessed trap bias for rare species visually.

To test for differences in community composition among traps, we used a

nonparametric analysis of similarities analysis (ANOSIM), visualized potential differences

with nonmetric multidimensional scaling (NMDS), and quantified the influence of trap type

for inferences on the regional patterns of community composition with hierarchical

clustering. For all analyses, we calculated Bray-Curtis dissimilarity matrices based on the

trap-specific (BG, CDC, net) abundances of all taxa within a site aggregated across sampling

nights. The ANOSIM analysis is a non-parametric test for differences in mosquito

communities among traps that compares the ranks of Bray-Curtis dissimilarity measures

from samples collected from the same vs. different traps [42]. To visualize this, we created

an ordination of traps and sites in mosquito community space for each region of the park.

Before all ordinations, we applied a Wisconsin transformation followed by a square root

transformation to the species matrices, which standardizes by species maxima and reduces

the influence of highly abundant taxa, respectively [43]. The ordinations converged on a

stable two-dimensional solution, based on stress values. We conducted all community

analyses in R using the vegan package [44].

Describing regional patterns of disease vectors

We additionally describe how known prime vectors for West Nile virus (*Cx. pipiens,*

Cx. quinquefasciatus, Cx. theileri, Cx. univittatus), Rift Valley fever (*Ae. dentatus, Ae.*

mcintoshi, Ae. ochraceus), Sindbis (*Cx. univittatus*), and Wesselsbron (none found) are

distributed across regions [21,26]. Additional known prime vectors for these infections

were not found in the study (Ae. caballus, *Ae. circumluteolus, Ae. juppi*). Chikungunya and

dengue fever outbreaks are less common in South Africa [14], but we describe the

distribution of their vector, *Ae. aegypti* [21] because additional known prime vectors in

Africa were not found (*Ae. africanus, Ae. albopictus, Ae. cordellieri, Ae. furcifer, Ae.*

luteocephalus, *Ae. neoafricanus*, *Ae. taylori*). We focus on the known prime vectors for a

conservative description, but additional mosquito species are considered suspected vectors

(reviewed in [26]). We use the numbers of *Cx. pipiens* complex to approximate the numbers

of *Cx. pipiens* and *Cx. quinquefasciatus; Cx. univittatus* complex for *Cx. univittatus; Ae.*

dentatus complex for *Ae. dentatus.* Although this assumption may hide epidemiologically

important variation, it is a valid approximation for these infections because multiple

members of the complex are known or suspected vectors. However, we do not plot the

distribution of malaria vectors because the most abundant species of *An. gambiae s.l.* in

KNP, *An. quadriannulatus*, is not a malaria vector [15].

Results

We collected 955 female mosquitoes, 946 (99.1%) of which were identified to the

level of species or species complex (Table S2). The most common species included

members of the *Culex univittatus* complex, *Aedes vexans* complex, and *Culex pipiens*

complex. We also collected over 50 *Anopheles gambiae* s.l., *Anopheles pretoriensis*, and

Culex theileri females.

Mosquito communities can be characterized with the net and CDC traps after multiple

trapping nights

Based on all traps together, mosquito community richness was sensitive to the

number of sampling nights (Fig 2; Fig S1). Taken across all sites in Malelane, 89% (24/27)

of the total number of unique species identified after four trapping nights were collected by

night 2 and 96% (26/27) were collected by night 3. In Shingwedzi, 79% (19/24) of the

species were collected by night 2 and 96% (23/24) by night 3. In Punda Maria, 53% (8/15)

of the species were collected by night 2 and 87% (13/15) by night 3. We note that these

percentages overestimate the percent of richness captured because species accumulation

curves suggest more than four nights of sampling are required to estimate total species

richness (Fig 2). The CDC and net trap together sampled all of the mosquito species

captured (range across regions, 93-100%), while the BG and GAT trap captured far fewer

species (Fig 2). Estimates of richness did not significantly differ between samples from the

net or CDC trap (Wilcoxon signed-rank test, p-value = 0.940), but estimates based on the

BG trap were lower than both the net and CDC trap (p-values <0.001).

Daily mosquito abundance estimates were comparable between the net and CDC traps but not

the BG and GAT trap

Most female individuals were collected in the CDC or the net trap, while the BG and

GAT trap captured far fewer individuals (Fig 3a). Together, the CDC and net trap sampled

96% of the individuals collected (range across regions, 94-99%). Estimates of mosquito

abundance did not significantly differ based on samples from the net or CDC trap (Wilcoxon

signed-rank test, p-value = 0.095), but estimates based on the BG trap were lower than

both the net and CDC trap (p-values <0.001; Fig 3b). Regression analyses showed that trap

choice influences comparisons between regions. All traps identified similar trends, with the

highest numbers captured in Malelane and the lowest numbers captured in Punda Maria

(Fig3c; Table S3). However, this spatial pattern was only significantly different for models

fit to the CDC data ($\beta_{1,1}$ for Punda Maria vs. Malelane = -1.88, p-value = 0.002). Unlike the

CDC trap, the BG and net trap were influenced by weather conditions (Table S3). The BG

and net trap collected higher numbers in warm, low wind conditions (Fig 3c). The net trap

also collected higher numbers in low relative humidity conditions. Results based on counts

aggregated from multiple traps were similar to results based on counts from either the CDC

or net trap alone (Fig 3c; Table S3).

Mosquito community composition was consistent between the net and CDC trap but not the

BG trap

Community composition was similar between the net and CDC trap, but not for the

BG trap (ANOSIM overall R = 0.126, p-value = 0.04; pairwise p-values: net vs. CDC p-value =

0.894; net vs. BG p-value = 0.023; CDC vs. BG p-value = 0.009). NMDS ordinations of traps

in species space reflect this relationship although some variation across regions of the park

does exist (Fig S2; Table S4). However, we did not find evidence for species-specific bias

between any of the traps, suggesting that differences in community composition in the BG

trap are driven by the relatively lower abundance collected in the trap (Fig 4; Fig S3). For

common species collected in the net vs. CDC trap, the number of individuals collected was

not significantly different between the traps (Fig 4a; No hypothesis tests for individual

species were significant). Rare species also did not show trap biases and include *Ae.*

aerarius, Ae. metallicus, Ae. unidentatus, An. maculipalpis, An. ziemanni, Cx. antennatus, Cx.

bitaenorhynchus, Cx. nebulosus, Lutzia tigripes, Mansonia africana, and *Uranotaenia balfouri*

(Fig S4). For comparisons with the BG trap, the net and CDC traps both collected higher

numbers of individuals, but there were no species or genus shifts driving this pattern (Fig

4b-c; Fig S3).

315   Although trap choice influences estimates of community composition (e.g.

316  community richness, ordinations), hierarchical cluster analysis suggests that may be less

317  important for comparisons between regions (Fig 5; Fig S5). Regional patterns in mosquito

318  communities were consistent across trap types, with samples from Malelane and Satara

319  being more similar than samples from Shingwedzi. Mosquito communities estimated from

320  the CDC and net trap were clustered by region (Fig S5), indicating that communities within

321  a region were more closely related to each other than communities between regions

322  regardless of the net or CDC trap. In contrast, samples from the BG trap were clustered

323  separately from the samples from the net and CDC trap due to the overall lower

324  abundances in the BG trap (Fig 5). We, therefore, describe the regional patterns of disease

325  vectors based on data from all traps together (Fig 6). Rift Valley fever vectors were most

326  common in Satara while West Nile virus and Sindbis vectors were more common in

327  Shingwedzi and Punda Maria. We did not find known vectors for Wesselsbron (*Ae. caballus,*

328  *Ae. circumluteolus*) within the park.

Discussion

331   Mosquito trapping is used for disease surveillance, biodiversity surveys, and

332  nuisance-reduction. In light of these multiple, non-exclusive aims, this study compared

333  traps based on four potential goals: collecting large numbers of mosquitoes; estimating

334  mosquito community composition, including vector or rare species; characterizing the

335  spatial patterns of abundance; and characterizing the spatial patterns of community

336  composition. We expected trade-offs among these aims, for example with traps specializing

337  on one vector to be less successful in estimating community richness (and vice versa, e.g.

[45]). In contrast to this expectation, our results indicated that the net and CDC trap

consistently performed best across multiple outcomes.

The CDC and net trap collected higher abundances and more unique species

compared to the BG and GAT traps, which collected no mosquitoes on most nights. This

result is different from trap comparison studies in Europe [7], the U.S. (BG: [13], GAT: [46]),

and South America (BG: [47], GAT: [48]), where both the BG and GAT traps have been

shown to perform well. One reason for their relatively low success within KNP could be the

diversity and types of species present within the park. The GAT trap has been designed to

capture container-breeding species, such as *Ae. aegypti* [48], and the BG trap performs well

in sampling *Ae. aegypti, Ae. albopictus*, and *Cx. pipiens* [7,49]. Although *Ae. aegypti* was

present in our study and previous studies [15], they were relatively rare (Table S1). The BG

trap's sampling efficiency for the two most-common species complexes in our dataset –

members of the *Ae. vexans* complex and *Cx. univitatus* complex – is either low [7] for the *Ae.*

vexans complex or previously unknown for the *Cx. univitatus* complex. An alternative

reason for the low success of the BG trap could be that it is negatively influenced by the

presence of alternative refugia and oviposition sites [50]. Habitat heterogeneity is likely to

be higher within KNP compared to other, primarily urban or suburban environments

where mosquito traps have been evaluated ([13, 46, 47], but see [7]). Additional

comparisons in urban environments in southern Africa are needed to distinguish these two

hypotheses.

By providing a detailed comparison between the net and CDC traps, our results also

suggest that estimates of community richness and diversity are comparable between them.

The CDC and net traps differed, however, in their inference about which regions had the

highest mosquito abundance because the net trap was more sensitive to weather

conditions. This difference is likely to be important for studies comparing vector

abundance across space or time and suggests that vector surveillance or control programs

should use caution in comparing patterns of abundance between studies that did not use

the same trap. However, the consistent patterns of community composition in the net and

CDC trap suggest that comparisons of vector communities between studies using the net

and CDC trap are possible.

The choice between the CDC and net trap should also consider other features of the

traps, such as ease of use and specimen quality. For morphological identification, the net

trap has the advantage that mosquito specimens can be collected with minimal damage,

which makes them easier to identify [14]. Therefore, studies requiring precise species

identification such as biodiversity assessments may prefer the net trap. However, for

sampling large numbers of sites or sites in remote locations, the CDC trap has an advantage

because the timing of when traps need to be cleared is more flexible compared to the net

trap, which has to be cleared at sunrise. These practical considerations may mean that the

CDC trap is better suited for large, comparative studies.

378   Conclusions

After assessing four different mosquito trapping methods in a natural savanna

ecosystem, we recommend the net trap, the CDC trap, or their combined use for outdoor

mosquito surveillance in southern Africa. These traps performed well based on four

evaluation criteria: collecting large numbers of mosquitoes; estimating mosquito

community composition, including vector or rare species; characterizing the spatial

patterns of abundance; and characterizing the spatial patterns of community composition.

This suggests they are appropriate for both biodiversity surveys and vector surveillance. As

such, this study provides a valuable proof-of-principle for characterizing the spatial

patterns of non-vectors as well as vectors for multiple diseases.

Declarations

Ethics approval: Not applicable

Consent for publication: Not applicable

Availability of data and materials: The dataset supporting the conclusions of this article are

available in the dryad digital repository and the free online mosquito databases, VectorMap

(http://vectormap.si.edu).

Competing Interests: The authors declare that they have no competing interests.

Funding: This study was supported by the Gratama Fund from University of Leiden (grant

number 2016.08) the Uyttenboogaart-Eliasen foundation for comparative entomology to

EEG (SUB.2016.12.08), and the RCN-IDEAS travel grant to EEG.

Author contribution statement: EG, BB, MS conceived and designed the analysis; EG, BB, DG,

MS collected the data; EG, MG, MS conducted identifications; EG, PB, MS performed analyses;

all authors wrote the paper.

Acknowledgements: We thank South African National Parks (SANParks) for their support in

conducting this study in Kruger and providing us with mosquito specimens to help with

identification. We thank Purvance H. Shikwambana and a great field team: Vicky Beckers,

Nina Haver, Louie Krol, Skhumbuza Mdletshe, Karabo Moloi, Nondumiso Myataza, and Gijs

van Nes. We thank Dr. Alan Kemp, Professor. A. Paulo Gouveia de Almeida, and Dr. Anthony

Cornel for their assistance with identifications. This study was supported by the Gratama

Fund from University of Leiden (grant number 2016.08) the Uyttenboogaart-Eliasen

foundation for comparative entomology to EEG (SUB.2016.12.08), and the RCN-IDEAS

travel grant to EEG.

References

1. Meyer SDB, Ritchie SA, Laurance SGW. Mosquito communities and disease risk influenced

by land use change and seasonality in the Australian tropics. Parasit Vectors. 2016;9:387.

2. Janko MM, Irish SR, Reich BJ, Peterson M, Doctor SM, Mwandagalirwa MK, et al. The links

between agriculture, *Anopheles* mosquitoes, and malaria risk in children younger than 5 years in

the Democratic Republic of the Congo: a population-based, cross-sectional, spatial study. Lancet

Planet Health. 2018;2:e74–82.

3. Diuk-Wasser MA, Brown HE, Andreadis TG, Fish D. Modeling the spatial distribution of

mosquito vectors for West Nile virus in Connecticut, USA. Vector Borne Zoonotic Dis.

2006;6:283–95.

4. Galardo AKR, Zimmerman RH, Lounibos LP, Young LJ, Galardo CD, Arruda M, et al.

Seasonal abundance of anopheline mosquitoes and their association with rainfall and malaria

along the Matapí River, Amapá, Brazil. Med Vet Entomol. 2009;23:335–49.

5. Dennett JA, Vessey NY, Parsons RE. A comparison of seven traps used for collection of

Aedes albopictus and *Aedes aegypti* originating from a large tire repository in Harris County

(Houston), Texas. J Am Mosq Control Assoc. 2004;20:342–9.

6. Hoel DF, Kline DL, Allan SA. Evaluation of six mosquito traps for collection of *Aedes*

albopictus and associated mosquito species in a suburban in north central Florida. J Am Mosq

Control Assoc. 2009;25:47–57.

7. Lühken R, Pfitzner WP, Börstler J, Garms R, Huber K, Schork N, et al. Field evaluation of

four widely used mosquito traps in Central Europe. Parasit Vectors. 2014;7:268.

8. L'Ambert G, Ferre J-B, Schaffner F, Fontenille D. Comparison of different trapping methods

for surveillance of mosquito vectors of West Nile virus in Rhone Delta, France. J Vector Ecol.

2012;37:269–75.

9. Jupp PG, McIntosh BM, Nevill EM. A survey of the mosquito and culicoides faunas at two

localities in the Karoo region of South Africa with some observations on bionomics.

Onderstepoort J Vet Res. 1980;47:1–6.

10. Crepeau TN, Healy SP, Bartlett-Healy K, Unlu I, Farajollahi A, Fonseca DM. Effects of

Biogents Sentinel Trap field placement on capture rates of adult Asian tiger mosquitoes, Aedes

albopictus. PLoS One. 2013;8:e60524.

11. Kraemer MUG, Sinka ME, Duda KA, Mylne AQN, Shearer FM, Barker CM, et al. The

global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. eLife.

2015;4:e08347.

12. Hay SI, Sinka ME, Okara RM, Kabaria CW, Mbithi PM, Tago CC, et al. Developing Global

Maps of the Dominant *Anopheles* Vectors of Human Malaria. PLoS Med. 2010;7:e1000209.

13. Farajollahi A, Kesavaraju B, Price DC, Williams GM, Healy SP, Gaugler R, et al. Field

Efficacy of BG-Sentinel and Industry-Standard Traps for *Aedes albopictus* (Diptera: Culicidae)

and West Nile Virus Surveillance. J Med Entomol. 2009;46:919–25.

14. Jupp PG. Mosquitoes of southern Africa. P.O.Box 178 Hartebeespoort 0216, South Africa:

Ekogilde Publishers; 1996.

15. Cornel AJ, Lee Y, Almeida APG, Johnson T, Mouatcho J, Venter M, et al. Mosquito

community composition in South Africa and some neighboring countries. Parasit Vectors.

2018;11:331.

16. Sikaala CH, Killeen GF, Chanda J, Chinula D, Miller JM, Russell TL, et al. Evaluation of

alternative mosquito sampling methods for malaria vectors in Lowland South - East Zambia.

Parasit Vectors. 2013;6:91.

17. Govella NJ, Chaki PP, Mpangile JM, Killeen GF. Monitoring mosquitoes in urban Dar es

Salaam: Evaluation of resting boxes, window exit traps, CDC light traps, Ifakara tent traps and

human landing catches. Parasit Vectors. 2011;4:40.

18. Maharaj R, Moonasar D, Baltazar C, Kunene S, Morris N. Sustaining control: lessons from

the Lubombo spatial development initiative in southern Africa. Malar J. 2016;15:409.

19. Sharp BL, Kleinschmidt I, Streat E, Maharaj R, Barnes KI, Durrheim DN, et al. Seven years

of regional malaria control collaboration Mozambique, South Africa, and Swaziland. Am J Trop

Med Hyg. 2007;76:42–7.

20. World Health Organization. Global Health Observatory (GHO) data. Malaria country

profiles. [Internet]. Available from: https://www.who.int/malaria/publications/country-

profiles/profile_zaf_en.pdf?ua=1

21. Burke A, Dandalo L, Munhenga G, Dahan-Moss Y, Mbokazi F, Ngxongo S, et al. A new

malaria vector mosquito in South Africa. Sci Rep. 2017;7:43779.

22. Coetzee M, Craig M, le Sueur D. Distribution of African malaria mosquitoes belonging to

the *Anopheles gambiae* complex. Parasitol Today. 2000;16:74–7.

23. Coetzee M, Hunt RH, Wilkerson R, Torre AD, Coulibaly MB, Besansky NJ. *Anopheles*

coluzzii and *Anopheles amharicus*, new members of the *Anopheles gambiae* complex. Zootaxa.

2013;3619:246–74.

24. Coetzee M, Fontenille D. Advances in the study of *Anopheles funestus*, a major vector of

malaria in Africa. Insect Biochem Mol Biol. 2004;34:599–605.

25. Jupp P. Mosquitoes as vectors of human disease in South Africa. South Afr Fam Pract.

2005;47:68–72.

26. Braack L, Gouveia de Almeida AP, Cornel AJ, Swanepoel R, de Jager C. Mosquito-borne

arboviruses of African origin: review of key viruses and vectors. Parasit Vectors. 2018;11:29.

27. Coetzee M, Hunt R, Braack L, Davidson G. Distribution of mosquitos belonging to the

Anopheles gambiae complex, including malaria vectors, south of latitude 15-degrees-S. South

Afr J Sci. 1993;89:227–31.

28. Charlwood JD. Some like it hot: a differential response to changing temperatures by the

malaria vectors *Anopheles funestus* and *An. gambiae* s.l. Peerj. 2017;5:e3099.

29. Kabaghe AN, Chipeta MG, Gowelo S, Mburu M, Truwah Z, McCann RS, et al. Fine-scale

spatial and temporal variation of clinical malaria incidence and associated factors in children in

rural Malawi: a longitudinal study. Parasit Vectors. 2018;11:129.

30. Hammami P, Tran A, Kemp A, Tshikae P, Kgori P, Chevalier V, et al. Rift Valley fever

vector diversity and impact of meteorological and environmental factors on Culex pipiens

dynamics in the Okavango Delta, Botswana. Parasit Vectors. 2016;9:434.

31. Pickett STA, Cadenasso ML, Benning TL. Biotic and abiotic variability as key determinants

of savanna heterogeneity at multiple spatiotemporal scales. The Kruger Experience: Ecology and

Management of Savanna Heterogeneity. Washington: Island Press; 2003. p. 22–40.

32. Ciota AT, Matacchiero AC, Kilpatrick AM, Kramer LD. The effect of temperature on life

history traits of Culex mosquitoes. J Med Entomol. 2014;51:55–62.

33. Hamer GL, Anderson TK, Donovan DJ, Brawn JD, Krebs BL, Gardner AM, et al. Dispersal

of adult *Culex* mosquitoes in an urban West Nile virus hotspot: A mark-capture study

incorporating stable isotope enrichment of natural larval habitats. PLoS Negl Trop Dis.

2014;8:e2768.

34. Muir LE, Kay BH. *Aedes aegypti* survival and dispersal estimated by mark-release-recapture

in northern Australia. Am J Trop Med Hyg. 1998;58:277–82.

35. Medeiros MCI, Boothe EC, Roark EB, Hamer GL. Dispersal of male and female *Culex*

quinquefasciatus and *Aedes albopictus* mosquitoes using stable isotope enrichment. PLoS Negl

Trop Dis. 2017;11:e0005347.

36. Verdonschot PFM, Besse-Lototskaya AA. Flight distance of mosquitoes (Culicidae): A

metadata analysis to support the management of barrier zones around rewetted and newly

constructed wetlands. Limnologica. 2014;45:69–79.

37. Gillies T, Coetzee M. A supplement of the Anophelinae of Africa south of the Sahara

(Afrotropical Region): Publications of the South African Institute for Medical Research.

1987;55:1–143.

38. Koekemoer LL, Kamau L, Hunt RH, Coetzee M. A cocktail polymerase chain reaction assay

to identify members of the *Anopheles funestus* (Diptera: Culicidae) group. Am J Trop Med Hyg.

2002;66:804–11.

39. Venter FJ, Scholes RJ, Eckhardt HC. The abiotic template and its associated vegetation

pattern. The Kruger Experience: ecology and management of savanna heterogeneity.

Washington: Island Press; 2003. p. 83–129.

40. R Development Core Team. R: A language and environment for statistical computing.

Vienna, Austria: R Foundation for Statistical Computing; 2014.

41. Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. J

Stat Softw. 2015;67:1–48.

42. Anderson MJ, Walsh DCI. PERMANOVA, ANOSIM, and the Mantel test in the face of

heterogeneous dispersions: What null hypothesis are you testing? Ecol Monogr. 2013;83:557–74.

43. Legendre P, Gallagher ED. Ecologically meaningful transformations for ordination of species

data. Oecologia. 2001;129:271–80.

44. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:

Community Ecology Package. R package version 2.5-3 [Internet]. 2018. Available from:

https://cran.r-project.org/package=vegan

45. Brown R, Hing CT, Fornace K, Ferguson HM. Evaluation of resting traps to examine the

behaviour and ecology of mosquito vectors in an area of rapidly changing land use in Sabah,

Malaysian Borneo. Parasit Vectors. 2018;11:346.

46. Burkett DA, Kelly R, Porter CH, Wirtz RA. Commercial mosquito trap and gravid trap

oviposition media evaluation, Atlanta, Georgia. J Am Mosq Control Assoc. 2004;20:233—238.

47. Degener CM, Eiras ÁE, Ázara TMF, Roque RA, Rösner S, Codeço CT, et al. Evaluation of

the Effectiveness of Mass Trapping With BG-Sentinel Traps for Dengue Vector Control: A

Cluster Randomized Controlled Trial in Manaus, Brazil. J Med Entomol. 2014;51:408–20.

48. Eiras AE, Buhagiar TS, Ritchie SA. Development of the gravid *Aedes* trap for the capture of

adult female container-exploiting mosquitoes (Diptera: Culicidae). J Med Entomol.

2014;51:200–9.

49. Meeraus WH, Armistead JS, Arias JR. Field comparison of novel and gold standard traps for

collecting *Aedes albopictus* in northern Virginia. J Am Mosq Control Assoc. 2008;24:244–8.

50. Ball TS, Ritchie SR. Sampling biases of the BG-Sentinel trap with respect to physiology,

age, and body size of adult *Aedes aegypti* (Diptera: Culicidae). J Med Entomol. 2010;47:649–56.

 Figures

547   Fig 1. Map of the spatial locations and weather conditions of trapping sites within KNP.

548   Colors represents the five regions where trapping occurred. From south to north, these

549   include the Malelane, Skukuza (no weather data), Satara, Shingwedzi, and Punda Maria

550   sections. Each dot in the map represents a unique water body sampled. Regions of the park

551   were characterized by distinct weather patterns (Table S1).

Fig 2. Richness (number of unique species) was sensitive to sampling effort and trap type.

Richness values in each region were aggregated across sites; figure S1 displays data for

each site within a region. Solid lines indicate that no traps were compromised at any site,

dashed lines indicate that one trapping site was not collected. Sites within the Satara and

Punda Maria region were only sampled for three nights due to rain.

Fig 3. Mosquito abundance. (A) The number of mosquitoes captured by trap type; dots

represent the number captured on each night at each site. (B) Histograms showing the

difference in the number of mosquitoes sampled between trap types paired by sampling

site. (C) Regression parameter estimates and standard errors from statistical models

characterizing the median number of mosquitoes sampled at sites relative to Malelane and

compared to the mean relative humidity (RH), temperature, or wind speed conditions.

Colors indicate whether data used for model fitting was based on one trap or aggregated

from multiple traps. Table S3 defines the parameter estimates and hypothesis tests.

Fig 4. There were no species-specific differences between (A) the net vs. cdc trap, while

species-specific differences between (B) the net vs. BG trap and the (C) CDC vs. BG trap

were driven by overall abundance in the net or CDC trap. Dots represent the difference in

the number of mosquitoes collected between pairs of traps based on the total number of

mosquitoes sampled across nights at each site. (A) Lines represent the median and

interquartile range of the data. Data displayed do not include sites where no individuals of

a given species were collected in either trap, but results remain consistent regardless of

whether these sites are or are not included. No hypothesis test for the individual species

was significant (p-values = 0.181, 0.174, 0.345, 0.143, 1, 0.134, 0.476, 0.346, 1, 0.498, 0.360,

0.152, 0.796, 0.931, 1, 0.719, 0.372, 0.423, 0.265, 0.725, 1, 0.850, 1). See Figure S3 for

species-specific comparisons with the BG trap and Table S4 for a summary table for the BG

trap.

Figure 5. Communities collected in the net and CDC trap are clustered by region. Traps and

regions are ordered according to the tree produced by clustering (Fig S5). Colors represent

species abundance, with color bins defining the 30th to 90th percentile in increments of 10.

592   Fig 6. The percentage of mosquitoes identified as primary vectors in each region.

593