Title

DYRK2 activates heat shock factor 1 promoting resistance to proteotoxic stress and tumour growth in triple-negative breast cancer

Authors/Affiliations

Rita Moreno[1], Sourav Banerjee[2], Angus W. Jackson[1], Jean Quinn[3], Gregg Baillie[3], Jack E. Dixon[2,4,5], Albena T. Dinkova-Kostova[1], Joanne Edwards[3] and Laureano de la Vega[1*]

[1]Division of Cellular Medicine, School of Medicine, University of Dundee, Scotland, UK.

[2]Department of Pharmacology, University of California San Diego, La Jolla, CA 92093-0721

[3]Unit of Gastrointestinal Oncology and Molecular Pathology, Institute of Cancer Sciences, College of Medical, Veterinary, and Life Sciences, University of Glasgow, Glasgow, United Kingdom

[4]Department of Cellular and Molecular Medicine, University of California San Diego, La Jolla, CA 92093, USA.

[5]Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, CA 92093, USA

*Correspondence: l.delavega@dundee.ac.uk

Summary

The proteotoxic-stress response pathway, controlled by heat shock factor 1 (HSF1), supports both cancer progression and chemoresistance and is an attractive target for cancer treatment. As developing HSF1 inhibitors has proved to be very challenging, the identification and targeting of upstream regulators of HSF1 presents a more tractable alternative strategy. Here we demonstrate that in triple negative breast cancer (TNBC) cells, the dual-specificity tyrosine-regulated kinase 2 (DYRK2) directly interacts with and phosphorylates HSF1, promoting its nuclear stability and transcriptional activity. Thus, DYRK2 depletion reduces HSF1 activity and sensitises TNBC cells to proteotoxic stress. Importantly, in tumours from TNBC patients, DYRK2 protein levels positively correlate with active HSF1 and associates with poor prognosis, suggesting that DYRK2 could be promoting TNBC. In agreement with this, DYRK2 depletion significantly reduces tumour growth in a TNBC xenograft model. These findings identify DYRK2 as both, an important modulator of the HSF1 transcriptional program, and a potential therapeutic target in TNBC.

Keywords

DYRK2; Proteotoxic stress; HSF1; Triple-Negative Breast Cancer; HSP70

Introduction

TNBC is a heterogeneous group of breast tumours defined by their lack of expression of estrogen (ER), progesterone (PR) and Her2 receptors, accounting for about 15-20% of all breast cancer cases[1]. TNBC tend to be diagnosed with higher grade cancer, have a greater rate of recurrence and metastases, and do not respond to ER- or HER2-targeted therapies, resulting in limited therapeutic options[2]. New strategies coupled with patient stratification

approaches to predict likely responders to new emerging treatments are urgently required to enhance the therapeutic options in TNBC.

Breast cancer cells, as most cancer cells, are exposed to sustained proteotoxic stress as a result of severe aneuploidy and high mutation load, which increases the amount of toxic, misfolded proteins in the cell[3, 4]. This proteotoxic stress is countered by the proteotoxic stress-response pathway, which promotes the correct folding of proteins via the heat shock factor 1 (HSF1) pathway (also known as the heat shock response). HSF1 therefore helps cancer cells to survive, supporting oncogenesis. This has been demonstrated by the reduced susceptibility of *Hsf1*-knockout mice to tumour formation driven either by *Ras* or *p53* mutations or by chemical carcinogens[5]. At basal conditions, HSF1 is maintained mainly in the cytoplasm and upon proteotoxic stress, HSF1 is activated and translocates to the nucleus[6]. HSF1 activity and stability are tightly controlled by multiple post-translational modifications[7]. Among these, phosphorylation of serine 320 and serine 326 are associated to enhanced transcriptional activity, stability and nuclear accumulation of HSF1[8-10]. Although the sequence of modification events leading to HSF1 activation or the interplay among the different phosphorylation events are unclear, the predominant view is that phosphorylation on S326 is critical and dominant over any inhibitory phosphorylation events, and thus HSF1 phosphorylated on S326 is a hallmark for HSF1 activation[11]. Once in the nucleus, HSF1 activates a transcriptional program including a number of heat shock proteins (HSPs), which function as molecular chaperones, facilitating the folding, transportation, ubiquitination, and proteasomal degradation of proteins[12, 13], protecting cells against proteotoxic stress and apoptosis[14]. Of the HSPs, *HSP70*, is one of the best-characterised HSF1 targets, and is expressed at high levels in a variety of cancers, correlating with growth, survival, invasion and

resistance to therapy[15]. For breast cancer in particular, high levels of HSF1 correlate with poor prognosis[16] and HSF1 depletion strongly reduces proliferation and viability of breast cancer cells, without affecting normal breast epithelial cells[5].

Although there is no direct evidence linking dual-specificity tyrosine-phosphorylation-regulated kinase 2 (DYRK2) to the HSF1 pathway in human cells, the yeast homolog of DYRK kinases, YAK1, phosphorylates and activates HSF1[17]. Thus, we hypothesized that DYRK2, by regulating the HSF1 pathway, could have a role in controlling proteotoxic stress responses and contributing to TNBC tumour progression.

Herein we found that DYRK2 phosphorylates and activates HSF1, promoting resistance to proteotoxic stress in TNBC cells, and tumour growth in TNBC xenograft models. We further report that the DYRK2-HSF1 link is also relevant in clinical TNBC samples. DYRK2 protein levels correlates with nuclear levels of HSF1, and associates with poor cancer-specific patient survival and high tumour recurrence. Together our data reveal DYRK2 as a kinase that promotes cancer cell survival by activating the HSF1 pathway, and support a tumour promoting role for DYRK2 in TNBC.

Results

1- DYRK2 phosphorylates HSF1

To test whether DYRK2 phosphorylates HSF1, we overexpressed DYRK2 and, by use of phosphospecific antibodies, we observed that the levels of endogenous HSF1 phosphorylated at S326 and S320 were increased (Fig. 1A); for this effect, the kinase activity of DYRK2 was required, as a kinase-dead version of DYRK2 (DYRK2-KD) did not induce HSF1 phosphorylation. To further test the role of DYRK2 kinase activity on HSF1 phosphorylation, we mutated the gatekeeper residue within DYRK2, creating an analogue-sensitive DYRK2

form (DYRK2-AS) that is selectively sensitive to PP1 inhibitors[18]. Using the DYRK2-AS form together with three different PP1 inhibitors, we observed that specific inhibition of DYRK2 activity, dose-dependently reduced the DYRK2-mediated phosphorylation of endogenous HSF1 (Fig. 1B). Additionally, harmine, a well characterised pan-specific DYRK inhibitor[19], was also able to reduce the phosphorylation of endogenous HSF1 mediated by DYRK2 overexpression (Fig. S1A).

To address if HSF1 is a direct substrate for DYRK2, we performed an *in vitro* kinase assay using recombinant His-HSF1 and GST-DYRK2. In close agreement with the cell culture experiments, we found that DYRK2 phosphorylates His-HSF1 at S326 and S320 *in vitro* (Fig. 1C). Furthermore, incubation of recombinant GST-DYRK2-AS with a PP1 inhibitor inhibited the phosphorylation of His-HSF1 on both S326 and S320 (Fig. S1B). Mass spectrometry analysis of a sample taken at an early time point (5 minutes) from the *in vitro* reaction showed that HSF1 was further phosphorylated at S307, T323 and S363 (Fig. S1C). Unfortunately, only antibodies against S307 were commercially available, and in our analysis, endogenous HSF1 was not phosphorylated at this site upon DYRK2 overexpression (not shown). We therefore focused on S320, S326, T323 and S363 in our subsequent studies.

Once we established that DYRK2 can phosphorylate HSF1 in cells and *in vitro*, we tested whether DYRK2 plays a role in HSF1 phosphorylation in response to proteotoxic stress in TNBC cell lines, using MDA-MB-231 and MDA-MB-468 as models. Using heat shock as a proteotoxic stress inducer, we showed that DYRK2 depletion leads to a reduction of HSF1 phosphorylation in response to heat shock (Fig. 1D). To test if this effect was specific for DYRK2, we silenced individually two other members of the DYRK family, DYRK1A and DYRK1B. We found that in our TNBC cell models, DYRK2 is the main DYRK member involved in HSF1 phosphorylation (Fig. 1E), although other members of the family can also modulate HSF1 to a lesser extent.

This is also the case in other cancer cell lines (Fig. S1D). In agreement with the role of DYRKs in heat shock-mediated HSF1 phosphorylation, incubation with harmine dramatically impaired HSF1 phosphorylation in response to heat shock in TNBC cells (Fig. 1F).

Interestingly, in addition to reducing HSF1 phosphorylation, DYRK2 knockdown appeared to reduce the levels of HSF1 under heat shock conditions (Fig. 1D, 1E, and Fig. S1D), which could be a cause or a consequence of the lower HSF1 phosphorylation levels. However, acute (1 hour) harmine-mediated DYRK inhibition reduced HSF1 phosphorylation without affecting HSF1 levels (Fig. 1F) which, based on the HSF1 relatively long half-life (>6 hours)[10, 20], suggests that reduction of HSF1 phosphorylation might precede or is uncoupled to HSF1 destabilisation.

2- DYRK2 interacts with HSF1 via two domains

Our results provide evidence to support a functional interaction between DYRK2 and HSF1. In order to test whether DYRK2 and HSF1 physically interact in cells, we performed co-immunoprecipitation. The interaction between the two proteins, which was not easily detectable at basal conditions, was increased upon exposure to proteotoxic stress (Fig. 2A and 2E). This weak interaction might reflect a transient binding between DYRK2 and HSF1, which is a common occurrence for kinases and their substrates. In order to map the interaction site(s), we used an *in vitro* interaction peptide array. By incubating recombinant GST-HSF1 with a membrane containing an array of overlapping peptides covering the entire DYRK2 protein, we identified two potential binding regions (Fig. 2B) within conserved aminoacid sequences (Fig. S2A). Binding region 1 (BR1) with sequence DDQGSYV, is located in a surface-exposed loop, and binding region 2 (BR2) with sequence TDA, falls within the unstructured C-terminus of DYRK2 (Fig. S2B). Although protein loops are not involved in

regulating kinase activity, they often play a crucial role in mediating protein-protein interactions[21]. In order to validate the functional relevance of these two regions, we created DYRK2 constructs harbouring mutations either in BR1 or BR2, or in both simultaneously. Based on our results, BR1 appeared to be more important for the ability of DYRK2 to phosphorylate HSF1 (Fig. 2C). Furthermore, when both DYRK2 binding regions were mutated, HSF1 phosphorylation was strongly reduced in both S320 and S326, but notably, the ability of DYRK2 to phosphorylate SIAH2[22], another known substrate, remained intact (Fig. 2D). This observation suggests that these two regions are not important for general DYRK2 kinase activity, but rather for its specific interaction with HSF1. As expected, the DYRK2 BR1/BR2 mutant (YV/AR-T/A) did not interact with HSF1 (Fig. 2E).

3- DYRK2 promotes HSF1 nuclear stability

As nuclear HSF1 is the active form of HSF1, we asked whether DYRK2 was affecting HSF1 nuclear levels. We tested in TNBC cells the role of DYRK2 on the nuclear levels of HSF1 at basal and after heat shock conditions by employing stable CRISPR-mediated knockout as well as siRNA-mediated knockdown. Both DYRK2 depletion strategies reduced HSF1 nuclear levels in basal conditions and after heat shock, as well as the levels of nuclear phospho-HSF1 in TNBC cells (Fig. 3A and Fig. S3A, S3B). This was also the case for other cancer cell lines (Fig. S3C and S3D). We used two independent gRNAs to knockout DYRK2 in the different TNBC cell lines, obtaining in both cases the same reduction in levels of nuclear HSF1 as with siDYRK2, suggesting a specific effect. Nevertheless, to further confirm the on-target effect of DYRK2 on HSF1 nuclear levels, we reconstituted DYRK2-KO cells with DYRK2, showing that the levels of nuclear HSF1 and phospho-HSF1 in DYRK2-KO cells were rescued (Fig. 3B). Moreover, DYRK2 overexpression resulted in increased HSF1 nuclear levels and phosphorylation (Fig. S3E).

In agreement with our previous results, acute harmine-mediated DYRK inhibition led to reduction in nuclear HSF1 phosphorylation without affecting nuclear HSF1 levels (Fig. 3C). To assess the effect of DYRK2 on HSF1 stability, we compared the HSF1 nuclear and cytosolic levels upon cycloheximide treatment in control and DYRK2 knocked-down TNBC cells. In cycloheximide chase experiments, DYRK2 depletion reduced the levels of nuclear but not cytosolic HSF1, suggesting that DYRK2 controls HSF1 nuclear stability (Fig. 3D).

The positive effect of DYRK2 on HSF1 stability could be either dependent or independent of DYRK2-mediated HSF1 phosphorylation. We reasoned that as phosphorylation on S326 and on S320 are known to promote HSF1 stability and nuclear accumulation respectively[8, 10], those two sites were most likely responsible for the positive effect of DYRK2 on HSF1 nuclear stability. Unexpectedly, our results showed that mutation of S326 and/or S320 were not sufficient to completely abolish the effect of DYRK2 on HSF1 stability (Fig. S3F). Nevertheless, additional mutation of the other two identified sites (T323 and S363) significantly reduced the effect of DYRK2 on HSF1 stability (Fig. 3E). These results demonstrate the relevance of the combined phosphorylation on HSF1 for the effect of DYRK2 on HSF1 stability, although we cannot completely rule out the participation of additional mechanisms unrelated to HSF1 phosphorylation.

4- DYRK2 affects the expression levels of the HSF1 target gene *HSP70*

Based on its positive effect on the levels of nuclear phosphorylated HSF1, we hypothesised that DYRK2 could be promoting HSF1 transcriptional activity. To test this hypothesis, we compared the levels of the prototypic HSF1 target gene, heat shock protein 70 (*HSP70*), in TNBC cells with and without DYRK2. DYRK2 knockout reduced significantly the induction of *HSP70* mRNA levels in response to heat shock (Fig. 4A and 4B) and in response to bortezomib,

another inducer of proteotoxic stress (Fig. S4A). Similarly, HSF1 knockout strongly affected the levels of *HSP70*, whereas HSF1/DYRK2 double knockout did not reduce the *HSP70* mRNA levels any further, confirming that the effect of DYRK2 on the expression of *HSP70* is mediated by HSF1 (Fig. 4A). Importantly the reduction on *HSP70* levels observed in DYRK2-KO cells was recovered by reconstituting them with DYRK2 (Fig. 4C). In addition, harmine reduced the *HSP70* mRNA levels in a DYRK2-dependent manner in TNBC as well as in other cancer cell lines (Fig. S4B).

5- DYRK2 reduces sensitivity to proteotoxic stress via HSF1

To assess the functional significance of the reduced HSF1 activity in DYRK2-impaired TNBC cells, we tested the effect of DYRK2 depletion on the sensitivity of these cells to proteotoxic stress. In agreement with our data on the importance of DYRK2 for HSF1 function and with the well-characterised protective role of HSF1 against proteotoxicity, DYRK2 depletion (both by siRNA and CRISPR-mediated knockout) sensitised TNBC cells to heat shock, as measured by levels of PARP cleavage, a marker of apoptosis (Fig 5A, 5B and Fig. S5A). The same result was observed in other cancer cell lines (Fig. S5B).

As expected, HSF1-deficient cells were more sensitive to heat shock than wild-type cells. Importantly, the effect of DYRK2 depletion on the enhanced sensitivity to proteotoxic stress on wild-type cells was largely abolished in TNBC HSF1-KO cells (Fig. 5C), confirming the involvement of HSF1. In agreement, harmine sensitised TNBC cells to heat shock in a HSF1-dependent manner (Fig. S5C). Together, these results strongly suggest that DYRK2 has a role in the HSF1-mediated resistance to proteotoxic stress.

6- DYRK2 levels correlate with HSF1 nuclear levels, and associate with prognosis and tumour recurrence in tissue from TNBC patients

Next, to address if DYRK2 regulates HSF1 *in vivo*, we investigated whether in human TNBC tumours, the protein levels of DYRK2 correlate with levels of nuclear HSF1, which is a surrogate for active HSF1. To answer this question, we analysed by immunohistochemistry (IHC) the levels and localisation of both DYRK2 and HSF1 in 120 samples from TNBC patients (antibodies validation in figure S6A). When samples were divided into high and low expression groups for either HSF1 or DYRK2 (H-score: < or >145 for cytoplasmic and nuclear DYRK2; < or > 156 for cytoplasmic HSF1; and < or > 100 for nuclear HSF1), nuclear HSF1 positively correlated with nuclear DYRK2, but not cytoplasmic DYRK2 (Fig. 6A and S6B). Interestingly, when cytoplasmic HSF1 was analysed, no association was observed with DYRK2. Our data further support the conclusion that DYRK2 positively regulates HSF1 nuclear levels in tissue from TNBC patients.

Based on this novel correlation between DYRK2 and HSF1 in TNBC human samples and on the known association of nuclear HSF1 with poor prognosis, we hypothesised that high levels of nuclear DYRK2 would associate with poor prognosis in TNBC. To test this hypothesis, we utilised a cohort of 750 breast cancer samples that had available a previously constructed tissue microarray, including 148 TNBC (TMA information in Fig. S6C). Human tumour samples were stratified on the basis of high or low H-score (H-score: low<145 WHUs and high>145 WHUs). DYRK2 protein levels were assessed in 3 individuate TMA cores and mean WHU employed for analysis (representative figures with high and low DYRK2 levels in Fig. S6D). In the full cohort, DYRK2 levels did not associate with survival (Fig. S6E). However, in TNBC samples, high nuclear DYRK2 levels significantly associated with reduced cancer-specific survival (Fig. 6B) but not with overall survival (Fig. S6F). Notably, the strength of this

association was highest in the TNBC subgroup of patients with ER, PR, HER2, and AR negative disease (Fig. 6C). In TN-AR negative samples, nuclear DYRK2 was observed as an independent factor in Cox Regression multivariate analysis when combined with other clinical parameters (Fig. S6G). Moreover, high DYRK2 levels were associated with shorter time to recurrence in both TNBC and TN-AR negative breast cancer (Fig. S6H).

These results establish DYRK2 protein levels as potential prognostic factor and promising novel therapeutic target in TNBC, especially in the TN/AR-negative subgroup of patients, for whom there is no targeted therapy available. Interestingly, several of the commonly used TNBC cell models (including the two cell lines used in this study, MDA-MB-231 and MDA-MB-468) are also AR negative[23], and thus there is full agreement between our cell culture and tissue analysis data.

7- The DYRK2-HSF1 axis promotes tumour growth in TNBC xenograft models.

Based on our cell culture and tissue data, we wondered whether targeting this newly identified DYRK2-HSF1 link could affect tumour growth *in vivo.* To answer this question, and as there are no specific DYRK2 inhibitors available, we evaluated the tumour formation capacity of WT and HSF1-KO TNBC cells after DYRK2 knockdown. Tumours derived from TNBC DYRK2-deficient cells had significantly slower growth rates and smaller weight than those formed by DYRK2-proficient cells (Fig. 7A and 7B), confirming the role of DYRK2 in promoting TNBC tumour growth *in vivo*. Similarly, tumours derived from TNBC HSF1-KO cells grew more slowly and were lighter than those derived from WT cells. In this case, the effect of knocking down DYRK2 was less pronounced than in tumours from WT cells (Fig. 7a and 7B). Notably, the fact that tumours derived from TNBC HSF1-KO cells grew much more slowly than those from WT cells, made it difficult to unambiguously conclude that DYRK2 depletion does not

affect tumour growth in HSF1-KO cells. Nonetheless, these experiments illustrate the importance of both DYRK2 and HSF1 for TNBC tumour growth and further show that, although HSF1 might not be the only mediator of the tumour-promoting activity of DYRK2, this kinase plays a major role in the growth of HSF1-proficient tumours. Overall, our data support the potential biological importance of the DYRK2-HSF1 axis in regulating cancer cell growth *in vivo*.

Discussion

The HSF1 pathway represents an attractive therapeutic target as it plays roles in both cancer progression and chemoresistance[5, 11, 16, 24, 25]. However, developing specific HSF1 inhibitors has proven to be challenging, and thus, an alternative approach is to understand and to target HSF1 upstream regulatory pathways. In this study, we shed light on the regulation of the HSF1 pathway by the kinase DYRK2. Other kinases have been shown to activate HSF1 by phosphorylating the same residues as DYRK2 (i.e. MEK1, mTOR and p38 MAPK phosphorylate S326[10, 26, 27] and PKA phosphorylates S320[8]), and thus we did not expect DYRK2 depletion to completely abolish HSF1 phosphorylation. Remarkably, DYRK2 depletion significantly reduced HSF1 phosphorylation and its nuclear levels, and was sufficient to impair HSF1 response and promote TNBC apoptosis. This finding suggests that either these other kinases are not relevant in TNBC cells, or they cannot completely compensate for the absence of DYRK2. The latter could be either because DYRK2 is necessary for a full HSF1 activation, or because the combined/sequential effect of multiple kinases is necessary, and thus elimination of any of the individual kinases involved would impair HSF1 response. Further studies are necessary to clarify this point, but nevertheless, our results indicate that DYRK2 is a major kinase controlling HSF1 activation in TNBC. Also, it is important to point out that DYRK2 is, to date,

the only known kinase that phosphorylates HSF1 on both S320 and S326, which are associated with its activation. Interestingly, DYRK2 can phosphorylate HSF1 on at least four sites *in vitro*, and the mutagenesis studies show that only when all four sites are mutated, the effect of DYRK2 on HSF1 stability is lost, suggesting that those sites play a role *in vivo*. Although how these sites control HSF1 stability is not known, the fact that DYRK2 might phosphorylate sites associated with both positive (S326, S320) and negative (S363) regulation of HSF1 might appear counter-intuitive. We propose that this conundrum could be explained by the current view that phosphorylation of S326 dominates over any other phosphorylation events. Thus, during heat shock, HSF1 undergoes both activating and inhibitory phosphorylation events, but as long as S326 is phosphorylated, the overall effect is HSF1 activation. Once S326 is dephosphorylated, the inhibitory phosphorylation events ensure that the HSF1 response is transient and does not persist.

We have shown that DYRK2 activates the pro-survival HSF1 pathway providing a support mechanism and a survival advantage to cancer cells. Importantly, this new link between DYRK2 and HSF1 appears to be relevant in TNBC patients as DYRK2 levels correlate positively with HSF1 nuclear levels, and negatively associates with cancer specific survival and time to recurrence, supporting a pro-tumoural role for DYRK2 in TNBC. Furthermore, we showed that DYRK2 predominantly affects nuclear HSF1, which is consistent with the observed increase in nuclear levels of DYRK2 in response to proteotoxic stress (Fig. 3A, 3B, S3A, S3B, S3C). This observation in cell lines, combined with the positive correlation between levels of nuclear DYRK2, nuclear HSF1 and association with patient prognosis observed in tumour samples suggest that the nuclear pool of DYRK2 is responsible for its HSF1-dependent tumour promoter role in TNBC.

Our results suggest that DYRK2 may be both a prognostic biomarker and a therapeutic target, which will be especially relevant in TN-AR negative (or quadruple negative) breast cancer patients, for whom there is no targeted therapy available. The development of new specific DYRK2 inhibitors and their validation *in vivo* (i.e. using xenograft and genetic models) is needed to further substantiate this hypothesis. In this context, it is encouraging to see in our study the dramatic inhibitory effect of the DYRK inhibitor harmine on HSF1 activation and *HSP70* expression. Harmine has been shown to delay tumour growth in a genetic model of pancreatic carcinogenesis[28], and although this drug inhibits various members of the DYRK family, our results support the relevance of the DYRK family as potential drug targets to impair cancer cell survival. In agreement, the drug curcumin inhibits DYRK2 activity and reduces TNBC tumour growth[29], further highlighting the importance of developing new specific DYRK2 inhibitors which might have potential clinical relevance.

Our data, together with previous reports showing that DYRK2 controls proteasome activity[30], presents DYRK2 as an apical upstream positive regulator of the two main pathways involved in alleviation of proteotoxic stress in cancer cells. Thus, DYRK2 might be an excellent target to tackle cancer cells dependent on such support mechanism. Based on this, combinatory therapies using inhibitors of either the proteasome or the HSF1 pathway together with DYRK2 inhibitors, are expected to have enhanced efficacy against TNBC.

Acknowledgments

We would like to thank the Glasgow Biorepository for providing the tissue. We are extremely grateful to the Medical Research Institute of the University of Dundee, Cancer Research UK

(C52419/A22869) (LDLV) and (CRUK/A18644) (ADK), Mary Kay Ash Breast Cancer grant 047.16 (JED) and UCSD Moores Cancer Centre Translational and Clinical Pilot Project award 2018 (JED) for financial support.

Author Contributions

Conceptualization, R.M., A.D-K., and L.V.; Formal Analysis, J.E.; Investigation, R.M., S.B., A.J., GB and J.Q.; Resources, J.E.D. and J.E.; Writing – Original Draft, L.V.; Writing – Review & Editing, R.M., S.B., J.E., A.D-K. and L.V.; Visualization, R.M.; Funding Acquisition, J.E.D., J.E. and L.V.

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. DYRK2 phosphorylates HSF1. A) 293T cells were transiently transfected to express GFP-tagged DYRK2 wild-type (WT) or a kinase dead (KD) version. After 48 hours, cells were lysed and the levels of endogenous HSF1 and phospho-HSF1 were analysed as indicated. **B)** 293T cells were transiently transfected with a GFP-tagged DYRK2 analog sensitive (AS) version. After 48 hours, cells were treated for a further 3 hours with increasing concentrations of three different PP1 inhibitors as indicated. Cells were lysed and the levels of endogenous HSF1 and phospho-HSF1 were analysed by western blot. **C)** *Upper panel*, purified recombinant His-HSF1 (1 µg) was incubated in kinase buffer with increasing concentrations of recombinant GST-DYRK2 at 30°C for 30 min. *Lower panel*, purified recombinant His-HSF1 (1 µg) was incubated with GST-DYRK2 (20 ng) at 30°C for various times as indicated. The reactions were

terminated by the addition of SDS gel loading buffer, the proteins were resolved by SDS-PAGE, and the levels of phosphorylated HSF1 were analysed. **D)** MDA-MB-231 or MDA-MB-468 cells were transfected with either siControl or siDYRK2. After 48 hours, cells were either incubated at 42 °C for one hour or kept at 37 °C. Cells were lysed in SDS buffer and the levels of the indicated proteins were analysed by western blotting. **E)** MDA-MB-231 or MDA-MB-468 cells were transfected with either siControl, siDYRK1A, siDYRK1B or siDYRK2. 48 hours later cells were incubated at 42 °C for one hour. Cells were lysed in SDS buffer and the levels of the indicated proteins were analysed by western blotting. **F)** MDA-MB-231 cells were treated with increasing concentrations of harmine. One hour later, cells were incubated at 42 °C. After one hour, cells were lysed in SDS buffer and the levels of the indicated proteins were analysed by western blotting. See also Figure S1.

Figure 2. DYRK2 interacts with HSF1 via two domains. A) 293T cells were transfected with the indicated plasmids. After 48 hours, cells were incubated at 42 °C for the indicated periods of time. A fraction of the cell lysates was tested for the correct expression of the transfected proteins (Input), while the remaining extracts were used for immunoprecipitation with anti-HA antibodies. After elution of bound proteins in 1 x SDS sample buffer, coprecipitated HSF1 was visualized by immunoblotting using an anti-GFP antibody as shown. **B)** A peptide array library covering the complete sequence of DYRK2 was incubated with recombinant GST-HSF1 protein (*upper panel*) or a GST control protein (*lower panel*) and the bound HSF1 protein was revealed by immunoblotting against GST as shown. The specific positive binding regions for HSF1 were indicated with a black box. Sequences of DYRK2 peptides within the binding regions (BR1 and BR2) interacting with HSF1 are shown in bold. **C)** 293T cells were transiently transfected with the GFP-tagged versions of either DYRK2-WT or DYRK2 constructs harbouring mutations on the BR1 or BR2 as indicated. After 48 hours, cells were lysed and the

levels of endogenous HSF1 and phospho-HSF1 were analysed as indicated. **D)** 293T cells were transiently transfected with the Flag-tagged versions of either DYRK2-WT or a DYRK2 mutant with mutations in both regions (DYRK2 YV/AR-T/A) together with an inactive form of SIAH2 (HA-SIAH2-RM). After 48 hours, cells were lysed and the levels of SIAH2 and phospho-SIAH2 were analysed as indicated. **E)** 293T cells were transfected with the indicated plasmids. After 48 hours, cells were incubated at 42 °C for the indicated periods of time. A fraction of the cell lysates was tested for the correct expression of the transfected proteins (input), while the remaining extracts were used for immunoprecipitation with anti-GFP antibodies. After elution of bound proteins in 1 x SDS sample buffer, coprecipitated HSF1 was visualized by immunoblotting using an anti-Flag antibody as shown. See also Figure S2

Figure 3. DYRK2 promotes HSF1 nuclear stability. A) Control (WT) and CRISPR-mediated DYRK2-KO MDA-MB-231 cells were incubated at 37 °C or at 42 °C for the indicated times. Nuclear and cytoplasmic fractions were analysed by western blot for the levels of the indicated proteins (L for long-exposure; S for short-exposure). **B)** DYRK2-KO 293T were transfected with either empty vector or with GFP-DYRK2. Control (WT), DYRK2-KO and DYRK2-GFP reconstituted DYRK2-KO cells were incubated at 37 °C or 42 °C for the indicated times. Cells were lysed, and nuclear fractions were analysed for the levels of the indicated proteins by western blotting. **C)** MDA-MB-231 cells were treated with either vehicle or 8 μM of Harmine. Two hours later, cells were incubated at 37 °C or 42 °C as indicated. After that, cells were lysed and nuclear and cytoplasmic fractions were analysed by western blotting (L for long-exposure; S for short-exposure). **D)** MDA-MB-231 cells were transfected with either siControl or siDYRK2, and 48 hours later cells were incubated with 20 μM of cycloheximide for the indicated times. Nuclear and cytosolic fractions were analysed for the levels of the indicated proteins by western blotting. **E)** 293T cells were transfected with either Flag-tagged

HSF1 or with Flag-tagged HSF1 mutated in S320A, T323I, S326A, S363A (HSF1-4mut) in combination with siControl or siDYRK2 as indicated. After 48 hours, cells were exposed to 42 °C. After one hour, cells were lysed and nuclear and cytosolic fractions were analysed for the levels of the indicated proteins by western blotting. See also Figure S3

Figure 4. DYRK2 affects the expression levels of the HSF1 target gene *HSP70*. A) *Upper panel,* control (WT), DYRK2-KO, HSF1-KO or HSF1/DYRK2-KO MDA-MB-231 cells. *Lower panel,* control (WT), DYRK2-KO, HSF1-KO or HSF1/DYRK2-KO MDA-MB-468 cells. In both cases, cells were incubated at 37 °C or 42 °C for one hour. The mRNA levels for *HSP70* (*HSPA1A*) were quantified using real-time PCR. The data were normalized using β-actin as an internal control. Data represent means ± SD (n=3) and are expressed relative to the control sample levels at 37 °C. *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001, ****P≤ 0.0001. **B)** Control (WT) and DYRK2-KO MDA-MB-468 cells were incubated at 37 °C, at 42 °C for 5, 15, 30 or 60 minutes, or at 42 °C for 60 minutes plus recovery time at 37 °C (up to 180 minutes). After that, mRNA levels for *HSP70* (*HSPA1A*) were quantified using real-time PCR. The data were normalized using β-actin as an internal control. Data represent means ± SD (n=3) and are expressed relative to the control samples at 37 °C. **C)** DYRK2-KO MDA-MB-231 were transfected with either empty vector or with GFP-DYRK2. Control (WT), DYRK2-KO and DYRK2-GFP reconstituted DYRK2-KO cells were incubated at 37 °C or 42 °C for the indicated times. After that, mRNA levels for HSP70 (HSPA1A) were quantified using real-time PCR. The data were normalized using β-actin as an internal control. Data represent means ± SD (n=3) and are expressed relative to the control samples at 37 °C. *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001, ****P≤ 0.0001. See also Figure S4

Figure 5. DYRK2 reduces sensitivity to proteotoxic stress via HSF1. A) Control (WT) and DYRK2-KO MDA-MB-231 cells were incubated at 37 °C (-) or at 45 °C for the indicated times,

followed by recovery at 37 °C. On the next day, cells were lysed and the levels of apoptosis were analysed by western blotting using an antibody that recognises cleaved PARP. **B and C)** MDA-MB-231 (B) or HSF1-KO MDA-MB-231 (C) cells transfected with either siControl or siDYRK2 were incubated at 37 °C or at 45 °C for the indicated times, followed by recovery at 37 °C. On the next day, cells were lysed and the levels of apoptosis were analysed by western blotting using an antibody that recognises cleaved parp. See also Figure S5

Figure 6. DYRK2 levels correlate with HSF1 nuclear levels and associates with poor prognosis and tumour recurrence in tissue from TNBC patients. A) Chi Square test correlations between nuclear/cytoplasmic HSF1, and nuclear/cytoplasmic DYRK2 in triple negative breast cancer. Significance p<0.05. **B)** Relationship between cytoplasmic, nuclear and total DYRK2 levels and cancer-specific survival in patients with triple negative invasive ductal breast cancer. "Low expression both" or "High expression both" means samples with low or high levels of both nuclear and cytoplasmic DYRK2; "High expression one" means samples with high levels of either nuclear or cytoplasmic DYRK2; "Low expression both/one high" means all the samples that do not have high levels of nuclear and cytoplasmic DYRK2. **C)** Relationship between cytoplasmic and nuclear DYRK2 levels and cancer-specific survival in patients with triple negative and AR negative invasive ductal breast cancer. See also Figure S6

Figure 7. The DYRK2-HSF1 axis promotes tumour growth in TNBC xenograft models. A) DYRK2 was knocked down in control or HSF1-KO MDA-MB-231 cells using two independent shRNAs targeting DYRK2, along with a scrambled shRNA control. The cells were then injected subcutaneously into NSG™ mice. Tumour volume was measured twice a week (n=8 per condition). **B)** Post 30 days of injection, tumours were resected, imaged and tumour weight was measured. ****p<0.0001, ***p<0.001, *p<0.05 (compared to scrambled control, two-way ANOVA, mean ± SD from n=8 mice each).

Material and Methods

Cell culture

All cell lines used in the study have been validated by STR profiling and were routinely tested

for mycoplasma. All cell lines were grown in DMEM containing 10% FBS at 37°C and 5% CO2.

Antibodies, plasmids and reagents

Antibodies recognizing Flag (M2) were obtained from Sigma-Aldrich (Dorset, UK); anti-HA (Y-

11), anti-Lamin B2 (C-20) and anti-Tubulin (TU-02) were obtained from Santa Cruz

Biotechnology (Dallas, Texas, USA). Anti-GFP was obtained from the MRC Protein

Phosphorylation and Ubiquitination Unit, (School of Life Science, University of Dundee, UK);

anti-cleaved PARP (19F4) was obtained from Cell Signalling (MA, USA) and anti-DYRK2 was

also obtained from both Cell Signalling and from the Protein Phosphorylation Unit (School of

Life Science, University of Dundee, UK); anti-HSF1 was obtained from Enzo Life Science (NY,

USA); anti-phospo-HSF1 (ser 326), anti-phospo-HSF1 (ser 320) and anti GST-HRP were

obtained from Abcam (Cambridge, UK). HRP-conjugated secondary antibodies were obtained

from Life Technologies (Carlsbad, California, USA). Dynabeads™ Protein G was obtained from

Thermo scientific (Waltham, MA USA).

Expression vectors for DYRK2-GFP, DYRK2-KD-GFP, DYRK2-Flag and SIAH2-RM-HA were a gift

from Marco A. Calzado (University of Cordoba, Spain); HSF1-GFP have been already

described[27]. All gRNAS were cloned into pLentiCRISPr-V2, which was a gift from Feng Zhang

(Addgene plasmid # 52961). Point mutants were produced by conventional point

mutagenesis. The DYRK2-analog sensitive was obtained by mutating the gatekeeper residue

F228 to Alanine. His-HSF1 has already been described[27] and GST-DYRK2 was obtained from

the MRC Protein Phosphorylation and Ubiquitination Unit (School of Life Science, University

of Dundee, UK). GFP-DYRK2 was cloned into the lentiviral 290-puro plasmid[31] obtaining a puro

resistant construct that was used for reconstitution experiments.

The siRNAs used against DYRK2 were the SMART pool: ON-Target Plus from Dharmacon (CO,

USA). The shRNAs used to target DYRK2 (shDYRK2-1 gggtagaagcggtattaaa and shDYRK2-2

ggagaaaacgtcagtgaaa) were expressed from the pLL3.7 vector

Bortezomib was obtained from Santa Cruz Biotechnology. DMSO was obtained from SIGMA-

Aldrich (MA, USA), Harmine was obtained from Tocris Bioscience (Bristol, UK) and the PP1

inhibitors were obtained from Cayman Chemicals (MI, USA).

Quantitative real time PCR (rt-qPCR)

RNA was extracted using RNeasy kit (QIAGEN). 500 ng of RNA per sample was reverse-

transcribed to cDNA using Omniscript RT kit (QIAGEN) supplemented with RNase inhibitor

(QIAGEN) according to the manufacturer's instructions. Resulting cDNA was analysed using

TaqMan Universal Master Mix II (Life technologies). Gene expression was determined using

an Applied Biosystems 7300 Real-Time PCR system by the comparative ΔΔCT method. All

experiments were performed at least in triplicates and data were normalized to the

housekeeping gene β-actin. Probes were obtained from Applied Biosystems. When

applicable, the differences between groups were determined by 2 way ANOVA. Analyses were

performed using GraphPad Prism (GraphPad Software); a P value of <0.05 was considered

significant. $*P \leq 0.05$, $**P \leq 0.01$, $***P \leq 0.001$.

Cell transfections

On the day prior to transfection, cells were plated to the required cell density (70-90%

confluency). Lipofectamine 2000 and Lipofectamine RNAiMAX (Invitrogen) were used for

plasmid DNA and siRNA respectively. The plasmid DNA/siRNA and lipofectamine were

individually diluted in Optimem (Gibco) and incubated for 10 minutes at room temperature.

Diluted DNA/siRNA was added to the diluted Lipofectamine solution (1:1 ratio) and further incubated for 15 minutes. DNA-lipid complex was added to the cells and incubated overnight in a humidified incubator at 37°C and 5% CO2. The next morning, the medium was replaced with fresh medium and cells were incubated 36 hours more prior lysis.

Lentivirus production

293T cells were transfected using Lipofectamine 2000 (Invitrogen) with the empty vector (pLL3.7) or the lentiviral shDYRK2-1 or shDYRK2-2 together with the packaging vectors (psPAX2 and pMD2.G) and cultivated in OptiMEM medium (Invitrogen). The next day the cells were further grown in DMEM complete medium and one day later the lentivirus-containing supernatant was collected, filtered and used to transduce cells.

Cell lysis protocol and western blotting

Cells were washed and harvested in ice-cold phosphate-buffered saline (PBS) and lysed in either SDS buffer or RIPA buffer [50 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 mM EDTA, 1% NP40, 0.5% sodium deoxycholate, 0.5 mM Na3VO4, 50 mM NaF, 2 µg/ml leupeptine, 2 µg/ml aprotinin, 0.05mM pefabloc]. Cells directly lysed in SDS were boiled for two minutes, sonicated and boiled again for another 5 minutes. Cells lysed in RIPA buffer were sonicated and lysates were cleared by centrifugation for 15 minutes at 4°C. Protein concentration was established using the BCA assay (Pierce). Supernatant was mixed with SDS sample buffer and boiled for 5 minutes. Equal amounts of protein were separated by SDS-PAGE, followed by semidry blotting to a polyvinylidene difluoride membrane (Thermo Scientific). After blocking of the membrane with 5% (w/v) TBST nonfat dry milk, primary antibodies were added. Appropriate secondary antibodies coupled to horseradish peroxidase were detected by enhanced chemiluminescence using Clarity™ Western ECL Blotting Substrate (BIO-Rad).

Subcellular fractionation (nuclear/cytoplasmic separation)

Cells were washed and harvested with ice-cold PBS. Pelleted cells were resuspended in 400µl of low-salt buffer A (10mM Hepes/KOH pH7.9, 10mM KCL, 0.1mM EDTA, 0.1mM EGTA, 1mM β-Mercaptoethanol). After incubation for 10 minutes on ice, 10µl of 10% NP-40 was added and cells were lysed by gently vortexing. The homogenate was centrifuged for 10 seconds at 13,200 rpm in a microfuge. The supernatant representing the cytoplasmic fraction was collected and the pellet containing the cell nuclei was washed 4 additional times in buffer A, then resuspended in 100µl high-salt buffer B (20mM Hepes/KOH pH7.9, 400mM NaCL, 1mM EDTA, 1mM EGTA, 1mM β-Mercaptoethanol). The lysates were sonicated and centrifuged at 4°C for 15 minutes at 13,200 rpm. The supernatant representing the nuclear fraction was collected. Protease and phosphatase inhibitors were freshly added to both buffers.

Immunoprecipitation

Cells were washed with PBS and lysed in IP buffer (50 mM Hepes pH 7.5, 50 mM NaCl, 1% (v/v) Triton X-100, 2 mM EDTA, 10 mM sodium fluoride, 0.5 mM sodium orthovanadate, leupeptine (10 µg/ml), aprotinin (10 µg/ml), and 1 mM PMSF), followed by a sonication step. The extract was centrifuged and the supernatant was transferred to a new tube. The immunoprecipitation was performed with 2 µg of precipitating antibodies together with 50 µl of Dynabeads™ Protein G. Tubes were rotated for 30 min on a spinning wheel at 4°C. The immunoprecipitates were washed 3x with PBS/0,01% Tween-20 and eluted by boiling in 1 x SDS sample buffer. Equal amounts of protein were separated by SDS-PAGE.

CRISPR-edited cell lines

The endogenous DYRK2 and/or HSF1 genes were knocked-out by transfecting cells with pLentiCRISPR-v2 (which codes for Cas9, and a puromycin cassette) containing gRNAs against the first exon of the short DYRK2 isoform or against the fourth exon of HSF1. For MDA-MB-231, HeLa and 293T DYRK2-KO cells the gRNA sequence used was GCTTGCCAGTGGTGCCAGAG

and for MDA-MB-468 DYRK2-KO cells the target sequence was CGCTCACGGACAGATCCAGG. Additionally, we also tested some of our results in MDA-MB-231 cells in which we almost completely remove the DYRK2 ORF by using two gRNAs (N-term sequence GCTTGCCAGTGGTGCCAGAG and C-term sequence GAAGCTGAGCTAGAAGGTGG). For HSF1-KO cells the gRNA sequence used was AAGTACTTCAAGCACAACAA. Control cells were transfected with the empty pLentiCRISPRV2 vector. After transfection, cells were exposed to 2 µg/ml of puromycin for two days followed by a medium exchange. Surviving cells were clonally selected (in the case of control cells were used as pool population) by serial dilution, and positive clones were identified by genomic analysis and western blot. At least two clones for each cell line were used for the experiments.

In vitro kinase assay

Bacterially expressed full length HSF1 His tagged was incubated with GST-DYRK2 in Kinase Buffer (25 mM TRIS pH 7.5, 5 mM β-glycerophosphate, 10 mM MgCl2 and freshly added 2mM DTT, 0.1 mM Na3VO4, 1 mM ATP) for different periods of time (5-60 min) at 30 °C. The reaction was inactivated by adding SDS loading buffer (50 mM Tris-HCl pH 6.8, 10%, SDS, 40% glycerin, 15% β-mercaptoethanol, 0.1 % bromophenol blue) and samples were boiled at 95 °C for 5 minutes and loaded in SDS-PAGE gels.

In vitro peptide binding

Overlapping dodecapeptides covering the entire DYRK2 protein were spotted in an automated process on cellulose membranes by using Fmoc-protection chemistry (Proteomics facility, CNB-CSIC, Spain). The membrane was blocked overnight in non-fat milk in TBS buffer and incubated for 6 h with 60 nmol of recombinant GST-HSF1 or GST proteins as a control. After extensive washing, anti-GST antibody coupled to horseradish peroxidase was added for

1 hour. After washing, blots were revealed by enhanced chemiluminescence using Clarity™ Western ECL Blotting Substrate (BIO-Rad).

Mass Spectrometry

Purified His-HSF1 (1ug) was incubated with GST-DYRK2 (20 ng) in 20 uL buffer containing 25mM Tris-HCl (pH7.5), 5 mM MgCl2, and 1 mM ATP for 5 minutes. The mixtures were reduced, alkylated and digested with trypsin (1:100) prepared in 100mM TEAB buffer (pH 8.5). The mixture was incubated at 37°C overnight before 8 uL of 10mM EDTA (in 0.1% TFA) was added. The peptide mixture was then subjected to nano-LC-MS/MS analysis according to a previous report[32]. Database search was carried out using Peaks 7.0 against UniProt-human (version 2018-05-28). False discovery rate was set at 1% (for peptide spectrum matches). The non-modified and modified (methionine oxidation and phosphorylation) peptides from HSF1 detected are shown in blue lines. The putative oxidation sites are labelled in yellow and phosphorylation sites in red.

Xenograft assays

Female NSG™ or Homozygous Foxn1nu (J:NU 007850) (Jackson Labs, USA) mice were housed and maintained at the University of California-San Diego (UCSD) in full compliance with policies of the Institutional Animal Core and Use Committee (IACUC). DYRK2 was knocked down in MDA-MB-231 (parental or HSF1 KO) cells with two independent shRNA oligoes as described previously[30]. MDA-MB-231 parental or HSF1 KO with either scrambled control or DYRK2 knock-down shRNA (1 or 2) were counted and suspended in sterile phosphate buffer saline containing 50% Matrigel (BD Bioscience). 300,000 cells (100 µl) were injected subcutaneously into the neck of each mouse. Tumour dimensions were measured twice per week using a digital caliper and tumour volume was calculated as (length × width^2)/2. After

the indicated days post-injection, mice were euthanized and tumours were excised and weighed.

Eight mice per group was the standard sample size for tumour xenograft experiments as defined by statistical power analyses (80-90% power; p<0.05) carried out using R packages. The investigators were not blinded to allocation during experiments and outcome assessment. Data were analysed using Graphpad Prism statistical package. All results are presented as mean ± SD unless otherwise mentioned

Patient cohort and Tissue microarrays

Patients presenting with invasive ductal breast cancer at Glasgow Royal Infirmary, Western Infirmary and Stobhill Hospital between 1995 and 1998, with formalin-fixed paraffin embedded blocks of the primary tumour available for evaluation were studied (n =850). The study was approved by the Research Ethics Committee of the West Glasgow University Hospitals NHS Trust. Clinicopathological data including age, tumour size, tumour grade, lymph node status, type of surgery and use of adjuvant treatment (chemotherapy, hormonal based therapy and/ or radiotherapy) were retrieved from the routine reports. Tumour grade was assigned according to the Nottingham Grading System. ER and PR status were assessed on tissue microarrays (TMAs) using immunohistochemistry (IHC) with Dako (Glostrup, Denmark) ER antibody and Leica (Wetzlar, Germany) PR antibody. Her2 status were assessed as described[33].

Tissue analysis

IHC was conducted in triplicate on tissue microarrays (TMAs). Prior to this, antibody specificity was confirmed using proficient and deficient cell lines (either knockdown or knockout cell lines) (Fig S6A). TMA sections (2.5μm) were dewaxed by immersion in histoclear then rehydrated using a series of alcohols. Heat induced antigen retrieval was performed in

a solution of either Tris-EDTA pH9 (DYRK2) or citrate buffer pH6 (HSF1) after which the sections were incubated in 3% hydrogen peroxide. Non-specific binding was blocked by incubation in 1.5% normal horse serum before being incubated with the optimum concentration of primary antibody overnight at 4°C. Antibody dilutions were prepared in antibody diluent (Agilent, London, UK). The primary antibodies and concentrations used are as follows; DYRK2 (Abgent AP7534a) at 1:200 and HSF1 (Cell Signaling 4256) at 1:500. Staining was visualized using ImmPRESS™ and ImmPACT ™DAB, (both Vector Labs). Tissue was counterstained using Harris Haematoxylin before being dehydrated and mounted using DPX. The stained TMA sections were scanned using a Hamamatsu NanoZoomer (Welwyn Garden City, Hertfordshire, UK) at x20 magnification. Visualization was carried out using Slidepath Digital Image Hub, version 4.0.1 (Slidepath, Leica Biosystems, Milton Keynes, UK). The weighted histoscore method was employed to assess protein expression [GB], with a second independent observer [JE] scoring 10% of the scores. Histoscores were calculated using the following formula: 0x % negative staining + 1x % of weakly positive staining + 2x % of moderately positive staining +3x % of strongly positive staining giving a range of 0 (minimum) to 300 (maximum). The results were considered discordant if the histoscores differed by more than 50 weighted histoscore units (WHU). Cytoplasmic and nuclear expression, if observed, was calculated separately.

Supplementary Figure legends

Figure S1. Related to Figure 1. DYRK2 phosphorylates HSF1. A) 293T cells were transiently transfected with empty vector or Flag-tagged DYRK2 as indicated. After 48 hours, cells were treated for a further 3 hours with vehicle or harmine (8 μM). Cells were lysed and the levels of endogenous HSF1 and phospho-HSF1 were analysed as indicated. **B)** 20 ng of recombinant

GST-DYRK2-WT (*left panel*) or GST-DYRK2-AS (*right panel*) were incubated with His-HSF1 (1

µg) and increasing concentrations of 1NM-PP1 in kinase buffer without ATP. After 5 minutes,

ATP was added and the reaction incubated at 30°C for 30 min. The reactions were terminated

by the addition of SDS gel loading buffer, the proteins were resolved by SDS-PAGE, and the

levels of phosphorylated HSF1 were analysed. **C)** Nano-LC-MS/MS analysis of His-HSF1

incubated with GST-DYRK2 for 5 minutes. Database search was carried out using Peaks 7.0

against UniProt-human(version 2018-05-28). False discovery rate was set at 1% (for peptide

spectrum matches). The non-modified and modified (methionine oxidation and

phosphorylation) peptides from HSF1 detected are shown in blue lines. The putative oxidation

sites are labled in yellow and phosphorylation sites in red. **D)** HeLa cells were transfected with

either siControl or siRNAs against all members of the DYRK family. 48 hours later cells were

incubated at 37 °C or 42 °C for one hour. Cells were lysed in SDS buffer and the levels of the

indicated proteins were analysed by western blotting.

Figure S2. Related to Figure 2. DYRK2 interacts with HSF1 via two domains. A) Species

alignment for the DYRK2 sequence containing the two identified binding regions, BR1 and BR2

(underlined). Asterisks mark conserved residues. **B)** Representation of the 3D structure of

DYRK2 obtained for UniProt. The square focused on the DDQGSYV region and shows in blue

the position of the different residues.

Figure S3. Related to Figure 3. DYRK2 promotes HSF1 nuclear stability. A) Control (WT) and

CRISPR-mediated DYRK2-KO MDA-MB-468 cells were incubated at 37 °C or at 42 °C for the

indicated times. Nuclear and cytosolic fractions were analysed by western blot for the levels

of the indicated proteins (L for long exposure; S for short exposure). **B, C and D)** MDA-MB-

468 cells (B), HeLa cells (C) or 293T cells (D) were transfected with either siControl or siDYRK2.

48 hours later cells were incubated at 37 °C or at 42 °C for the indicated times. Nuclear

fractions were analysed by western blot for the levels of the indicated proteins. **E)** 293T cells were transfected with either empty vector or GFP-DYRK2. After 48 hours, cells were incubated at 37 °C or 42 °C for the indicated periods of time. Nuclear fractions were analysed by western blot for the levels of the indicated proteins. **F)** 293T cells were transfected with either Flag-tagged HSF1 or with the indicated Flag-tagged mutant HSF1 in combination with siControl or siDYRK2. After 48 hours, cells were exposed to 42 °C. After one hour, cells were lysed and analysed for the levels of the indicated proteins by western blotting

Figure S4. Related to Figure 4. DYRK2 affects the expression levels of the HSF1 target gene *HSP70*. A) Control (WT) and DYRK2-KO MDA-MB-231 cells (*left panel*), or Control (WT) and DYRK2-KO MDA-MB-468 cells (*right panel*), were incubated with either vehicle or 20nM Bortezomib (Brtz) for four hours. The mRNA levels for *HSP70* (*HSPA1A*) were quantified using real-time PCR. The data were normalized using β-actin as an internal control. Data represent means ± SD (n=3) and are expressed relative to the control samples treated with vehicle. *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001, ****P≤ 0.0001. **B)** Control (WT) or DYRK2-KO MDA-MB-231 cells (*left panel*), or MDA-MB-468 cells (*right panel*), or 293T cells (*lower panel*) were incubated with increasing concentrations of harmine. After 2 hours, the mRNA levels for *HSP70* (*HSPA1A*) were quantified using real-time PCR. The data were normalized using β-actin as an internal control. Data represent means ± SD (n=3) and are expressed relative to the control samples treated with vehicle. *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001, ****P≤ 0.0001.

Figure S5. Related to Figure 5. DYRK2 reduces sensitivity to proteotoxic stress via HSF1. A and B) Control (WT) and DYRK2-KO MDA-MB-468 cells (A) or 293T cells (B) were incubated at 37 °C or at 45 °C for the indicated times, followed by recovery at 37 °C. On the next day, cells were lysed and the levels of apoptosis were analysed by western blotting using an antibody that recognises cleaved PARP. **C)** Control (WT) (*left panel*) or HSF1-KO *(right panel)* MDA-MB-

231 cells were treated with vehicle or increasing concentrations of harmine. After two hours, cells were incubated at 37 °C or at 45 °C for the indicated times, followed by recovery at 37 °C. On the next day, cells were lysed and the levels of apoptosis were analysed by western blotting using an antibody that recognises cleaved PARP.

Figure S6. Related to Figure 6. DYRK2 levels correlate with HSF1 nuclear levels, prognosis and tumour recurrence in tissue from TNBC patients. A) DYRK2 and HSF1 antibody validation using Formalin-Fixed, Paraffin-Embedded (FFPE) cell pellets. **B)** Chi Square test correlation between low and high HSF1 and DYRK2 protein expression in triple negative breast cancer samples. Significance $p < 0.05$. **C)** Tissue microarray details. **D)** Representative samples showing high and low DYRK2 staining. **E)** Relationship between cytoplasmic, nuclear and total DYRK2 levels and cancer-specific survival in patients with invasive ductal breast cancer (total cohort). **F)** Relationship between nuclear DYRK2 levels and overall survival in patients with triple negative (left panel) or triple negative and AR negative (right panel) invasive ductal breast cancer. **G)** Univariate and multivariate survival analysis for patients with triple negative and AR negative breast cancer to assess the relationship between clinico-pathological characteristics, DYRK2 and cancer specific survival. **H)** Relationship between nuclear DYRK2 levels in tumour cells and time to recurrence in patients with triple negative (left panel) or triple negative and AR negative (right panel) invasive ductal breast cancer.

References

Bianchini G, Balko JM, Mayer IA, Sanders ME, Gianni L. Triple-negative breast cancer: challenges and opportunities of a heterogeneous disease. Nat Rev Clin Oncol 2016; 13: 674-690.

Andreopoulou E, Schweber SJ, Sparano JA, McDaid HM. Therapies for triple negative breast cancer. Expert Opin Pharmacother 2015; 16: 983-998.

Donnelly N, Storchova Z. Causes and consequences of protein folding stress in aneuploid cells. Cell Cycle 2015; 14: 495-501.

Ohashi A, Ohori M, Iwai K, Nakayama Y, Nambu T, Morishita D *et al*. Aneuploidy generates proteotoxic stress and DNA damage concurrently with p53-mediated post-mitotic apoptosis in SAC-impaired cells. Nat Commun 2015; 6: 7668.

Dai C, Whitesell L, Rogers AB, Lindquist S. Heat shock factor 1 is a powerful multifaceted modifier of carcinogenesis. Cell 2007; 130: 1005-1018.

Baler R, Dahl G, Voellmy R. Activation of human heat shock genes is accompanied by oligomerization, modification, and rapid translocation of heat shock transcription factor HSF1. Mol Cell Biol 1993; 13: 2486-2496.

Dayalan Naidu S, Dinkova-Kostova AT. Regulation of the mammalian heat shock factor 1. FEBS J 2017; 284: 1606-1627.

Murshid A, Chou SD, Prince T, Zhang Y, Bharti A, Calderwood SK. Protein kinase A binds and activates heat shock factor 1. PLoS One 2010; 5: e13830.

Guettouche T, Boellmann F, Lane WS, Voellmy R. Analysis of phosphorylation of human heat shock factor 1 in cells experiencing a stress. BMC Biochem 2005; 6: 4.

Tang Z, Dai S, He Y, Doty RA, Shultz LD, Sampson SB *et al*. MEK guards proteome stability and inhibits tumor-suppressive amyloidogenesis via HSF1. Cell 2015; 160: 729-744.

Mendillo ML, Santagata S, Koeva M, Bell GW, Hu R, Tamimi RM *et al*. HSF1 drives a transcriptional program distinct from heat shock to support highly malignant human cancers. Cell 2012; 150: 549-562.

Brandvold KR, Morimoto RI. The Chemical Biology of Molecular Chaperones--Implications for Modulation of Proteostasis. J Mol Biol 2015; 427: 2931-2947.

Doyle SM, Genest O, Wickner S. Protein rescue from aggregates by powerful molecular chaperone machines. Nat Rev Mol Cell Biol 2013; 14: 617-629.

Jolly C, Morimoto RI. Role of the heat shock response and molecular chaperones in oncogenesis and cell death. J Natl Cancer Inst 2000; 92: 1564-1572.

Rerole AL, Jego G, Garrido C. Hsp70: anti-apoptotic and tumorigenic protein. Methods Mol Biol 2011; 787: 205-230.

Santagata S, Hu R, Lin NU, Mendillo ML, Collins LC, Hankinson SE *et al*. High levels of nuclear heat-shock factor 1 (HSF1) are associated with poor prognosis in breast cancer. Proc Natl Acad Sci U S A 2011; 108: 18378-18383.

Lee P, Cho BR, Joo HS, Hahn JS. Yeast Yak1 kinase, a bridge between PKA and stress-responsive transcription factors, Hsf1 and Msn2/Msn4. Mol Microbiol 2008; 70: 882-895.

Bishop AC, Ubersax JA, Petsch DT, Matheos DP, Gray NS, Blethrow J *et al*. A chemical switch for inhibitor-sensitive alleles of any protein kinase. Nature 2000; 407: 395-401.

Gockler N, Jofre G, Papadopoulos C, Soppa U, Tejedor FJ, Becker W. Harmine specifically inhibits protein kinase DYRK1A and interferes with neurite formation. FEBS J 2009; 276: 6324-6337.

Kourtis N, Moubarak RS, Aranda-Orgilles B, Lui K, Aydin IT, Trimarchi T *et al*. FBXW7 modulates cellular stress response and metastatic potential through HSF1 post-translational modification. Nat Cell Biol 2015; 17: 322-332.

Papaleo E, Saladino G, Lambrughi M, Lindorff-Larsen K, Gervasio FL, Nussinov R. The Role of Protein Loops and Linkers in Conformational Dynamics and Allostery. Chem Rev 2016; 116: 6391-6423.

Perez M, Garcia-Limones C, Zapico I, Marina A, Schmitz ML, Munoz E *et al*. Mutual regulation between SIAH2 and DYRK2 controls hypoxic and genotoxic signaling pathways. J Mol Cell Biol 2012; 4: 316-330.

Speers C, Zhao SG, Chandler B, Liu M, Wilder-Romans K, Olsen E *et al*. Androgen receptor as a mediator and biomarker of radioresistance in triple-negative breast cancer. NPJ Breast Cancer 2017; 3: 29.

Desai S, Liu Z, Yao J, Patel N, Chen J, Wu Y *et al*. Heat shock factor 1 (HSF1) controls chemoresistance and autophagy through transcriptional regulation of autophagy-related protein 7 (ATG7). J Biol Chem 2013; 288: 9165-9176.

Shah SP, Lonial S, Boise LH. When Cancer Fights Back: Multiple Myeloma, Proteasome Inhibition, and the Heat-Shock Response. Mol Cancer Res 2015; 13: 1163-1173.

Chou SD, Prince T, Gong J, Calderwood SK. mTOR is essential for the proteotoxic stress response, HSF1 activation and heat shock protein synthesis. PLoS One 2012; 7: e39679.

Dayalan Naidu S, Sutherland C, Zhang Y, Risco A, de la Vega L, Caunt CJ *et al*. Heat Shock Factor 1 Is a Substrate for p38 Mitogen-Activated Protein Kinases. Mol Cell Biol 2016; 36: 2403-2417.

Luna J, Boni J, Cuatrecasas M, Bofill-De Ros X, Nunez-Manchon E, Gironella M *et al*. DYRK1A modulates c-MET in pancreatic ductal adenocarcinoma to drive tumour growth. Gut 2018.

Banerjee S, Ji C, Mayfield JE, Goel A, Xiao J, Dixon JE *et al*. Ancient drug curcumin impedes 26S proteasome activity by direct inhibition of dual-specificity tyrosine-regulated kinase 2. Proc Natl Acad Sci U S A 2018.

Guo X, Wang X, Wang Z, Banerjee S, Yang J, Huang L *et al*. Site-specific proteasome phosphorylation controls cell proliferation and tumorigenesis. Nat Cell Biol 2016; 18: 202-212.

de la Vega L, Grishina I, Moreno R, Kruger M, Braun T, Schmitz ML. A redox-regulated SUMO/acetylation switch of HIPK2 controls the survival threshold to oxidative stress. Mol Cell 2012; 46: 472-483.

Chakraborty P, Huang JTJ, Hiom K. DHX9 helicase promotes R-loop formation in cells with impaired RNA splicing. Nat Commun 2018; 9: 4346.

Bartlett JM, Going JJ, Mallon EA, Watters AD, Reeves JR, Stanton P *et al*. Evaluating HER2 amplification and overexpression in breast cancer. J Pathol 2001; 195: 422-428.

Fig 1

A

GFP-HSF1 + + + +
HA-DYRK2 + + + +
42 °C (min) - 5 15 30

IP: HA — GFP, HA
Input — GFP, HA, Tubulin

B

GST-HSF1

GST

BR1
C12 N-G-G-Y-D-**D-D-Q-G-S-Y-V**
C13 Y-D-**D-D-Q-G-S-Y-V**-Q-V-P

BR2
G13 K-L-R-T-N-L-A-Q-M-**T-D-A**
G14 T-N-L-A-Q-M-**T-D-A**-N-G-N
G15 A-Q-M-**T-D-A**-N-G-N-I-Q-
G16 **T-D-A**-N-G-N-I-Q-Q-R-T-V

C

GFP-DYRK2 - + - - - - - - -
GFP-DYRK2-TDA/TAA - - + - - - - - -
GFP-DYRK2-TDA/ADA - - - + - - - - -
GFP-DYRK2-Q/R - - - - + - - - -
GFP-DYRK2-DD/AA - - - - - + - - -
GFP-DYRK2-YV/AR - - - - - - + - -
GFP-DYRK2-GS/AA - - - - - - - + -
GFP-DYRK2-YV/AR-T/A - - - - - - - - +

HSF1
p-HSF1 (S326)
p-HSF1 (S320)
GFP
Tubulin

D

Flag-DYRK2 - + -
Flag-DYRK2-YV/AR-T/A - - +
HA-SIAH2-RM + + +

Flag
HA
p-SIAH2-S28
Tubulin

E

Flag-HSF1 + + + + + + +
GFP-DYRK2 - + + + - - -
GFP-DYRK2-YV/AR-T/A - - - - + + +
42 °C (min) - - 5 15 - 5 15

IP: GFP — Flag, GFP
Input — Flag, GFP, Tubulin

Fig 2

Fig 3

Fig 4

A

45°C (min)   -   5   15   30   45

Cleaved Parp
DYRK2
Tubulin

B

45°C (min)   -   5   15   30   45   60

Cleaved Parp
DYRK2
Tubulin

C

45°C (min)   -   10   20   30

Cleaved Parp
DYRK2
HSF1
Tubulin

Fig 5

A

	Nuclear HSF1	Cytoplasmic HSF1
Nuclear DYRK2	p-value= 0,034	p-value= 0,670
Cytoplasmic DYRK2	p-value= 0,177	p-value= 0,385

B

C

Fig 6

Fig 7

Fig S1

S2A

<pre>
                                  BR1   BR2
                          ** ********* ***   ** * ********
Homo sapiens   NNGGYD<u>DDQGSYV</u>QVPH   NLAQ<u>MTDA</u>NGNIQ
Pan troglodytes   NNGGYDDDQGSYVQVPH   NLAQMTDANGNIQ
Macaca mulatta   NNGGYDDDQGSYVQVPH   NLAQMTDANGNIQ
Canis lupus familiaris   NNGGYDDDQGSYVQVPH   NLAQMTDANGNIQ
Bos Taurus   NNGGYDDDQGSYVQVPH   NLAQMTDANGNIQ
Mus musculus   NNGGYDDDQGSYVQVPH   NLAQMTDANGNIQ
Rattus norvegicus   NNGGYDDDQGSYVQVPH   NLAQMTDANGNIQ
Gallus gallus   NNCGYDDDQGSYIQVPH   NLAQMTDANGNIQ
Xenopus tropicalis   NNGGYDDDQGSYVQVPH   NLTQITDANGNIQ
Danio rerio   NNGGYDDDQGSYIHVPH   NLAQMTDANGNIQ
                          * **   * *   *
Drosophila melanogaster   NNSEYDNEQGAYIHVPH
Anopheles gambiae   GNSDYDNEQGSYIHIAH
Caenorhabditis elegans   NNGGYDDENGSYQLVVH
</pre>

S2B

Fig S2

Fig S3

S4A

S4B

S5A

45°C (min)   -   5   15   30   45

Cleaved Parp

DYRK2

Tubulin

S5B

45°C (min)   -   30   45   60

Cleaved Parp

DYRK2

Tubulin

S5C

Cleaved Parp

Tubulin

MB-MDA-231 WT   MB-MDA-231 HSF1-KO

Fig S5

S6A

DYRK2 levels

MDA-MB-231 DYRK2-KO

x40

x20

MDA-MB-231 Control

x40

x20

Negative control

x20

HSF1 Levels

HeLa

Control

HSF1-KO

S6B

	Patients (n=120)	Low cytoplasmic HSF1	High cytoplasmic HSF1	P-value
DYRK2 cytoplasmic (low/high)	36/84	21/15	56/28	0.385
DYRK2 nuclear (low/high)	100/20	65/35	12/8	0.670
	Patients (n=120)	Low nuclear HSF1	High nuclear HSF1	P-value
DYRK2 cytoplasmic (low/high)	36/84	22/14	40/44	0.177
DYRK2 nuclear (low/high)	100/20	56/44	6/14	0.034

Fig S6

S6C

Patient Cohort Characteristics	Number of samples
Age (<50/>50)	208/507
Histological tumour type (other/lobular/ductal)	38/48/629
Tumour size (<20mm/20mm-50mm/>50mm/missing)	398/278/38/1
Tumour grade (1/2/3/missing)	128/320/265/2
Nodal status (negative/positive/missing)	404/301/10
ER status (negative/positive)	226/489
PR status (negative/positive)	376/339
HER2 status (negative/positive/missing)	593/115/7
Ki67 (<15%/>15%/missing)	473/208/34
Molecular subtype (luminal A/ luminal B/ triple negative/ER negative and HER2 positive/missing)	314/163/148/64/26
AR status (negative/positive/missing)	186/478/51
Endocrine therapy (no/yes)	276/439
Chemotherapy (no/yes/missing)	425/288/2
Radiotherapy (no/yes/missing)	374/339/2
Recurrence (no/local/distant/both/missing)	559/39/105/7/5
Survival status (alive/cancer death/death other cause)	425/155/135

S6D

Low DYRK2 levels

High DYRK2 levels

Fig S6

S6E

S6F

Fig S6

S6G

Patient Cohort Characteristics	Univariate analysis		Multivariate analysis	
	P value	Hazard ratio (95% CI)	P value	Hazard ratio (95% CI)
Age (<50/>50)	0.615	0.803(0.341-1.890)		
Histological tumour type (other/lobular/ductal)	0.353	5.412(0.154-190)		
Tumour size (<20mm/20mm-50mm/>50mm/missing)	0.023	2.387(1.129-5.047)		
Tumour grade (1/2/3/missing)	0.745	1.157(0.481-2.782)		
Nodal status (negative/positive/missing)	0.001	5.932(2.167-16.242)	0.006	6.004(1.675-21.811)
Ki67 (<15%/>15%/missing)	0.762	1.169(0.435-3.218)		
Chemotherapy (no/yes/missing)	0.707	1.209(0.443-3.302)		
Radiotherapy (no/yes/missing)	0.537	1.330(0.537-3.295)		
Recurrence (no/local/distant/both/missing)	<0.001	3.887(2.578-5.860)	<0.001	4.287(2.460-7.470)
Cytoplasmic DYRK2 (low/high)	0.067	3.955(0.909-17.203)		
Nuclear DYRK2 (low/high)	0.023	3.115(1.167-8.315)	0.013	3.920(1.338-11.487)
Total DYRK2 (low/high)	0.009	2.564(1.264-5.203)		

S6H

Fig S6