TITLE PAGE

Title

Complete genome assembly of clinical multidrug resistant *Bacteroides fragilis* isolates enables
comprehensive identification of antimicrobial resistance genes and plasmids.

Running title

Genomes and plasmids of six clinical MDR *B. fragilis*

Author names

Thomas V. Sydenham[1,2,3]*, Søren Overballe-Petersen[4], Henrik Hasman[4], Hannah Wexler[5],
Michael Kemp[1,2], Ulrik S. Justesen[1,2]

Affiliations

[1]Research Unit of Clinical Microbiology, Department of Clinical Research, University of Southern
Denmark, Odense, Denmark; [2]Department of Clinical Microbiology, Odense University Hospital,
Odense, Denmark; [3]Department of Clinical Microbiology, Lillebaelt Hospital, Vejle, Denmark;
[4]Bacteria, Parasites & Fungi, Statens Serum Institut, Copenhagen, Denmark;[5] GLAVA Health Care
System and David Geffen School of Medicine at UCLA, Los Angeles, CA, United States.

Corresponding author

Thomas V. Sydenham, Thomas.sydenham@rsyd.dk

ORCIDs

Thomas V. Sydenham: 0000-0003-1058-2449

Henrik Hasman: 0000-0001-6314-2709

Michael Kemp: 0000-0001-5989-0421

Ulrik S. Justesen: 0000-0002-6130-1902

KEYWORDS

Bacteroides fragilis; antimicrobial resistance; genome sequencing; plasmid; oxford nanopore;
hybrid assembly; insertion sequences

REPOSITORIES

Sequence files (MinION reads de-multiplexed with Deepbinner and basecalled with Albacore in
fast5 format and Illumina MiSeq reads in fastq format) and final genome assemblies have been
deposited to NCBI/ENA/DDBJ under Bioproject accessions PRJNA525024, PRJNA244942,
PRJNA244943, PRJNA244944, PRJNA253771, PRJNA254401, and PRJNA254455

ABSTRACT

Bacteroides fragilis constitutes a significant part of the normal human gut microbiota and can also
act as an opportunistic pathogen. Antimicrobial resistance and the prevalence of antimicrobial
resistance genes are increasing, and prediction of antimicrobial susceptibility based on sequence
information could support targeted antimicrobial therapy in a clinical setting. Complete identification
of insertion sequence (IS) elements carrying promoter sequences upstream of resistance genes is
necessary for prediction of antimicrobial resistance. However, *de novo* assemblies from short
reads alone are often fractured due to repeat regions and the presence multiple copies of identical
IS elements. Identification of plasmids in clinical isolates can aid in the surveillance of the

dissemination of antimicrobial resistance and comprehensive sequence databases support
microbiome and metagenomic studies. Here we test several short-read, hybrid and long-lead
assembly pipelines by assembling the type strain *B. fragilis* CCUG4856T
(=ATCC25285=NCTC9343) with Illumina short reads and long reads generated by Oxford
Nanopore Technologies (ONT) MinION sequencing. Hybrid assembly with Unicycler, using quality
filtered Illumina reads and Filtlong filtered and Canu corrected ONT reads produced the assembly
of highest quality. This approach was then applied to six clinical multidrug resistant *B. fragilis*
isolates and, with minimal manual finishing of chromosomal assemblies of three isolates, complete,
circular assemblies of all isolates were produced. Eleven circular, putative plasmids were identified
in the six assemblies of which only three corresponded to a known cultured *Bacteroides* plasmid.
Complete IS elements could be identified upstream of antimicrobial resistance genes, however
there was not complete correlation between the absence of IS elements and antimicrobial
susceptibility. As our knowledge on factors that increase expression of resistance genes in the
absence of IS elements is limited, further research is needed prior to implementing antimicrobial
resistance prediction for *B. fragilis* from whole genome sequencing.

IMPACT STATEMENT

Bacterial whole genome sequencing is increasingly used in public health, clinical, and research
laboratories for typing, identification of virulence factors, phylogenomics, outbreak investigation
and identification of antimicrobial resistance genes. In some settings, diagnostic microbiome
amplicon sequencing or metagenomic sequencing directly from clinical samples is already
implemented and informs treatment decisions. The prospect of prediction of antimicrobial
susceptibility based on resistome identification holds promises for shortening time from sample to
report and informing treatment decisions. Databases with comprehensive reference sequences of
high quality are a necessity for these purposes. *Bacteroides fragilis* is an important part of the
human commensal gut microbiota and is also the most commonly isolated anaerobic bacterium
from non-faecal clinical samples but few complete genome assemblies are available through public
databases. The fragmented assemblies from short read de novo assembly often negate the
identification of insertion sequences upstream of antimicrobial resistance gens, which is necessary
for prediction of antimicrobial resistance from whole genome sequencing. Here we test multiple
assembly pipelines with short read Illumina data and long read data from Oxford Nanopore
Technologies MinION sequencing to select an optimal pipeline for complete genome assembly of
B. fragilis. However, *B. fragilis* is a highly plastic genome with multiple inversive repeat regions,
and complete genome assembly of six clinical multidrug resistant isolates still required minor
manual finishing for half the isolates. Complete identification of known insertion sequences and
resistance genes was possible from the complete genome. In addition, the current catalogue of
Bacteroides plasmid sequences is augmented by eight new plasmid sequences that do not have
corresponding, complete entries in the NCBI database. This work almost doubles the number of
publicly available complete, finished chromosomal and plasmid *B. fragilis* sequences paving the
way for further studies on antimicrobial resistance prediction and increased quality of microbiome
and metagenomic studies.

ABBREVIATIONS

AMR, antimicrobial resistance; WGS, whole genome sequencing; IS, insertion sequence; ONT,
Oxford Nanopore Technologies;

DATA SUMMARY

1. Sequence read files (Oxford Nanopore (ONT) fast5 files and Illumina fastq files) as well as the
final genome assemblies have been deposited to NCBI/ENA/DDBJ under Bioproject
accessions PRJNA525024, PRJNA244942, PRJNA244943, PRJNA244944, PRJNA253771,
PRJNA254401, and PRJNA254455.

2. Fastq format of demultiplexed ONT reads trimmed of adapters and barcode sequences are available at doi.org/10.5281/zenodo.2677927
3. Genome assemblies from the assembly pipeline validation are available at doi: doi.org/10.5281/zenodo.2648546.
4. Genome assemblies corresponding to each stage of the process of the assembly are available at doi.org/10.5281/zenodo.2661704.
5. Full commands and scripts used are available from GitHub: https://github.com/thsyd/bfassembly as well as a static version at doi.org/10.5281/zenodo.2683511

INTRODUCTION

Bacteroides fragilis is a Gram-negative anaerobic bacterium that is commensal to the human gut but can act as an opportunistic pathogen; it is the most commonly isolated anaerobic bacteria from non-faecal clinical samples (1). Antimicrobial resistance rates are increasing for *B. fragilis*, especially for carbapenems and metronidazole, two widely used antimicrobials for treatment of severe infections and anaerobe bacteria (2,3). Antimicrobial susceptibility testing of anaerobes using agar dilution or gradient strip methods can be costly and labour intensive and despite efforts to validate disk diffusion as a less expensive option, turn-around time will still be least 18 hours and validation for individual species will be required (4).

Antimicrobial resistance prediction from bacterial whole genome sequences, from cultured isolates as well as metagenomes, could be implemented in clinical microbiology in the near future, with the potential for improved sample-to-report turnover time and possibly eliminating the need for phenotypical testing for individual species (5–8). For a few species, prediction of antimicrobial resistance from WGS has been validated, but for the majority of clinical relevant species challenges still remain (6,9,10).

Based on DNA-DNA hybridisation studies, *B. fragilis* can be divided into two DNA homology groups (division I and II), whose ribosomal contents are so different that the two divisions can be distinguished by mass spectrometry routinely used to identify isolates in clinical laboratories (11). *B. fragilis* division I carry the chromosomal cephalosporinase gene *cepA* whilst *B. fragilis* division II harbour the chromosomal metallo-β-lactamase gene *cfiA* (also known as *ccrA*) (12,13). The *cfiA* gene can confer resistance to carbapenems, a class of antimicrobials usually reserved for patients with severe sepsis or infections with multidrug-resistant bacteria. But expression levels are partly controlled by insertion sequence (IS) elements carrying promotor sequences inserted upstream of the gene and only 30-50% of clinical isolates that harbour *cfiA* display phenotypically reduced susceptibility to carbapenems (3). The same pattern of expression control can be observed for genes associated with resistance to metronidazole (*nim* genes) and clindamycin (*erm* genes) (1).

In 2014 we observed that identification of IS elements upstream of known antimicrobial resistance genes in *B. fragilis* was hampered in short read *de novo* assemblies even though the genes could be identified (14). This occurred because contigs were often terminated close to the start of the resistance genes, presumably due to the proliferation of multiple copies of the same IS elements throughout the *B. fragilis* genomes. Genome assemblies from short read sequencing technologies alone most often result in fragmented assemblies because of repetitive regions and genome elements with multiple occurrences in the chromosomes and plasmids (15,16). Therefore, we could not predict antimicrobial resistance (AMR) phenotypes in *B. fragilis* using only short reads for WGS since IS element identification is a prerequisite for correct genotype-phenotype associations. Long read sequencing technologies are increasingly being utilised to increase the contiguity of bacterial genome assemblies and often result in complete, closed chromosomes and plasmids (17–20). This provides possibilities for comprehensive identification of IS elements, insights into genome structures and characterisation of other mobilisable elements and associated genes.

Complete identification and characterisation of plasmids in sequenced isolates would allow for
improved analysis of the plasmid-mediated spread of antimicrobial resistance.

Bioinformatic analysis of WGS data depends heavily on high-quality reference databases.
Anaerobes make up most of the bacterial human commensal microbiota but are most likely
underrepresented in public databases of whole genomes from cultured isolates. The NCBI
Genome database (accessed 31-03-2019) contains genome sequences of 191,411 bacteria of
which 13,483 are marked as complete assemblies. Only seven of these are *Bacteroides fragilis*
(21–27). In comparison there are 776 assemblies of *E. coli* marked as complete and 398 of *S.*
aureus. Improving the representation of complete assemblies of *B. fragilis* in the public genome
databases will support the development of antimicrobial resistance prediction from WGS as well as
microbiome and metagenomic analysis projects.

The aims of this study were to select an optimal assembly software pipeline for complete, circular
assembly of *Bacteroides fragilis* and demonstrate the utility of complete assembly for both plasmid
identification and comprehensive detection of genes and IS elements associated with antimicrobial
resistance. We assembled the *B. fragilis* CCUG4856T (= ATCC25285 = NCTC9343) reference
strain utilising long reads generated with the MinION sequencer from Oxford Nanopore Technology
(ONT) and high-quality Illumina short reads and selected the best assembly pipeline by comparing
assemblies to the Sanger sequenced reference NCTC9343 (RefSeq accesion GCF_000025985.1).
The best assembly pipeline was then applied to six clinical multi-drug resistant *B. fragilis* isolates
from our 2014 study (14).

METHODS

Culture conditions and DNA extraction

Bacteroides fragilis CCUG4856T and the six strains described in our previous study were included
(14,21). Strains were stored at -80° in beef extract broth with 10% glycerol (SSI Diagnostica) and
cultured on solid chocolate agar with added vitamin K and cysteine (SSI Diagnostica) for 48 hrs in
an anaerobic atmosphere at 35 °C. Ten µl of culture was transferred to 14 ml saccharose serum
broth (SSI Diagnostica) and incubated for 18 hrs under the same conditions. DNA was then
extracted using the Genomic-Tip G/500 kit (Qiagen) following the manufacturers protocol for Gram
negative bacteria and eluted into 5 mM Tris pH 7.5 0.5 mM EDTA buffer. Quality control was
performed by measuring fragment length on a TapeStation 2500 (Genomic DNA ScreenTape,
Agilent), purity on the NanoDrop (ThermoFisher Scientific) and concentration on the Qubit (dsDNA
BR kit; Invitrogen). The eluted DNA was then stored at -20 °C.

Illumina library preparation, sequencing and quality control

The strains had previously been sequenced and assembled using Illumina short reads for our
previous study (14), but to minimise biological disparities we opted to re-sequence with Illumina
using the same DNA extraction prepared for long read sequencing. Paired-end libraries were
generated using the Nextera XT DNA sample preparation kit (Illumina) according to the
manufacturer's protocol. DNA was sequenced on a MiSeq sequencer (Illumina) with 150 bp reads
for a theoretical read depth of 100x. Read quality metrics were evaluated using FastQC
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and fastp v0.19.6 (28). Filterbytile
from the BBmap package (http://sourceforge.net/projects/bbmap/) was used for removing low-
quality reads based on positional information on the sequencing flowcell and TrimGalore
(http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), with settings --qual 20 and --
length 126, provided additional adapter and quality trimming. FastQ files were then randomly
down-sampled to < 100x crude read depth using an estimated genome size of 5.3 Mb, as higher
read depths tend to reduce assembly quality (29).

Nanopore library preparation and MinION sequencing

Sequencing libraries were prepared using the Rapid Barcoding kit (SQK-RPB004; Oxford
Nanopore Technologies) following the manufacturers protocol (version
RPB_9059_v1_revC_08Mar2018) with SPRI bead clean up (AMPure XT beads; Beckman Coulter)
as described. Sequencing was performed as multiplex runs on a MinION connected to a Windows
PC with MinKnow v1.15.1 using FLO-MIN106 R9.4 flowcells. Raw fast5 files were transferred to
the Computerome high performance cluster (https://www.computerome.dk/) for analysis. Four
sequencing runs were performed, as the first two runs did not provide enough data for complete
assembly of all isolates (see results section).

Fast5 demultiplexing, base-calling, quality control and filtering

The raw fast5 files were demultiplexed with Deepbinner v0.2.0 and base-called using Albacore
v2.3.3, retaining only those barcodes Deepbinner and Albacore agreed upon for minimal barcode
misclassification (30). Porechop v0.2.4 (https://github.com/rrwick/Porechop) with the --
discard_middle option was used for adapter and barcode trimming and read statistics were
collected using NanoPlot (31). Filtlong v0.2.0 (https://github.com/rrwick/Filtlong) was used to filter
the long reads by either removing the worst 10% or by retaining 500Mbs in total, which ever option
resulted in fewer reads.

Assembly validation

To select and validate the optimum assembly pipeline *Bacteroides fragilis* CCUG4856T was
assembled using a variety of well-known assemblers and polishing tools (Table 1). Each
assembler was run with the Filtlong filtered reads as input or the filtered reads corrected with Canu
1.8 (with standard settings, corMinCoverage=0, or coroutCoverage=999). Canu was also tested
with the unfiltered reads as input. Hybrid assemblers used the filtered long reads and the filtered,
trimmed and down-sampled Illumina reads. Unicycler includes polishing with Racon and Pilon. For
assemblers other than Unicycler, Racon polishing with ONT reads was run for one or two rounds
and Pilon was run until no changes were made or for a maximum of six rounds. Racon polishing
with Illumina reads was run for one round.

The original Sanger sequenced *Bacteroides fragilis* NCTC9343 (=CCUG4856T) (21) downloaded
from NCBI RefSeq (accession GCF_000025985.1) was used as reference sequence for the
assembly comparisons and Quast v5.0.2 was used for assembly summary statistics, indel count,
and K-mer-based completion (32). BUSCO v3.0.2b with the bacteroidetes_odb9 dataset, CheckM
v1.0.12, and Prokka v1.13.3 were used to assess gene content (33–35). Average nucleotide
identity was calculated using https://github.com/chjp/ANI/blob/master/ANI.pl and ALE v0.9, which
uses a likelihood based approach to assess the quality of different assemblies, was also used to
score the assemblies (36,37). Ranking of assemblies was based on number of contigs, number of
circular contigs, closeness to total length compared to the reference genome, number of local
misassembles, number of mismatches per 100 kb, number of indels per 100kb, average nucleotide
identity (ANI), CheckM and BUSCO scores, and the total ALE score (a higher score is better).
Please see https://github.com/thsyd/bfassembly for full bioinformatics methods.

Genome assembly of MDR *B. fragilis* isolates

The assembly strategy deemed to produce the highest quality genome for CCUG4856T was
chosen for initial assembly of the six MDR *B. fragilis* isolates. Manual finishing of incomplete
assemblies was performed using Bandage for visualisation of assembly graphs and BLASTn
searches (38). Minimap2 and BWA MEM were used to map reads to the assemblies for coverage
graphs (39,40). Long read assembly with Flye was compared to the Unicycler assembly and used
to guide and validate the manual finishing results. Circlator's *fixstart* task was used to fix the start
position of the manually finished genomes to be at the *dnaA* gene (41).

The assembled genomes were submitted to NCBI GenBank and annotated with PGAP (42).
ABRicate v0.8.10 (https://github.com/tseemann/ABRicate) (with options --minid 40 --mincov 25)

was used to screen for antimicrobial resistance genes with the ResFinder (database date 19-08-
2018), NCBI Bacterial Antimicrobial Resistance Reference Gene Database (database date 19-09-
2018), and CARD (v2.0.3) databases, supplemented with nucleotide sequences for the multidrug
efflux-pump genes *bexA* (GenBank: AB067769.1:3564..4895) and *bexB* (GenBank:
AY375536.1:4599..5963) (43,44). IS elements were identified using ABRicate with data from the
IS-finder database (http://www-is.biotoul.fr/, Update: 2018-07-25) (45).

Identification of plasmids and mobile genetic elements

The PLSDB web server (https://ccb-microbe.cs.uni-saarland.de/plsdb/) (data v. 2019_03_05)
contains bacterial plasmid sequences retrieved from the NCBI and was used for screening and
identifying putative plasmids sequences (46). Only hits to accessions from cultured organisms
were included. Putative plasmids not identified using PLSDB, were evaluated by the read depth
relative to the chromosome (higher relative read depth indicates plasmid sequence) and Pfam
families covering known plasmid replication domains from Table 1 in reference (47) were
downloaded from the Pfam database (Pfam 32.0, https://pfam.xfam.org/) and used for screening
putative plasmids with ABRicate.

RESULTS

Sequencing data quality

For Illumina data, a median of 3,465,082 reads (interquartile range [IQR]: 3,177,493-5,001,077)
were generated for each isolate (Supplementary Table S1)). After filtering, adapter-removal and
down sampling a median of 449,022,741 bases (IQR: 433,517,549-530,257,210) were available
per isolate with 87-96% Q30 bases corresponding to calculated read depths of 75-103%. The
%GC content of the reads for each isolate (median 42.9%, range: 42.6-43.3%) were very
consistent and within the expected range for the *Bacteroides* genus (40-48%) (48) .

Isolates were sequenced in runs multiplexed with other isolates not included in this study. Based
on initial test assemblies using Unicycler without filtering or Canu correction (not shown) it was
concluded that data from the first ONT sequencing runs were to be supplemented by additional
runs to increase the chance of complete assembly of all isolates. Concatenating reads from runs, a
median of 75,598 reads [IQR: 50,210-112,065] with a median length of 2,938-4,393 bases were
generated for each isolate (Supplementary Table S1). Filtering with Filtlong and correction with
Canu resulted in a median of 8,515 reads (IQR: 6,226-10,370) with median lengths of 6,181-
38,588 for each isolate as input for the assemblies.

Selecting the optimal assembly pipeline

141 assemblies of *B. fragilis* CCUG4856T were generated using the various assemblers and
polishing steps (Supplementary Table S2). Compared to the reference genome, Unicycler
assemblies were of the highest quality (**Table 2**). Unicycler, with any of the read input options,
produced two circular contigs of the expected lengths, and the differences between the various
Unicycler assemblies were minimal (**Table 3**). Assemblies with Canu corrected reads showed
slightly higher genome fractions and average nucleotide identities to the reference and fewer
mismatches and indels, when compared to Unicycler alone. Unicycler assemblies corrected with
Racon using Illumina reads worsened slightly overall with 0.04-0.19 more indels and 0.14-0.25
more mismatches per 100 kbp. Based on this initial evaluation, the assembly pipeline using Canu
corrected reads with default options was chosen (Assembly "OF.CS" in **Table 3**). This would
reduce the number of long reads, compared to Canu correction with corMinCoverage=0, or
coroutCoverage=999, and thereby lead to a faster run-time for Unicycler.

The hybrid Unicycler assembly of CCUG4856T with standard Canu corrected ONT reads consists
of two circular contigs of 5,205,133 and 36,560 bp in length. The plasmid is the same length as
plasmid pBF9343 from the reference assembly GCF_000025985.1 and the chromosome is seven

bases shorter. Alignments of the Sanger sequenced assembly GCF_000025985.1 with the hybrid
Unicycler assembly show an 88,045 bp inversion in the hybrid assembly compared to the Sanger
assembly (Figure 1). This inversion is present in all the best assemblies, including assemblies
derived from solely ONT sequences or Illumina sequences (Supplementary Figure S1) as well as
two additional assemblies of NCTC9343/ATCC25285 from PacBio and Illumina sequences
downloaded from NCBI RefSeq (Supplementary Figure S2).

Complete assembly of six multidrug resistant isolates

Unicycler, using filtered and trimmed Illumina reads and the Filtlong filtered and Canu corrected
ONT reads from the first sequencing runs, generated complete, continuous, circular assemblies for
two of the six isolates (BFO18 and BFO67) (Figure 2). For the assemblies that were not complete
with sequencing data from the first MinION runs, increasing the amount of ONT data resulted in
fewer contigs overall, except for BF067, where the additional data from the second sequencing run
led to a fragmented assembly and manual finishing was necessary. Performing assembly of isolate
S01 without Canu correction of the ONT reads from the first sequencing resulted in a closed
chromosome and performing Canu correction of reads resulted in a fragmentation of the
chromosome. This was ameliorated by including more ONT data. By manual finishing using read
mapping and additional assembly with Flye, the remaining three assemblies were circularised.
Chromosomes varied in length from 5,141,257 – 5,504,076 bp. Alignment of ONT and Illumina
reads to the chromosome assemblies showed even coverage for both sequencing technologies
(Supplementary Figure S3). For BFO85 a >100% relative read depth increase was observed at
approximately 25kb-38kb. This could represent a 12 kb repeat region that was not resolved in the
assembly. Seven (47%) of the 15 PGAP annotated CDS' in the 13kb region were annotated as
hypothetical proteins. None of the annotated CDS' represented mobilisable proteins.

Eleven putative plasmid sequences were identified

A total of 11 putative circular plasmids were identified in the six *B. fragilis* isolates (**Table 4**). Zero
to three putative plasmids were identified per isolate with lengths varying from 2,782 to 85,671bp.

The PLSDB database contains NCBI RefSeq plasmid sequences marked as complete. Three of
the 11 putative plasmid sequences were found to match (ID > 98%) a sequence in PLSDB (Table
4). These three all matched the cryptic plasmid pBFP35 (49). The NCBI Nucleotide database was
queried using BLASTn with the remaining unidentified putative plasmid sequences (50). BFO18
putative plasmid sequence pBFO18_1 (7,221 bp) resembles plasmid pIP421, a 7.2kb plasmid with
metronidazole resistance gene *nimD* and IS*1169.* Partial sequences in NCBI GenBank spanning
the *nimD* gene, IS element and *RepA* (GenBank Y10480.1 and X86702.1) showed 99 %ID to their
alignment to pBFO18_1 (not shown) (51,52). Strain S01 putative plasmid sequence pBFS01_2
(8,331 bp) showed 99.87 %ID to the 1486bp partial sequence of *B. fragilis* plasmid pBF388c
(GenBank AM042593.1), a 8.3kb conjugative plasmid harbouring *nimE* and IS*Bf6* (53).

None of the three putative plasmid sequences of strain BFO18 could be identified using the
PLSDB but querying the NCBI nucleotide database using BLASTn revealed hits for all three. The
hits corresponded to circularised sequences (%ID: 99.56-99.96, %COV: 100) assembled from
mobilome metagenomic sequencing of the uncultured caecum content from a rat trapped at
Bispebjerg Hospital in Copenhagen, Denmark (two hours' drive from Odense University Hospital
where BFO18 was isolated from a patient's blood culture) (Supplementary Table S3) (47,54).
BLASTn searches of the remaining unidentified putative plasmids from the other strains did not
reveal complete hits.

Using ABRicate with the plasmid replication domains collected from the Pfam database, all
putative plasmids, except pBF017_1 and pBFS01_1, were found to have recognised replicon
domains (Table 4). DNA fragments of sizes matching pBFO17_1 and pBFS01_1 were detected by
PFGE of S1 endonuclease restriction enzyme treated plasmid DNA extracts (Supplementary

Figure S4) and the circular structures of the two sequences lacking a predicted replication domain,
were confirmed manually by visually inspecting BLASTn mapping of ONT sequences longer than
10 kbp to the assembled plasmid sequences with CLC Genomics Workbench 10 (Qiagen). Eleven
and 22 ONT reads spanned the complete lengths of pBFO17_1 and pBFS01_1 respectively and
contained no other elements. pBFO17_1 and pBFS01_1 demonstrate a degree of similarity of
close to 100%, except for an approximate total of 7,500 bp transposase and prophage sequences
in pBF017_1 (Figure 3). No alignment to chromosomal sequences of any of the included *B. fragilis*
isolates was observed using progressiveMauve (not shown) (55).

The GC content of pBFO17_1 and pBFS01_1 are 36.78% and 36.04% respectively. These lie
within the range for the *Bacteroides* genus but differ from the expected value for *B. fragilis* (43%),
which could indicate that the putative plasmids do not originate from *B. fragilis* (56). After
supplementing the PGAP annotations with RAST annotation (57), 63% (pBFO17_1) and 59%
(pBFSO1_1) of CDS' remained annotated as hypothetical or as domain of unknown function. Of
the annotated CDS' the majority were associated with mobilisable features, plasmids and phages
such as *parA* and *parB*, DNA partitioning proteins, conjugative transposon proteins, transposases,
DNA binding motif domain containing proteins, and reverse transcriptase protein. The results
above support the assembly data suggesting these two sequences are in fact plasmids.

Detection of antimicrobial resistance genes and Insertion sequence elements

We used ABRicate to screen assemblies for AMR genes (ResFinder, NCBI and CARD databases
supplemented with sequences for *bexA* and *bexB*) and IS elements (IS-finder database); several
AMR genes, possible homologs to known AMR genes and IS elements adjunct to the AMR genes
were detected (Table 5). Of note, isolate BFO17 contains two homologs of the metronidazole
resistance gene *nimJ* (with a 100% consensus) and two isolates, S01 and BFO85, harbour two
homologs of the tetracycline resistance gene *tetQ*. Homologs to *bexA* and *bexB* were identified
with 73.53-99.12 %ID and were all confirmed with BLASTx searches against the NCBI nr
database, as was done in our previous study (14). Partial hits for *ugd* was observed for several
isolates, but with low %ID and %COV, and possible represent identification of conserved domains,
but not *ugd* homologs. Increased expression of the *cfiA* metallo-beta-lactamase gene, *nim*-family
5-nitroimidazole genes and *erm* genes is partly regulated through IS elements containing promoter
sequences. Full length IS elements could be identified upstream of 11 (79%) of 14 *cfiA*, *nim* and
erm genes and upstream of two of three *CfxA4* genes and the *OXA-347* gene identified in BFO42.
The described *Bacteroides fragilis* promotors TAnnTTTG (-7) and TG or TTG or TGTG (-33) (58)
were searched for manually but could not be identified upstream of the two *cfiA* genes in isolates
BFO67 and BFO85 or the *ermB* gene in BFO85 for which no IS elements could be detected
upstream (not shown).

Correlation between identified genes and IS elements and phenotypical resistance

As in our previous study, the *cfiA* gene was identified in the five meropenem resistant isolates
(Table 5). All the *cfiA* genes were found on the chromosomal sequences. Complete IS elements
were identified upstream of the *cfiA* genes in BFO17, BFO18 and S01, but not in BFO67 or
BFO85. MICs for meropenem and imipenem were lower for these two isolates. *Nim* genes (-*A, -D,*
-E and -*J*) could be found in the four metronidazole resistant isolates, all with complete IS elements
upstream. Three of the *nim* genes were found on putative plasmids of the respective isolates. The
four clindamycin-resistant isolates all carried *erm*-genes with upstream IS elements. A transposase
was inserted in the *ermF-gene* in isolate BFO18, splitting it in two and the same isolate
demonstrated a lower clindamycin MIC (6 mg/L) than the other three clindamycin resistant isolates.

DISCUSSION

Hybrid genome assembly produces high quality *B. fragilis* genomes

The primary aim of this study was to select and validate an assembly method to reliably complete
chromosome and plasmid assembly of *B. fragilis* genomes. From 141 assembly variations, a
hybrid approach using Filtlong filtered and Canu corrected ONT reads with quality filtered Illumina
reads as input to Unicycler produced a complete, closed assembly of *B. fragilis* CCUG4856T with
high similarity to the reference assembly of the original Sanger sequenced reference assembly. An
88kb inversion was observed when comparing the two assemblies. Cerdeño-Tárraga and
colleagues observed difficulties in resolving certain regions of the Sanger sequenced assembly of
NCTC9343 due to invertible regions with flanking inverted repeat sequences (21). The observed
inversion in the hybrid Unicycler assembly, could be due to a) a superior assembly where the
longer ONT reads have overcome the shortcomings of the shorter Sanger sequences, b) an
incorrect assembly by Unicycler, c) a biological difference that has occurred over time between the
strain stored at NCTC and CCUG, or d) a biological difference that occurred during the culturing of
the strain, with dominance of a clone with the inversion, prior to DNA extraction as part of this
study. The observations that the inversion is also present in all the best assemblies from this study
and assemblies from two other research institutions support the conclusions that the current hybrid
Unicycler assembly represents the true orientation of the 88kb sequence.

Complete genome assembly of three of the six multidrug resistant isolates required manual
finishing

The assemblies of BFO18, S01, and BFO42 were completed by Unicycler without manual
intervention, but the chromosomes of BFO17, BFO67, and BFO85 could only be closed by
performing manual steps. The manual finishing steps are time consuming, difficult to replicate and
are easily biased. In order to be implemented in routine clinical laboratories, large scale,
automated, complete assembly of prokaryote genomes require robust methods with minimal
human interaction. Genome assembly using another long-read assembler, Flye, supported the
results of the manual finishing for two of three isolates. Flye is better at resolving repeats than
miniasm, the long read assembler included in the Unicycler pipeline (59). One option could be to
include the long-read assembly from Flye, in place of that of miniasm, to guide bridge building for
the higher quality Illumina-only contigs produced in the first steps of Unicycler. To resolve repeats it
is often necessary to have long reads that span the repeat. In prokaryotes repeats over 10kb are
not unusual and they are often spanned by the ONT reads generated, even by novice researchers.
But repeat regions of up to 120kb and duplications of 200kb have been described in some
prokaryotes (17,18,60). ONT sequencing runs will routinely result in many reads that span the
majority of repeats, but to obtain ONT reads that span specific 120-200kb repeats in a genome of
interest still requires skill and a certain amount of luck. Protocols for ONT sequencing have been
described that result in read lengths of over 2 Mb, but this requires skilled and experienced
researchers and lab technicians and demands high amounts of very high quality input DNA and
essentially sequencing of only one isolate per MinION flowcell (61).

ONT read depth did not serve as an indicator of whether the Unicycler assemblies would result in
closed chromosomal contigs in this study. Final ONT read depth, prior to Filtlong filtering and Canu
correction, ranged from 23-371x, but a high read depth alone, was not an indicator of closed
contigs. The three assemblies BF017, BFO67, and BFO85 required manual finishing to complete
the assemblies and had ONT raw read depths of 99-137x. After Filtlong filtering and Canu
correction the median read lengths were 21,932-29,893b and read length N50 was 25,765-34,815b
for the three isolates (Supplementary Table S1). Canu correction improved the Unicycler assembly
of *B. fragilis* CCUG4856T by nearly all parameters. But whilst Canu correction of the data from the
first sequencing run resulted in the complete assembly of BFO67, the assembly of S01 worsened
slightly. Increasing the amount of ONT data for BFO67 fragmented the complete chromosome.
However, increasing the ONT read depth did decrease the number of contigs per isolate in our
study overall.

Defining an optimal approach for complete prokaryote genome assembly is a continuous process,
as sequencing technologies and assembly software develop and mature. Ring and colleagues
found that Canu correction prior to Unicycler hybrid assembly was superior to other hybrid
assembly or long read assembly approaches for assembly of *Bordetella pertussis* genomes that
contain long duplicated regions (18). Unicycler also performs well in other studies comparing
assessing genome assemblers for bacterial genome and plasmid assembly (19). De Maio and
colleagues recently published a preprint comparing hybrid assembly strategies for 20
Enterobacteriaceae isolates (20). In their dataset, simply randomly subsampling ONT reads to an
approximate read depth of 100x was slightly superior to applying Canu correction or Filtlong
filtering prior to Unicycler assembly. For 85% of isolates the expected number of circular contigs
were all assembled. For only one additional isolate Canu correction or Filtlong filtering resulted in
the assembly of the expected number of circular contigs. Manual steps, including down sampling
ONT reads or removing the Canu correction are options to consider, if chromosomes are not
complete and circularised after initial Unicycler assembly, providing ONT read depth of 100x is
available.

We chose to benchmark a selection of widely used genome assemblers for short read, long-read
and hybrid bacterial genome assembly as well as polishing tools for long read assemblies, but
many other options have been published. Most assemblers and polishing tools were run using
default parameters, and it is possible that further optimisation of settings for the individual software
packages might have improved assemblies further than was demonstrated here. As sequencing
technologies and assembly software continues to improve, continued validation of pipelines is
advisable. Software such as poreTally provides user friendly options for benchmarking genome
assembly pipelines prior to implementation (62).

Bacteroides plasmids are not well represented in public databases

A secondary aim of this study was to identify plasmids in the hybrid assemblies. Automated tools
have been developed and validated for identification of plasmids from genome assemblies or read
data, but they are dependant of collated databases of known plasmid sequences. As such, tools
such as PlasmidFinder or mlplasmids can be applied for plasmid identification for
Enterobacteriaceae or *Enterococcus faecium*, but *B. fragilis* is not supported at the time of writing
(63,64). Therefore, we evaluated putative plasmid sequences by sequence identity and length
comparison using the PLSDB webpage, identifying plasmid replication domains, and using
circularisation and relative coverage as indicators that a sequence represents a plasmid in a given
isolate.

Only four of the twelve plasmid sequences from the seven isolates could be identified using the
PLSDB and three of these were the same plasmid, pBFP35. Two other putative plasmids,
pBFO18_1 and pBFS01_2 were likely plasmids pBF388c and pIP421 based on the partial
sequences from these plasmids and plasmid length. This still leaves half of the circularised,
putative plasmids unidentified. The two longer putative plasmids, pBFO17_1 and pBFS01_1,
displayed a high degree of similarity, a GC% out of the normal range for *B. fragilis*, and a relative
read depth of double the reads compared to the chromosome. Most annotated CDS' were
associated with mobilisable elements, but no known plasmid replication domains could be
identified. From the sequencing data alone, we cannot conclude that they represent true plasmids,
however the findings above and manual inspection of long read mapping support that inference.

There are only 14 complete plasmid sequences from cultured *Bacteroides* isolates in the PLSDB
v2019_03_05, which is based on the NCBI RefSeq database. Many other *Bacteroides* plasmids
have been partially described, and some are represented by partial sequences or marked as contig
level in the NCBI nucleotide database (65–68). Metagenomic sequencing and genome assembly
projects are expanding the public sequence databases and screening the NCBI nucleotide
database, sequences with a high degree of similarity to the putative plasmid sequences from one

patient isolate (BFO18) could be found. These originated from a rat caecum metagenomic plasmid
sequencing project from Copenhagen, a few hours' drive from Odense University Hospital. To
understand and perform surveillance of the dissemination of plasmids there is a need for increased
submissions of high quality, annotated and phenotypically validated sequences of bacterial isolates
including plasmids. This study adds significantly to the number of complete plasmid sequences
associated with *Bacteroides.*

Complete assembly allows comprehensive identification of resistance determinants in *B.*
fragilis

We also intended to comprehensively identify resistance genes and IS elements in the hybrid
genome assemblies. Using ABRicate with several resistance gene databases and IS-element
nucleotide sequences, the findings of our previous study were confirmed and enhanced.
Assemblies from Illumina sequencing alone would only allow partial IS element identification (14).
Now, with the complete assemblies, comprehensive identification of known IS elements upstream
of the relevant resistance genes could be completed. In our first study we used ResFinder with the
available database at that time. Now, by including several databases, and lowering the %ID
threshold, the number of genes identified increased. Additionally, for as a result of the complete
genome assembly of BFO17, we could now identify two copies of *nimJ*, while only one copy was
identified in the short read draft assembly of the same isolate in the previous study. Husain and
colleagues identified the presence of three copies of *nimJ* in strain HMW615, when describing the
nimJ gene (69). We confirmed this finding by running ABRicate on the HMW615 assembly as done
with the isolates of this study (not shown). Interestingly, RAST annotates a third *nim* gene
(nucleotide positions 1,359,590..1,360,093) in the Unicycler hybrid assembly of BFO17, and the
PGAP annotation includes an additional annotation of a pyridoxamine 5'-phosphate oxidase family
gene (nucleotide positions 940,032..940,505), the family that includes the *nim*-genes. It is possible
that one or more novel homologs of the *nim* are present in BFO17.

IS elements could be identified upstream of most relevant resistance genes. However, in three
cases no IS element was present upstream of a resistance gene, even though the isolates
displayed phenotypical resistance associated with increased expression of the specific gene.
Known *B. fragilis* promoter sequences could not be identified upstream of the genes "missing"
upstream IS elements, however *B. fragilis* promotors are still not completely described, so it is
possible there are other unknown variants.

By selecting an optimal genome assembly strategy for *B. fragilis*, supplemented with minimal
manual finishing efforts, and applying this to six multidrug resistant isolates, the number of
complete *B. fragilis* genomes and plasmids in the public databases has now almost doubled. The
future aim of performing antimicrobial resistance prediction based solely on WGS information for *B.*
fragilis demands near-complete genomes for identification of IS elements upstream of resistance
genes. However, we must caution that the absence of an IS element upstream of *cfiA* does not
always correlate to susceptibility to carbapenems. Future studies are needed to address this, and
utilising complete genome assembly for genome wide association studies is one approach that
could be pursued. Technologies that provide a single solution for real-time, high-quality sequencing
of long reads will be essential for implementing near real-time diagnostics of infectious diseases
and characterisation of pathogens.

AUTHOR STATEMENTS

Authors and contributors

The study was conceptualised by T.V.S. and U.S.J. Funding was secured by T.V.S., M.K., H.H.
and U.S.J. Data curation and investigation was performed by T.V.S.. Formal analysis was done by
T.V.S. and S.O-P.. Resources were provided by M.K., T.V.S., H.H. and U.S.J.. U.S.J, H.H. and

M.K. supervised the work. T.V.S wrote the original draft and edited the manuscript. U.S.J., M.K.,
T.V.S., H.H., S.O-P. and H.M.W revised the manuscript.

Conflicts of interest

The authors declare that there are no conflicts of interest

Funding information

Funding supporting this work was provided from the Danish Medical Research Grant [case no.
2013-5480/912523-108] as well as internal funds from the Department of Clinical Microbiology,
Odense University Hospital, Odense, Denmark. Thomas V. Sydenham's salary as a PhD student
was funded by the University of Southern Denmark and he received a travel grant from the Henrik
& Emilie Ovesen Foundation.

Ethical approval

Isolates were obtained as part of routine clinical care, and details about the isolates have
previously been published. No ethical approvals were required.

Acknowledgements

We are very grateful to Professor Henrik Westh, Department of Clinical Microbiology, Hvidovre
Hospital, Denmark, for allowing use of their Computerome High Performance Computing cluster
account for data analysis and to Mikala Wang, Dept. of Clinical Microbiology, Aarhus University
hospital, for gifting us the MDR *B. fragilis* strain isolated at her department. We thank Valentina
Galata, Saarland University, Saarbrücken, Germany (first author of the paper describing PLSDB)
and Natalie Ring, University of Bath, Bath, United Kingdom (first author of reference (18)) for kind
and helpful answers to questions via e-mail.

531 **References**

1. Wexler HM. Bacteroides: the good, the bad, and the nitty-gritty. Clin Microbiol Rev. 2007
Oct;20(4):593–621. doi: 10.1128/CMR.00008-07

2. Nagy E, Urbán E, Nord CE. Antimicrobial susceptibility of Bacteroides fragilis group isolates in
Europe: 20 years of experience. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect
Dis. 2011 Mar;17(3):371–9. doi: 10.1111/j.1469-0691.2010.03256.x

3. Ferløv-Schwensen SA, Sydenham TV, Hansen KC, Hoegh SV, Justesen US. Prevalences of
antimicrobial resistance and the cfiA resistance gene in danish bacteroides fragilis group
isolates since 1973. Int J Antimicrob Agents. 2017 Jun 27; doi:
10.1016/j.ijantimicag.2017.05.007

4. Nagy E, Justesen US, Eitel Z, Urbán E. Development of EUCAST disk diffusion method for
susceptibility testing of the Bacteroides fragilis group isolates. Anaerobe. 2015 Feb;31:65–71.
doi: 10.1016/j.anaerobe.2014.10.008

5. Zankari E, Hasman H, Kaas RS, Seyfarth AM, Agersø Y, Lund O, et al. Genotyping using
whole-genome sequencing is a realistic alternative to surveillance based on phenotypic
antimicrobial susceptibility testing. J Antimicrob Chemother. 2013 Apr;68(4):771–7. doi:
10.1093/jac/dks496

6. Stoesser N, Batty EM, Eyre DW, Morgan M, Wyllie DH, Del Ojo Elias C, et al. Predicting
antimicrobial susceptibilities for Escherichia coli and Klebsiella pneumoniae isolates using
whole genomic sequence data. J Antimicrob Chemother [Internet]. 2013 May 30 [cited 2013
Aug 19]; doi: 10.1093/jac/dkt180

7. Boolchandani M, D'Souza AW, Dantas G. Sequencing-based methods and resources to
study antimicrobial resistance. Nat Rev Genet. 2019 Mar 18;1. doi: 10.1038/s41576-019-
0108-4

8. Didelot X, Bowden R, Wilson DJ, Peto TEA, Crook DW. Transforming clinical microbiology
with bacterial genome sequencing. Nat Rev Genet. 2012 Sep;13(9):601–12. doi:
10.1038/nrg3226

9. Davies TJ, Stoesser N, Sheppard AE, Abuoun M, Fowler PW, Swann J, et al. Reconciling the
potentially irreconcilable? Genotypic and phenotypic amoxicillin-clavulanate resistance in
Escherichia coli. bioRxiv. 2019 Jan 7;511402. doi: 10.1101/511402

10. Ellington MJ, Ekelund O, Aarestrup FM, Canton R, Doumith M, Giske C, et al. The role of
whole genome sequencing in antimicrobial susceptibility testing of bacteria: report from the
EUCAST Subcommittee. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis. 2017
Jan;23(1):2–22. doi: 10.1016/j.cmi.2016.11.012

11. Nagy E, Becker S, Sóki J, Urbán E, Kostrzewa M. Differentiation of division I (cfiA-negative)
and division II (cfiA-positive) Bacteroides fragilis strains by matrix-assisted laser
desorption/ionization time-of-flight mass spectrometry. J Med Microbiol. 2011 Nov;60(Pt
11):1584–90. doi: 10.1099/jmm.0.031336-0

12. Rogers MB, Parker AC, Smith CJ. Cloning and characterization of the endogenous
cephalosporinase gene, cepA, from Bacteroides fragilis reveals a new subgroup of Ambler
class A beta-lactamases. Antimicrob Agents Chemother. 1993 Nov;37(11):2391–400. doi:
10.1128/AAC.37.11.2391

13. Rasmussen BA, Gluzman Y, Tally FP. Cloning and sequencing of the class B beta-lactamase
gene (ccrA) from Bacteroides fragilis TAL3636. Antimicrob Agents Chemother. 1990
Aug;34(8):1590–2.

14. Sydenham TV, Sóki J, Hasman H, Wang M, Justesen US, ESGAI (ESCMID Study Group on
Anaerobic Infections). Identification of antimicrobial resistance genes in multidrug-resistant
clinical Bacteroides fragilis isolates by whole genome shotgun sequencing. Anaerobe. 2015
Feb;31:59–64. doi: 10.1016/j.anaerobe.2014.10.009

15. Ricker N, Qian H, Fulthorpe RR. The limitations of draft assemblies for understanding
prokaryotic adaptation and evolution. Genomics. 2012 Sep;100(3):167–75. doi:
10.1016/j.ygeno.2012.06.009

16. Page AJ, De Silva N, Hunt M, Quail MA, Parkhill J, Harris SR, et al. Robust high-throughput
prokaryote de novo assembly and improvement pipeline for Illumina data. Microb Genomics
[Internet]. 2016 Aug 25 [cited 2019 Apr 17];2(8). doi: 10.1099/mgen.0.000083

17. Schmid M, Frei D, Patrignani A, Schlapbach R, Frey JE, Remus-Emsermann MNP, et al.
Pushing the limits of de novo genome assembly for complex prokaryotic genomes harboring
very long, near identical repeats. Nucleic Acids Res. 2018 Sep 28;46(17):8953–65. doi:
10.1093/nar/gky726

18. Ring N, Abrahams JS, Jain M, Olsen H, Preston A, Bagby S. Resolving the complex
Bordetella pertussis genome using barcoded nanopore sequencing. Microb Genomics. 2018
Nov 21; doi: 10.1099/mgen.0.000234

19.   Wick RR, Judd LM, Gorrie CL, Holt KE. Completing bacterial genome assemblies with
multiplex MinION sequencing. Microb Genomics [Internet]. 2017 [cited 2018 Apr 3];3(10). doi:
10.1099/mgen.0.000132

20.   Maio ND, Shaw LP, Hubbard A, George S, Sanderson N, Swann J, et al. Comparison of long-
read sequencing technologies in the hybrid assembly of complex bacterial genomes. bioRxiv.
2019 Jan 28;530824. doi: 10.1101/530824

21.   Cerdeño-Tárraga AM, Patrick S, Crossman LC, Blakely G, Abratt V, Lennard N, et al.
Extensive DNA inversions in the B. fragilis genome control variable gene expression.
Science. 2005 Mar 4;307(5714):1463–5. doi: 10.1126/science.1107008

22.   Kuwahara T, Yamashita A, Hirakawa H, Nakayama H, Toh H, Okada N, et al. Genomic
analysis of Bacteroides fragilis reveals extensive DNA inversions regulating cell surface
adaptation. Proc Natl Acad Sci U S A. 2004 Oct 12;101(41):14919–24. doi:
10.1073/pnas.0404172101

23.   Patrick S, Blakely GW, Houston S, Moore J, Abratt VR, Bertalan M, et al. Twenty-eight
divergent polysaccharide loci specifying within- and amongst-strain capsule diversity in three
strains of Bacteroides fragilis. Microbiology. 2010 Nov;156(Pt 11):3255–69. doi:
10.1099/mic.0.042978-0

24.   Nikitina AS, Kharlampieva DD, Babenko VV, Shirokov DA, Vakhitova MT, Manolov AI, et al.
Complete Genome Sequence of an Enterotoxigenic Bacteroides fragilis Clinical Isolate.
Genome Announc [Internet]. 2015 May 7;3(3). doi: 10.1128/genomeA.00450-15

25.   Risse J, Thomson M, Patrick S, Blakely G, Koutsovoulos G, Blaxter M, et al. A single
chromosome assembly of Bacteroides fragilis strain BE1 from Illumina and MinION nanopore
sequencing data. GigaScience [Internet]. 2015 Dec 4 [cited 2017 Jun 22];4. doi:
10.1186/s13742-015-0101-6

26.   Soki, Jozsef. Bacteroides fragilis S14 genome sequencing and assembly (Data accessed on
NCBI RefSeq database accession GCF_001682215.1). 2015.

27.   Ho P-L, Yau C-Y, Wang Y, Chow K-H. Determination of the mutant–prevention concentration
of imipenem for the two imipenem–susceptible Bacteroides fragilis strains, Q1F2 (cfiA-
positive) and ATCC 25282 (cfiA-negative). Int J Antimicrob Agents. 2018 Feb 1;51(2):270–1.
doi: 10.1016/j.ijantimicag.2017.08.004

28.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018 Sep 1;34(17):i884–90. doi: 10.1093/bioinformatics/bty560

29.   Desai A, Marwah VS, Yadav A, Jha V, Dhaygude K, Bangar U, et al. Identification of
Optimum Sequencing Depth Especially for De Novo Genome Assembly of Small Genomes
Using Next Generation Sequencing Data. PLoS ONE [Internet]. 2013 Apr 12 [cited 2014 May
22];8(4). doi: 10.1371/journal.pone.0060204

30.   Wick RR, Judd LM, Holt KE. Deepbinner: Demultiplexing barcoded Oxford Nanopore reads
with deep convolutional neural networks. bioRxiv. 2018 Sep 14;366526. doi: 10.1101/366526

31.   De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack: visualizing and
processing long-read sequencing data. Bioinformatics. 2018 Aug 1;34(15):2666–9. doi:
10.1093/bioinformatics/bty149

32. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. Bioinformatics. 2013 Apr 15;29(8):1072–5. doi: 10.1093/bioinformatics/btt086

33. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res. 2015 Jul 1;25(7):1043–55. doi: 10.1101/gr.186072.114

34. Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, et al. BUSCO
Applications from Quality Assessments to Gene Prediction and Phylogenomics. Mol Biol Evol.
2018 Mar 1;35(3):543–8. doi: 10.1093/molbev/msx319

35. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014 Jul
15;30(14):2068–9. doi: 10.1093/bioinformatics/btu153

36. Richter M, Rosselló-Móra R. Shifting the genomic gold standard for the prokaryotic species
definition. Proc Natl Acad Sci. 2009 Oct 22;pnas.0906412106. doi:
10.1073/pnas.0906412106

37. Clark SC, Egan R, Frazier PI, Wang Z. ALE: a generic assembly likelihood evaluation
framework for assessing the accuracy of genome and metagenome assemblies.
Bioinformatics. 2013 Feb 15;29(4):435–43. doi: 10.1093/bioinformatics/bts723

38. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo
genome assemblies. Bioinforma Oxf Engl. 2015 Oct 15;31(20):3350–2. doi:
10.1093/bioinformatics/btv383

39. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018 Sep
15;34(18):3094–100. doi: 10.1093/bioinformatics/bty191

40. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinforma Oxf Engl. 2009 Jul 15;25(14):1754–60. doi: 10.1093/bioinformatics/btp324

41. Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. Circlator: automated
circularization of genome assemblies using long sequencing reads. Genome Biol. 2015 Dec
29;16:294. doi: 10.1186/s13059-015-0849-0

42. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al. NCBI
prokaryotic genome annotation pipeline. Nucleic Acids Res. 2016 19;44(14):6614–24. doi:
10.1093/nar/gkw569

43. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.
Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother. 2012
Nov;67(11):2640–4. doi: 10.1093/jac/dks261

44. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017:
expansion and model-centric curation of the comprehensive antibiotic resistance database.
Nucleic Acids Res. 2017 04;45(D1):D566–73. doi: 10.1093/nar/gkw1004

45. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference centre for
bacterial insertion sequences. Nucleic Acids Res. 2006 Jan 1;34(Database issue):D32-36.
doi: 10.1093/nar/gkj014

46.   Galata V, Fehlmann T, Backes C, Keller A. PLSDB: a resource of complete bacterial
plasmids. Nucleic Acids Res. 2019 Jan 8;47(D1):D195–202. doi: 10.1093/nar/gky1050

47.   Jørgensen TS, Xu Z, Hansen MA, Sørensen SJ, Hansen LH. Hundreds of Circular Novel
Plasmids and DNA Elements Identified in a Rat Cecum Metamobilome. PLoS ONE [Internet].
2014 Feb 4 [cited 2019 Mar 12];9(2). doi: 10.1371/journal.pone.0087924

48.   Shah HN. The Genus Bacteroides and Related Taxa. In: Balows A, Trüper HG, Dworkin M,
Harder W, Schleifer K-H, editors. The Prokaryotes: A Handbook on the Biology of Bacteria:
Ecophysiology, Isolation, Identification, Applications [Internet]. New York, NY: Springer New
York; 1992 [cited 2019 Mar 21]. p. 3593–607. doi: 10.1007/978-1-4757-2191-1_34

49.   Sóki J, Wareham DW, Rátkai C, Aduse-Opoku J, Urbán E, Nagy E. Prevalence, nucleotide
sequence and expression studies of two proteins of a 5.6kb, class III, Bacteroides plasmid
frequently found in clinical isolates from European countries. Plasmid. 2010 Mar;63(2):86–97.
doi: 10.1016/j.plasmid.2009.12.002

50.   Database resources of the National Center for Biotechnology Information. Nucleic Acids Res.
2016 Jan 4;44(Database issue):D7–19. doi: 10.1093/nar/gkv1290

51.   Trinh S, Haggoud A, Reysset G, Sebald M. Plasmids pIP419 and pIP421 from Bacteroides:
5-nitroimidazole resistance genes and their upstream insertion sequence elements. Microbiol
Read Engl. 1995 Apr;141 (Pt 4):927–35. doi: 10.1099/13500872-141-4-927

52.   Haggoud A, Trinh S, Moumni M, Reysset G. Genetic analysis of the minimal replicon of
plasmid pIP417 and comparison with the other encoding 5-nitroimidazole resistance plasmids
from Bacteroides spp. Plasmid. 1995 Sep;34(2):132–43. doi: 10.1006/plas.1995.9994

53.   Sóki J, Gal M, Brazier JS, Rotimi VO, Urbán E, Nagy E, et al. Molecular investigation of
genetic elements contributing to metronidazole resistance in Bacteroides strains. J Antimicrob
Chemother. 2006 Feb;57(2):212–20. doi: 10.1093/jac/dki443

54.   Hartmeyer GN, Sóki J, Nagy E, Justesen US. Multidrug-resistant Bacteroides fragilis group on
the rise in Europe? J Med Microbiol. 2012 Dec;61(Pt 12):1784–8. doi:
10.1099/jmm.0.049825-0

55.   Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment with Gene
Gain, Loss and Rearrangement. PLOS ONE. 2010 Jun 25;5(6):e11147. doi:
10.1371/journal.pone.0011147

56.   Nishida H. Comparative Analyses of Base Compositions, DNA Sizes, and Dinucleotide
Frequency Profiles in Archaeal and Bacterial Chromosomes and Plasmids [Internet].
International Journal of Evolutionary Biology. 2012 [cited 2019 Apr 26]. doi:
10.1155/2012/342482

57.   Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, et al. RASTtk: a modular and
extensible implementation of the RAST algorithm for building custom annotation pipelines and
annotating batches of genomes. Sci Rep. 2015 Feb 10;5:8365. doi: 10.1038/srep08365

58.   Bayley DP, Rocha ER, Smith CJ. Analysis of cepA and other Bacteroides fragilis genes
reveals a unique promoter structure. FEMS Microbiol Lett. 2000 Dec 1;193(1):149–54. doi:
10.1111/j.1574-6968.2000.tb09417.x

59. Lin Y, Yuan J, Kolmogorov M, Shen MW, Chaisson M, Pevzner PA. Assembly of long error-
prone reads using de Bruijn graphs. Proc Natl Acad Sci. 2016 Dec 27;113(52):E8396–405.
doi: 10.1073/pnas.1604560113

60. Kamath GM, Shomorony I, Xia F, Courtade TA, Tse DN. HINGE: long-read assembly
achieves optimal repeat resolution. Genome Res. 2017;27(5):747–56. doi:
10.1101/gr.216465.116

61. Payne A, Holmes N, Rakyan V, Loose M. Whale watching with BulkVis: A graphical viewer for
Oxford Nanopore bulk fast5 files. bioRxiv. 2018 May 3;312256. doi: 10.1101/312256

62. de Lannoy C, Risse J, de Ridder D. poreTally: run and publish de novo nanopore assembler
benchmarks. Bioinformatics [Internet]. [cited 2019 Apr 26]; doi:
10.1093/bioinformatics/bty1045

63. Carattoli A, Zankari E, García-Fernández A, Larsen MV, Lund O, Villa L, et al. In Silico
Detection and Typing of Plasmids using PlasmidFinder and Plasmid Multilocus Sequence
Typing. Antimicrob Agents Chemother. 2014 Jul 1;58(7):3895–903. doi: 10.1128/AAC.02412-
14

64. Arredondo-Alonso S, Rogers MRC, Braat JC, Verschuuren TD, Top J, Corander J, et al.
mlplasmids: a user-friendly tool to predict plasmid- and chromosome-derived sequences for
single species. Microb Genomics [Internet]. 2018 [cited 2019 May 9];4(11). doi:
10.1099/mgen.0.000224

65. Nguyen M, Vedantam G. Mobile genetic elements in the genus Bacteroides, and their
mechanism(s) of dissemination. Mob Genet Elem. 2011;1(3):187–96. doi:
10.4161/mge.1.3.18448

66. Shkoporov AN, Khokhlova EV, Kulagina EV, Smeianov VV, Kuchmiy AA, Kafarskaya LI, et al.
Analysis of a novel 8.9kb cryptic plasmid from Bacteroides uniformis, its long-term stability
and spread within human microbiota. Plasmid. 2013 Mar 1;69(2):146–59. doi:
10.1016/j.plasmid.2012.11.002

67. McNulty NP, Wu M, Erickson AR, Pan C, Erickson BK, Martens EC, et al. Effects of diet on
resource utilization by a model human gut microbiota containing Bacteroides cellulosilyticus
WH2, a symbiont with an extensive glycobiome. PLoS Biol. 2013;11(8):e1001637. doi:
10.1371/journal.pbio.1001637

68. Pierce JV, Bernstein HD. Genomic Diversity of Enterotoxigenic Strains of Bacteroides fragilis.
PLoS ONE [Internet]. 2016 Jun 27 [cited 2019 Mar 26];11(6). doi:
10.1371/journal.pone.0158171

69. Husain F, Veeranagouda Y, Hsi J, Meggersee R, Abratt V, Wexler HM. Two multidrug-
resistant clinical isolates of Bacteroides fragilis carry a novel metronidazole resistance nim
gene (nimJ). Antimicrob Agents Chemother. 2013 Aug;57(8):3767–74. doi:
10.1128/AAC.00386-13

70. Ruan J, Li H. Fast and accurate long-read assembly with wtdbg2. bioRxiv. 2019 Jan
26;530972. doi: 10.1101/530972

71. Li H. Minimap and miniasm: fast mapping and de novo assembly for noisy long sequences.
Bioinformatics. 2016 Jul 15;32(14):2103–10. doi: 10.1093/bioinformatics/btw152

72. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat
graphs. Nat Biotechnol. 2019 Apr 1;1. doi: 10.1038/s41587-019-0072-8

73. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome
Res. 2017 May 1;27(5):722–36. doi: 10.1101/gr.215087.116

74. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New
Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J Comput Biol.
2012 May;19(5):455–77. doi: 10.1089/cmb.2012.0021

75. Antipov D, Korobeynikov A, McLean JS, Pevzner PA. hybridSPAdes: an algorithm for hybrid
assembly of short and long reads. Bioinformatics. 2016 Apr 1;32(7):1009–15. doi:
10.1093/bioinformatics/btv688

76. Souvorov A, Agarwala R, Lipman DJ. SKESA: strategic k-mer extension for scrupulous
assemblies. Genome Biol. 2018 Oct 4;19(1):153. doi: 10.1186/s13059-018-1540-z

77. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies
from short and long sequencing reads. PLOS Comput Biol. 2017 Aug 6;13(6):e1005595. doi:
10.1371/journal.pcbi.1005595

78. Loman NJ, Quick J, Simpson JT. A complete bacterial genome assembled *de novo* using
only nanopore sequencing data. Nat Methods. 2015 Aug;12(8):733–5. doi:
10.1038/nmeth.3444

79. Vaser R, Sovic I, Nagarajan N, Sikic M. Fast and accurate de novo genome assembly from
long uncorrected reads. Genome Res. 2017 Jan 18;gr.214270.116. doi:
10.1101/gr.214270.116

80. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An Integrated
Tool for Comprehensive Microbial Variant Detection and Genome Assembly Improvement.
PLOS ONE. 2014 Nov 19;9(11):e112963. doi: 10.1371/journal.pone.0112963

81. Krumsiek J, Arnold R, Rattei T. Gepard: a rapid and sensitive tool for creating dotplots on
genome scale. Bioinforma Oxf Engl. 2007 Apr 15;23(8):1026–8. doi:
10.1093/bioinformatics/btm039

82. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. Bioinformatics.
2011 Apr 1;27(7):1009–10. doi: 10.1093/bioinformatics/btr039

83. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster
version of the PHAST phage search tool. Nucleic Acids Res. 2016 08;44(W1):W16-21. doi:
10.1093/nar/gkw387

 TABLES AND FIGURES

Genome assembler and version	Link	Reference
Wtdbg2 v2.3	https://github.com/ruanjue/wtdbg2	(70)
Miniasm v0.3r179	https://github.com/lh3/miniasm	(39,71)
Flye v2.3.7	https://github.com/fenderglass/Flye	(59,72)
Canu v1.8	https://github.com/marbl/canu	(73)
Spades (including Hybridspades) v3.13.0	https://github.com/ablab/spades	(74,75)
Skesa v2.3.0	https://github.com/ncbi/SKESA	(76)
Unicycler v0.4.7	https://github.com/rrwick/Unicycler	(77)
Assembly polishing tools		
Nanopolish v0.10.2	https://github.com/jts/nanopolish	(78)
Racon v1.3.1	https://github.com/isovic/racon	(79)
Pilon v1.22	https://github.com/broadinstitute/pilon	(80)

797 Table 1 - Genome assemblers and polishing tools tested

Assembly	Contigs	Largest contig	Total length	Mis-assemblies	Genome fraction (%)	Mismatches per 100 kbp	Indels per 100 kbp	Average nucleotide identity	CheckM Completeness	BUSCOs: complete and single-copy/ complete and duplicate/ fragment (of 443)	Prokka genes	Prokka rRNA	Prokka tRNA	Total ALE score
GCF_000025985.1	2	5,205,140	5,241,700	0	100.000	0	0	100.0000	99.26	442/0/1	4439	19	73	-17071758.95
Skesa	46	553,341	5,201,945	3	99.237	0.23	0.15	99.9980	99.26	440/2/1	4391	2	62	-20926329.69
Spades	23	1,779,941	5,212,217	4	99.396	0.44	0.17	99.9866	99.26	440/2/1	4407	3	56	-19676529.39
Canu.OF.CO.RO2.RI.PI3	2	5,247,938	5,350,432	8	99.972	4.94	15.9	99.9752	99.26	442/0/1	4634	19	73	-19283611.73
Flye.OF.CS.PI5.RI	5	2,282,650	5,269,269	4	99.917	1.07	6.24	99.9781	99.26	441/1/1	4476	19	73	-18222322.23
Miniasm.OF.CM.RO2.PI5	3	5,204,445	5,277,434	2	99.972	5.21	17.75	99.9691	98.88	442/0/1	4607	19	73	-17789234.97
Wtdbg2.OF.CO.RO2.PI6.RI	3	5,192,352	5,234,448	7	99.723	3.23	3.04	99.9807	99.26	442/0/1	4437	19	73	-18750266.21
SpadesHybrid.CS	5	3,093,122	5,242,724	7	99.987	1.89	0.53	99.9856	99.26	440/2/1	4441	19	73	-18535980.68
Unicycler.OF.CS	2	5,205,133	5,241,693	2	99.972	0.84	0.48	99.9997	99.26	442/0/1	4435	19	73	-17200232.52

799 Table 2. Selected quality indicators for the best genome assembly of *B. fragilis* CCUG4856T per assembly pipeline. RefSeq accession
800 GCF_000025985.1 was used as reference. OF: ONT reads filtered with Filtlong, CS: Canu corrected standard settings, CM: Canu
801 corrected with option corMinCoverage=0, CO: Canu corrected with option coroutCoverage=999, RO2: Two rounds of Racon polishing
802 with ONT reads, RI: Racon polishing with Illumina reads, PI[n]: Pilon polishing with Illumina reads, [n] rounds. Full results are available in
803 Supplementary Table S2.

Assembly	Total length (bp)	Largest contig (bp)	Local mis-assem blies	Genome fraction (%)	Mismatc hes per 100 kbp	Indels per 100 kbp	K-mer-based compl. (%)	K-mer-based misjoins	Average nucletide identity	Prokka CDS	Prokka genes	Total ALE score
GCF_0000 25985.1	5,241,700	5,205,140	0	100.00 0	0	0	100.00	0	100.00000	4346	4439	-17071758.95
OF	5,241,602	5,205,042	3	99.970	1.11	0.65	99.96	0	99.99896	4343	4436	-17245134.52
OF.RI	5,241,606	5,205,046	3	99.970	1.09	0.67	99.96	3	99.99887	4345	4438	-17247815.86
OF.CS	5,241,693	5,205,133	2	99.972	0.84	0.48	99.97	1	99.99974	4342	4435	-17200232.52
OF.CS.RI	5,241,698	5,205,138	2	99.972	0.88	0.52	99.96	1	99.99968	4346	4439	-17206271.66
OF.CM	5,241,691	5,205,131	2	99.972	0.88	0.5	99.96	1	99.99966	4343	4436	-17201292.44
OF.CM.RI	5,241,696	5,205,136	2	99.972	0.95	0.55	99.97	1	99.99975	4343	4436	-17193184.79
OF.CO	5,241,693	5,205,133	2	99.972	0.84	0.48	99.97	1	99.99974	4342	4435	-17200232.52
OF.CO.RI	5,241,698	5,205,138	2	99.972	0.88	0.52	99.96	1	99.99968	4346	4439	-17206271.66

Table 3. Hybrid Unicycler assemblies of *B. fragilis* CCUG4856T. RefSeq accession GCF_000025985.1 was used as reference. OF: ONT
reads filtered with Filtlong; CS: Canu corrected standard settings; CM: Canu corrected with option corMinCoverage=0; CO: Canu
corrected with option coroutCoverage=999; RI: Racon polishing with Illumina reads. Unicycler performs assembly polishing with Racon
(ONT reads) and Pilon. Full results are available in Supplementary Table S2.

Figure 1. Dot plot matrix of the alignment of the reference assembly and the hybrid Unicycler
assembly using Gepard v1.40 (81). The *B. fragilis* NCTC9343 (RefSeq GCF_000025985.1)
reference assembly derived from Sanger sequencing is on the x-axis and the hybrid Unicycler
assembly on the y-axis. On this otherwise near perfect alignment with high similarity, an 88,045 bp
inversion with 100% ID is observed at nucleotide positions 2,941,962..3,030,006 on the Unicycler
assembly (2,005,742..2,093,786 on the reference sequence) (indicated by the blue arrow).

Strain	Best Spades graph	Hybrid assembly after first nanopore sequencing run. No Canu correction	Hybrid assembly after first nanopore sequencing run with Canu correction	Hybrid assembly with data from supplementary runs with Canu correction	After manual finishing	Final assemblies contig statistics		
						No.	Length	Relative read depth
CCUG 4856T						1	5,205,133	1.00x
						2	36,560	7.42x
BFO17						1	5,474,541	1.00x
						2	85,671	1.85x
						3	5,594	23.03x
BFO18						1	5,302,644	1.00x
						2	7,221	25.98x
						3	4,137	50.80x
						4	2,782	59.91x
S01						1	5,325,251	1.00x
						2	78,085	2.29x
						3	8,331	20.99x
						4	5,595	22.67x
BFO42						1	5,141,257	1.00x
						2	8,306	40.06x
						3	5,594	40.15x
BFO67						1	5,478,614	1.00x
						2	6,129	94.15x
BFO85						1	5,504,076	1.00x

Figure 2. Evolution of genome assemblies with added data and manual finishing. The best SPAdes
assembly graphs by Unicycler with short reads only are shown on the far left. Supplying ONT
reads improved the assemblies overall, but only three were circularised with singular chromosome
contigs with data from the initial MinION sequencing runs. Adding additional ONT data and
correcting reads with Canu did not improve assemblies for all isolates. Manual finishing was
necessary to finish assemblies for three isolates. Assembly graph images generated with
Bandage. Read information can be found in Supplementary Table S1.

| Strain | Sequence | Length (bp) | Relative read depth | GC% | PLSDB results | | | | Plasmid replicon family (%COV, %ID) |
					Best hit accession no.	Plasmid hit name	%ID	Length of the sequence of best hit (bp)	
CCUG4856T	Chr.	5,205,133	1.00x	43.19	-	-	-	-	-
	pBF9343	36,560	7.42x	32.19	NC_006873.1	pBF9343	100	36,560	Rep_3 (100/100)
BFO17	Chr.	5,474,541	1.00x	43.51	-	-	-	-	-
	pBFO17_1	85,671	1.85x	36.78	NC_006873.1	pBF9343	80.7	36,560	none
	pBFO17_2	5,594	23.03x	39.65	NC_011073.1	pBFP35	99.9	5,594	Rep_1 (100/100)
BFO18	Chr.	5,302,644	1.00x	43.34	-	-	-	-	-
	pBFO18_1	7,221	25.98x	42.32	NC_015168.1	pBACSA02	85.6	19,280	Rep_3 (99.69/99.69)
	pBFO18_2	4,137	50.80x	45.40	NC_019534.1	pBFUK1	92.2	12,817	Rep_3 (100.00/98.24)
	pBFO18_3	2,782	59.91x	41.45	NC_005026.1	pBI143	94.6	2,747	RepL (89.66/49.22)[a]
S01	Chr.	5,325,251	1.00x	43.57	-	-	-	-	-
	pBFS01_1	78,085	2.29x	36.04	NC_006873.1	pBF9343	80.7	36,560	none
	pBFS01_2	8,331	20.99x	41.17	NC_015166.1	pBACSA03	95.6	6,277	Rep_3 (100.00/97.85)
	pBFS01_3	5,595	22.67x	39.62	NC_011073.1	pBFP35	99.9	5,594	Rep_1 (100.00/99.48)
BFO42	Chr.	5,141,257	1.00x	43.35	-	-	-	-	-
	pBFO32_1	8,306	40.06x	43.34	KJ830768.1	pBF69566b	96.0	11,019	RHH_1 (92.94/64.63) Rep_3 (93.64/68.31)
	pBFO32_2	5,594	40.15x	39.63	NC_011073.1	pBFP35	99.9	5,594	Rep_1 (100.00 /99.48)
BFO67	Chr.	5,478,614	1.00x	43.85	-	-	-	-	-
	pBFO67_1	6,129	94.15x	41.67	NC_011073.1	pBFP35	76.9	5,594	Rep_3 (100.00/99.69)
BFO85	Chr.	5,504,076	1.00x	43.60	-	-	-	-	-

Table 4. Putative plasmid sequences of the complete *B. fragilis* assemblies. Putative plasmid sequences from the hybrid assemblies of *B.*
fragilis CCUG4856T and the six MDR *B. fragilis* isolates were screened using the PLSDB. The best hit to plasmids from cultured isolates
is shown. Only three putative plasmids from the MDR *B. fragilis* isolate assemblies could be identified with confident %ID. For most
sequences, plasmid replication family proteins were identified in the putative plasmids using ABRicate with a database of sequences

downloaded from the Pfam database, strengthening the interpretation that the circularised putative plasmid sequences do in fact
represent plasmids harboured by the isolates. Notes: [a]Annotated as RepA protein in the PGAP annotation. Abbreviations: %ID, %COV,
no.; number, Chr.; chromosome.

Figure 3. Linear representation of an alignment of putative circular plasmid sequences pBFO17_1 pBFS01_1. Comparison of the putative circular plasmids pBFO17_1 and pBFS01_1 (reverse complement for better visualisation) using EasyFig (82). EasyFig uses BLAST to identify sequences of similarity. Sequence similarities of >98% is indicated by full colouring, a darker colour indicates a higher %ID. Products of annotated CDS' are shown. CDS' annotated as hypothetical or Domain of Unknown Function are coloured white. The two sequences show a very high degree of similarity. pBFO17_1 is 7,586 bp longer than pBFS01_1. This is mainly due to the insertion of a reverse transcriptase (pBFO17_1, 11367..13034) (disrupting a DNA methylase), the insertion of prophage from position 56125 to 61162) (identified as an incomplete prophage using PHASTER (83)) and an IS*1380* family-like transposase (67933..69237). The regions pBFO17_1 50711..52501 and pBFS01_1 32248..30304 are not similar. Possibly, the insertion of two transposases in pBFO17_1 have excised most of the ParB-family DNA partitioning protein in the corresponding sequence range in pBFS01_1.

Antimicrobial susceptibility[a]				Antimicrobial resistance genes and IS elements[b]					
Strain	Anti­mi­crobi­al	Etest MIC (mg/L)	Result	Gene	Upstream IS element	Sequenc e[c]	%ID	%COV	Associated resistance to drug class
BFO17	MEM	>32	R	cfiA11	IS614B	Chr	100.00	99.20	Carbapenem
	IPM	>32	R						
	MTZ	>32	R	nimJ	IS614B	Chr	99.40	100.00	Nitroimidazole
				nimJ	IS614B	Chr c	99.40	100.00	Nitroimidazole
	CLI	0.094	S						
	PTZ	>256	R						
				tetQ		Chr	99.34	99.34	Tetracycline
				cfxA4		Chr	85.71	100.00	Cephamycin
				bexB		Chr c	91.21	100.00	Fluoroquinolone
				bexA		Chr	73.77	99.02	Fluoroquinolone
BFO18	MEM	>32	R	cfiA2_1	ISBf12	Chr	100.00	100.00	Carbapenem
	IPM	16	R				100.00	100.00	
	MTZ	16	R	nimD	IS1169	2	99.19	100.00	Nitroimidazole
	CLI	6	R	ermF[d]	IS4351	Chr c	99.83	72.03	Clindamycin
					ISBthe1[d]	Chr c	70.97	97.19	
				erm(F)[d]		Chr c	99.58	29.71	
				lnu(AN2)		Chr c	100.00	100.00	Clindamycin
	PTZ	>256	R						
				ugd		Chr	65.69	53.04	Polymyxin
				bexA		Chr c	73.60	99.02	Fluoroquinolone
				bexB		Chr	91.14	100.00	Fluoroquinolone
				tet(Q)		Chr c	99.79	100.00	Tetracycline
				mef(En2)		Chr c	99.83	100.00	Macrolides
S01	MEM	>32	R	cfiA13_1	IS1187	Chr	99.20	100.00	Carbapenem
	IPM	16	R						
	MTZ	64	R	nimE	ISBf6	3	100.00	100.00	Nitroimidazole
	CLI	>32	R	erm(F)	IS1187	Chr	99.50	100.00	Clindamycin
	PTZ	6	S						
				tetQ		Chr	90.02	99.95	Tetracycline
				tet(Q)		Chr c	99.84	100.00	Tetracycline
				bexB		Chr c	91.06	100.00	Fluoroquinolone
				bexA		Chr	74.03	98.80	Fluoroquinolone
BFO42	MEM	0.094	S						
	IPM	0.25	S						
	MTZ	8	R	nimA	ISBf13	2	98.64	96.61	Nitroimidazole
	CLI	>256	R	erm(F)	IS613	Chr	99.50	100.00	Clindamycin

				Gene	IS element	Location	%ID	%COV	Antimicrobial
				lnu(AN2)		Chr	100.00	100.00	Clindamycin
	PTZ	0.38	S						
				ugd		Chr	70.38	31.45	
				cepA-49		Chr$_c$	100.00	100.00	Cephalosporin
				mef(En2)		Chr	99.83	100.00	Macrolide
				ugd		Chr	71.15	31.11	Polymyxin
				tetQ		Chr$_c$	100.00	100.00	Tetracycline
				bexB		Chr	99.12	100.00	Fluoroquinolone
				ere(D)		Chr	96.66	100.00	Erythromycin
				aadS		Chr$_c$	99.88	100.00	Aminoglycoside
				OXA-347	IS*613*	Chr$_c$	100.00	100.00	Penicillin, cephalosporin
				bexA		Chr	75.09	99.62	Fluoroquinolone
BFO67	MEM	8	R	*cfiA13_1*	None	Chr	100.00	100.00	Carbapenem
	IPM	0.5	S						
	MTZ	0.19	S						
	CLI	0.38	S						
	PTZ	2	S						
				cfxA2	IS*Bf11*	Chr	99.69	100.00	Cephamycin
				mef(En2)		Chr	99.75	100.00	Macrolide
				lnu(AN2)		Chr	100.00	100.00	Clindamycin
				ugd		Chr$_c$	66.76	56.30	Polymyxin
				tet(Q)		Chr	100.00	100.00	Tetracycline
				bexB		Chr$_c$	90.92	100.00	Fluoroquinolone
				bexA		Chr	73.90	99.02	Fluoroquinolone
BFO85	MEM	32	R	*cfiA2_1*	None	Chr	100.00	100.00	Carbapenem
	IPM	1	S						
	MTZ	0.25	S						
	CLI	>256	R	*ermB*		Chr$_c$	99.19	98.66	Clindamycin
	PTZ	2	S						
				ugd		Chr	69.84	31.45	Polymyxin
				tetQ		Chr$_c$	90.02	99.95	Tetracycline
				aadE		Chr$_c$	100.00	100.00	Aminoglycoside
				aad9		Chr$_c$	100.00	100.00	Aminoglycoside
				bexB		Chr$_c$	90.92	100.00	Fluoroquinolone
				bexA		Chr	73.53	99.02	Fluoroquinolone
				cfxA2	IS*614*	Chr$_c$	100.00	100.00	Cephamycin
				tet(Q)		Chr$_c$	99.84	100.00	Tetracycline

Table 5. Antimicrobial susceptibility and resistance genes and IS elements for the six MDR *B.*
fragilis strains. Identified genes are displayed next to the relevant antimicrobials. Identified IS
elements in correct orientation (opposite strand) directly upstream of the genes are included. The
%ID and %COV refer to the gene hit. Hits with %ID or %COV <98% were confirmed with BLASTx

searches. The hits for *ugd* represent possible homologs for genes coding for PmrE, which is
involved in polymyxin resistance in Gram-negative bacteria. Full ABRicate results with nucleotide
positions and information on the IS elements is available the Supplementary Tables S4 AMR-IS-
results.xlsx. Notes: [a] Results from previously published work following EUCAST breakpoints (14). [c]
A [c] denotes complement strand. [d] A transposase has inserted itself, splitting the *ermF* gene in two.
Abbreviations: %ID; percent identity, %COV; coverage percentage, Chr; chromosome.

Strain	Best Spades graph	Hybrid assembly after first nanopore sequencing run. No Canu correction	Hybrid assembly after first nanopore sequencing run with Canu correction	Hybrid assembly with data from supplementary runs with Canu correction	After manual finishing	Final assemblies contig statistics		
						No.	Length	Relative read depth
CCUG 4856T						1	5,205,133	1.00x
						2	36,560	7.42x
BFO17						1	5,474,541	1.00x
						2	85,671	1.85x
						3	5,594	23.03x
BFO18						1	5,302,644	1.00x
						2	7,221	25.98x
						3	4,137	50.80x
						4	2,782	59.91x
S01						1	5,325,251	1.00x
						2	78,085	2.29x
						3	8,331	20.99x
						4	5,595	22.67x
BFO42						1	5,141,257	1.00x
						2	8,306	40.06x
						3	5,594	40.15x
BFO67						1	5,478,614	1.00x
						2	6,129	94.15x
BFO85						1	5,504,076	1.00x