Discovery of an essential mycobacterial polar growth factor involved in cell envelope

integrity and lipomannan and lipoarabinomannan transport

Kuldeepkumar R. Gupta[a], Kathryn C. Rahlwes[b], Chunyan Wang[c], Jun Liu[c], Yasu S.

Morita[b], and E. Hesper Rego[a,1]

[a]Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven,

CT 06519

[b]Department of Microbiology, University of Massachusetts, Amherst, MA 01003

[c]Department of Microbial Pathogenesis, Microbial Sciences Institute, Yale University

School of Medicine, New Haven, CT, USA 06519

[1]To whom correspondence should be addressed: hesper.rego@yale.edu

ABSTRACT

Mycobacteria grow and divide differently compared to well-studied model bacteria. They

insert new cell envelope at their poles instead of along their side walls; and, they lack

obvious homologs of many well-conserved cell growth and division proteins. Furthermore,

while mycobacteria share several similarities to Gram-positive bacteria, unlike these

organisms they possess an outer membrane, which is abundant in long-chain fatty acids

and several glycolipids and lipoglycans. A better understanding of the unique factors that

make this unusual structure could lead to therapeutic targets for pathogenic

mycobacteria. Here, we study a gene of previously unknown function – *msmeg_0317* -

predicted to be essential and associated with mycobacterial cell growth and division

proteins. We find that transcriptional depletion of *msmeg_0317* leads to loss of polar

growth, disruption of the mycobacterial outer membrane, and cell death. Surprisingly, we also observe that depletion results in the accumulation of cell-associated lipoglycans lipomannan and lipoarabinomannan (LM/LAM), while overexpression of CwdA leads to increased shedding of LM/LAM. LM/LAM have been extensively studied in relation to infection but their role in bacterial physiology is less clear. Altogether, our data suggest that MSMEG_0317, renamed CwdA, is involved in the transport of LM/LAM to the mycobacterial outer membrane, and reveal unexpected connections between the correct localization of LM/LAM, polar growth, and the structural integrity of the mycobacterial cell envelope.

IMPORTANCE. Some of the most successful antibiotics target bacterial cell envelope synthesis. However, the cell envelope of mycobacteria is significantly different from that of other, more well-studied, bacteria. Its core structure consists of a covalently-linked network of peptides, carbohydrates, and fatty acids. Additionally, there are several lipids non-covalently interspersed throughout. Many of the enzymes which synthesize these lipids are known, but how they are transported remains largely unclear. Here, we discover that an essential protein, CwdA, is involved in the transport of LM/LAM, abundant lipoglycans in the mycobacterial cell envelope. Depletion of CwdA leads to the loss of polar growth and the disintegration of the mycobacterial outer membrane. These results suggest that the proteins which transport molecules across the mycobacterial cell envelope may represent an abundant source of novel drug targets for the treatment of mycobacterial infections, like tuberculosis.

INTRODUCTION

Mycobacterium tuberculosis (Mtb), the etiological agent of human tuberculosis (TB), is responsible for approximately 1.4 million deaths each year. One of the pathogen's distinguishing features is its unusual cell envelope. Like nearly all other bacterial species, the plasma membrane is surrounded by peptidoglycan, a rigid mesh-like structure made up of carbohydrate chains crosslinked by peptide bridges. However, in contrast to the peptidoglycan of other well-characterized bacteria, mycobacterial peptidoglycan is covalently linked to the highly branched hetero-polysaccharide arabinogalactan, which is, itself, covalently bound to extremely long-chained fatty acids called mycolic acids. Electron microscopy has revealed that the outer cell envelope is a lipid bilayer, and, as such, it is referred to as the outer membrane, or, alternatively, the mycomembrane (1, 2). In addition to this core structure, several lipids are abundantly and non-covalently interspersed across the plasma membrane and outer membrane (3, 4). This complex cell envelope is a double-edged sword: it is a formidable barrier to many antibiotics, yet provides a number of potentially targetable structures. Indeed, two of the four first-line TB antibiotics target cell-envelope biosynthesis.

In addition to their unusual cell envelope, mycobacteria differ from other well-studied rod-shaped bacteria in important ways. Notably, mycobacteria elongate from their poles rather than along their side walls. The details of polar growth are poorly understood (5-7). Many key proteins important for cell elongation in other species appear to be missing in mycobacteria (6, 8). Likewise, it is almost completely unknown how proteins involved in synthesizing the different layers, and free lipids, work together to build this complex cell envelope.

Given the unusual physiology of mycobacteria, perhaps it is unsurprising that

greater than 25% of the genes encoded by the mycobacterial genome are annotated as

having unknown function (9). In our previous studies on the mycobacterial division

complex, 'the divisome', we precipitated known divisome and elongasome components

from *M. smegmatis* (Msm), and identified multiple candidate mycobacterial-specific

divisome proteins in the precipitate, (8, 10). One of these was MSMEG_0317, a protein

of unknown function that is highly conserved in actinobacteria. The Mtb homolog of

msmeg_0317, *rv0227c*, is predicted to be essential (11, 12). Here, we sought to

understand the role of MSMEG_0317 during mycobacterial growth and division.

MSMEG_0317 belongs to the DUF3068-domain super-family of proteins, which

are exclusively found in actinobacteria. One member of this family in corynebacteria has

channel activity, so this domain has been renamed PorA (13, 14). In addition, by structure-

based homology prediction, MSMEG_0317 shares limited homology to mycobacterial

lipoproteins, LppX and LrpG, which transport lipids, phthiocerol dimycocerosates (PDIMs)

and triacylglycerol (TAG), respectively, to the outer mycobacterial membrane (15, 16).

We find that MSMEG_0317 localizes to sites of new cell wall synthesis and is

indeed essential for cell viability in Msm. Transcriptional depletion leads to loss of cellular

elongation, reductive division, and eventually cell lysis. Surprisingly, we find that

lipomannan (LM), and lipoarabinomannan (LAM) – two large lipoglycans abundant in the

mycobacterial cell envelope - accumulate as MSMEG_0317 is depleted and are shed into

the culture media more abundantly upon MSMEG_0317 overexpression. By cryo-electron

microscopy, we find that the cell envelopes of depleted cells are devoid of outer

membrane and are highly disrupted. Together, these results show that MSMEG_0317

plays an important role in LM/LAM transport to the outer membrane, polar elongation,

and the structural organization of the mycobacterial cell wall. As such, we propose to

rename *msmeg_0317*, *cwdA*, for <u>c</u>ell <u>w</u>all <u>d</u>isruption, mutant A.

RESULTS

***cwdA* is essential for polar growth**

To determine the essentiality of *cwdA*, we used an allele swapping strategy (17).

Briefly, in a strain of Msm whose only copy of *cwdA* was at the L5 phage integration site

(18), we exchanged *cwdA* for either an empty vector or for another copy of itself (Fig. 1A).

Consistent with *cwdA* being essential for cell growth, we observed approximately 1000-

fold fewer colonies when we exchanged *cwdA* with an empty vector compared to

exchanging it for another copy of itself (Fig. 1A).

To further confirm the essentially of *cwdA*, we constructed a strain in which the

only copy of the gene was tetracycline inducible (Fig. 1B). Removal of anhydrotetracycline

(ATC) from the culture media prevented cell growth (Fig. 1B). By time-lapse microscopy

we observed that, while cells expressing CwdA became longer, on average, as cell

density increased within the microfluidic device (Fig. 2A,C & Movie S1), cells depleted for

CwdA stopped elongating but continued to divide (reductive division), became wider, and,

in many instances, eventually lysed (Fig. 2B,C & Movie S2). Together these data show

that *cwdA* is essential for polar elongation in Msm.

CwdA localizes to the sites of cell wall synthesis

As our data suggested CwdA is important for polar growth, we hypothesized that

it would localize to the poles, which are the sites of new cell wall synthesis in

mycobacteria. To test this hypothesis, we made a translational fusion to CwdA with mRFP

expressed by its native promoter. By allele swapping, we find that *cwdA-mrfp* restores

bacterial growth (Fig. S1) and thus encodes a functional *cwdA*. Fluorescence microscopy

at a single time-point showed that CwdA-mRFP localizes to mid-cell and to the poles (Fig.

3A). In polar growing bacteria, the site of division eventually becomes the site of

elongation. Thus, in addition to localizing to the poles, elongation-complex proteins will

also localize to the site of division early during the next cell cycle, making it difficult to

distinguish true septal localization from static images. To disentangle whether CwdA also

localizes to the septum, we visualized CwdA-mRFP by time-lapse microscopy. By

constructing kymographs averaged over many cells, we compared the fluorescence

probability distribution over the course of the cell cycle to the earliest known markers for

the division complex (FtsZ-mCherry2B) and the elongation complex (eGFP-Wag31).

CwdA-mRFP localizes to the site of division after FtsZ, but before Wag31 (Fig. 3B). These

data are consistent with CwdA being a member of both the mycobacterial division and

elongation complexes, but its essential function is as part of the elongation complex.

CwdA is involved in LM/LAM transport

The corynebacterial homolog of *cwdA* was recently implicated in LM/LAM

biosynthesis (19), as a deletion mutant of the *Corynebacterium glutamicum* homolog

resulted in disappearance of LAM and accumulation of truncated LM, which was clearly

detectable by faster migration on SDS-PAGE. To test if this is also the case in

mycobacteria, we used the same electrophoretic approach to analyze LM/LAM, and thin

layer chromatography (TLC) to examine potential changes in other lipid species upon

depletion of CwdA, before and after the onset of cell death (Fig. S2B). In contrast to the

results obtained in *C. glutamicum*, we observe a dramatic and transient increase in the

total amount of cell-associated of LM/LAM during depletion (Figs. 4A & S2A), up until the

onset of cell death. There were no obvious changes in the migration of LM/LAM on SDS-

PAGE, implying that the impact of CwdA depletion on LM/LAM biosynthesis was minimal.

Additionally, other lipids, including the precursor of LM/LAM biosynthesis such as

Ac_1PIM_2, showed no reproducible change (Figs. 4B,C & Fig. S2A).

These data are not consistent with the proposed biosynthetic function of the

corynebacterial CwdA homolog. Instead, the transient increase of LM/LAM prior to cell

death suggests a potential role for CwdA in cell envelope trafficking of these lipoglycans.

LM/LAM are thought to be embedded in the plasma membrane, and also transported to

the outer membrane (20, 21). Furthermore, LM/LAM in the outer membrane are likely to

be shed into the extracellular milieu (22-24). Interestingly, sequence-based homology

modeling indicated that CwdA shares some homology to LppX and LprG, proteins known

to transport lipids from the inner membrane to the outer membrane in mycobacteria.

Therefore, we hypothesized that depletion of CwdA results in the transient increase of

cell-associated LM/LAM due to the disruption of LM/LAM trafficking. Based on this

hypothesis, we reasoned that over-expression of CwdA may facilitate more trafficking,

which could deplete cell-associated LM/LAM and increase shedding. To test this, we

constructed strains which inducibly expressed CwdA from a multi-copy plasmid.

Consistent with our hypothesis, upon induction with ATC we observed less LM/LAM

associated with Msm cells (Figs. S3A & S4) and more LAM secreted in the culture

supernatant (Fig. S3B). These data suggest a model whereby CwdA is involved, directly

or indirectly, in the transport of LM/LAM to the outer membrane of mycobacteria.

CwdA is important for cell envelope integrity

Our model predicts that CwdA-depleted cells should have greatly altered cell

envelopes. Thus, we visualized the cell envelope of wild type and mutant cells by high

resolution cryo-electron microscopy (cryo-EM). As has been previously observed (1, 2),

wild type Msm cells exhibit a distinct plasma membrane and outer membrane that is

approximately 50-nm thick (Fig. 5A-C). Consistent with our model, we find that cells

depleted for CwdA have largely unstructured cell walls, and appear devoid of electron

density in the outer membrane (Fig. 5D-F). We also observe dramatic fraying of the outer

membrane, and large invaginations of plasma membrane (Fig. 6). Thus, CwdA is

important for maintaining the structural organization of the mycobacterial cell envelope.

Collectively, our data show that CwdA is an essential polar growth factor involved

in transporting, either directly or indirectly, LM/LAM to the outer membrane. Further, our

data suggest that the correct localization of LM/LAM is important for the structural

organization and integrity of the mycobacterial cell envelope.

DISCUSSION

The extraordinary diversity of the bacterial kingdom has led to a plethora of

bacterial cell shapes and geometries of cell growth and division. Unlike well-studied

model bacteria, several organisms, including the major human pathogen *Mycobacterium*

tuberculosis, grow from their poles rather than their side-walls. Our understanding of polar

growth is at a nascent stage compared to our knowledge of side-wall growth.

Mycobacteria, and related organisms, have an additional challenge of orchestrating the

biosynthesis and transport of several additional covalently and non-covalently attached

lipids and glycans to their thick layer of peptidoglycan. How these organisms coordinate

the biogenesis of this complex structure to elongate and divide is almost completely

unknown.

Here, we have studied a highly conserved mycobacterial protein, MSMEG_0317,

in Msm and propose to rename it CwdA. As predicted (19), we find that the gene which

encodes this protein is essential in Msm. The Mtb homolog*, rv0227c,* is almost certainly

essential as well, as no transposon insertions have been found in this gene, even in data

sets which combine insertion counts from high density mutagenesis across multiple

independent experiments (11).

Our data suggest that CwdA's essential function occurs during polar elongation,

as cells depleted for CwdA stop elongating, and CwdA-mRFP localizes to the poles.

Interestingly, CwdA also localizes to the site of division, yet depleted cells continue to

divide. This is consistent with recent work from several groups, which suggests that the

essentiality of certain cell envelope structures is different for elongation and division. For

example, *C. glutamicum* cells treated with ethambutol, which inhibits the polymerization

of arabinan, a core structure in both arabinogalactan and LAM, stop elongating, but

continue to divide (25, 26).

What is the function of CwdA? Our work suggests that CwdA is involved in the

transport of LM/LAM to the outer membrane, either directly or indirectly, and is important

for maintaining the structural integrity of the mycobacterial cell envelope. Upon depletion,

the levels of several other lipids remain constant or show no reproducible change,

suggesting that CwdA may be acting on LM/LAM directly, though our lipid profiling was

not exhaustive and we cannot rule out the possibility that another structural lipid is

affected by CwdA depletion. However, supporting the notion that CwdA specifically acts

upon LM/LAM, we observe a temporal correlation, up until the time of cell lysis, between

the accumulation of LM/LAM and the disruption of the envelope. There has been a good

deal of speculation as to the localization of LM/LAM in the mycobacterial envelope, but

several groups have now shown that at least a fraction of LM/LAM resides in the outer

membrane (23, 24). Our data are consistent with a model in which LM/LAM are structural

components of the outer membrane. However, mutants of Msm that produce diminished

amounts or truncated versions of LM/LAM are viable, suggesting that the presence of

LM/LAM in the outer membrane may not be essential for viability (27, 28). Why, then, is

cwdA essential? We speculate that the accumulation of these large lipoglycans in the

periplasm of CwdA-depleted cells causes envelope disruption and cell rupture.

One implication from our work is that the transport or correct localization of lipids

to the outer membrane is needed for polar elongation, but not division, suggesting that

there could be various feedback loops to ensure complete construction of the elaborate

mycobacterial cell envelope for insertion of new material to elongate. These checkpoints

may be missing during division. This notion is supported by other mutants defective in

transporting important outer membrane components like trehalose monomycolate, which,

like the *cwdA* mutant, undergo reductive cell division (29).

The mycobacterial cell envelope remains an enduring target for the development

of new TB therapeutics. The standard TB therapy uses a combination of four drugs; two

of these drugs target cell wall synthesis. In addition to the core, and covalently linked

structure of peptidoglycan, arabinogalactan, and mycolic acids, the mycobacterial cell

envelope is also rich with many other lipids. The functional role of these lipids in bacterial

cell physiology is relatively unexplored but they are likely important, if not essential,

components of the cell envelope. Our data argue that a more complete understanding of

the transport of these molecules could lead to new targets for the development of anti-TB

therapeutics.

Materials and Methods

Bacterial strains and culture conditions: Msm mc^{2}155 was grown in

Middlebrook 7H9 broth supplemented with 0.05% Tween80, 0.2% glycerol, 5 gm/L

albumin, 2 gm/L dextrose and 0.003 gm/L catalase or plated on LB agar. *Escherichia coli*

DH5α cells were grown in LB broth or on LB agar plates. Concentrations of antibiotics

used for Msm is as follows: 20 μg/ml zeocin, 25 μg/ml kanamycin, 50 μg/ml hygromycin,

and 20 μg/ml nourseothricin. Concentrations of antibiotics used for *E. coli* used for *E. coli*

is as follows: 40 μg/ml zeocin, 50 μg/ml kanamycin, 100 μg/ml hygromycin, and 40 μg/ml

nourseothricin.

Plasmids and strains construction: Before deleting the native copy of

msmeg_0317, a merodiploid strain was created by inserting a second copy of

msmeg_0317 gene under pTetO promoter using a Kan^R L5 integrating vector.

Subsequently, in the merodiploid strain, an in-frame deletion was made by replacing the

native copy of *msmeg_0317 gene* with a zeocin resistance cassette flanked by loxP sites

using recombineering (18). The plasmid constructs were made using isothermal Gibson

assembly whereby insert and vector backbone shared 20-25 bp of homology.

msmeg_0317 and its fluorescent variant were cloned in a Nat[R] L5 integrating vector for

allele exchange at the L5 site. The *msmeg_0317* depletion strain was made by

transforming an episomal Hyg[R] marked vector constitutively expressing TetR repressor

into a strain expressing a single copy of *msmeg_0317* controlled by the pTetO promoter.

mRFP was fused to the c-terminal of *msmeg_0317* gene cloned with 200 bp upstream

sequence to capture the native promoter. *ftsZ-mCherry*2B and *eGFP-wag31* genes were

cloned under medium strength promoter ptb21 and integrated at tweety (tw) and L5 phage

integration sites, respectively.

MSMEG_0317 Depletion: To transcriptionally deplete *msmeg_0317* in Msm, the

ATC-inducible Tet-ON system was used. The only copy of *msmeg_0317* was driven by

the pTetO promoter, while the TetR repressor was constitutively expressed in trans from

an episomal vector. MSMEG_0317 was depleted by removing ATC from the medium and

cells were grown for 18 hours. Subsequently, MSMEG_0317 depleted cells were re-

diluted in fresh medium, also without ATC, and samples at different timepoints were taken

and processed for cyro-electron microscopy. Alternatively, to avoid changes in LM/LAM

abundance that have been found to correlated with cell density and growth phase (21),

at 18 hours of depletion, half of the culture was removed for lipid extraction and analysis,

and replaced with fresh media. Lipid analysis was performed at timepoints before and

after the decrease in OD indicating cessation of growth (Fig. S3).

Time-lapse microscopy: An inverted Nikon Ti-E microscope was used for the

time-lapse imaging. An environmental chamber was maintained at 37°C. Cells were

cultivated in an B04 microfluidics plate from CellAsic; continuously supplied with fresh

medium; and imaged every 15 minutes using a 1.4 N.A. phase contrast objective.

Kymograph analysis. Time-lapse images were analyzed in open source image

analysis software Fiji (30) and a custom MATLAB program was used to generate

kymographs. Specifically, for a single cell, in Fiji, a 5-pixel wide segmented line was drawn

from the new pole to the old pole at each time point during the cell cycle. This was

repeated on 20-50 cells. These line profiles were then imported into MATLAB, where a

custom script was used to generate an average kymograph by 2D interpolation of the

individual kymographs.

Lipid extraction and analysis. Extraction, purification and analysis of lipids were

as described previously (31). Briefly, crude lipids were extracted from equal wet cell pellet

weights of the MSMEG_0317 depletion strains after 18, 24, 27, 30, and 33 hrs with or

without ATC. After lipid extraction using chloroform/methanol, LM and LAM were

extracted from the delipidated pellet by incubation with phenol/water (1:1) for 2 hrs at

55°C. Phospholipids and PIMs extracted by chloroform/methanol were further purified by

n-butanol/water phase partitioning, and separated by high performance thin layer

chromatography (HPTLC) (Silica gel 60, EMD Merck) using chloroform/methanol/ 13 M

ammonia/1 M ammonium acetate/ water (180:140:9:9:23). Phospholipids were visualized

via cupric acetate staining. PIMs were visualized with orcinol staining as described (32).

LM/LAM samples were separated by SDS-PAGE (15% gel) and visualized using ProQ

Emerald 488 glycan staining kit (Life Technologies). To detect LM/LAM in culture

supernatants, the supernatants were initially treated with a final concentration of 50 μg/ml

Proteinase K for 4 hours at 50°C. The treated supernatants were electrophoresed on 15%

SDS-PAGE. LM/LAM were blotted onto nitrocellulose at 20 V for 45 minutes using semi-

dry transfer method. Post-transfer, the membrane was blocked by 5% milk in Tris-

buffered saline supplemented with 0.05% Tween-20 (TBST). The blocked membrane was

then probed overnight with CS-35 antibody (BEI Resources, NIH) at 1:250 dilution at 4°C.

The membrane was washed with TBST five times for five minutes each. Post-washing, it

was probed with anti-mouse secondary for one hour at room temperature. Membrane

was then washed five times for five minutes with TBST. Thermo Scientific's west dura

chemiluminescent reagent was used to develop the membrane.

Sample preparation and image collection for Cryo-EM: Wild type and

MSMEG_0317 depleted Msm were pelleted, washed twice with 1X phosphate buffered

saline (PBS), and suspended in a very small volume of PBS to get a very dense

suspension of bacterial cells. The dense culture was subsequently mixed with 10-nm

colloidal gold and deposited onto fresh glow-discharged holey carbon grids. The grids

were then blotted with filter paper manually and rapidly frozen in liquid ethane. The frozen

grids were transferred under -170 °C into a 300kV Titan Krios equipped with K2 Summit

direct detector, a quantum energy filter (Gatan, Pleasanton, CA) and an FEI volta phase

plate (FEI). The data was collected using SerialEM (33) and Digital Micrograph

software. The motion correction for each movie was done with MotionCor2 (34). The

microscope was operated at 300kV with a magnification of 3,600x and 26,000 x for low-

mag map and high-mag map respectively. The grey levels of each micrograph are

obtained using MATLAB.

Acknowledgements

We thank Cara Boutte and Maikel Boot for helpful suggestions and for critical reading of

the manuscript. Work carried out in this manuscript was supported by career award at

scientific interface (CASI) from Burroughs Welcome Fund and Searle and Pew funds to

E.H.R.

References:

1. **Hoffmann C, Leis A, Niederweis M, Plitzko JM, Engelhardt H.** 2008. Disclosure of the mycobacterial outer membrane: cryo-electron tomography and vitreous sections reveal the lipid bilayer structure. Proc Natl Acad Sci U S A **105:**3963-3967.
2. **Zuber B, Chami M, Houssin C, Dubochet J, Griffiths G, Daffe M.** 2008. Direct visualization of the outer membrane of mycobacteria and corynebacteria in their native state. J Bacteriol **190:**5672-5680.
3. **Jackson M.** 2014. The mycobacterial cell envelope-lipids. Cold Spring Harb Perspect Med **4**.
4. **Jankute M, Cox JA, Harrison J, Besra GS.** 2015. Assembly of the Mycobacterial Cell Wall. Annu Rev Microbiol **69:**405-423.
5. **Kieser KJ, Rubin EJ.** 2014. How sisters grow apart: mycobacterial growth and division. Nat Rev Microbiol **12:**550-562.
6. **Hett EC, Rubin EJ.** 2008. Bacterial growth and cell division: a mycobacterial perspective. Microbiol Mol Biol Rev **72:**126-156, table of contents.
7. **Baranowski C, Rego EH, Rubin EJ.** 2019. The Dream of a Mycobacterium. Microbiol Spectr **7**.
8. **Wu KJ, Zhang J, Baranowski C, Leung V, Rego EH, Morita YS, Rubin EJ, Boutte CC.** 2018. Characterization of Conserved and Novel Septal Factors in Mycobacterium smegmatis. J Bacteriol **200**.
9. **Doerks T, van Noort V, Minguez P, Bork P.** 2012. Annotation of the M. tuberculosis hypothetical orfeome: adding functional information to more than half of the uncharacterized proteins. PLoS One **7:**e34302.
10. **Rego EH, Audette RE, Rubin EJ.** 2017. Deletion of a mycobacterial divisome factor collapses single-cell phenotypic heterogeneity. Nature **546:**153-157.
11. **DeJesus MA, Gerrick ER, Xu W, Park SW, Long JE, Boutte CC, Rubin EJ, Schnappinger D, Ehrt S, Fortune SM, Sassetti CM, Ioerger TR.** 2017. Comprehensive Essentiality Analysis of the Mycobacterium tuberculosis Genome via Saturating Transposon Mutagenesis. MBio **8**.

12.   **Zhang YJ, Ioerger TR, Huttenhower C, Long JE, Sassetti CM, Sacchettini JC, Rubin**
EJ. 2012. Global assessment of genomic regions required for growth in Mycobacterium
tuberculosis. PLoS Pathog **8:**e1002946.

13.   **Soltan Mohammadi N, Mafakheri S, Abdali N, Barcena-Uribarri I, Tauch A, Benz R.**
2013. Identification and characterization of the channel-forming protein in the cell wall of
Corynebacterium amycolatum. Biochim Biophys Acta **1828:**2574-2582.

14.   **Abdali N, Younas F, Mafakheri S, Pothula KR, Kleinekathofer U, Tauch A, Benz R.**
2018. Identification and characterization of smallest pore-forming protein in the cell wall
of pathogenic Corynebacterium urealyticum DSM 7109. BMC Biochem **19:**3.

15.   **Sulzenbacher G, Canaan S, Bordat Y, Neyrolles O, Stadthagen G, Roig-Zamboni V,**
Rauzier J, Maurin D, Laval F, Daffe M, Cambillau C, Gicquel B, Bourne Y, Jackson
M. 2006. LppX is a lipoprotein required for the translocation of phthiocerol
dimycocerosates to the surface of Mycobacterium tuberculosis. EMBO J **25:**1436-1444.

16.   **Martinot AJ, Farrow M, Bai L, Layre E, Cheng TY, Tsai JH, Iqbal J, Annand JW,**
Sullivan ZA, Hussain MM, Sacchettini J, Moody DB, Seeliger JC, Rubin EJ. 2016.
Mycobacterial Metabolic Syndrome: LprG and Rv1410 Regulate Triacylglyceride Levels,
Growth Rate and Virulence in Mycobacterium tuberculosis. PLoS Pathog **12:**e1005351.

17.   **Pashley CA, Parish T.** 2003. Efficient switching of mycobacteriophage L5-based
integrating plasmids in Mycobacterium tuberculosis. FEMS Microbiol Lett **229:**211-215.

18.   **van Kessel JC, Hatfull GF.** 2007. Recombineering in Mycobacterium tuberculosis. Nat
Methods **4:**147-152.

19.   **Cashmore TJ, Klatt S, Yamaryo-Botte Y, Brammananth R, Rainczuk AK,**
McConville MJ, Crellin PK, Coppel RL. 2017. Identification of a Membrane Protein
Required for Lipomannan Maturation and Lipoarabinomannan Synthesis in
Corynebacterineae. J Biol Chem **292:**4976-4986.

20.   **Rahlwes KC, Sparks IL, Morita YS.** 2019. Cell Walls and Membranes of Actinobacteria.
Subcell Biochem **92:**417-469.

21.   **Dhiman RK, Dinadayala P, Ryan GJ, Lenaerts AJ, Schenkel AR, Crick DC.** 2011.
Lipoarabinomannan localization and abundance during growth of Mycobacterium
smegmatis. J Bacteriol **193:**5802-5809.

22.   **Pitarque S, Larrouy-Maumus G, Payre B, Jackson M, Puzo G, Nigou J.** 2008. The
immunomodulatory lipoglycans, lipoarabinomannan and lipomannan, are exposed at the
mycobacterial cell surface. Tuberculosis (Edinb) **88:**560-565.

23.   **Gaur RL, Ren K, Blumenthal A, Bhamidi S, Gibbs S, Jackson M, Zare RN, Ehrt S,**
Ernst JD, Banaei N. 2015. Correction: LprG-Mediated Surface Expression of
Lipoarabinomannan Is Essential for Virulence of Mycobacterium tuberculosis. PLoS
Pathog **11:**e1005336.

24.   **Shukla S, Richardson ET, Athman JJ, Shi L, Wearsch PA, McDonald D, Banaei N,**
Boom WH, Jackson M, Harding CV. 2014. Mycobacterium tuberculosis lipoprotein
LprG binds lipoarabinomannan and determines its cell envelope localization to control
phagolysosomal fusion. PLoS Pathog **10:**e1004471.

25.   **Zhou X, Rodriguez-Rivera FP, Lim HC, Bell JC, Bernhardt TG, Bertozzi CR, Theriot**
JA. 2019. Sequential assembly of the septal cell envelope prior to V snapping in
Corynebacterium glutamicum. Nat Chem Biol **15:**221-231.

26.   **Schubert K, Sieger B, Meyer F, Giacomelli G, Bohm K, Rieblinger A, Lindenthal L,**
Sachs N, Wanner G, Bramkamp M. 2017. The Antituberculosis Drug Ethambutol
Selectively Blocks Apical Growth in CMN Group Bacteria. MBio **8**.
27.   **Fukuda T, Matsumura T, Ato M, Hamasaki M, Nishiuchi Y, Murakami Y, Maeda Y,**
Yoshimori T, Matsumoto S, Kobayashi K, Kinoshita T, Morita YS. 2013. Critical roles
for lipomannan and lipoarabinomannan in cell wall integrity of mycobacteria and
pathogenesis of tuberculosis. MBio **4**:e00472-00412.
28.   **Kaur D, McNeil MR, Khoo KH, Chatterjee D, Crick DC, Jackson M, Brennan PJ.**
2007. New insights into the biosynthesis of mycobacterial lipomannan arising from
deletion of a conserved gene. J Biol Chem **282**:27133-27140.
29.   **Fay A, Czudnochowski N, Rock J, Johnson JR, Krogan NJ, Rosenberg O, Glickman**
MS. 2019. Two accessory proteins govern MmpL3 mycolic acid transport in mycobacteria.
bioRxiv doi:10.1101/581447:**581447**.
30.   **Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,**
Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V,
Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-
image analysis. Nat Methods **9**:676-682.
31.   **Rahlwes KC, Puffal J, Morita YS.** 2019. Purification and Analysis of Mycobacterial
Phosphatidylinositol Mannosides, Lipomannan, and Lipoarabinomannan. Methods Mol
Biol **1954**:59-75.
32.   **Sena CB, Fukuda T, Miyanagi K, Matsumoto S, Kobayashi K, Murakami Y, Maeda**
Y, Kinoshita T, Morita YS. 2010. Controlled expression of branch-forming
mannosyltransferase is critical for mycobacterial lipoarabinomannan biosynthesis. J Biol
Chem **285**:13326-13336.
33.   **Mastronarde DN.** 2005. Automated electron microscope tomography using robust
prediction of specimen movements. J Struct Biol **152**:36-51.
34.   **Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA.** 2017.
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
microscopy. Nat Methods **14**:331-332.
35.   **Larrouy-Maumus G, Gilleron M, Skovierova H, Zuberogoitia S, Brennan PJ, Puzo**
G, Jackson M, Nigou J. 2015. A glycomic approach reveals a new mycobacterial
polysaccharide. Glycobiology **25**:1163-1171.
434
435

436
Fig. 1. *cwdA* is essential in Msm. (A) L5-integrating vectors carrying either *cwdA* or a
gene encoding TetR were transformed into a strain whose only copy of *cwdA* was at the
L5 integration site, and transformants were counted. Bars represent means of biological
triplicates and are representative data of three independently performed experiments. (B)
Optical density over time of a strain whose only copy of *cwdA* is tetracycline inducible,
with and without ATC (+/-ATC). Points are means of three biological replicates. Error bars
are too small to be visible. Experiment was performed at least 4 times on separate days,
and representative data are shown. ATC=anhydrotetracycline. Time on the *x*-axis
corresponds to time after ATC was removed.
446
447

Fig. 2. CwdA-depletion causes reductive cell division and eventual cell lysis. (A,B)
The strain shown in Fig. 1B was visualized by time-lapse microscopy in the presence (A)
or absence (B) of ATC. Cells were loaded into a microfluidic device 18 hours after the
removal of ATC (B) or a mock control (A). Representative frames are 3 hours apart. (C)
Birth lengths and cell widths over time of depletion (orange = +ATC; red = -ATC). Open
circles represent individual cells (length: N = 261 (length) & 253 (width) for +ATC; N =
355 (length) & 254 (width) -ATC). Closed circles are not fits to the data but represent the
mean of cells in 2.25-hr (length) or 2.75-hr (width) bins. The experiment was performed
twice on separate days. In +ATC cultures, cell density became too high to measure length
and width past ~33 hrs. ATC=anhydrotetracycline

459
460
461

Fig. 3. CwdA localizes to the sites of division and elongation. (A) Merged phase-contrast and fluorescence images of Msm cells expressing CwdA-mRFP, eGFP-Wag31 and FtsZ-mCherry2b. (B) Average fluorescent distributions over time (kymographs) from cells aligned from new to old poles and from birth to division (N=20 CwdA-mRFP; N=48 FtsZ-mCherry2B; N=20 eGFP-Wag31).

469
470
Fig. 4. LM/LAM accumulate as CwdA is depleted. In a strain whose only copy of *cwdA*
is ATC-inducible, (A) LM/LAM, (B) non-polar and polar PIMs, (C) and plasma membrane
lipids were analyzed by (A) SDS-PAGE and (B-C) TLC during CwdA depletion (without
ATC). Cell pellets were normalized by wet weight to account for differences in cell density.
Two other biological replicates are shown in Fig. S2. CL, cardiolipin; PE,
phosphatidylethanolamine; PI, phosphatidylinositol.
477
478

480 **Fig. 5. Depletion of CwdA leads to loss of outer membrane.** The cell envelope of a
481 representative wild type Msm cell (A-C) and a cell depleted for CwdA for 24 hours were
482 (D-F) visualized (A,B,D,E), and measured (C,F) by cryo-electron microscopy. PM =
483 plasma membrane; OM = outer membrane. Scale bar = 100 nm.

486
487 **Fig. 6. Cryo-electron microscopy of CwdA-depleted cells** (A-H) The cell envelopes of
488 CwdA-depleted cells show plasma membrane invaginations (A&E; C&G; D&H) and outer
489 membrane wall fraying (B-D; F-H) (arrows). (A-C; E-G) 24 hours; (D&H) 33 hours of
490 depletion. Scale bar = 100 nm.
491
492
493

Supplemental Information

Figure S1: CwdA-mRFP encodes a functional copy of CwdA

Figure S2: Two additional biological replicates of LM/LAM analysis during CwdA depletion

Figure S3: Detection of cell-associated and shed LM/LAM during CwdA over-expression

Figure S4: An additional biological replicate of cell-associated LM/LAM and other lipids

during CwdA over expression

Movie S1. Cells whose only copy of *cwdA* is ATC-inducible are imaged over time by

phase-contrast microscopy in the presence of ATC. Cells were treated in parallel to those

shown in Movie S2, except ATC was included in the culture media at all times.

Movies S2. Cells whose only copy of *cwdA* is ATC-inducible, are imaged over time by

phase-contrast microscopy in the absence of ATC. Cells were loaded into the device 18

hours after the removal of ATC.

Table S1: A list of strains used in this study.

Strain	Usage
Mycobacterium smegmatis mc^2 155	Wild type strain, lab stock
mc2 155 L5::pTetO-*cwdA*-kan^R	Merodiploid strain used for deleting native copy of *cwdA*
mc2 155 ΔcwdA L5::pTetO-*cwdA*-kan^R	Parental strain used for allele exchange of *cwdA* at L5 site
mc2 155 ΔcwdA L5::pTetO-cwdA/pTetR	ATC inducible *cwdA* strain where *cwdA* is under TetO promoter and the strain carries constitutive TetR repressor; used for transcriptional depletion of *cwdA*
mc2 155 ΔcwdA L5::pTetO-MS0317-myc/pTetR	ATC inducible C-terminal myc-tagged *cwdA* strain where *cwdA* is under TetO promoter and the strain carries

	constitutive TetR repressor; used for transcriptional depletion of *cwdA*
mc2 155 /pTetOR-*cwdA*-myc	ATC inducible C-terminal myc-tagged *cwdA* strain; used for over-expression of CwdA protein
mc2 155 Δ*cwdA* pL5:pNative-*cwdA*-mRFP-myc	Strain carrying C-terminal mRFP tagged CwdA protein expressed from native promoter to track the localization of CwdA protein
mc2155 L5::pG-MCK-ptb21-egfp-wag31 tweety::ptb21-ftsZ-mCherry2B	Strain used to track the localization of Wag31 and FtsZ proteins (Rego et al 2017, *Nature*)

513 Table S2: A list of primers used in this study

MS_0317_FP	GCTTAATTAAGAAGGAGATATACATAT GAACCGCGCTGTGGCGCTGCGTATCG	Forward primer to amplify *cwdA* from Msm genome
MS_0317_RP	GGTCCCCAATTAATTAGCTAAAGCTTTC AGATCGGTCGGTCCGGTGGCAGATCG	*Reverse* primer to amplify *cwdA* from Msm genome
MS_0317_1	GGAGCGCAGGTAGTGCCCGGTTCTCG	Forward primer to amplify the upstream region for *cwdA* knock-out construct
MS_0317_2	GGCGTTACCCAACTTAATCGCCTTGCA GCTCACCCCAGAAGTCCACATGCCGCG	Reverse primer to amplify the upstream region for *cwdA* knock-out construct
MS_0317_3	CGCGGCATGTGGACTTCTGGGGTGAGC TGCAAGGCGATTAAGTTGGGTAACGCC	forward primer to amplify zeo^R cassette for

		cwdA knock-out construct
MS_0317_4	GCACCTGGATGCCCTGCGTATCGAA GAAACAGCTATGACCATGATTACGCC	reverse primer to amplify zeo^R cassette for *cwdA* knock-out construct
MS_0317_5	GGCGTAATCATGGTCATAGCTGTTTC TTCGATACGCAGGGCATCCAGGTGC	Forward primer to amplify the downstream region for *cwdA* knock-out construct
MS_0317_6	AGACTCCAGTGGCCCTGCAACAACCGC	Reverse primer to amplify the downstream region for *cwdA* knock-out construct
MS0317_Nat_Fwd	CGATCCGCATGCTTAATTAAG AAGGAGATATACTAATGAACCGCG CTGTGGCGCTGCGT	Forward primer for sub-cloning *cwdA* into L5 integrating pTetO-Nat^R vector
MS0317_Nat_Rev	CCTCTAGGGTCCCCAATTAATT AGCTAAAGCTTTCAGATCGGTC GGTCCGGTGGCAGATCG	Reverse primer for sub-cloning *cwdA* into L5 integrating pTetO-Nat^R vector
MS0317_KO_Fwd	GGATCACGATCACCGCCACCAC GTAGC	Forward sequencing primer to confirm deletion of *cwdA* gene
MS0317_KO_Rev	CCAGAACACCAGCGGAGCCACC GGAGTCC	Reverse sequencing primer to confirm

		deletion of *cwdA* gene
MS0317-Nat_myc-Rev	TCCCCAATTAATTAGCTAAAGCTT TCACAGGTCTTCCTCGCTGATCAGC TTCTGCTCGATCGGTCGGTCCGGTGGC	Reverse primer for attaching myc tag at C-terminus of *cwdA*
pNative317-RFP_1	CGATCCGCATGCTTAATTAAGAAGGAG ATATACTAATGAACCGCGCTGTGGCGCTGCGT	Gibson primers used for making *pNative_ cwdA _mrfp* construct
pNative317-RFP_2	CCTTGATGACGTCCTCGGAGGAGGC CGAGCCGCCGATCGGTCGGTCCGGT GGCAGATCGG	Gibson primers used for making *pNative_ cwdA _mrfp* construct
pNative317-RFP_3	CCGATCTGCCACCGGACCGACCGAT CGGCGGCTCGGCCTCCTCCGAGGACGTCATCAAGG	Gibson primers used for making *pNative_ cwdA _mrfp* construct
pNative317-RFP_4	GGTCCCCAATTAATTAGCTAAAGCTT TCACAGGTCTTCCTCGCTGATC AGCTTCTGCTCGGCGCCGG TGGAGTGGCGGCCCTCGGCGC	Gibson primers used for making *pNative_ ms_0317_mrfp* construct

Supplemental Figures

Fig. S1. CwdA-mRFP encodes a functional copy. L5-allele swapping with *cwdA* or *cwdA-mRFP* expressed by the native promoter. Bars represent means of 3 biological replicates. Error bars indicate standard deviation.

A

Replicate 1
hours without ATC

+ATC 18 24 27 30 33 marker +ATC 18 24 27 30 33

Replicate 2
hours without ATC

LAM
LM

Ac_2PIM_2
Ac_1PIM_2
Ac_2PIM_6
Ac_1PIM_6

CL
PE
PI
Ac_1PIM_2

B

OD_{600nm}

hours without ATC

527
528
Fig. S2. Two additional biological replicates of LM/LAM analysis during CwdA
depletion. (A) As in Main Text Figure 4, LM/LAM and other lipids were extracted and
separated by SDS-PAGE and TLC in cells expressing (+ATC) and depleted for (-ATC)
CwdA. Two biological replicates are shown. (B) The optical density of the cultures was
measured as CwdA was depleted. To avoid changes in LM/LAM due to cell density, at
the indicated timepoints, half of the culture was taken for lipid analysis, and replaced with
fresh media. Filled circles indicate means; error bars indicate standard deviation.
536

537
538
539
Fig. S3. Detection of cell-associated and shed LM/LAM during CwdA over-
expression. In cells carrying an ATC-inducible copy of *cwdA* on a multicopy plasmid,
(A) cell-associated LM/LAM were extracted, separated via SDS-PAGE, and visualized
by ProQ Emerald with and without ATC. (B) The culture supernatants of those cells
were also probed for LAM by western blot using an anti-LAM antibody (CS-35, BEI
resources) specific to the arabinan moiety of LAM. The high molecular weight species
marked by * likely corresponds to the recently described LAM-like polysaccharide (35).
547

548
549

550
551
552
553

Fig. S4. An additional biological replicate of cell-associated LM/LAM and other lipids during CwdA over expression. As in Fig S3, cell-associated were separated and visualized by SDS-PAGE. Other lipids were also separated and visualized by TLC.