**Title: An essential periplasmic protein coordinates lipid trafficking and is required for asymmetric polar growth in mycobacteria**

**Authors:** Kuldeepkumar R. Gupta[a], Celena M. Gwin[a], Kathryn C. Rahlwes[b], Kyle J. Biegas[d,e], Chunyan Wang[a,c], Jin Ho Park[a], Jun Liu[a,c], Benjamin M. Swarts[d,e], Yasu S. Morita[b], and E. Hesper Rego[a,1]

**Affiliations**

[a] Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT 06519

[b] Department of Microbiology, University of Massachusetts, Amherst, MA 01003

[c] Microbial Sciences Institute, Yale University, West Haven, CT 06516

[d] Department of Chemistry and Biochemistry, Central Michigan University, Mount Pleasant, MI 48859

[e] Biochemistry, Cell, and Molecular Biology Program, Central Michigan University, Mount Pleasant, MI 48859

[1]To whom correspondence should be addressed: hesper.rego@yale.edu

**ABSTRACT**

Mycobacteria, including the human pathogen *Mycobacterium tuberculosis*, grow by

inserting new cell wall material at their poles. This process and that of division are

asymmetric, producing a phenotypically heterogeneous population of cells that respond

non-uniformly to stress (Aldridge et al., 2012; Rego et al., 2017; Richardson et al., 2016).

Surprisingly, deletion of a single gene – *lamA* – leads to more symmetry, and to a

population of cells that is more uniformly killed by antibiotics (Rego et al., 2017). How

does LamA create asymmetry? Here, using a combination of quantitative time-lapse

imaging, bacterial genetics, and lipid profiling, we find that LamA recruits essential

proteins involved in cell wall synthesis to one side of the cell – the old pole. One of these

proteins, MSMEG_0317, here renamed PgfA, was of unknown function. We show that

PgfA is a periplasmic protein that interacts with MmpL3, an essential transporter that flips

mycolic acids in the form of trehalose monomycolate (TMM), across the plasma

membrane. PgfA interacts with a TMM analog suggesting a direct role in TMM transport.

Yet our data point to a broader function as well, as cells with altered PgfA levels have

differences in the abundance of other lipids and are differentially reliant on those lipids for

survival. Overexpression of PgfA, but not MmpL3, restores growth at the old poles in cells

missing *lamA*. Together, our results suggest that PgfA is a key regulator of polar growth

and cell envelope composition in mycobacteria, and that the LamA-mediated recruitment

of this protein to one side of the cell is a required step in the establishment of cellular

asymmetry.

**INTRODUCTION**

     *Mycobacterium tuberculosis* (Mtb), the etiological agent of human tuberculosis (TB), is responsible for approximately 1.4 million deaths each year. One of the pathogen's distinguishing features is its unusual cell envelope. Like nearly all other bacterial species, the plasma membrane is surrounded by peptidoglycan, a rigid mesh-like structure made up of carbohydrate chains crosslinked by peptide bridges. However, in contrast to the peptidoglycan of other well-characterized bacteria, mycobacterial peptidoglycan is covalently linked to the highly branched hetero-polysaccharide arabinogalactan, which is, itself, covalently bound to extremely long-chained fatty acids called mycolic acids. Collectively, this structure is known as the mycolyl-arababinogalactan-peptidoglycan complex or mAGP. Electron microscopy has revealed that the outer most layer is a lipid bilayer, and, as such, it is referred to as the outer membrane, or, alternatively, the mycomembrane (Hoffmann et al., 2008; Zuber et al., 2008). In addition to this core structure, several lipids, lipoglycans, and glycolipids, are abundantly and non-covalently interspersed across the plasma membrane and mycomembrane (Jackson, 2014; Jankute et al., 2015). This complex cell envelope is a double-edged sword: it is a formidable barrier to many antibiotics yet provides several potentially targetable structures. Indeed, two of the four first-line TB antibiotics target cell envelope biosynthesis.

     In addition to their unusual cell envelope, mycobacteria differ from other well-studied rod-shaped bacteria in important ways. Notably, mycobacteria elongate by adding new material at their poles rather than along their side walls. Over the course of a cell cycle, one pole grows more than the other, giving rise to an asymmetric growth pattern (Aldridge et al., 2012). Importantly, closely related organisms like corynebacteria that

have similar cell wall architecture grow more evenly from their poles, suggesting that

asymmetry is not simply a consequence of polar growth and may be actively created by

mycobacteria (Rego et al., 2017). In fact, while the molecular details of asymmetric polar

growth are not yet well understood (Baranowski et al., 2019; Kieser and Rubin, 2014), we

have discovered that LamA, a protein of unknown function specific to the mycobacterial

genus, is involved (Rego et al., 2017). Deletion of *lamA* results in more growth from the

pole formed from the previous round of division, "the new pole", and less from the

established growth pole, "the old pole", leading to less asymmetry (Fig. 1A) (Rego et al.,

2017).

To understand LamA's role in mycobacterial division and elongation, we identified

multiple putative LamA-interacting proteins of known and unknown function (Rego et al.,

2017). One of these proteins, MSMEG_0317, is predicted to be associated with several

other divisome proteins (Wu et al., 2018) (Fig. 1A). Attempts to delete *msmeg_0317* from

the *Mycobacterium smegmatis* chromosome have been unsuccessful (Cashmore et al.,

2017), and transposon insertion mapping has predicted the *M. tuberculosis* homolog,

*rv0227c*, to be essential (DeJesus, M. A. et al., 2017; Zhang et al., 2012). MSMEG_0317

belongs to the DUF3068-domain super-family of proteins, which are exclusively found in

actinobacteria. One member of this family in corynebacteria has channel activity, so this

domain has been renamed PorA (Abdali et al., 2018; Soltan Mohammadi et al., 2013). In

addition, by structure-based homology prediction, MSMEG_0317 shares limited

homology to CD36, which transports long-chained fatty acids into eukaryotic cells (Patel

et al., 2022). In corynebacteria, the putative homolog of MSMEG_0317, LmcA, is thought

to have lipid binding activity (Patel et al., 2022), and is involved in an ill-defined step of

lipoglycan synthesis (Cashmore et al., 2017; Patel et al., 2022).

Here, we sought to understand the role of MSMEG_0317 in relation to LamA

during mycobacterial growth and division. We find that in *M. smegmatis*, a model

mycobacterial species, MSMEG_0317 is essential for polar growth, and localizes to the

old pole in a LamA-dependent manner. There, it localizes with MmpL3, the essential

mycolic acid flippase, to build the mycomembrane. Our data suggest that one function of

MSMEG_0317 is to traffic mycolic acids in the periplasmic space, and that its

overexpression is sufficient to restore growth at the old pole in cells missing *lamA*. With

only 21% sequence identity to MSMEG_0317, the putative corynebacterial homolog

LmcA cannot substitute for MSMEG_0317, suggesting that these proteins are not

orthologs, and that the function of MSMEG_0317 is specific to mycobacteria. Together,

these results implicate MSMEG_0317 in mycolic acid trafficking, and argue that it is an

important regulator of cell wall composition and incorporation at one site of growth – the

old pole – in mycobacteria. As such, we propose to rename MSMEG_0317, PgfA, for

Polar growth factor A.

**RESULTS**

**PgfA is essential for polar growth and localizes to the sites of new cell wall**

**synthesis**

To verify the essentiality of *pgfA* in *M. smegmatis*, we used an allele swapping

strategy (Pashley and Parish, 2003). Briefly, in a strain whose only copy of *pgfA* was at

the L5 phage integration site (Lewis and Hatfull, 2000; van Kessel and Hatfull, 2007), we

exchanged *pgfA* for either an empty vector or for another copy of itself (Fig. 1B).

Consistent with *pgfA* being essential for cell growth, we observed approximately 1000-

fold fewer colonies when we exchanged *pgfA* with an empty vector compared to

exchanging it for another copy of itself (Fig. 1B).

To visualize the morphology of *pgfA*-depleted cells, we constructed a strain in

which the only copy of the gene was tetracycline inducible (Fig. 1C; Figure 1 – figure

supplement 1A). Removal of anhydrotetracycline (ATC) from the culture media prevented

cell growth (Figure 1 – figure supplement 1A). By time-lapse microscopy we observed

that, while cells expressing PgfA became longer, on average, as cell density increased

within the microfluidic device (Fig. 1C; Figure 1 – figure supplement 1B) cells depleted for

PgfA stopped elongating but continued to divide (reductive division), became wider, and,

in many instances, eventually lysed (Fig. 1C, Figure 1 – figure supplement 1B).

To determine where and when PgfA functions in the cell, we fused PgfA to mRFP

and expressed the resulting chimera from the native *pgfA* promoter. By allele swapping,

we find that P$_{native}$-*pgfA-mrfp* restores bacterial growth (Figure 1 – figure supplement 1C)

and thus encodes a functional PgfA. Fluorescence microscopy at a single time-point

showed that PgfA-mRFP localizes to mid-cell and to the poles (Fig. 1D, Figure 1 – figure

supplement 2A). In polar growing bacteria, the site of division eventually becomes the site

of elongation. Thus, in addition to localizing to the poles, elongation-complex proteins can

also appear at mid-cell before daughter cell separation is clearly observed. This makes it

difficult to determine true septal-associated localization from a single time point. To

disentangle whether PgfA, in addition to localizing to the poles, also localizes to the

**Figure 1. PgfA is an essential polar growth factor that localizes asymmetrically. (A)** *top:* Graphical depiction of growth pattern in WT and Δ*lamA* cells. Green=old cell wall material. Grey= new cell wall material. *bottom*: LamA is a membrane protein that co-immuno-precipitates with PonA1, a bifunctional penicillin binding protein, and MSMEG_0317/PgfA, a protein of unknown function. **(B)** Schematic and results of allelic exchange experiment. Vectors with *pgfA* or without *pgfA* (*tetR*) were transformed into a strain whose only copy of *pgfA* was at the L5 integration site. Transformants carrying the incoming vectors were counted by colony forming units (CFU). **(C)** A strain whose only copy of PgfA is tetracycline-inducible was imaged over time with (+pgfA) or without (-pgfA) anhydrotetracycline (ATC). Cells were loaded into a microfluidic device 18 hours after the removal of ATC (bottom) or a mock control (top). **(D)** Cells whose only copy of PgfA was fused to mRFP, were imaged over time by phase and fluorescence microscopy in a microfluidic device with constant perfusion of media. **(E)** *Top*. Individual cells (N=25) were followed from birth to division and the fluorescence was measured from new to old pole. Each resulting kymograph was interpolated over cell length and time and then averaged together. Using this analysis, we find that PgfA is first at the old poles (pink triangles), partially re-localizes to the septum (yellow triangles) during cell division, and then disappears from the site of division before the next cell cycle (blue triangles) to establish asymmetry in the next generation. *Bottom*. A depiction of this localization pattern is shown as a cartoon.

septum during division, we visualized PgfA-mRFP by time-lapse microscopy (Fig. 1D).

For a single cell, we measured the fluorescence distribution over time as a function of

both cell cycle time and cell length. As we wanted to compare across strains, we averaged

many of these individual trajectories together. The resulting 'average' kymograph

represents the probability of finding a fluorescent protein in a particular cellular location

at a particular stage of the cell cycle (Fig. 1E). Using this analysis, we compared the

spatiotemporal localization of PgfA-mRFP to the earliest known markers for the division

complex (FtsZ-mCherry2B) and the elongation complex (eGFP-Wag31) (Figure 1 – figure

supplement 2B). We find that PgfA localizes primarily to the old pole before the onset of

division. During division, the fluorescence intensity at the old pole becomes slightly less

intense as PgfA-mRFP re-localizes to the septum (Fig. 1D, E). This event occurs during

the latter stages of FtsZ recruitment, but before the arrival of Wag31 to the mid-cell,

suggesting that PgfA is a late divisome-associated protein (Figure 1 – figure supplement

2). At the end of division, PgfA-mRFP disappears from the septal site such that the new

daughter cells are once again born with an asymmetric distribution of PgfA (Fig. 1D, E).

Taken together, these data are consistent with PgfA being a member of both the

mycobacterial division and elongation complexes and show that PgfA is essential for polar

growth.

**PgfA and MmpL3 are recruited to the old pole by LamA to build the mycomembrane**

The manner of cell death suggested a defect in the cell envelope of PgfA-depleted

cells. To resolve the layers of the mycobacterial cell envelope in detail we used cryo-

electron microscopy (cryo-EM) to visualize frozen-hydrated cells. As has been previously

observed (Hoffmann et al., 2008; Zuber et al., 2008), wild type *M. smegmatis* cells exhibit

distinct plasma and outer membranes, both clearly observed as bilayers, separated by

approximately 50-nm (Fig. 2A). In cells depleted of PgfA, the outer membrane is frayed

(Fig. 2A and Figure 2 – figure supplement 1) and largely devoid of electron density (Fig.

2A). Thus, PgfA is important for maintaining the structural organization of

mycomembrane.

**Figure 2. LamA recruits PgfA and MmpL3 to the old pole to build the mycomembrane.**
**(A)** The cell envelope of a representative wild type Msm cell and a cell depleted of PgfA for 24 hours were visualized and measured by cryo-electron microscopy. PM = plasma membrane; OM = outer membrane. Scale bar = 100 nm. **(B)** The lysates from strains expressing PgfA-FLAG with or without MmpL3-msfGFP were immuno-precipitated (IP) using GFP-trap beads, and the elution was probed for the presence of PgfA-FLAG via anti-FLAG Western blot. **(C)** *left*: Fluorescent protein fusions to either PgfA or MmpL3 were imaged by time-lapse microscopy in wildtype, Δ*lamA,* and complement cells (Δ*lamA* +pNative-*lamA*) N=25. The resulting images were then processed as described in Figure 1D. *right:* The same data as on the left is plotted at a single time point to show the decrease in old pole accumulation of MmpL3 and PgfA in the absence of *lamA.* **(D)** Cells were depleted using CRISPRi for *pgfA* or *mmpL3* and a representative timepoint is shown. Cell wall material is excreted from the poles and septa during depletion (white triangles).

Many of the cytoplasmic and periplasmic enzymes involved in mycomembrane

synthesis have been identified. However, the molecular details of how precursors are

trafficked from the plasma membrane to build the complex structure are almost

completely unknown. We do know that one key step involves the essential protein

MmpL3, which transports trehalose monomycolate (TMM) across the plasma membrane

to the periplasm (Su, C. C. et al., 2021; Xu et al., 2017). There, TMM is trafficked, through

unknown mechanisms, to the antigen 85 enzymes (Backus et al., 2014), and incorporated

onto mAGP or made into trehalose dimycolate (TDM), a component of the outer

mycomembrane leaflet (Chiaradia et al., 2017). Importantly, MmpL3 has become an anti-

TB therapeutic target, as multiple compounds have been found to inhibit its function

(Adams et al., 2021; La Rosa et al., 2012; Umare et al., 2021).

Using an *E. coli* two-hybrid approach, MmpL3 from *M. tuberculosis* was recently

found to interact with several factors involved in cell wall synthesis, growth, and division,

including Rv0227c (64% identity to *M. smegmatis* PgfA) (Belardinelli et al., 2019). To

verify that PgfA and MmpL3 interact in intact mycobacterial cells, we constructed strains

that expressed fusions to these proteins to enable co-immunoprecipitation. Precipitating

MmpL3-GFP with a nanobody against GFP, we found PgfA-FLAG in the elution only when

MmpL3-GFP was present (Fig. 2B). Taken together, these results show that MmpL3 and

PgfA interact in the cell.

As our initial interest in PgfA was prompted by its connection to LamA, we

wondered if PgfA and MmpL3 would localize differently in Δ*lamA* cells. Thus, we

visualized fluorescent fusions to PgfA and MmpL3 in strains with and without *lamA* and

analyzed the fluorescence distributions over time. MmpL3-msfGFP displayed the same

spatial and temporal localization pattern as PgfA-mRFP, again suggesting they are part

of the same complex (Fig. 2C). Surprisingly, in Δ*lamA* cells, PgfA-mRFP and MmpL3-

msfGFP became dramatically less abundant at the old pole, with no change in abundance

at the new pole. The loss of polar PgfA and MmpL3 in Δ*lamA* could be complemented by

integrating *lamA* at a single site on the chromosome (Fig. 2C). The abundance of

msfGFP-MurJ, which transports peptidoglycan precursors, was not as dramatically

changed at the old pole in Δ*lamA* cells, showing that the LamA-dependent loss of MmpL3

and PgfA was specific and not a general loss of elongation-complex proteins (Figure 2 –

figure supplement 2).

If PgfA and MmpL3 are functioning together, then cells depleted of either essential

protein may have the same terminal phenotype. To assay this, we created CRISPRi -

guides to both *mmpL3* and *pgfA* and visualized the morphology of cells over time as the

depletion was induced (Rock et al., 2017). As *pgfA* is predicted to be the first gene in a

two gene operon (Martini et al., 2019), we were concerned about polar effects of the

CRISPRi depletion (Rock et al., 2017). To address this, we integrated another copy of

MSMEG_0315, the second gene in the operon, at a phage site expressed by its native

promoter. Consistent with the notion that PgfA and MmpL3 function in the same pathway,

we found that cells depleted for *mmpL3* phenocopied those depleted for *pgfA* in that they

become progressively shorter and wider (Figure 2 - video supplement 1). By phase

contrast microscopy, we also observed cell wall material excreted from the poles and the

septa in both *mmpL3*- and *pgfA*- depleted cells (Fig. 2D). These observations are

reminiscent of cells treated with ethambutol (Wuo et al., 2022), a drug that inhibits the

synthesis of arabinan, the anchor point for mycolic acids, leading to accumulation and

shedding of TMM (Kilburn and Takayama, 1981; Mikusova et al., 1995). Indeed, inhibition

or depletion of MmpL3 leads to TMM accumulation (Degiacomi et al., 2017; Fay et al.,

2019a; Su, Chih-Chia et al., 2019; Tahlan et al., 2012).

Collectively, our data are consistent with a model in which LamA recruits PgfA and

MmpL3 to the old pole, either directly or indirectly, where they function to construct the

mycomembrane. Further, our data suggest that cells depleted for PgfA accumulate TMM,

possibly to toxic levels.

**PgfA is a periplasmic protein that binds TMM**

As our data suggested that PgfA-depleted cells accumulated TMM, we wondered

if PgfA was directly or indirectly interacting with TMM. A recent proteomics study identified

PgfA as a putative TMM-interacting protein (Kavunja et al., 2020). To verify this, we tested

whether PgfA could be captured from live *M. smegmatis* cells using N-x-AlkTMM-C15, a

synthetic photo-cross-linkable TMM analogue containing an alkyne "click" chemistry

handle that enables specific detection and/or enrichment of cross-linked protein

interactors (Kavunja et al., 2020). *M. smegmatis* cells expressing PgfA-3xFLAG were

incubated with N-x-AlkTMM-C15 and exposed to UV irradiation to effect cross-linking.

Then, cell lysates were collected and subjected to "click" reaction to dual label N-x-

AlkTMM-C15-modified proteins with fluorophore and biotin affinity tags. Biotinylated

proteins were captured on avidin beads, eluted, and analyzed by anti-FLAG Western blot.

We found that PgfA was enriched exclusively in N-x-AlkTMM-C15-treated, UV-exposed

*M. smegmatis*, demonstrating direct interaction between PgfA and the TMM analogue

(Fig. 3A and Figure 3 – figure supplement 1)

Several accessory proteins work with MmpL3 and/or are required to transport

TMM across the plasma membrane. However, to date, all the known accessory proteins

reside in the cytoplasm or have

globular domains on the

cytoplasmic side of the plasma

membrane(Fay et al., 2019b). The

topology of the predicted PgfA

structure, shows a beta barrel-like

domain anchored by one or,

possibly two, transmembrane

helices(Patel et al., 2022).

Prediction of the orientation with

respect to the membrane is

ambiguous (Figure 3 – figure

supplement 2A). To map the

topology of PgfA, we fused

mGFPmut3 to either the N- or the

C-terminal side of PgfA (Fig. 3B).

As GFP does not fluoresce in the

mycobacterial periplasm,[36] we

reasoned that fluorescence would

indicate cytoplasmic localization

of GFP. Importantly, both fusion

**Figure 3. PgfA is a periplasmic protein that binds TMM and is involved in TMM trafficking. (A)** Cells expressing PgfA-3xFLAG were cultured with N-x-AlkTMM-C15 (100 μM), UV-irradiated, and lysed. Lysates were reacted with azido-TAMRA-biotin reagent (AzTB) by Cu-catalyzed azide-alkyne cycloaddition (CuAAC "click" reaction), and analyzed before (input) and after (output) avidin bead enrichment by anti-FLAG Western blot. The full analysis, including Coomassie and in gel-fluorescence scanning is displayed in Figure 3 – figure supplement 1. Data are representative of two independent experiments. **(B)** PgfA is fused to GFPmut3 at either its N- or C- terminus and integrated into Msm as the sole copy of PgfA. Cells carrying these fusions are imaged by fluorescence microscopy. Images are displayed on identical intensity scales to allow direct comparison. **(C)** Using CRISPRi, cells were depleted of MmpL3 or PgfA at the indicated timepoints and the cell envelopes were analyzed for TMM and TDM by TLC. **(D)** Cells were induced to overexpress PgfA at the indicated timepoints, and their cell envelopes were analyzed by TLC for TMM and TDM. Data in C and D are representative from at two least independent experiments, all with similar results.

proteins can replace wild type PgfA at high efficiency, and thus encode functional PgfA

(Figure 3 – figure supplement 2B). Using fluorescence microscopy, we find that both the

N- and C-terminal fusions fluoresce, although the N-terminal fusion is significantly dimmer

(Fig. 3B). Nevertheless, collectively, these data show that PgfA is an inner membrane

protein with a large periplasmic domain and that it binds TMM.

**PgfA abundance correlates to altered TMM/TDM ratios**

If PgfA is involved in the trafficking of TMM, then its depletion, as with depletion of

MmpL3, should result in an altered TMM/TDM ratio (Degiacomi et al., 2017; Fay et al.,

2019; Su et al., 2019; Tahlan et al., 2012). Total cell-associated amounts of TMM and

TDM can be visualized by thin layer chromatography (TLC). As expected, depletion of

*mmpL3* leads to accumulation of TMM and a co-incident reduction of TDM (Fig. 3C). To

test if this is also the case for PgfA, we repeated the same procedure on cells depleted

of *pgfA*. In this case, we find that TMM accumulates while TDM levels are more stable

(Fig. 3C). These data are consistent with a model in which TMM accumulates in a different

subcellular compartment in *pgfA*-depleted cells, compared to *mmpL3*-depleted cells.

Specifically, it suggests that in *pgfA*-depleted cells, at least some of the TMM pool may

still be accessible to antigen 85 enzymes.

To further test the notion that PgfA is involved in TMM trafficking in the periplasm,

we created a strain which inducibly overexpressed the protein from a multi-copy plasmid.

Again, consistent with the notion that PgfA is involved in trafficking TMM, we find that

inducing the overexpression of PgfA results in increased TDM levels (Fig. 3D). Together

these data support a model in which PgfA functions as part of the TMM transport pathway

in the periplasm.

**The function of PgfA is specific to mycobacteria and its abundance negatively**

**correlates with the abundance of lipoglycans LM/LAM**

Our data indicate a role for PgfA in the transport of TMM. In contrast, the putative

corynebacterial homolog of *pgfA* has been proposed to function in the biosynthesis of two

large lipoglycans abundant in the mycobacterial cell envelope, lipomannan and

lipoarabinomannan (LM/LAM) (Cashmore et al., 2017). A deletion mutant of *NCgl2760*,

the presumed ortholog of *pgfA* in *C. glutamicum,* resulted in disappearance of LAM and

accumulation of truncated LM, which was clearly detectable by faster migration on SDS-

PAGE (Cashmore et al., 2017). To test if this is also the case in mycobacteria, we used

the same electrophoretic approach to analyze LM/LAM, and thin layer chromatography

(TLC) to examine potential changes in other lipid species upon depletion of *pgfA*. In

contrast to the results obtained in *C. glutamicum*, we observe a dramatic and transient

increase in the total amount of cell-associated LAM during depletion, before cells began

dying (Fig. 4A, Figure 4 – figure supplement 1). There were no obvious changes in the

migration of LM/LAM on SDS-PAGE, implying that the impact of PgfA depletion on

LM/LAM biosynthesis was minimal. Additionally, other lipids, including the precursor of

LM/LAM biosynthesis such as $AcPIM_2$, show no reproducible change (Figs. 4B,C; Figure

4 – figure supplement 1).

These data led us to hypothesize that PgfA and NCgl2760 are not orthologs,

despite the fact that there is synteny in their chromosomal location. However, while the

**Figure 4. LM/LAM levels negatively correlate with abundance of PgfA.** In a strain whose only copy of *pgfA* is ATC-inducible, **(A)** LM/LAM, **(B)** non-polar and polar PIMs, **(C)** and plasma membrane lipids were analyzed by **(A)** SDS-PAGE and **(B-C)** TLC during PgfA depletion (without ATC). Cell pellets were normalized by wet weight to account for differences in cell density. To avoid changes in LM/LAM due to cell density, at the indicated timepoints, half of the cultures were taken for lipid analysis, and replaced with fresh media. Two other biological replicates are shown in Figure 4 – figure supplement 1. CL, cardiolipin; PE, phosphatidylethanolamine; PI, phosphatidylinositol. **(D)** Results of exchanging the indicated *pgfA* and *C.glutamicum NCgl2760* alleles for a wildtype copy of *pgfA*. Correct transformants are positive for the incoming selective marker and negative for the outgoing (KanR, NatS). **(E)** Transposon insertions in Δ*lamA* comparted to WT. The only two genes with significantly different number of insertions (reduced) were *lamA* itself and *sucT*, a gene that codes for a protein known to modify LAM and arabinogalactan. **(F)** In cells carrying an ATC-inducible copy of *pgfA* on a multicopy plasmid, cell-associated LM/LAM were extracted, separated via SDS-PAGE, and visualized by ProQ Emerald with and without ATC at the indicated timepoints.

genes surrounding *NCgl2760* and *pgfA* share moderate to high degrees of sequence

identity (~30-50%), *NCgl2760* and *pgfA* are the clear outliers, with only 21% sequence

identity. In fact, a recent study solved the structure of the periplasmic domain of

MSMEG_0317 (PgfA) and compared it to the predicted AlphaFold  (Jumper et al., 2021;

Varadi et al., 2022) structure of NCgl2760 and found differences in the size and shape of

the beta-barrel pocket (Patel et al., 2022). Additionally, NCgl2760 is missing the long C-

terminal cytoplasmic tail that is present in PgfA. Thus, we wondered if NCgl2760 can

substitute for PgfA in *M. smegmatis*. To test this, we first asked if PgfA's cytoplasmic tail

is important for its function using our allele swapping strategy (Fig. 1B). We find that

transformation of an allele deleted for this tail is inefficient (Figure 4 – figure supplement

2A), and that the few transformants that do survive have altered morphology (Figure 4 –

figure supplement 2B). Thus, the long C-terminal tail of PgfA is necessary for its function.

We next tried to swap alleles of NCgl2760 with or without the C-terminal PgfA tail.

Transformation of both genes was inefficient (Figure 4 – figure supplement 2C) and none

of the surviving transformants had correctly swapped out the parental *pgfA* gene (Fig.

4D). Thus, *NCgl2760*, even when coupled to the important cytoplasmic tail of PgfA,

cannot substitute for *pgfA* in *M. smegmatis*. These data indicate that PgfA and NCgl2760

may not be orthologs and could have diverged through evolution to perform different

functions.

Why does depletion of PgfA result in an accumulation of LM/LAM? To investigate

this, we decided to leverage Δ*lamA* cells, which naturally have less PgfA and MmpL3

(Fig. 2C). PgfA and MmpL3 have been described as highly vulnerable drug targets, *i.e.* a

small change in their abundance causes growth arrest or cell death (Bosch et al., 2021).

How do Δ*lamA* cells grow at a normal rate with fewer of these essential proteins? We

reasoned that changes in gene essentiality in Δ*lamA* may uncover compensatory

mechanisms that promote survival. Thus, we created transposon libraries in Δ*lamA* and

wild type cells and compared insertions across the genomes. There were very few

changes: aside from Δ*lamA* itself, the only gene that sustained significantly fewer

insertions in Δ*lamA* across replicates was MSMEG_3187 or *sucT* (Fig. 4E, Supplemental

Table 1). SucT succinylates arabinogalactan and LAM, a modification that changes the

structural properties of the mycomembrane (Palcekova et al., 2019). These data suggest

that in cells deleted for *lamA*, in which the levels of PgfA and MmpL3 are lower, the

presence of lipoglycans or their modifications become more important. In fact, we find

that cells overexpressing PgfA, which results in more TDM (Fig. 3D), also have lower

levels of cell-associated LAM (Fig. 4F).

Taken, together, our data suggest that PgfA and the presumed corynebacterial

ortholog, Ncgl2760, do not have the same function. Further, our data uncover a previously

unknown connection between TMM transport, or assembly of the mycomembrane, and

LM/LAM. We find that the levels of LAM, and to a lesser extent LM, are negatively

correlated with the levels of PgfA-mediated TMM-trafficking, suggesting that there may

be competing pathways for the transport or synthesis of these molecules, a notion

supported genetically by Tn-seq. Alternatively, altered PgfA-mediated TMM trafficking

may lead to differences in LM/LAM accessibility to extraction methods.

**PgfA is necessary and sufficient for old pole growth in Δ*lamA* and the molecular**

**requirements for growth between the poles are different**

As we have shown, PgfA and MmpL3 are necessary for polar growth (Fig.1A, 2D).

Additionally, in Δ*lamA* cells, which grow less from the old pole, PgfA and MmpL3 are less

**Figure 5. PgfA is sufficient to restore old pole growth in Δ*lamA* cells.** **(A)** Cells were stained with amine-reactive dye, loaded into a microfluidic device, and imaged while perfusing dye-free media. By following the amount of unlabeled cell wall material, growth at the old and new poles was measured over time. The amount of growth incorporated per cell cycle is compared across WT cells, Δ*lamA* cells, and Δ*lamA* cells expressing a second copy of *pgfA* or *mmpL3* driven by a strong promoter**.** (N=51 WT; 44 Δ*lamA*; 36 Δ*lamA*+*pgfA;* 35 Δ*lamA*+*mmpL3*). Bars represent medians of the population. **(B)** Cells were stained with a fluorescent d-alanine analog, HADA, fixed, and imaged by fluorescence microscopy. MicrobeJ was used to segment the cells and measure intensity profiles along the long axis of the cell. Profiles were then aligned by the brightest pole and ordered by length as a proxy for cell cycle. (N=194 WT; N=259 Δ*lamA;* N=335 Δ*lamA*+*lamA)* **(C)** A model for asymmetric growth. PgfA is recruited by LamA to the old pole, where, together with other proteins, it functions to organize the mycobacterial cell envelope and promote fast cell elongation at one side of the cell.

abundant at this site (Fig. 2C). We wondered if loss of old pole growth in Δ*lamA* cells

could be directly attributed to the lowered abundance of either PgfA or MmpL3 at this site.

To test if PgfA and/or MmpL3 are sufficient to restore old pole growth in Δ*lamA*, we

created strains that expressed a second copy of these genes driven by a strong ribosomal

promoter. By quantifying polar growth with an established pulse-chase assay (Aldridge et

al., 2012; Rego et al., 2017), we find that Δ*lamA* cells encoding an extra copy of *pgfA*, but

not *mmpL3*, grow more from the old pole (Fig. 5A). These data suggest that the function

of PgfA or its interaction with other factors is necessary and sufficient for growth from the

old pole in *M. smegmatis*.

Intriguingly, overexpression of PgfA does not lead to more growth at the new pole

(Fig. 5A). This mirrors the observation that in Δ*lamA* cells - which grow more from the

new pole - neither MmpL3 nor PgfA are more abundant at this pole (Fig. 2C). Instead, the

levels of MurJ-msfGFP become increased (Figure 2 – figure supplement 2), leading us to

hypothesize that peptidoglycan synthesis increases at this site in the absence of *lamA*.

To test this, we stained cells with the fluorescent D-Ala analogue HADA, which is

incorporated into peptidoglycan through the cytoplasmic route of synthesis(García-

Heredia et al., 2018). In agreement with the increased MurJ abundance at the new pole,

we find that Δ*lamA* cells have increased HADA staining at the new pole (Fig. 5B). These

data suggest that the molecular requirements for growth between the new and old poles

may be different: with PgfA being rate limiting at the old pole and PG synthesis being rate

limiting at the new pole.

**DISCUSSION**

Our understanding of the mechanisms that govern polar growth and division are

at a nascent stage compared to our understanding of these processes in model rod

shaped bacteria  (Baranowski et al., 2019; Kieser and Rubin, 2014). This is unfortunate

because the enzymes that create and remodel the cell envelope are a rich source of anti-

bacterial drug targets. In addition, the unusual mode of asymmetric polar growth - coupled

with the complexity of constructing the multi-layered mycobacterial cell envelope - means

that the details of growth and division in mycobacteria are almost certainly different from

laterally growing rod-shaped bacteria.

Here, we investigate the function of a mycobacterial specific factor LamA. We had

previously shown that LamA is, in part, responsible for asymmetric polar growth. Cells

missing LamA grow more symmetrically and are uniformly susceptible to several drugs

(Rego et al., 2017). How does LamA create asymmetry? We make inroads into answering

this question by investigating the cellular function of MSMEG_0317, another protein of

unknown function, which was predicted to interact with LamA. We show that

MSMEG_0317, renamed PgfA, is involved in the trafficking of mycolic acids in the

periplasmic space. PgfA interacts with a TMM analog, consistent with a role in directly

trafficking TMM. Additionally, our data point to a more global function for PgfA in

regulating the composition of the cell envelope. Specifically, we find that PgfA levels

negatively correlate with the abundance of LAM, and to some extent LM, two large and

abundant lipoglycans in the cell envelope. These data suggest that there may be

compensatory or competing mechanisms for the transport and/or synthesis of TMM and

LM/LAM (Fig. 5C). Further strengthening that argument, we find that Δ*lamA* cells, which

have less PgfA, are more reliant on SucT - an enzyme that succinylates LAM and

arabinogalactan (Bhamidi et al., 2008; Palcekova et al., 2019). It may be that TMM and

LM/LAM are simply competing for space in the outer leaflet of the mycomembrane.

However, to date, altered LM/LAM levels have not been reported for other TMM-trafficking

mutants, suggesting that there may be more active regulation controlling the levels or

transport of these two molecules. Indeed, others have speculated that succinylation itself

may act to negatively regulate mycoloylation, as succinylated arabinan chains are found

to be unmycoloylated (Bhamidi et al., 2008).

Deletion of *lamA* results in less PgfA and MmpL3 at one side of the cell, the 'old

pole', leading to less growth from that side of the cell. Increased expression of PgfA but

not MmpL3 restores growth at that side of the cell, suggesting that PgfA's role in

coordinating lipid trafficking, or the presence of the protein itself, is rate limiting for growth

at the old pole (Fig. 5C). Further research will be needed to understand how a TMM-

binding protein influences the rate of cell wall insertion at the old pole. Indeed, insertion

of new cell wall material requires the concerted effort of many synthetic enzymes,

including those that polymerize and crosslink peptidoglycan. In mycobacteria, the

bifunctional class A PBPs (aPBPs) are the dominant enzymes required for cell elongation.

In other organisms, aPBPs are activated by inner or outer membrane-anchored proteins

(Fenton et al., 2018; Paradis-Bleau et al., 2010; Typas et al., 2010), though no activator

of PonA1 - the major aPBP in mycobacteria - has been found. Thus, it is intriguing to

speculate that PgfA – a protein involved in mycomembrane precursor transport - may be

needed to activate PG synthesis at the old pole in mycobacteria.

What are the requirements for growth at the new pole – the pole newly formed by

division? In Δ*lamA*, this pole grows more without an increase in PgfA or MmpL3; instead,

enzymes involved in PG synthesis are more abundant at this site in Δ*lamA*. These

observations suggest that the requirements for growth between the two poles are

fundamentally different, at least for a certain time after the onset of pole elongation.

Indeed, new poles will eventually become old poles in the next round of division. We see

that PgfA and MmpL3 localize to the "new old poles" after division, establishing

asymmetry in the next generation. How does LamA affect the growth of the poles in

opposite directions – inhibiting one but activating another? Future research is needed,

but our data suggest that the new and old poles of mycobacteria may be distinct sites,

with different molecular factors and requirements for growth. Our results suggest that

LamA influences these differences to create asymmetry in individual cells, leading to a

heterogeneous population better able to survive certain stresses, like antibiotics.

**Acknowledgements**

We thank members of the Rego lab for helpful suggestions. This work was supported by

the National Institutes of Health, R01AI148255 to E.H.R, who was also supported by the

Searle and Pew Foundations. B.M.S. was supported by the National Science Foundation

(CAREER Award 1654408). C.W and J.L. were supported by the NIH grants

R01AI087946 and R01GM110243.

**Materials and Methods**

Table 1: Strains used in this study

| Reagent type (species) or resource | Designation | Source or reference | Identifiers | Additional Information |
|---|---|---|---|---|
| strain (*Mycobacterium smegmatis*) | | | *Mycobacterium smegmatis* mc$^2$155 | Wild type *M. smegmatis* |
| strain (*M. smegmatis*) | KG26 | this work | mc2155Δmsmeg_0317 L5::ptetO-msmeg_0317 | Parental swap strain |
| Strain ( *M. smegmatis*) | HR342 | Rego et al 2016 | mc2155ΔMSMEG_4265 | LamA knockout strain |
| strain (*M. smegmatis*) | KG97 | This work | mc2 155 Δmsmeg_0317 L5::ptetO-msmeg_0317-myc /pTetR | Strain used for the Anhydrotetracyline inducible depletion of PgfA |
| strain (*M. smegmatis*) | KG81 | This work | mc2 155 Δmsmeg_0317 pL5:pNative- msmeg_0317 -mRFP-myc | Strain used for determining PgfA localization |
| strain (*M. smegmatis*) | HR388 | Rego et al 2016 | mc2155 L5::pG-MCK-ptb21-egfp-wag31 tweety::ptb21-ftsZ-mCherry2B | Strain used to determine localization of FtsZ & Wag31 |
| strain (*M. smegmatis*) | CMG184 | This work | mc2155 tw:: ptb21 mmpL3$_{TB}$-msfGFP | Strain used to determine localization of MmpL3 |

| strain (*M. smegmatis*) | CMG530 | This work | mc2155 Δ msmeg_0317 L5:: pTetO- msmeg_0317 - 3X-Flag tw:: ptb21-mmpL3$_{TB}$-msfGFP | Strain used for co-immunoprecipitation of MmpL3 and PgfA |
|---|---|---|---|---|
| strain (*M. smegmatis*) | KG167 | This work | mc2155 Δ msmeg_0317 L5:: pTetO- msmeg_0317 - 3X-Flag | Control strain used for co-immunoprecipitation of MmpL3 and PgfA |
| strain (*M. smegmatis*) | CMG184 | This work | mc2155 tw:: ptb21 mmpL3$_{TB}$-msfGFP | Strain used to determine localization of MmpL3 |
| strain (*M. smegmatis*) | CMG481 | This work | mc2155 ΔlamA::zeo L5:: pNative-lamA-strep tw:: ptb21 mmpL3$_{TB}$-msfGFP | Strain used to determine localization of MmpL3 in a ΔlamA background with LamA complemented under its native promoter |
| strain (*M. smegmatis*) | CMG557 | This work | mc2155 L5:: pTetO-MS0317-myc #1 | Strain used for measuring polar growth in a PgfA merodiploid background |
| strain (*M. smegmatis*) | KG175 | This work | mc2155 ΔlamA L5::pTetO-ms0317-myc | Strain used for measuring polar growth in a PgfA merodiploid and LamA knockout background |
| strain (*M. smegmatis*) | CMG585 | This work | mc2155 L5:: pTetO mmpL3$_{TB}$-myc | Strain used for measuring polar growth in an MmpL3 merodiploid background |
| strain (*M. smegmatis*) | CMG587 | This work | mc2155 ΔlamA L5:: pTetO mmpL3$_{TB}$-myc | Strain used for measuring polar growth in an MmpL3 merodiploid and LamA knockout background |
| strain (*M. smegmatis*) | KG154 | This work | mc2 155 ΔlamA Δmsmeg_0317 pL5:pNative- msmeg_0317 -mRFP-myc | Strain used for PgfA localization in LamA knockout background |

| | | | | |
|---|---|---|---|---|
| strain (*M. smegmatis*) | KG157 | This work | mc2 155 ΔlamA Δmsmeg_0317 pL5:pNative- msmeg_0317 -mRFP-myc pL5 pNative-lamA-strep | Strain used for PgfA localization in a background where lamA was complemented in LamA knockout |
| strain (*M. smegmatis*) | CMG206 | This work | mc2155 tw:: ptb21 msfGFP-MurJ$_{TB}$ | Strain used for localization of MurJ |
| strain (*M. smegmatis*) | CMG199 | This work | mc2155 ΔlamA tw:: ptb21 msfGFP-murJ$_{TB}$ | Strain used for localization of MurJ in a LamA knockout background |
| strain (*M. smegmatis*) | CMG485 | This work | mc2155 ΔlamA::zeo L5:: pNative LamA strep tw:: ptb21 msfGFP-MurJ$_{TB}$ | Strain used for localization of MurJ in a LamA knockout background where LamA was complemented under its native promoter |
| strain (*M. smegmatis*) | KG244 | This work | mc2 155 L5::pLJR962-mmpL3-sgRNA | CRISPR inducible MmpL3 depletion strain |
| strain (*M. smegmatis*) | KG289/ CMG549 | This work | mc2 155 L5::pLJR962-msmeg_0317-sgRNA:: pTweety-pNative-msmeg_0315 | CRISPR inducible PgfA depletion strain |
| strain (*M. smegmatis*) | KG47 | This work | mc2 155 Δmsmeg_0317 L5::pTetO-msmeg_0317-mGFPmut3-myc | C-terminal GFP tagged PgfA strain |
| Strain (*M. smegmatis*) | KG73 | This work | mc2 155 Δmsmeg_0317 L5:: pTetO-mGFP-mut3-MS0317-myc | N-terminal GFP tagged PgfA strain |
| Strain (*M. smegmatis*) | KG167 | This work | mc2 155 Δmsmeg_0317 L5:: pTetO-MS0317-3XFLAG | Strain carrying MSMEG_0317-3X-FLAG, used for TMM pulldown |

| | | | | |
|---|---|---|---|---|
| Strain (*M. smegmatis*) | KG60 | This work | mc2 155 /pTetOR-msmeg_0317-myc | Strain carrying tetracycline inducible over-expression plasmid for PgfA |
| Strain (*M. smegmatis*) | KG286 | This work | mc2 155 Δmsmeg_0317 L5:: pTetO-msmeg ΔC | Strain carrying msmeg_0317 without c-terminal cytoplasmic tail |
| Plasmid (*E. coli*) | KG147 | This work | DH5 α /pL5-pTetO-*Ncgl2760*-myc | Plasmid carrying corynebacterial *ncgl2760* gene |
| Plasmid (*E. coli*) | KG284 | This work | DH5 α /pL5-pTetO-*Ncgl2760*-msmeg_0317-C-term | Plasmid carrying corynebacterial *ncgl2760* gene fused to cytosolic C-terminal of *msmeg_0317* gene |
| Plasmid (*E. coli*) | CMG541 | This work | Dh5a / pTweety pNative Msm_0315-strep | Plasmid carrying mycobacterial gene MSMEG_0315 with a C-terminal strep tag |

**Bacterial strains and culture conditions**: *M. smegmatis* mc$^2$155 was grown in Middlebrook 7H9 broth supplemented with 0.05% Tween80, 0.2% glycerol, 5 gm/L albumin, 2 gm/L dextrose and 0.003 gm/L catalase or plated on LB agar. *Escherichia coli* DH5α cells were grown in LB broth or on LB agar plates. Concentrations of antibiotics used for *M. smegmatis* is as follows: 20 μg/ml zeocin, 25 μg/ml kanamycin, 50 μg/ml hygromycin, and 20 μg/ml nourseothricin. Concentrations of antibiotics used for *E. coli* used for *E. coli* is as follows: 40 μg/ml zeocin, 50 μg/ml kanamycin, 100 μg/ml hygromycin, and 40 μg/ml nourseothricin.

**Plasmid and strain construction**: Strains used in this study are listed in Table 1.

Oligos and primers are listed in Supplemental Table 2. In brief, before deleting the native

copy of *msmeg_0317*, a merodiploid strain was created by inserting a second copy of

*msmeg_0317* gene under pTetO promoter using a Kan[R] L5 integrating vector.

Subsequently, in the merodiploid strain, an in-frame deletion was made by replacing the

native copy of *msmeg_0317 gene* with a zeocin resistance cassette flanked by loxP sites

using recombineering  (van Kessel and Hatfull, 2007). *msmeg_0317* and its variants,

including the fluorescent protein fusions, were cloned in a Nat[R] L5 integrating vector for

allele exchange at the L5 site. The *msmeg_0317* depletion strain was made by

transforming an episomal Hyg[R] marked vector constitutively expressing TetR repressor

into a strain expressing a single copy of *msmeg_0317* controlled by the pTetO promoter.

For expression of *msmeg_0317* driven by the native promoter, 200bp upstream of

msmeg_0317 chromosomal locus was used. For the MmpL3 and MurJ fluorescent

fusions and co-IP experiments, the genes were cloned from the *M. tuberculosis* genome,

expressed from the ptb21 promoter and integrated in single copy at the tweety phage

integration site. All plasmid constructs were made using isothermal Gibson assembly

whereby insert and vector backbone shared 20-25 bp of homology.

**MSMEG_0317 Depletion:**

*Promoter depletion.* To transcriptionally deplete *msmeg_0317* in Msm, the ATC-inducible

Tet-ON system was used. The only copy of *msmeg_0317* was driven by the pTetO

promoter, while the TetR repressor was constitutively expressed in trans from an

episomal vector. MSMEG_0317 was depleted by removing ATC from the medium and

cells were grown for 18 hours. Subsequently, MSMEG_0317 depleted cells were re-

diluted in fresh medium, also without ATC, and samples at different timepoints were taken

and processed for cyro-electron microscopy. Alternatively, to avoid changes in LM/LAM

abundance that have been found to correlated with cell density and growth phase (21),

at 18 hours of depletion, half of the culture was removed for lipid extraction and analysis,

and replaced with fresh media. Lipid analysis was performed at timepoints before and

after the decrease in OD indicating cessation of growth (Fig. S3).

*CRISPRi depletion.* To transcriptionally deplete *mmpL3* and *msmeg_0317* using

CRISPRi, both dCas9 and the respective guide RNAs were cloned in an L5 integrating

plasmid pIJR962 (Rock et al., 2017). The CRISPRi silencing of *mmpL3/msmeg_0317* was

induced by adding ATC to a final concentration of 100 ng/ul. The induction was carried

out for the indicated timepoints.

**Time-lapse and fluorescence microscopy**: An inverted Nikon Ti-E microscope

was used for the time-lapse imaging. An environmental chamber (Okolabs) was

maintained at 37°C. Cells were cultivated in an B04 microfluidics plate from CellAsic,

continuously supplied with fresh medium, and imaged every 15 minutes using a 60x 1.4

N.A. Plan Apochromat phase contrast objective (Nikon). Fluorescence was excited using

the Spectra X Light Engine (Lumencor), separated using single- or multi- wavelength

dichroic mirrors, filtered through single bandpass emission filters, and detected with an

sCMOS camera (ORCA Flash 4.0). Filters are: GFP (Ex: 470/24; Em: 515/30 or 525/50);

mRFP (Ex: 550/15; Em: 595/44 or 630/75). To reduce phototoxicity exposure times were

kept below 100ms for excitation with 470nm and 300ms for excitation with 550nm.

**Kymograph analysis and Image analysis.** Time-lapse images were analyzed in

open-source image analysis software Fiji (30) and a custom MATLAB program was used

to generate kymographs. Source code will be uploaded to Github. Specifically, for a single

cell, in Fiji, a 5-pixel wide segmented line was drawn from the new pole to the old pole at

each time point during the cell cycle. This was repeated on 20-50 cells. These line profiles

were then imported into MATLAB, where a custom script was used to generate an

average kymograph by 2D interpolation of the individual kymographs. For the

demographs shown in Fig 5, cells were segmented and intensity profiles measured using

MicrobeJ. Output intensity profiles were then transferred to Matlab, sorted by cell length,

and aligned by the brightest pole for visualization.

**N-x-AlkTMM-C15 mediated protein capture:**

*Preparation of cell lysate. M. smegmatis* expressing 3x FLAG-tagged

MSMEG_0317 starter culture was generated by inoculating a single colony from a freshly

streaked LB agar plate supplemented with zeocin and nourseothricin (20 ug/ml each) into

10 ml liquid 7H9 medium in a sterile culture tube. The starter culture was grown until mid-

logarithmic phase and then diluted to OD600 0.2 with 7H9 medium to a final volume of

80 ml. The culture was split into two equal volumes in 125ml sterilized culture flasks. To

one flask, N-x-AlkTMM-C15 was added to a final concentration of 100 $\mu$M with (final

DMSO concentration of 2%), while the other flask was left untreated as a DMSO control

(final DMSO concentration of 2%). Both flasks were incubated with shaking until OD600

0.8 was reached. The cells were pelleted by centrifugation at 6,500 xg at 4 °C for 10 min.

The cell pellets were washed twice with PBS, re-suspended in 6 ml PBS, and split into

two equal volumes. One aliquot was exposed to UV irradiation for 30 min with a 5-watt

365 nm UV bench lamp (UVP) while the other was left unexposed as a control. The cell

pellets were collected by centrifugation and re-suspended in 600 $\mu$l lysis buffer (2 mg/ml

lysozyme, 1 mM phenylmethylsulfonyl fluoride (PMSF) in 1x PBS), transferred to

scintillation vials, and incubated at 37 °C for 2 h. The mixtures were transferred to 1.5 ml

screw-cap vials containing 0.25 ml of 0.1 mm zirconia/silica beads (BioSpec Products)

and subjected to bead beating 3x for 1 min each using a FastPrep-24 bead beater (MP

Biomedicals). The lysate was transferred to a scintillation vial, then SDS was added to a

final concentration of 2% from a 20% SDS stock. Cell extracts were incubated at 60 °C

for 2 h with constant stirring. The lysates were transferred to microcentrifuge tubes and

centrifuged at 3,200 xg for 10 min at 4 °C. The supernatant was collected and stored in

separate tubes at 4 °C until use.

CuAAC and affinity enrichment. To reduce SDS concentration, 500 $\mu$l of

methanol/chloroform (2:1 v/v) was added to 184 $\mu$l of cell lysate. The resulting protein

precipitate was centrifuged at 18,000 xg at 4 °C for 10 minutes and the supernatant was

discarded. The precipitate was air dried and solubilized using 184 $\mu$l of 0.5% SDS/0.05%

LDAO buffer. Copper catalyzed azide-alkyne cycloaddition (CuAAC) was carried out by

sequential addition of 1 mM AzTB (azido-TAMRA-biotin (4 $\mu$l), Click Chemistry Tools), 60

mM sodium ascorbate (4 $\mu$l), 6.4 mM TBTA ligand (4 $\mu$l), and 50 mM copper sulfate (4 $\mu$l)

to give a final volume of 200 $\mu$l. The final reagent concentration was 20 $\mu$M AzTB; 1.2 mM

sodium ascorbate; 128 $\mu$M TBTA; and 1 mM copper(II) sulfate. The mixture was

thoroughly mixed by pipetting up and down and the reaction was incubated for 2 h in the

dark with constant agitation at 37 °C. Excess AzTB was removed by precipitating proteins

in methanol/chloroform (2:1 v/v), discarding supernatant, and solubilizing protein pellets

in 200 $\mu$l 0.5% SDS/0.05% LDAO buffer as described above. Protein concentration was

determined by Bradford assay. 15 $\mu$l of each sample was saved as input sample. The

remaining sample was mixed with 40 μl Pierce avidin-agarose beads (ThermoFisher

Scientific) that had been pre-washed 3x with 0.5% SDS/0.05% LDAO buffer (100 μl). The

bead mixture was incubated at room temperature with constant rotation for 2 h. The beads

were washed 3x with 0.5% SDS/0.05% LDAO buffer (100 μl) followed by 3x with PBS

(100 μl) with centrifugation at 1,000 xg for 1 min between each wash. Bound proteins

were eluted by boiling at 95 °C for 15 min in 30 μl of 4x sample buffer.

SDS-PAGE and Western blot. 5 μg of input and 10 μl of output was resolved by

SDS-PAGE by 4-20% acrylamide gels and analyzed by in-gel fluorescence using a

Typhoon FLA 7000 (GE Healthcare Life Science) using the rhodamine channel to detect

TAMRA-labeled proteins.

The gel was fixed for 15 min (40% ethanol and 10% acetic acid in DI water), rinsed

3x with DI water 10 minutes each and stained overnight with agitation in QC colloidal

Coomassie stain (Bio-Rad). The gel was rinsed with DI water 3x for 10 min and imaged

using a ChemiDoc Touch Imaging System (Bio-Rad) and processed by Image Lab

software (Bio-Rad).

The above conditions were used to generate samples for Western blot analysis. 5

μg input controls and 10 μl eluted proteins were resolved by SDS-PAGE. After gel

electrophoresis, proteins were transferred onto an Immun-Blot PVDF membrane (Bio-

Rad). The PVDF membrane was equilibrated in ethanol and blotting filter paper

(ThermoFisher Scientific) was equilibrated in transfer buffer (25 mM Tris, 193 mM glycine,

and 20% ethanol) before placement in a transfer cassette. Proteins were transferred

electrophoretically at a constant voltage of 25 V for 7 min using a Trans-Blot Turbo

Transfer System (Bio-Rad). After transfer, the membrane was blocked overnight at 4 °C

in 5% dry milk in Tris-buffered saline containing Tween 20, pH 7.6 (50 mM Tris, 0.5 M

NaCl, 0.02% Tween 20 (TBST)). Anti-FLAG-HRP (ThermoFisher Scientific) was used at

1:1,000 dilution using 2% dry milk in TBST. The membrane was incubated with antibody

at 4 °C overnight with constant rocking. The membrane was washed 3x for 10 min each

with TBST. The membrane was treated with SuperSignal West Pico PLUS

chemiluminescent reagent (ThermoFisher Scientific) and chemiluminescence was

detected on a ChemiDoc Touch Imaging system (Bio-Rad).

**Tn-seq.** Transposon libraries in wild type and Δ*lamA* (HR342) were prepared as

and sequenced described elsewhere (Dragset et al., 2019). This was done independently

on two separate days for a total of two biological replicates in each strain. The TRANSIT

and TPP python packages were used to map insertions to the mc$^2$155 genome and

quantitatively compare insertions across stains, using the 'resampling' option (DeJesus,

Michael A. et al., 2015).

**Co-immunoprecipitation:**

Mycobacterial cultures were grown to mid-log phase as described above. Protocol for

crosslinking was adapted from Belardinelli et al, Sci Rep, 2019 (Belardinelli et al., 2019).

Cells were washed with 1x phosphate buffered saline (PBS) once. Pellets were

resuspended in 1mL of 1x PBS with 1.25mM dithiobis (succinimidyl propionate) (DSP)

and incubated for 30 minutes at 37˚C for crosslinking. After incubation, cells were pelleted

at 10,000xg for 5 minutes at room temperature and the supernatant was discarded. The

pellet was resuspended in lysis buffer (50mM Tris-HCl, pH 7.4; 150mM NaCl; 10ug/mL

DNase I; one tablet Roche cOmplete EDTA free protease inhibitor cocktail; and and 0.5%

Igepal Nonidet P40 Substitute) and lysed with a BeadBug$^{TM}$ Microtube Homogenizer at

4,000rpm six times for 30 seconds each, icing in between. Lysed cells were spun down

at 15,000xg for 15 minutes at 4˚C, and the supernatant was transferred to a clean

eppendorf tube. Lysates were incubated, where indicated, with either GFP-Trap Magnetic

Agarose (Chromotek) or magnetic a-FLAG M2 beads (Sigma-Aldrich) and incubated at

4˚C overnight, rotating. After incubation, samples were spun down at 2,500xg for 1 minute

at room temperature and flow through was discarded. Beads were washed three times

with non-detergent wash buffer (10mM Tris-HCl, pH 7.4; 150mM NaCl, 0.5mM EDTA).

GFP-Trap samples were eluted with 2x Laemmli Buffer (Bio-Rad) prepared with 50mM

(DTT) and boiled at 95˚C for 5 minutes. FLAG M2 beads were eluted twice with 3xFLAG

peptide (Sigma-Aldrich) for 30 minutes rotating at 4˚C. All samples not already treated

with Laemmli Buffer + DTT were prepared for western blotting by addition of Laemmli

Buffer + DTT and boiled at 95˚C for 5 minutes, to reverse all crosslinks.

*Western blot verification*

Samples were run on NuPAGE™ (Thermo Fisher) 4-12% Bis-Tris gels in 1x MOPS-SDS

running buffer. Proteins were transferred to nitrocellulose membranes and probed with

1:1,000 a-FLAG primary (Sigma-Aldrich, clone M2) and 1:5,000 a-mouse secondary

(Thermo Fisher, Superclonal™ A28177). SuperSignal™ West Femto Extended Duration

Substrate (Thermo Fisher) was used for chemiluminescent visualization on an Amersham

ImageQuant 800 system.

**Lipid extraction and analysis.**

*TMM/TDM.* Crude lipids were extracted from equal wet cell pellet weights of the

either *mmpL3* depletion or *msmeg_0317* depleted cells after 9,12, and 15 hours with or

without ATC with 2:1 chloroform/methanol mixture for 12 hours. The organic layer was

separated from the cell debris centrifugation at 4000g. The organic extract was air dried

overnight. The dried extract was dissolved in 50 μl of 2:1 chloroform/methanol. TMMs and

TDMs were separated on by high performance thin layer chromatography (HPTLC) (Silica

gel 60, EMD Merck) using chloroform/methanol/ water (25:4:9:0.4). TMMs/TDMs were

visualized by spraying the TLC sheet with 20% 1-napthaol in 5% sulfuric acid and charring

the plate at 110°C Trehalose containing lipids (TMMs/TDMs) appear as purple bands

after charring.

*LM/LAM.* Extraction, purification and analysis of lipids were as described

previously (Rahlwes et al., 2019). Briefly, crude lipids were extracted from equal wet cell

pellet weights of the MSMEG_0317 depletion strains after 18, 24, 27, 30, and 33 hrs with

or without ATC. After lipid extraction using chloroform/methanol, LM and LAM were

extracted from the delipidated pellet by incubation with phenol/water (1:1) for 2 hrs at

55°C. Phospholipids and PIMs extracted by chloroform/methanol were further purified by

*n*-butanol/water phase partitioning, and separated by high performance thin layer

chromatography (HPTLC) (Silica gel 60, EMD Merck) using chloroform/methanol/ 13 M

ammonia/1 M ammonium acetate/ water (180:140:9:9:23). Phospholipids were visualized

via cupric acetate staining. PIMs were visualized with orcinol staining as described(Sena

et al., 2010). LM/LAM samples were separated by SDS-PAGE (15% gel) and visualized

using ProQ Emerald 488 glycan staining kit (Life Technologies). To detect LM/LAM in

culture supernatants, the supernatants were initially treated with a final concentration of

50 μg/ml Proteinase K for 4 hours at 50°C. The treated supernatants were

electrophoresed on 15% SDS-PAGE. LM/LAM were blotted onto nitrocellulose at 20 V

for 45 minutes using semi-dry transfer method. Post-transfer, the membrane was blocked

by 5% milk in Tris-buffered saline supplemented with 0.05% Tween-20 (TBST). The

blocked membrane was then probed overnight with CS-35 antibody (BEI Resources, NIH)

at 1:250 dilution at 4°C. The membrane was washed with TBST five times for five minutes

each. Post-washing, it was probed with anti-mouse secondary for one hour at room

temperature. Membrane was then washed five times for five minutes with TBST. Thermo

Scientific's west dura chemiluminescent reagent was used to develop the membrane.

**Sample preparation and image collection for Cryo-EM.** Wild type and

MSMEG_0317 depleted *M. smegmatis* cells were pelleted, washed twice with 1X

phosphate buffered saline (PBS), and suspended in ~20µl PBS. The culture was

subsequently deposited onto freshly glow-discharged holey carbon grids. The grids were

then blotted with filter paper manually for about 4 seconds and rapidly frozen in liquid

ethane. The frozen grids were transferred into a 300kV Titan Krios electron microscope

(ThermoFisher Scientific) equipped with a K2 Summit direct detector and a quantum

energy filter (Gatan, Pleasanton, CA). Cryo-EM movie stacks were collected using

SerialEM (Mastronarde, 2005). MotionCor2 (Zheng et al., 2017) was used for drift

correction of the cryo-EM movie stacks. The grey levels of each micrograph are obtained

using MATLAB.

**Experimental Replicates.** Biological replicates are defined as independent

cultures grown in parallel or on separate days. Technical replicates are defined at the

same culture, measured independently.  All the experiments were performed at least

twice - often three or more times - with biological replicates.

656  **Data and Materials Availability.** Data generated in this study are included in the

657 manuscript and supporting files. Request for strains can be addressed to the

658 corresponding author: hesper.rego@yale.edu

Reference List

Abdali, N., Younas, F., Mafakheri, S., Pothula, K.R., Kleinekathöfer, U., Tauch, A., and Benz, R. (2018). Identification and characterization of smallest pore-forming protein in the cell wall of pathogenic Corynebacterium urealyticum DSM 7109. BMC Biochem *19,* 3.

Adams, O., Deme, J.C., Parker, J.L., CRyPTIC Consortium, Fowler, P.W., Lea, S.M., and Newstead, S. (2021). Cryo-EM structure and resistance landscape of M. tuberculosis MmpL3: An emergent therapeutic target. Structure *29,* 1182-1191.e4.

Aldridge, B.B., Fernandez-Suarez, M., Heller, D., Ambravaneswaran, V., Irimia, D., Toner, M., and Fortune, S.M. (2012). Asymmetry and aging of mycobacterial cells lead to variable growth and antibiotic susceptibility. Science *335,* 100-104.

Backus, K.M., Dolan, M.A., Barry, C.S., Joe, M., McPhie, P., Boshoff, H.I.M., Lowary, T.L., Davis, B.G., and Barry, C.E. (2014). The Three Mycobacterium tuberculosis Antigen 85 Isoforms Have Unique Substrates and Activities Determined by Non-active Site Regions. J Biol Chem *289,* 25041-25053.

Baranowski, C., Rego, E.H., and Rubin, E.J. (2019). The Dream of a Mycobacterium. Microbiol. Spectr. *7,* 10.1128/microbiolspec.GPP3-2018.

Belardinelli, J.M., Stevens, C.M., Li, W., Tan, Y.Z., Jones, V., Mancia, F., Zgurskaya, H.I., and Jackson, M. (2019). The MmpL3 interactome reveals a complex crosstalk between cell envelope biosynthesis and cell elongation and division in mycobacteria. Sci. Rep. *9,* 10728-8.

Bhamidi, S., Scherman, M.S., Rithner, C.D., Prenni, J.E., Chatterjee, D., Khoo, K.H., and McNeil, M.R. (2008). The identification and location of succinyl residues and the characterization of the interior arabinan region allow for a model of the complete primary structure of Mycobacterium tuberculosis mycolyl arabinogalactan. J. Biol. Chem. *283,* 12992-13000.

Bosch, B., DeJesus, M.A., Poulton, N.C., Zhang, W., Engelhart, C.A., Zaveri, A., Lavalette, S., Ruecker, N., Trujillo, C., Wallach, J.B*., et al.* (2021). Genome-wide gene expression tuning reveals diverse vulnerabilities of M. tuberculosis. Cell *184,* 4579-4592.e24.

Cashmore, T.J., Klatt, S., Yamaryo-Botte, Y., Brammananth, R., Rainczuk, A.K., McConville, M.J., Crellin, P.K., and Coppel, R.L. (2017). Identification of a Membrane Protein Required for Lipomannan Maturation and Lipoarabinomannan Synthesis in Corynebacterineae. J. Biol. Chem. *292,* 4976-4986.

Chiaradia, L., Lefebvre, C., Parra, J., Marcoux, J., Burlet-Schiltz, O., Etienne, G., Tropis, M., and Daffé, M. (2017). Dissecting the mycobacterial cell envelope and defining the composition of the native mycomembrane. Scientific Reports *7,* 12807-12.

Degiacomi, G., Benjak, A., Madacki, J., Boldrin, F., Provvedi, R., Palù, G., Kordulakova, J., Cole, S.T., and Manganelli, R. (2017). Essentiality of mmpL3 and impact of its silencing on Mycobacterium tuberculosis gene expression. Scientific Reports *7,* 43495.

DeJesus, M.A., Gerrick, E.R., Xu, W., Park, S.W., Long, J.E., Boutte, C.C., Rubin, E.J., Schnappinger, D., Ehrt, S., Fortune, S.M., Sassetti, C.M., and Ioerger, T.R. (2017). Comprehensive Essentiality Analysis of the Mycobacterium tuberculosis Genome via Saturating Transposon Mutagenesis. MBio *8,* 10.1128/mBio.02133-16.

DeJesus, M.A., Ambadipudi, C., Baker, R., Sassetti, C., and Ioerger, T.R. (2015).
TRANSIT - A Software Tool for Himar1 TnSeq Analysis. PLOS Computational Biology
*11,* e1004401.
Dragset, M.S., Ioerger, T.R., Zhang, Y.J., Mærk, M., Ginbot, Z., Sacchettini, J.C., Flo,
T.H., Rubin, E.J., and Steigedal, M. (2019). Genome-wide Phenotypic Profiling
Identifies and Categorizes Genes Required for Mycobacterial Low Iron Fitness. Sci Rep
*9,*
Fay, A., Czudnochowski, N., Rock, J.M., Johnson, J.R., Krogan, N.J., Rosenberg, O.,
and Glickman, M.S. (2019a). Two Accessory Proteins Govern MmpL3 Mycolic Acid
Transport in Mycobacteria. mBio *10,* 10.1128/mBio.00850-19.
Fay, A., Czudnochowski, N., Rock, J.M., Johnson, J.R., Krogan, N.J., Rosenberg, O.,
and Glickman, M.S. (2019b). Two Accessory Proteins Govern MmpL3 Mycolic Acid
Transport in Mycobacteria. mBio *10,* 10.1128/mBio.00850-19.
Fenton, A.K., Manuse, S., Flores-Kim, J., Garcia, P.S., Mercy, C., Grangeasse, C.,
Bernhardt, T.G., and Rudner, D.Z. (2018). Phosphorylation-dependent activation of the
cell wall synthase PBP2a in Streptococcus pneumoniae by MacP. Proceedings of the
National Academy of Sciences - PNAS *115,* 2812-2817.
García-Heredia, A., Pohane, A.A., Melzer, E.S., Carr, C.R., Fiolek, T.J., Rundell, S.R.,
Lim, H.C., Wagner, J.C., Morita, Y.S., Swarts, B.M., and Siegrist, M.S. (2018).
Peptidoglycan precursor synthesis along the sidewall of pole-growing mycobacteria.
eLife *7,* e37243.
Hoffmann, C., Leis, A., Niederweis, M., Plitzko, J.M., and Engelhardt, H. (2008).
Disclosure of the mycobacterial outer membrane: cryo-electron tomography and
vitreous sections reveal the lipid bilayer structure. Proc. Natl. Acad. Sci. U. S. A. *105,*
3963-3967.
Jackson, M. (2014). The mycobacterial cell envelope-lipids. Cold Spring Harb Perspect.
Med. *4,* 10.1101/cshperspect.a021105.
Jankute, M., Cox, J.A., Harrison, J., and Besra, G.S. (2015). Assembly of the
Mycobacterial Cell Wall. Annu. Rev. Microbiol. *69,* 405-423.
Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A.*, et al.* (2021). Highly accurate
protein structure prediction with AlphaFold. Nature *596,* 583.
Kavunja, H.W., Biegas, K.J., Banahene, N., Stewart, J.A., Piligian, B.F., Groenevelt,
J.M., Sein, C.E., Morita, Y.S., Niederweis, M., Siegrist, M.S., and Swarts, B.M. (2020).
Photoactivatable Glycolipid Probes for Identifying Mycolate-Protein Interactions in Live
Mycobacteria. J. Am. Chem. Soc. *142,* 7725-7731.
Kieser, K.J., and Rubin, E.J. (2014). How sisters grow apart: mycobacterial growth and
division. Nat. Rev. Microbiol. *12,* 550-562.
Kilburn, J.O., and Takayama, K. (1981). Effects of ethambutol on accumulation and
secretion of trehalose mycolates and free mycolic acid in Mycobacterium smegmatis.
Antimicrob. Agents Chemother. *20,* 401-404.
La Rosa, V., Poce, G., Canseco, J.O., Buroni, S., Pasca, M.R., Biava, M., Raju, R.M.,
Porretta, G.C., Alfonso, S., Battilocchio, C.*, et al.* (2012). MmpL3 is the cellular target of
the antitubercular pyrrole derivative BM212. Antimicrob. Agents Chemother. *56,* 324-
331.

Lewis, J.A., and Hatfull, G.F. (2000). Identification and characterization of
mycobacteriophage L5 excisionase. Mol. Microbiol. *35,* 350-360.
Martini, M.C., Zhou, Y., Sun, H., and Shell, S.S. (2019). Defining the Transcriptional and
Post-transcriptional Landscapes of Mycobacterium smegmatis in Aerobic Growth and
Hypoxia. Front Microbiol *10,*
Mastronarde, D.N. (2005). Automated electron microscope tomography using robust
prediction of specimen movements. J Struct Biol *152,* 36-51.
Mikusova, K., Slayden, R.A., Besra, G.S., and Brennan, P.J. (1995). Biogenesis of the
mycobacterial cell wall and the site of action of ethambutol. Antimicrob. Agents
Chemother. *39,* 2484-2489.
Palcekova, Z., Angala, S.K., Belardinelli, J.M., Eskandarian, H.A., Joe, M., Brunton, R.,
Rithner, C., Jones, V., Nigou, J., Lowary, T.L.*, et al.* (2019). Disruption of the SucT
acyltransferase in Mycobacterium smegmatis abrogates succinylation of cell envelope
polysaccharides. J. Biol. Chem. *294,* 10325-10335.
Paradis-Bleau, C., Markovski, M., Uehara, T., Lupoli, T.J., Walker, S., Kahne, D.E., and
Bernhardt, T.G. (2010). Lipoprotein Cofactors Located in the Outer Membrane Activate
Bacterial Cell Wall Polymerases. Cell *143,* 1110-1120.
Pashley, C.A., and Parish, T. (2003). Efficient switching of mycobacteriophage L5-
based integrating plasmids in Mycobacterium tuberculosis. FEMS Microbiology Letters
*229,* 211-215.
Patel, O., Brammananth, R., Dai, W., Panjikar, S., Coppel, R.L., Lucet, I.S., and Crellin,
P.K. (2022). Crystal structure of the putative cell-wall lipoglycan biosynthesis protein
LmcA from Mycobacterium smegmatis. Acta Crystallogr. D. Struct. Biol. *78,* 494-508.
Rahlwes, K.C., Puffal, J., and Morita, Y.S. (2019). Purification and Analysis of
Mycobacterial Phosphatidylinositol Mannosides, Lipomannan, and Lipoarabinomannan.
In Bacterial Polysaccharides: Methods and Protocols, Brockhausen, Inka ed., (New
York, NY: Springer New York) pp. 59-75.
Rego, E.H., Audette, R.E., and Rubin, E.J. (2017). Deletion of a mycobacterial divisome
factor collapses single-cell phenotypic heterogeneity. Nature *546,* 153-157.
Richardson, K., Bennion, O.T., Tan, S., Hoang, A.N., Cokol, M., and Aldridge, B.B.
(2016). Temporal and intrinsic factors of rifampicin tolerance in mycobacteria. Proc.
Natl. Acad. Sci. U. S. A. *113,* 8302.
Rock, J.M., Hopkins, F.F., Chavez, A., Diallo, M., Chase, M.R., Gerrick, E.R., Pritchard,
J.R., Church, G.M., Rubin, E.J., Sassetti, C.M., Schnappinger, D., and Fortune, S.M.
(2017). Programmable transcriptional repression in mycobacteria using an orthogonal
CRISPR interference platform. Nat. Microbiol. *2,* 16274.
Sena, C.B.C., Fukuda, T., Miyanagi, K., Matsumoto, S., Kobayashi, K., Murakami, Y.,
Maeda, Y., Kinoshita, T., and Morita, Y.S. (2010). Controlled Expression of Branch-
forming Mannosyltransferase Is Critical for Mycobacterial Lipoarabinomannan
Biosynthesis. J Biol Chem *285,* 13326-13336.
Soltan Mohammadi, N., Mafakheri, S., Abdali, N., Bárcena-Uribarri, I., Tauch, A., and
Benz, R. (2013). Identification and characterization of the channel-forming protein in the
cell wall of Corynebacterium amycolatum. Biochimica Et Biophysica Acta (BBA) -
Biomembranes *1828,* 2574-2582.

Su, C.C., Klenotic, P.A., Cui, M., Lyu, M., Morgan, C.E., and Yu, E.W. (2021).
Structures of the mycobacterial membrane protein MmpL3 reveal its mechanism of lipid
transport. PLoS Biol. *19,* e3001370.
Su, C., Klenotic, P.A., Bolla, J.R., Purdy, G.E., Robinson, C.V., and Yu, E.W. (2019).
MmpL3 is a lipid transporter that binds trehalose monomycolate and
phosphatidylethanolamine. Proceedings of the National Academy of Sciences - PNAS
*116,* 11241-11246.
Tahlan, K., Wilson, R., Kastrinsky, D.B., Arora, K., Nair, V., Fischer, E., Barnes, S.W.,
Walker, J.R., Alland, D., Barry, C.E., and Boshoff, H.I. (2012). SQ109 Targets MmpL3,
a Membrane Transporter of Trehalose Monomycolate Involved in Mycolic Acid Donation
to the Cell Wall Core of Mycobacterium tuberculosis. Antimicrob Agents Chemother *56,*
1797-1809.
Typas, A., Banzhaf, M., van den Berg van Saparoea, Bart, Verheul, J., Biboy, J.,
Nichols, R.J., Zietek, M., Beilharz, K., Kannenberg, K., von Rechenberg, M.*, et al.*
(2010). Regulation of Peptidoglycan Synthesis by Outer-Membrane Proteins. Cell *143,*
1097-1109.
Umare, M.D., Khedekar, P.B., and Chikhale, R.V. (2021). Mycobacterial Membrane
Protein Large 3 (MmpL3) Inhibitors: A Promising Approach to Combat Tuberculosis.
ChemMedChem *16,* 3136-3148.
van Kessel, J.C., and Hatfull, G.F. (2007). Recombineering in Mycobacterium
tuberculosis. Nat. Methods *4,* 147-152.
Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan,
D., Stroe, O., Wood, G., Laydon, A.*, et al.* (2022). AlphaFold Protein Structure
Database: massively expanding the structural coverage of protein-sequence space with
high-accuracy models. Nucleic Acids Research *50,* D439-D444.
Wu, K.J., Zhang, J., Baranowski, C., Leung, V., Rego, E.H., Morita, Y.S., Rubin, E.J.,
and Boutte, C.C. (2018). Characterization of Conserved and Novel Septal Factors in
Mycobacterium smegmatis. J. Bacteriol. *200,* 10.1128/JB.00649-17. Print 2018 Mar 15.
Wuo, M.G., Dulberger, C.L., Brown, R.A., Sturm, A., Ultee, E., Bloom-Ackermann, Z.,
Choi, C., Garner, E.C., Briegel, A., Hung, D.T., Rubin, E.J., and Kiessling, L.L. (2022).
Antibiotic action revealed by real-time imaging of the mycobacterial membrane.
Xu, Z., Meshcheryakov, V.A., Poce, G., and Chng, S.S. (2017). MmpL3 is the flippase
for mycolic acids in mycobacteria. Proc. Natl. Acad. Sci. U. S. A. *114,* 7993-7998.
Zhang, Y.J., Ioerger, T.R., Huttenhower, C., Long, J.E., Sassetti, C.M., Sacchettini, J.C.,
and Rubin, E.J. (2012). Global assessment of genomic regions required for growth in
Mycobacterium tuberculosis. PLoS Pathog. *8,* e1002946.
Zheng, S.Q., Palovcak, E., Armache, J., Verba, K.A., Cheng, Y., and Agard, D.A.
(2017). MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy. Nat Methods *14,* 331-332.
Zuber, B., Chami, M., Houssin, C., Dubochet, J., Griffiths, G., and Daffe, M. (2008).
Direct visualization of the outer membrane of mycobacteria and corynebacteria in their
native state. J. Bacteriol. *190,* 5672-5680.
835
836

**Figure 1 – figure supplement 1. (A)** Optical density over time of a strain whose only
copy of *pgfA* is tetracycline inducible, with and without ATC (+/-ATC). Points are means
of three biological replicates. Error bars are too small to be visible. Experiment was
performed at least 4 times on separate days, and representative data are shown.
ATC=anhydrotetracycline. Time on the *x*-axis corresponds to time after ATC was
removed. **(B)** Birth lengths and cell widths over time of depletion (orange = +ATC; red =
-ATC). Open circles represent individual cells (length: N = 261 (length) & 253 (width) for
+ATC; N = 355 (length) & 254 (width) -ATC). Closed circles are not fits to the data but
represent the mean of cells in 2.25-hr (length) or 2.75-hr (width) bins. The experiment
was performed twice on separate days. In +ATC cultures, cell density became too high
to measure length and width past ~33 hrs. ATC=anhydrotetracycline **(C)** L5-allele
swapping with *pgfA* or *pgfA-mRFP* expressed by the native promoter. Bars represent
means of 3 biological replicates. Error bars indicate standard deviation.

851
852 **Figure 1 – figure supplement 2** (A) Merged phase-contrast and fluorescence images of
853 Msm cells expressing PgfA-mRFP, eGFP-Wag31 and FtsZ-mCherry2b. (B) Average
854 fluorescent distributions over time (kymographs) from cells aligned from new to old poles
855 and from birth to division (N=20 PgfA-mRFP; N=48 FtsZ-mCherry2B; N=20 eGFP-
856 Wag31).
857

858
859
860 **Figure 2 – figure supplement 1.** The cell envelopes of PgfA-depleted show outer
861 membrane wall fraying at 24 hours left) and 33 hours (right) of depletion.

862
863
864 **Figure 2 – figure supplement 2.** Kymographs of a fluorescent fusion to MurJ in WT and
865 in Δ*lamA* cells. N=25.
866

| | (A) Coomassie | | (B) In-gel TAMRA fluorescence | | (C) Anti-FLAG Western blot | |
|---|---|---|---|---|---|---|
| | Input | Output | Input | Output | Input | Output |

| Lane | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| N-x-AlkTMM-C15 | - | - | + | + | - | - | + | + | - | - | + | + | - | - | + | + | - | - | + | + | - | - | + | + |
| UV (365nm) | - | + | - | + | - | + | - | + | - | + | - | + | - | + | - | + | - | + | - | + | - | + | - | + |
| CuAAC (AzTB) | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |

867
868
**Figure 3– figure supplement 1.** Full results of experiment present in Main text Figure
3A. Photoactivatable TMM analogue interacts with MSMEG_0317 in live *M. smegmatis*.
KG167 expressing 3x FLAG-tagged MSMEG_0317 was cultured with N-x-AlkTMM-C15
(100 µM), UV-irradiated, and lysed. Lysates were reacted with azido-TAMRA-biotin
reagent (AzTB) by Cu-catalyzed azide-alkyne cycloaddition (CuAAC "click" reaction), and
analyzed before (input) and after (output) avidin bead enrichment. Input and output
samples were analyzed by (A) Coomassie staining, (B) in-gel fluorescence scanning, and
(C) anti-FLAG Western blot. Data are representative of two independent experiments.
High-intensity band at low molecular weight in the Coomassie-stained gels represents
lysozyme used in the cell lysis procedure.
879

881
882
883
884 **Figure 3 – figure supplement 2. (A)** Output from TOPCONS (https://topcons.cbr.su.se/),
885 which aggregates the results of several transmembrane prediction algorithms. The input
886 is MSMEG_0317, or PgfA. **(B)** Results of allele swapping with the indicated PgfA fusion
887 proteins.
888

891
**Figure 4 – figure supplement 1.** Two additional biological replicates of LM/LAM analysis
during PgfA depletion. **(A)** As in Main Text Figure 4A , LM/LAM and other lipids were
extracted and separated by SDS-PAGE and TLC in cells expressing (+ATC) and depleted
for (-ATC) PgfA. Two biological replicates are shown. (B) For the same replicates optical
density was measured during depletion. To avoid changes in LM/LAM due to cell density,
at the indicated timepoints, half of the cultures were taken for lipid analysis, and replaced
with fresh media.

899
900

901
902 **Figure 4 – figure supplement 2. (A)** Results of exchanging the indicated *pgfA* alleles for
903 a wildtype copy of *pgfA*. Bars represent medians. **(B)** Cell morphology analysis of a
904 surviving transformant. Lines represent medians. **(C)** Results of exchanging the indicated
905 *pgfA* alleles for a wildtype copy of *pgfA*. Bars represent medians.
906
907
908 **Figure 2 – video supplement 1:** Phase time-lapse microscopy of cells carrying ATC-
909 inducible CRISRPi guides targeting either *pgfA* (left) or *mmpL3* (right). Exposure to ATC
910 occurs at the beginning of the video.
911
912