Discovery of an essential mycobacterial polar growth factor involved in cell envelope

integrity and LM/LAM transport

Kuldeepkumar R. Gupta[a], Kathryn C. Rahlwes[b], Chunyan Wang[c], Jun Liu[c], Yasu S.

Morita[b], and E. Hesper Rego[a,1]

[a]Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven,

CT 06519

[b]Department of Microbiology, University of Massachusetts, Amherst, MA 01003

[c]Department of Microbial Pathogenesis, Microbial Sciences Institute, Yale School of

Medicine, New Haven, CT, USA 06519

[1]To whom correspondence should be addressed: hesper.rego@yale.edu

ABSTRACT. Mycobacteria grow and divide differently compared to well-studied model

bacteria. They insert new cell envelope at their poles instead of along their side walls;

and, they lack obvious homologs of well-conserved cell growth and division proteins.

Furthermore, while mycobacteria are classified as Gram-positive, unlike other bacteria in

this taxon, they possess an outer membrane abundant in long-chain fatty acids and

several glycolipids. A better understanding of the unique factors that make this unusual

structure could lead to therapeutic targets for pathogenic mycobacteria. Here, we study

a gene of previously unknown function - *msmeg_0317* - predicted to be essential and

associated with mycobacterial cell growth and division proteins. We find that

transcriptional depletion of *msmeg_0317* leads to loss of polar growth, disruption of the

outer mycobacterial cell wall, and cell death. Surprisingly, we also observe that depletion

results in the accumulation of glycolipids lipomannan and lipoarabinomannan (LM/LAM)

in the periplasm. LM/LAM have been extensively studied in relation to infection but their

role in bacterial physiology is less clear. Altogether, our data suggest that MSMEG_0317,

renamed CwdA, is involved in the transport of LM/LAM to the mycobacterial surface, and

reveal unexpected connections between the correct localization of LM/LAM, polar growth,

and the structural integrity of the mycobacterial cell envelope.

Introduction

Mycobacterium tuberculosis (Mtb), the agent of human tuberculosis (TB), is

responsible for approximately 1.5 million deaths each year. One of the pathogen's

distinguishing features is its unusual cell envelope. Like nearly all other bacterial species,

the plasma membrane is surrounded by peptidoglycan, a rigid mesh-like structure made

up of sugars and peptides. However, in contrast to the peptidoglycan of other well-

characterized bacteria, mycobacterial peptidoglycan is covalently linked to the highly

branched hetero-polysaccharide arabinogalactan, which is, itself, covalently bound to

extremely long chained fatty acids called mycolic acids. In addition to this core structure,

several lipids and glycolipids, are abundantly and non-covalently interspersed across the

plasma membrane and mycobacterial cell wall (1, 2). This multi-laminate cell envelope is

a double-edged sword: it is a formidable barrier to many antibiotics, yet provides a number

of potentially targetable structures. Indeed, two, of the four, first-line TB antibiotics target

cell-wall biosynthesis.

In addition to its unusual cell envelope, mycobacterial physiology is distinctly

different than other model bacterial species. Notably, mycobacteria elongate from their

poles rather than along their side walls. The details of polar growth are poorly understood

(3-5). Many key proteins important for cell elongation in other species appear to be

missing in mycobacteria (4, 6). Likewise, it is almost completely unknown how proteins

involved in synthesizing the different layers, and free (glyco)lipids, work together to build

the complex cell envelope.

Given its unusual physiology, perhaps it is unsurprising that greater than 25% of

the mycobacterial genome encodes genes of unknown function (7). In our previous

studies on the mycobacterial division complex, 'the divisome', we precipitated known

divisome and elongasome components from *M. smegmatis* (Msm), and identified multiple

candidate mycobacterial-specific divisome proteins in the precipitate, (6, 8). One of these

was MSMEG_0317, a protein of unknown function that is highly conserved in

actinobacteria. The Mtb homolog of *msmeg_0317*, *rv0227c*, is predicted to be essential

(9, 10). Here, as we sought to understand the role of MSMEG_0317 during mycobacterial

growth and division.

MSMEG_0317 belongs to the DUF3068-domain super-family of proteins, which

are exclusively found in actinobacteria. One member of this family in corynebacteria has

channel activity, so this domain has been renamed PorA (11). In addition, by structure-

based homology prediction, MSMEG_0317 shares limited homology to mycobacterial

lipoproteins, LppX and LprG, which transport lipids, phthiocerol dimycocerosates (PDIM)

and triacylglycerol (TAG), respectively, to the outer mycobacterial cell wall (12, 13).

We find that MSMEG_0317 localizes to sites of new cell wall synthesis and is

indeed essential for cell viability in Msm. Transcriptional depletion leads to loss of cellular

elongation, reductive division, and eventually cell lysis. Surprisingly, we find that

lipomannan (LM), and lipoarabinomannan (LAM) – two large glycolipids abundant

throughout the mycobacterial cell envelope - accumulate as *msmeg_0317* is depleted,

and are shed into the culture media more abundantly upon MSMEG_0317

overexpression (1, 14). By cryo-electron microscopy, we find that the cell envelopes of

depleted cells are devoid of outer cell wall and are highly disrupted. Together, these

results show that MSMEG_0317 plays an important role in LM/LAM transport to the outer

mycobacterial cell wall, polar elongation, and the structural organization of the

mycobacterial cell wall. As such, we propose to rename *msmeg_0317*, *cwdA*, for <u>c</u>ell <u>w</u>all

<u>d</u>isruption, mutant <u>A</u>.

Results

***cwdA* is essential for polar growth and cell wall integrity**

To determine the essentiality of *cwdA*, we used an allele swapping strategy (15).

Briefly, in a strain of Msm whose only copy of *cwdA* was at the L5 phage integration site

(16), we exchanged *cwdA* for either an empty vector or for another copy of itself (Fig. 1A).

Consistent with *cwdA* being essential for cell growth, we observed approximately 1000-

fold fewer colonies when we exchanged *cwdA* with an empty vector compared to

exchanging it for another copy of itself (Fig. 1A).

To further confirm the essentially of *cwdA* we constructed a strain in which the only

copy of the gene was tetracycline inducible (Fig. 1B). Removal of anhydrotetracycline

(ATC) from the culture media prevented cell growth (Fig. 1B). By time-lapse microscopy

we observed that, while cells expressing *cwdA* became longer, on average, as cell density

increased within the microfluidic device (Fig. 2A, C & Movie S1), cells depleted for *cwdA*

stopped elongating but continued to divide (reductive division), became wider, and, in

many instances, eventually lysed (Fig. 2B,C & Movie S2). Together these data show that

cwdA is essential for polar elongation in Msm.

CwdA localizes to the sites of cell wall synthesis

As our data suggested CwdA is important for polar growth, we hypothesized that

it would localize to the poles, which are the site of new cell wall synthesis in mycobacteria.

To test this hypothesis, we made a translational fusion to CwdA with mRFP expressed by

its native promoter. By allele swapping, we find that *cwdA-mrfp* restores bacterial growth

(Fig. S1) and thus encodes a functional *cwdA*. Fluorescence microscopy at a single time-

point showed that CwdA-mRFP localizes to mid-cell and to the poles (Fig. 3A). In polar

growing bacteria, the site of division eventually becomes the site of elongation. Thus, in

addition to localizing to the poles, elongation-complex proteins will also localize to the site

of division early during the cell cycle. To disentangle whether CwdA is also part of the

division complex we visualized CwdA-mRFP by time-lapse microscopy. By constructing

kymographs averaged over many cells, we compared the fluorescence distribution over

the course of the cell cycle to the earliest known marker for the division (FtsZ-mCherry2B)

complex and the elongation complex (eGFP-Wag31). CwdA-mRFP localizes to the site

of division after FtsZ (Fig. 3A, B), but before Wag31 (Fig. 3C). These data are consistent

with CwdA being a member of both the mycobacterial division and elongation complexes,

but its essential function is as part of the elongation complex.

CwdA is involved in LM/LAM transport and cell wall integrity

The corynebacterial homolog of *cwdA* was recently implicated in LM/LAM

biosynthesis (17), as a deletion mutant of the *Corynebacterium glutamicum* homolog

results in truncated LM. To test if this is also the case in mycobacteria we used thin layer

chromatography (TLC) to quantify the amounts of different lipid species in cells depleted

for CwdA. In contrast to the results obtained in *C. glutamicum*, we observe a dramatic

increase in the total amount of cell-associated of LM/LAM during depletion (Fig. 4A), while

other lipids (Fig. 4 B, C & Fig. S4) and sugars (Fig. S2) remain constant. Likewise, when

we inducibly over-expressed *cwdA* from a multicopy plasmid, we observed less LM/LAM

associated with Msm cells (Fig. S3A & S5).

These data are not consistent with the proposed function of the corynebacteria

cwdA homolog. As, by sequence-based homology modeling, CwdA shares some

homology to LppX and LprG, proteins known to transport lipids from the inner membrane

to the outer membrane in mycobacteria, we hypothesized that CwdA could be involved in

transporting LM/LAM to the outer membrane where it is likely to be shed (18-20).

Consistent with this hypothesis, upon over-expression of CwdA we observed more

LM/LAM in the culture supernatant (Fig. S3B). These data show that CwdA is involved in

transporting LM/LAM to the cell surface of mycobacteria (Fig. S3B).

CwdA is important for outer cell wall integrity.

Given the abrupt and violent nature of lysis when cells are depleted for CwdA, we

hypothesized that CwdA-depleted cells had gross alterations to their cell envelope. Thus,

we visualized the cell envelope of wild type and mutant cells by high resolution cryo-

electron microscopy (cryo-EM). As has been previously observed (21, 22), wild type Msm

cells exhibit a distinct plasma membrane and outer membrane that is approximately 50-

nm thick (Fig. 5A-C). Surprisingly, we find that cells depleted for CwdA have largely

unstructured cell walls, and appear devoid of electron density in the outer cell wall (Fig. 5

D-F). Early during depletion, before cell death, we observe dramatic fraying of the outer

membrane, and large invaginations of plasma membrane (Fig. 6). Thus, CwdA is

important for maintaining the structural integrity of the outer mycobacterial cell wall.

Collectively, our data show that CwdA is an essential polar growth factor involved

in transporting LM/LAM to the outer wall. Further, our data suggest that the correct

localization of LM/LAM is important for the structural organization and integrity of the

mycobacterial cell envelope.

Discussion

The extraordinary diversity of the bacterial kingdom has led to a plethora of

bacterial cell shapes and geometries of cell growth and division. Unlike well-studied

model bacteria, several organisms, including the major human pathogen *M. tuberculosis,*

grow from their poles rather than their side-walls. Our understanding of polar growth is at

a nascent stage compared to our knowledge of side-wall growth. Mycobacteria, and

related organisms, have an additional challenge of orchestrating the biosynthesis and

transport of several additional covalently and non-covalently attached lipids and sugars

to their thick layer of peptidoglycan. How these organisms coordinate the biogenesis of

this complex structure to elongate and divide is almost completely unknown.

Here, we have identified a highly conserved actinobacterial protein,

MSMEG_0317, in *M. smegmatis* and propose to rename it CwdA. We show that this gene

is essential in Msm. The Mtb homolog, *rv0227c*, is almost certainly essential as well, as

no transposon insertions have been found in this gene, even in data sets which combine

insertion counts from high density mutagenesis across multiple independent experiments

(9).

Our data suggest that CwdA's essential function occurs during polar elongation,

as cells depleted for CwdA stop elongating, and CwdA-mRFP localizes to the poles.

Interestingly, CwdA also localizes to the site of division, yet depleted cells continue to

divide. This is consistent with recent work from several groups, which suggests that the

essentiality of certain cell envelope structures is different for elongation and division. For

example, *C. glutamicum* cells treated with ethambutol, which inhibits the polymerization

of Arabian, a core structure in both arabinogalactan and LAM, stop elongating, but

continue to divide (23, 24).

What is the function of CwdA? Our work suggests that CwdA is involved in the

transport of LM/LAM to the cell surface either directly or indirectly, and for maintaining the

structural integrity of that mycobacterial cell wall. Upon depletion, the levels of several

other lipids remain constant, suggesting that *cwdA* may be acting on LM/LAM directly,

though our lipid profiling was not exhaustive and we cannot rule out the possibility that

another structural lipid is affected by CwdA depletion. However, supporting the notion that

CwdA specifically acts upon LM/LAM, we observe a temporal correlation, up until the time

of cell lysis, between the accumulation of LM/LAM and the disruption of the envelope.

This leads the question: Could LM/LAM be essential components of the outer cell

envelope? There has been a good deal of speculation as to the localization of LM/LAM in

the mycobacterial envelope, but several groups have now shown that at least a fraction

of LM/LAM resides in the outer cell wall (19, 20). Our data are consistent with this model.

In addition, our data suggest that either LM/LAM are important structural components of

the outer cell wall or the accumulation of these large glycolipids in the periplasm leads to

envelope disruption.

One implication from our work is that the transport of lipids to the cell surface is

needed for polar elongation, but not division, suggesting that there could be various

feedback loops to ensure complete construction of the elaborate mycobacterial cell

envelope for insertion of new material to elongate. These checkpoints appear missing

during division. This notion is supported by other mutants defective in transporting

important cell wall components like trehalose monomycolate, which, like the *cwdA*

mutant, undergo reductive cell division (25).

The mycobacterial cell envelope remains an enduring target for the development

of new TB therapeutics. The standard TB therapy uses a combination of four drugs; two

of these drugs target cell wall synthesis. In addition to the core, and covalently linked

structure of peptidoglycan, arabinogalactan, and mycolic acids, the mycobacterial cell

envelope is also rich with many other lipids. The functional role of these lipids in bacterial

cell physiology is relatively unexplored but they are likely important, if not essential,

components of the cell envelope. Our data argue that a more complete understanding of

the transport of these molecules could lead to new targets for the development of anti-TB

therapeutics.

Materials and Methods

Bacterial strains and culture conditions: *M. smegmatis* mc^{2}155 was grown in

Middlebrook 7H9 broth supplemented with 0.05% Tween80, 0.2% glycerol, 5 gm/L

albumins, 2 gm/L dextrose and 0.003 gm/L catalase or plated on LB agar. *Escherichia*

coli DH5α cells were grown in LB broth or on LB agar plates. Concentrations of antibiotics

used for *M. smegmatis* is as follows: 20 μg/ml zeocin, 25 μg/ml kanamycin, 50 μg/ml

hygromycin, and 20 μg/ml nourseothricin. Concentrations of antibiotics used for *E. coli* is

as follows: 40 μg/ml zeocin, 50 μg/ml kanamycin, 100 μg/ml hygromycin, and 40 μg/ml

nourseothricin.

Plasmids and strains construction: Before deleting the native copy of

msmeg_0317/cwdA, a merodiploid strain was created by inserting a second copy of

msmeg_0317/cwdA gene under pTetO promoter using a Kan^R L5 integrating vector.

Subsequently, in the merodiploid strain, an in-frame deletion was made by replacing the

native copy of *msmeg_0317/cwdA gene* with a zeocin resistance cassette flanked by lox

sites using recombineering (16). The plasmid constructs were made using isothermal

Gibson assembly whereby insert and vector backbone shared 20-25 bp of homology.

msmeg_0317/cwdA and its fluorescent variant were cloned in a Nat^R L5 integrating vector

for allele exchange at L5 site. The *msmeg_0317/cwdA* depletion strain was made by

transforming an episomally Hyg^R marked vector constitutively expressing TetR repressor

into a strain expressing *msmeg_0317* gene under pTetO promoter. mRFP was fused to

the c-terminal of *msmeg_0317* gene cloned with 200 bp upstream sequence to capture

the native promoter. *ftsZ-mCherry*2B and *eGFP-wag31* genes were cloned under medium

strength promoter ptb21 and integrated at tweety (tw) and L5 phage integration sites,

respectively.

MSMEG_0317 Depletion: To transcriptionally deplete *msmeg_0317/cwdA* in *M.*

smegmatis, anhydrotetracycline-inducible Tet-ON system was used. The

msmeg_0317/cwdA gene was expressed from pTetO promoter, while the TetR repressor

was constitutively expressed in trans from an episomal vector. MSMEG_0317 was

depleted by removing anhydrotetracyline from the medium and cells were grown for 18

hours. Subsequently, MSMEG_0317 depleted cells were re-diluted in fresh medium.

Alternatively, depletion was achieved by removing and adding fresh medium every three

hours from 18hr-old depletion cultures.

Time-lapse microscopy: An inverted Nikon Ti-E microscope was used for the

time-lapse imaging. An environmental chamber was maintained at 37°C. Cells were

cultivated in an B04 microfluidics plate from CellAsic; continuously supplied with fresh

medium; and imaged every 15 minutes using phase contrast.

Kymograph analysis. Time-lapse images were analyzed in open source image

analysis software Fiji (26) and a custom MATLAB program was used to generate

kymographs. Specifically, for a single cell, in Fiji, a 5-pixel wide segmented line was drawn

from the new pole to the old pole at each time point during the cell cycle. This was

repeated on 20-50 cells. These line profiles were then imported into MatLab, where a

custom script was used to generate an average kymograph by 2D interpolation of the

individual kymographs.

Lipid extraction and analysis. Crude lipids were extracted from the

MSMEG_0317*/cwdA* depletion strains after 18, 24, 27, 30, and 33 hrs without

anhydrotetracycline, following a previously published protocol(27). Briefly, LM and LAM

were extracted from the delipidated pellet by incubation with phenol/water (1:1) for 2 hrs

at 55°C. Phospholipids and PIMs were separated by high performance thin layer

chromatography (HPTLC) silica gel 60 (EMD Merck) using chloroform/methanol/ 13M

ammonia/1M ammonium acetate/ water (180:140:9:9:23). Phospholipids were visualized

via cupric acetate staining. Carbohydrates extracted in the aqueous phase of the butanol-

water phase partitioning were separated by HPTLC using the solvent n-butanol/

isoproponal/ water (13:12:4). Carbohydrates and PIMs were visualized with orcinol

staining as described (Sena 2010). LM/LAM samples were separated by SDS-PAGE

(15% gel) and visualized using ProQ Emerald 488 glycan staining kit (Life Technologies)

as described in (28). To detect LM/LAM in culture supernatants, the supernatants were

initially treated with a final concentration of 50 μg/ml Proteinase K for 4 hours at 50°C.

The treated supernatants were electrophoresed on 15% SDS-PAGE. LM/LAM were

blotted onto nitrocellulose at 20 V for 45 minutes using semi-dry transfer method. Post-

transfer, the membrane was blocked by 5% milk. The blocked membrane was then

probed with CS-35 antibody (BEI Resources, NIH) at 1:250 dilution at 4°C overnight. The

membrane was washed with 1X TBST five times for five minutes each. Post-washing, it

was probed with anti-mouse secondary for one hour at room temperature. Membrane

was then washed five times for five minutes with 1X TBST. Thermo Scientific's west dura

chemiluminescent reagent was used to develop the membrane.

Sample preparation and image collection for Cryo-EM: Wild type and

MSMEG_0317/CwdA depleted Msm were pelleted, washed twice with 1X phosphate

buffered saline (PBS), and suspended in a very small volume of PBS to get a very dense

suspension of bacterial cells. The dense culture was subsequently mixed with 10-nm

colloidal gold and deposited onto fresh glow-discharged holey carbon grids. The grids

were then blotted with filter paper manually and rapidly frozen in liquid ethane. The frozen

grids were transferred under -170 °C into a 300kV Titan Krios equipped with K2 Summit

direct detector, a quantum energy filter (Gatan, Pleasanton, CA) and an FEI volta phase

plate (FEI). The data was collected using SerialEM (29) and Digital Micrograph

softwares. The motion correction for each movie was done with MotionCor2 (30). The

microscope was operated at 300kV with a magnification of 3,600x and 26,000 x for low-

mag map and high-mag map respectively. The grey levels of each micrograph are

obtained using MatLab.

Acknowledgements

We thank Cara Boutte and Maikel Boot for helpful suggestions and for critical reading of

the manuscript. Work carried out in this manuscript was supported by career award at

scientific interface (CASI) from Burroughs Welcome Fund and Searle and Pew funds to

E.H.R.

References:

292
1.   Jackson M (2014) The mycobacterial cell envelope-lipids. *Cold Spring Harb Perspect Med*
4(10).
2.   Jankute M, Cox JA, Harrison J, & Besra GS (2015) Assembly of the Mycobacterial Cell
Wall. *Annu Rev Microbiol* 69:405-423.
3.   Kieser KJ & Rubin EJ (2014) How sisters grow apart: mycobacterial growth and division.
Nat Rev Microbiol 12(8):550-562.
4.   Hett EC & Rubin EJ (2008) Bacterial growth and cell division: a mycobacterial
perspective. *Microbiol Mol Biol Rev* 72(1):126-156, table of contents.
5.   Baranowski C, Rego EH, & Rubin EJ (2019) The Dream of a Mycobacterium.
Microbiology spectrum 7(2).
6.   Wu KJ*, et al.* (2018) Characterization of Conserved and Novel Septal Factors in
Mycobacterium smegmatis. *J Bacteriol* 200(6).

7.   Doerks T, van Noort V, Minguez P, & Bork P (2012) Annotation of the M. tuberculosis
hypothetical orfeome: adding functional information to more than half of the
uncharacterized proteins. *PLoS One* 7(4):e34302.
8.   Rego EH, Audette RE, & Rubin EJ (2017) Deletion of a mycobacterial divisome factor
collapses single-cell phenotypic heterogeneity. *Nature* 546(7656):153-157.
9.   DeJesus MA*, et al.* (2017) Comprehensive Essentiality Analysis of the Mycobacterium
tuberculosis Genome via Saturating Transposon Mutagenesis. *mBio* 8(1).
10. Zhang YJ*, et al.* (2012) Global assessment of genomic regions required for growth in
Mycobacterium tuberculosis. *PLoS Pathog* 8(9):e1002946.
11. Soltan Mohammadi N*, et al.* (2013) Identification and characterization of the channel-
forming protein in the cell wall of Corynebacterium amycolatum. *Biochimica et biophysica*
acta 1828(11):2574-2582.
12. Sulzenbacher G*, et al.* (2006) LppX is a lipoprotein required for the translocation of
phthiocerol dimycocerosates to the surface of Mycobacterium tuberculosis. *The EMBO*
journal 25(7):1436-1444.
13. Martinot AJ*, et al.* (2016) Mycobacterial Metabolic Syndrome: LprG and Rv1410 Regulate
Triacylglyceride Levels, Growth Rate and Virulence in Mycobacterium tuberculosis. *PLoS*
Pathog 12(1):e1005351.
14. Mishra AK, Driessen NN, Appelmelk BJ, & Besra GS (2011) Lipoarabinomannan and
related glycoconjugates: structure, biogenesis and role in Mycobacterium tuberculosis
physiology and host-pathogen interaction. *FEMS Microbiol Rev* 35(6):1126-1157.
15. Pashley CA & Parish T (2003) Efficient switching of mycobacteriophage L5-based
integrating plasmids in Mycobacterium tuberculosis. *FEMS Microbiol Lett* 229(2):211-
215.
16. van Kessel JC & Hatfull GF (2007) Recombineering in Mycobacterium tuberculosis. *Nat*
Methods 4(2):147-152.
17. Cashmore TJ*, et al.* (2017) Identification of a Membrane Protein Required for Lipomannan
Maturation and Lipoarabinomannan Synthesis in Corynebacterineae. *J Biol Chem*
292(12):4976-4986.
18. Pitarque S*, et al.* (2008) The immunomodulatory lipoglycans, lipoarabinomannan and
lipomannan, are exposed at the mycobacterial cell surface. *Tuberculosis (Edinburgh,*
Scotland) 88(6):560-565.
19. Gaur RL*, et al.* (2015) Correction: LprG-Mediated Surface Expression of
Lipoarabinomannan Is Essential for Virulence of Mycobacterium tuberculosis. *PLoS*
Pathog 11(12):e1005336.
20. Shukla S*, et al.* (2014) Mycobacterium tuberculosis lipoprotein LprG binds
lipoarabinomannan and determines its cell envelope localization to control
phagolysosomal fusion. *PLoS Pathog* 10(10):e1004471.
21. Zuber B*, et al.* (2008) Direct visualization of the outer membrane of mycobacteria and
corynebacteria in their native state. *J Bacteriol* 190(16):5672-5680.
22. Hoffmann C, Leis A, Niederweis M, Plitzko JM, & Engelhardt H (2008) Disclosure of the
mycobacterial outer membrane: cryo-electron tomography and vitreous sections reveal the
lipid bilayer structure. *Proc Natl Acad Sci U S A* 105(10):3963-3967.
23. Zhou X*, et al.* (2019) Sequential assembly of the septal cell envelope prior to V snapping
in Corynebacterium glutamicum. *Nature chemical biology* 15(3):221-231.

24. Schubert K, *et al.* (2017) The Antituberculosis Drug Ethambutol Selectively Blocks Apical
Growth in CMN Group Bacteria. *mBio* 8(1).
25. Fay A, *et al.* (2019) Two accessory proteins govern MmpL3 mycolic acid transport in
mycobacteria. *bioRxiv*:581447.
26. Schindelin J, *et al.* (2012) Fiji: an open-source platform for biological-image analysis. *Nat*
Methods 9(7):676-682.
27. Sena CB, *et al.* (2010) Controlled expression of branch-forming mannosyltransferase is
critical for mycobacterial lipoarabinomannan biosynthesis. *J Biol Chem* 285(18):13326-
13336.
28. Rahlwes KC, *et al.* (2017) The cell envelope-associated phospholipid-binding protein
LmeA is required for mannan polymerization in mycobacteria. *J Biol Chem*
292(42):17407-17417.
29. Mastronarde DN (2005) Automated electron microscope tomography using robust
prediction of specimen movements. *J Struct Biol* 152(1):36-51.
30. Zheng SQ, *et al.* (2017) MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* 14(4):331-332.

Fig. 1. *cwdA* is essential in *M. smegmatis*. (A) L5-integrating vectors carrying either a

copy of *cwdA* or a gene encoding *tetR* were transformed into a strain whose only copy of

cwdA was at the L5 integration site, and transformants were counted. Bars represent

means of biological triplicates and are representative data of 3 independently performed

experiments. (B) Optical density over time of a strain whose only copy of *cwdA* is

tetracycline inducible (+/-ATC). Points are means of three biological replicates. Error bars

are too small to be visible. Experiment was performed at least 4 times on separate days,

and representative data are shown. ATC=anhydrotetracycline. Time on the x-axis

corresponds to time after ATC was removed.

Fig. 2. *cwdA*-depletion causes reductive cell division and eventual cell lysis. (A,B)
The strain shown in Fig. 1B was visualized by time-lapse microscopy in the presence (A)
or absence of ATC (B). Cells were loaded into a microfluidic device 18 hours after the
removal of ATC (B) or a mock control (A). Representative frames are 3 hours apart. (C)
Birth lengths and cell widths over time of depletion (orange = +ATC; red = -ATC). Open
circles represent individual cells (length: N = 261 (length) & 253 (width) for +ATC; N =
355 (length) & 254 (width) -ATC). Closed circles are not fits to the data but represent the
mean of cells in 2.25-hr (length) or 2.75-hr (width) bins. The experiment was performed
twice on separate days. In +ATC cultures, cell density became too high to measure length
and width past ~33 hrs.

Fig. 3. CwdA localizes to the sites of division and elongaton. (A) Merged phase-
contrast and fluorescence images of Msm cells expressing CwdA-mRFP, eGFP-Wag31
and FtsZ-mCherry2b. (B) Average fluorescent distributions over time (kymographs) from
cells aligned from new to old poles and from birth to division (N=20 CwdA-mRFP; N=48
FtsZ-mCherry2B; N=20 eGFP-Wag31).

Fig. 4. LM/LAM accumulate as *cwdA* is depleted. (A-C) In cells depleted for *cwdA*, LM/LAM are separated by SDS-PAGE (A); non-polar and polar PIMs (B), and plasma-membrane lipids (C) are measured by TLC.

Fig. 5. Depletion of CwdA leads to loss of outer cell wall. The cell envelope of a
representative wild type *M. smegmatis* cell (A-C) and a cell depleted for CwdA for 24
hours (D-F) visualized (A,B,D,E) and measured (D,F) by cryo-electron microscopy. PM =
plasma membrane; OM = outer membrane.

429
430

Fig. 6. Cryo-Electron Microscopy of CwdA-depleted cells (A-H) The cell envelopes of CwdA-depleted cells, showing plasma membrane invaginations (A,E,C&H) and outer membrane wall fraying (B-H). (A,E) 24 hrs; (B,F) 24 hrs; (C,G) 24 hrs; (D,H) 33 hrs of depletion.

439 **Supplemental Figures**
440

441
442
443 **Fig. S1. CwdA-mRFP encodes a functional copy.** L5-allele swapping with *cwdA* or
444 *cwdA-mRFP* expressed by the native promoter. Bars represent means of 3 biological
445 replicates. Error is standard deviation.
446

447
Fig. S2. TLC of unidentified sugars during *cwdA* depletion. This TLC shows that during
CwdA depletion the levels of monosaccharides and disaccharides remain constant.
450

451
452

453
454 **Fig. S3.** Detection of LM/LAM during *cwdA* over-expression. A) Detection of cell
455 associated LM/LAM using SDS-PAGE in *cwdA* over-expression. B) Detection of LAM in
456 the culture supernatants using western blot during *cwdA* over-expression using CS-35
457 antibody. CS-35 antibody is specific to arabinan moiety, hence only LAM is detected in
458 the western blot experiments.

Replicate 1   Replicate 2

control   hours without ATC   marker   control   hours without ATC

18 24 27 30 33   18 24 27 30 33

LAM

LM

Ac_2PIM_2
Ac_1PIM_2
Ac_2PIM_6
Ac_1PIM_6

CL
PE

PI
Ac_1PIM_2

484
485
486 **Fig. S4. Accumulation of LM/LAM during *cwdA* depletion**. Two independent biological
487 replicates showing that *cwdA* depletion specifically causes an accumulation of LM/LAM
488 before cell death is triggered. Levels of the rest of the cell wall associated lipids remains
489 constant during *cwdA* depletion.
490
491
492
493

494
495
496 **Fig. S5. Detection of cell wall associated lipid during *cwdA* over-expression.**
497 LM/LAM levels associated with mycobacterial cells decrease upon *cwdA* over-
498 expression. However, the levels of the rest of the lipids remain constant.
499
500