1

PilT and PilU are homohexameric ATPases that coordinate to retract type IVa pili

Jennifer L. Chlebek[1], Hannah Q. Hughes[1], Aleksandra S. Ratkiewicz[2], Rasman Rayyan[2],

Joseph Che-Yen Wang[3], Brittany E. Herrin[1], Triana N. Dalia[1], Nicolas Biais[2], and Ankur B.

Dalia[1,*]

[1]Department of Biology, Indiana University, 1001 E. 3rd Street, Bloomington, IN 47405.

[2]Biology Department, CUNY Brooklyn College, 2900 Bedford Avenue, Brooklyn, NY 11210 and
Graduate Center of CUNY, 365 5th Avenue, NY 10016.

[3]Electron Microscopy Center, Indiana University, Bloomington, IN 47405.

*Correspondence to: ankdalia@indiana.edu.

Short Title: PilU is a PilT-dependent type IVa pilus retraction ATPase

Abstract

Bacterial type IV pili are critical for diverse biological processes including horizontal gene transfer, surface sensing, biofilm formation, adherence, motility, and virulence. These dynamic appendages extend and retract from the cell surface. In type IVa pilus systems, extension occurs through the action of an extension ATPase, PilB, while optimal retraction requires the action of a retraction ATPase, PilT. Many type IVa systems also encode a homolog of PilT called PilU. However, the function of this protein has remained unclear because *pilU* mutants largely lack obvious phenotypes. Here, we study the type IVa competence pilus of *Vibrio cholerae* as a model system to define the role of PilU. We show that the ATPase activity of PilU is critical for optimal pilus retraction in PilT ATPase mutants. PilU does not, however, contribute to pilus retraction in Δ*pilT* strains. Thus, these data suggest that PilU is a *bona fide* retraction ATPase that supports pilus retraction in a PilT-dependent manner. We also found that a Δ*pilU* mutant exhibited a reduction in the force of retraction suggesting that PilU is important for generating maximal retraction forces. Additional *in vitro* and *in vivo* data show that PilT and PilU act as independent homo-hexamers that likely form a complex to facilitate pilus retraction. Finally, we demonstrate that the role of PilU as a PilT-dependent retraction ATPase is conserved in *Acinetobacter baylyi*, suggesting that the role of PilU described here may be broadly applicable to diverse type IVa pilus systems.

Author Summary

Almost all bacterial species use thin surface appendages called pili to interact with their environments. These structures are critical for the virulence of many pathogens and represent one major way that bacteria share DNA with one another, which contributes to the spread of antibiotic resistance. To carry out their function, pili dynamically extend and retract from the

43   bacterial surface. Here, we show that retraction of pili in some systems is determined by the

44   combined activity of two motor ATPase proteins.

Introduction

Type IV pili are ubiquitous surface appendages in Gram-negative bacteria that promote diverse

activities including attachment, virulence, biofilm formation, horizontal gene transfer, and

twitching motility [1-5]. These structures can dynamically extend and retract from the cell

surface, which is often critical for their function. A detailed mechanistic understanding of pilus

dynamic activity, however, remains lacking.

Type IV pili are composed almost exclusively of a single protein called the major pilin, which

forms a helical fiber that extends from the cell surface [6]. Extension and retraction of the pilus is

hypothesized to occur through the interaction of cytoplasmic hexameric ATPases with the inner

membrane platform of the pilus machine, whereby ATP hydrolysis likely facilitates turning of the

platform in order to incorporate or remove major pilin subunits from the pilus fiber [7-9]. All type

IV pilus systems encode a predicted ATPase (PilB) that facilitates pilus extension [7, 10-12].

The type IVa pilus systems also contain a retraction ATPase (PilT) which depolymerizes the

pilus fiber and recycles pilin subunits into the inner membrane. Many type IVa pilus systems

also encode another putative ATPase called PilU. PilU is a homolog of the retraction ATPase

PilT, however, the function of this protein has remained unclear in most pilus systems because

pilU mutants largely lack obvious phenotypes. A notable exception is *Pseudomonas aeruginosa*,

where *pilU* mutants exhibit reduced twitching motility [13].

The facultative bacterial pathogen *Vibrio cholerae* uses a type IV competence pilus for DNA

uptake during natural transformation. We have recently demonstrated that the type IV

competence pili of *V. cholerae* can be fluorescently labeled by using a technique [14-16] in

which an amino acid of the major pilin subunit PilA is replaced with a cysteine (PilA^{S67C}) [17],

which allows for subsequent labeling with a fluorescently-labeled thiol-reactive maleimide dye

(AF488-mal). This labeling approach does not impede dynamic pilus activity and allows for the

direct observation and measurement of pilus extension and retraction by time-lapse

epifluorescence microscopy. Using this labeling technique, we have recently demonstrated that

the retraction of DNA-bound type IV competence pili is required to initiate the process of DNA

uptake during natural transformation [17].

Here, we employ our pilus labeling approach and complementary molecular methods to address

the role that PilU plays in type IV competence pilus retraction, and we extend our findings to the

type IVa competence pilus of *Acinetobacter baylyi*. Our results indicate that PilT and PilU form

independent homo-hexamers and that the ATPase activity of PilU promotes type IV pilus

retraction in a PilT-dependent manner.

Results

PilU promotes type IV competence pilus retraction in a PilT-dependent manner

In order to study the function and role of ATPases in the retraction of type IV competence pili in

V. cholerae, we utilize two main methods: (1) our labeling approach to directly measure pilus

retraction via epifluorescence time-lapse microscopy and (2) assessing natural transformation,

which is dependent on competence pilus retraction [17, 18]. Also, as for other pili [19-21],

competence pili can mediate cell-to-cell interactions, which causes hyperpiliated strains to

aggregate and pellet to the bottom of culture tubes [22]. Consistent with prior work [17, 18, 23-

28], mutant strains that lack the retraction ATPase PilT are poorly transformable, exhibit

markedly reduced rates of pilus retraction, aggregate, and are hyperpiliated (≥3 pili per cell)

compared to the parent strain (**Fig 1A-D**). Qualitatively, the number of cells that exhibit dynamic

pilus activity is reduced in the Δ*pilT* strain as observed by time-lapse epifluorescence

microscopy, which is consistent with the hyperpiliated phenotype of this mutant. The residual

retraction in a Δ*pilT* strain is not due to PilU because a Δ*pilT* Δ*pilU* double mutant strain displays

the same phenotype as a Δ*pilT* strain (**Fig 1A-D**). By contrast, Δ*pilU* mutants are largely

indistinguishable from the parent, other than a ~1.3-fold reduction in the rate of pilus retraction

(**Fig 1B**). Thus, these data support a model where PilT plays the dominant role in pilus

retraction, which is consistent with prior reports [17, 18, 26].

PilT is a predicted ATPase and its ATPase activity has been shown to be important in the

function of other type IV pilus systems [12, 29]. Therefore, we hypothesized that the ATPase

activity of PilT would be required for retraction of the competence pilus. We found, however, that

strains harboring mutations in PilT that ablate the ATPase activity of this protein—due to a

mutation in either an invariantly conserved lysine in the Walker A motif required for ATP binding

(*pilT^{K136A}*), a mutation in a conserved glutamic acid in the Walker B motif required for ATP

hydrolysis (*pilT^{E204A}*), or a double mutant containing both mutations (*pilT$^{E204A/K136A}$*)— all

transform similarly to the parent strain and retract competence pili at rates that are at most ~4-

fold lower than the parent (**Fig 1A-B**). This is in stark contrast to a Δ*pilT* strain where the

transformation frequency is reduced ~4-logs and retraction is >50-fold slower compared to the

parent strain (**Fig 1A-B**). These results were quite surprising and suggested that either (1) the

ATPase activity of the retraction machinery is not critical for promoting optimal pilus retraction or

(2) that another retraction ATPase may promote retraction in the absence of PilT ATPase

activity.

We tested the latter model and specifically assessed whether PilU could promote retraction in

PilT ATPase-deficient mutants. Indeed, we found that *pilT^{K136A}*Δ*pilU*, *pilT^{E204A}*Δ*pilU* and

*pilT$^{E204A/K136A}$*Δ*pilU* strains all display reduced transformation, are hyperpiliated, aggregate, and

exhibit markedly reduced retraction rates similar to a Δ*pilT* Δ*pilU* strain (**Fig 1A-D**). This is

consistent with PilU promoting retraction in the *pilT^{K136A}*, *pilT^{E204A}* and *pilT^{E204A/K136A}*

backgrounds. Importantly, the *pilT^{K136A}*Δ*pilU*, *pilT^{E204A}*Δ*pilU* and *pilT^{E204A/K136A}*Δ*pilU* strains could

be complemented via ectopic expression of *pilT* or *pilU* (**S1 Fig**). Since loss of *pilT* alone (i.e.

Δ*pilT*) results in a phenotype comparable to a Δ*pilT* Δ*pilU* mutant, this result indicates that PilU

is not sufficient to drive pilus retraction in the absence of PilT (**Fig 1A-D**). Consistent with this,

ectopic expression of *pilU* does not rescue the transformation deficit of a Δ*pilT* Δ*pilU* mutant (**S1**

Fig). Together, these data suggest that PilU supports pilus retraction in a PilT-dependent

manner.

Interestingly, the *pilT^{E204A}* strain is hyperpiliated (**Fig 1D**). Qualitatively, the frequency of cells

that exhibit pilus retraction in this mutant is much lower than the parent and comparable to the

Δ*pilT* Δ*pilU* strain (as assessed by epifluorescence time-lapse microscopy), which may explain

this hyperpiliated phenotype; however, when pilus retraction is observed, the rate of retraction is

closer to that of the parent than the Δ*pilT* Δ*pilU* mutant (**Fig 1B**). Considering the canonical

functions of the Walker A and B motifs in ATP-binding and hydrolysis, respectively, the Walker

B *pilT^{E204A}* mutant should be able to bind to ATP but not hydrolyze it. Because the ATP-binding

deficient *pilT^{K136A}* Walker A mutant exhibited parental levels of piliation (**Fig 1D**), we

hypothesized that the hyperpiliation observed in *pilT^{E204A}* may be due to the ATP-binding state

of this mutant. Consistent with this, we found that the Walker A and B double mutant

(*pilT^{E204A/K136A}*) exhibits parent levels of piliation in contrast to the *pilT^{E204A}* single mutant (**Fig**

1D). These results indicate that the phenotype of the ATP-binding deficient Walker A mutant

(*pilT^{K136A}*) is dominant over the ATP-hydrolysis deficient *pilT^{E204A}* Walker B mutation. Both of

these mutations should result in loss of ATPase activity. Therefore, to avoid phenotypic

consequences of variable states of ATP binding associated with the *pilT^{E204A}* mutant, we utilized

the *pilT^{K136A}* mutant as an ATPase-defective allele of PilT moving forward.

PilU is required for forceful retraction

The data thus far suggest that PilU is sufficient but not necessary to mediate pilus retraction and

natural transformation. However, PilU is present in many type IVa pilus systems, so the

question remains: Why is PilU so widely conserved if it is not necessary for pilus retraction?

Recent work in *Pseudomonas aeruginosa* indicates that PilU may be required for forceful

retraction [30]. Our data thus far indicate that pilus retraction in the *pilU* mutant is largely

indistinguishable from the parent (**Fig 1B**), however, these measurements are based on free pili

that are not retracting under load. We hypothesized that PilU might be required for forceful

retraction, such as when pili are attached to a surface. To test this, we assessed pilus retraction

using a micropillar assay in which binding of pili to micropillars and subsequent retraction results

in measurable bending and displacement of the micropillars. These data can then be used to

calculate the rates and force of pilus retraction [31]. In this assay, the Δ*pilU* mutant exhibited a

significant reduction in retraction force (~2-fold) and speed (~3-fold) compared to the parent

strain, despite the fact that this strain still has functional PilT-dependent retraction (**Fig 1B,E-F**).

This reduction in the force of retraction in the Δ*pilU* mutant is comparable to the reduction in

force of retraction observed in the Δ*pilT* mutant (**Fig 1E**). Thus, these data indicate that PilU is

necessary for generating maximal retraction forces in the *V. cholerae* competence pilus.

The ATPase activities of PilT and PilU define the rate of type IV competence pilus
retraction

Just like PilT, PilU is a AAA family ATPase with a conserved lysine in its Walker A motif.

Because we observed that PilU supports pilus retraction in the *pilT*^{K136A} mutant (**Fig 1B**), we

hypothesized that the ATPase activity of PilU may be important for supporting pilus retraction.

To test the relative roles of each of these ATPases in pilus retraction, we mutated the conserved

lysine in the Walker A motifs of PilT and/or PilU. The $pilU^{K134A}$ strain retracts ~1.5-fold slower

than the parent strain, similar to the $\Delta pilU$ mutant (**Fig 2A**). There may be a limited effect of the

$pilU^{K134A}$ mutation on pilus retraction because this strain still has a functional PilT ATPase.

However, we hypothesized that PilU ATPase activity may be critical for retraction in a

background where PilT ATPase activity is also inactivated. Consistent with this, we found that a

$pilT^{K136A}$ $pilU^{K134A}$ strain is indistinguishable from a $\Delta pilT$ strain for pilus retraction, piliation,

aggregation, and transformation (**Fig 2A-C**). Together, these data suggest that PilU ATPase

activity promotes type IV competence pilus retraction.

The data above suggest that the ATPase activities of PilT and PilU are critical for pilus

retraction. These ATPases are predicted to be motor proteins that facilitate the depolymerization

of pilus fibers. Thus, if we alter the ATPase rates of these motors, we hypothesized that this

should alter the rates of pilus retraction. To test this hypothesis, we mutated a leucine in the

Walker B domain of PilT and PilU to a cysteine, which was previously shown to reduce the rate

of ATP hydrolysis of *Neisseria gonorrhoeae* PilT ~2-fold [26]. When these Walker B mutations

were tested in isolation they did not alter the rate of pilus retraction (Comparing Parent strain to

$pilT^{L201C}$ and $pilU^{L199C}$) (**Fig 2A**). This may be due to the fact that these cells still have one fully

functional retraction ATPase (i.e. $pilT^{L201C}$ still has a fully functional PilU and $pilU^{L199C}$ still has a

fully functional PilT). We hypothesized that these Walker B "slow" ATPase mutations, however,

may have an impact on pilus retraction if they represented the sole functional retraction ATPase

in the cell. To that end, we generated both a $pilT^{L201C}$ $\Delta pilU$ strain and a $pilT^{K136A}$ $pilU^{L199C}$ strain.

In both cases, these strains exhibited reduced retraction rates relative to the appropriate parent

strain: a ~1.3-fold reduction for PilT^{L201C} (comparing $pilT^{L201C}\Delta pilU$ to $\Delta pilU$) and an ~1.5-fold

reduction for PilU^{L199C} (comparing $pilT^{K136A}pilU^{L199C}$ to $pilT^{K136A}$) (**Fig 2A**). This provides genetic

evidence that the ATPase activities of PilT and PilU are linked to the retraction rate of the type

IV competence pilus. Furthermore, it provides additional supporting evidence that PilU promotes

pilus retraction through its ATPase activity.

PilT and PilU directly interact with one another

All of the data presented above indicate that PilT and PilU can both support pilus retraction.

Therefore, we hypothesized that PilT and PilU may interact to form a complex that supports

pilus retraction. To address this, we utilized a Bacterial Adenylate Cyclase Two-Hybrid (BACTH)

assay [32, 33]. As expected for predicted hexameric proteins like PilT and PilU, the BACTH

analysis showed PilT-PilT and PilU-PilU self-interactions (**Fig 3** and **S2 Fig**). Additionally, the

BACTH assay showed that PilT and PilU interact with one another and that the Walker A

mutations of either protein (PilT^{K136A} and PilU^{K134A}) did not disrupt this interaction (**Fig 3** and **S2**

Fig). Thus, our BACTH results suggest that PilT and PilU may form a complex to mediate pilus

retraction. These results do not, however, indicate whether PilU and PilT form heteromeric

complexes or homomeric complexes to facilitate pilus retraction. We test this further below.

PilT and PilU form independent homohexameric rings in vitro and in vivo

PilT in other systems has been shown to form a hexameric complex [34], and by homology we

hypothesized that PilU also forms a hexamer. Based on transformation frequency assays, N-

terminally tagged ^{6X}His-PilT and ^{6X}His-PilU were fully functional (**S3A Fig**). Therefore, we

purified ^{6X}His-PilT and ^{6X}His-PilU to characterize these proteins *in vitro*. Negative stain

transmission electron microscopy revealed that both ^{6X}His-PilT and ^{6X}His-PilU formed hexameric

ring complexes (**Fig 4A**). These results indicate that these proteins are capable of forming

independent homohexameric complexes *in vitro*. PilT and PilU have both sequence (~65%

similarity) and structural similarity to one another (**Fig 4A**), thus it is also possible that these

proteins can be incorporated into one ring and form heterohexameric complexes that facilitate

pilus retraction *in vivo*.

In order to test if PilT and PilU form heterohexamers or independent homohexamers to facilitate

pilus retraction *in vivo*, we hypothesized that overexpression of ATPase-inactive mutants (i.e.

PilT^{K136A} and PilU^{K134A}) in backgrounds that have native PilT and PilU would allow us to

distinguish between these two models. For example, if these proteins form heteromeric

complexes, then we would predict that overexpression of either ATPase-inactive protein

(PilT^{K136A} or PilU^{K134A}) should result in complexes that are mostly composed of ATPase-inactive

proteins and result in inhibition of pilus retraction (yielding a hyperpiliated phenotype) and

reduce rates of natural transformation (**Fig 4B**). If these proteins solely form homomeric

complexes, however, then we would predict that overexpression of only one ATPase-inactive

protein (PilT^{K136A} or PilU^{K134A}) should only inhibit the activity of one motor complex (PilT or PilU,

respectively) and should not have a dramatic effect on pilus retraction or natural transformation

because the other retraction motor would still be intact (**Fig 4B**).

In order for us to successfully test this question, we needed to: (1) be able to ectopically

overexpress ATPase deficient alleles of PilT or PilU, (2) demonstrate that ATPase deficient

alleles can interact with their ATPase sufficient counterparts and form proper hexamers, and (3)

show that ectopic expression of ATPase deficient alleles of PilT or PilU can inhibit the activity of

their respective natively expressed ATPase. We test each of these in turn below. First, we

confirmed that we could ectopically overexpress PilT and PilU. Using functional 3xFLAG-tagged

versions of PilT and PilU (**S3A-B Fig**), we show that our chromosomal P_{tac} regulated constructs

resulted in dramatic overexpression of these proteins (**S3C Fig**). Importantly, overexpression of

WT PilT and PilU did not affect pilus activity or natural transformation (**S1 and S3B,D Figs**).

Next, we purified ^{6X}His-PilT^{K136A} and ^{6X}His-PilU^{K134A} and demonstrated that they formed ring

structures *in vitro* (**S3E Fig**), indicating that these proteins should form complexes with ATPase

sufficient alleles. This is also supported by our BACTH data (**Fig 3**). Finally, we wanted to

confirm that overexpressing the ATPase-inactive PilT^{K136A} or PilU^{K134A} could effectively impede

the function of the natively expressed ATPase-active PilT or PilU. To test this, we genetically

isolated either PilT or PilU as the sole functional retraction ATPase (i.e. PilT activity is isolated in

a Δ*pilU* background, while PilU activity is isolated in a *pilT^{K136A}* background) and then

overexpressed the cognate ATPase-inactive allele (P$_{tac}$-*pilT^{K136A}* or P$_{tac}$-*pilU^{K134A}*, respectively).

We assessed piliation, aggregation, and natural transformation as indicators of optimal pilus

retraction. We found that, as expected, overexpression of ATPase-inactive alleles was sufficient

to disrupt the activity of the natively expressed ATPase (**Fig 4C-D** - compare Δ*pilU* to Δ*pilU* P$_{tac}$-

pilT^{K136A} and *pilT^{K136A}* to *pilT^{K136A}* P$_{tac}$-*pilU^{K134A}*). Also, as expected, overexpression of both

ATPase-deficient alleles together (P$_{tac}$-*pilT^{K136A}* and P$_{tac}$-*pilU^{K134A}*) in a background that has

natively expressed PilT and PilU, results in hyperpiliation, aggregation and reduced natural

transformation (**Fig 4C-D**), which further demonstrates that overexpression of these ATPase-

deficient proteins inhibits the activity of their natively expressed ATPase–sufficient versions.

The experiments above indicated that it would be feasible to test whether PilT and PilU formed

homomeric or heteromeric complexes *in vivo* as described in our model (**Fig 4B**). When we

overexpressed only one ATPase-deficient protein (P$_{tac}$-*pilT^{K136A}* or P$_{tac}$-*pilU^{K134A}*) in a background

that has natively expressed PilT and PilU, we found that piliation and transformation were

indistinguishable from the parent (**Fig. 4C-D**). Thus, these data are consistent with PilT and PilU

forming independent homohexamers to facilitate pilus retraction.

Using our purified PilT and PilU, we also attempted to directly measure the ATPase activity of

these proteins. Unfortunately, the ATPase activity of these purified proteins was

indistinguishable from contaminating ATPases and/or below our detection limit (~1 nmol

pi/min/mg protein). Poor ATPase activity in purified retraction ATPases is not unique to our

system [12, 29, 35]. This may be due to the fact that the ATPase activities of these proteins are

stimulated *in vivo* via interaction with the proteins that compose the pilus machine, as previously

suggested [12]. Regardless, our previous *in vivo* data (**Fig 1A-D**) are consistent with the

pilT^{K136A} mutant being ATPase deficient.

The presence of PilT is not required for PilU to interact with the pilus platform

PilC.

Our data thus far indicate that PilU requires the presence of PilT (whether it is a functional

ATPase or not) to support optimal pilus retraction. The extension and retraction ATPases are

hypothesized to interact with the platform protein PilC to mechanically facilitate polymerization

and depolymerization of the pilus fiber [7, 36, 37]. Because PilU is dependent on the presence

of PilT to mediate optimal retraction, we hypothesized that PilT can interact with PilC, but that

PilU cannot. To test this, we utilized the BACTH assay to test the interactions of PilT and PilU

with PilC. This assay showed that PilT and PilU could both interact with PilC, and that the

Walker A mutations of these ATPases (PilT^{K136A} and PilU^{K134A}) did not affect these interactions

(**Fig 3**). The interaction between PilC and PilT supports the canonical model that retraction

ATPases interact with PilC to facilitate depolymerization of the pilus fiber. Interestingly, PilC and

PilU interact independently of PilT. Thus, the simple model of PilU requiring PilT for interaction

with the platform PilC appears to be incorrect.

The function of PilU as a PilT-dependent retraction ATPase is conserved

As mentioned earlier, many type IVa pilus systems encode PilU in addition to the canonical

retraction ATPase PilT. Therefore, we sought to assess whether PilU could facilitate PilT-

dependent retraction in other type IV pilus systems. To test this, we studied the type IVa

competence pili of *Acinetobacter baylyi*.

The competence pili of *A. baylyi* likely facilitate DNA uptake in a retraction-dependent manner,

similar to the competence pili of *V. cholerae*. Thus, we employed natural transformation assays

to dissect the role that PilT and PilU play in *A. baylyi* competence pilus retraction. These assays

showed that the PilT and PilU of *A. baylyi* behave largely like the PilT and PilU of *V. cholerae*.

The *pilU* mutant of *A. baylyi* displays a significant (~13-fold) reduction in natural transformation

(**Fig 5**), unlike what is observed in *V. cholerae* (**Fig 1A**). This may be because DNA uptake in *A.*

baylyi requires more forceful retraction or because PilU plays a more important role in pilus

retraction in this species. Most importantly, however, we found that PilU facilitates natural

transformation (and therefore pilus retraction) in backgrounds where the ATPase activity of PilT

is inactivated, but not in backgrounds where *pilT* is absent (**Fig 5** – compare *pilT*^{K136A} to *pilT*^{K136A}

Δ*pilU* and Δ*pilT*). Also, a strain where the ATPase activity of both PilT and PilU were inactivated

(*pilT*^{K136A} *pilU*^{K137A}) resulted in the loss of transformation equivalent to a retraction deficient

background (e.g. Δ*pilTU*) or a strain lacking competence pili (Δ*comP*). Importantly, strains

lacking *pilT* (i.e. Δ*pilT* and Δ*pilTU*) could only be complemented via ectopic expression of *pilT*

but not *pilU*, while strains that contained an ATPase defective allele of *pilT* (i.e. *pilT*^{K136A} Δ*pilU* or

pilT^{K136A} *pilU*^{K137A}) could be complemented by ectopic expression of *pilT* or *pilU* (**S4 Fig**) These

data are consistent with what was observed for *V. cholerae* competence pili (**Fig 1A and S1**

Fig) and suggest that PilU ATPase activity facilitates pilus retraction in a PilT-dependent

manner in diverse type IVa pilus systems.

Discussion

This study sheds light on the function of PilU as a retraction ATPase of type IVa competence pili

in *V. cholerae*. Specifically, our data indicate that PilU can mediate pilus retraction in a PilT-

dependent manner likely through its ATPase activity. Our results also show that PilT and PilU

form independent homohexamers that likely interact to form a complex that supports pilus

retraction. Furthermore, we demonstrate that the role of PilU in supporting pilus retraction is

conserved in at least one other type IVa pilus system.

Our results indicate that PilU can only facilitate pilus retraction in the presence of PilT,

regardless of whether that PilT is a functional ATPase. These results are also supported by a

recent preprint from the Blokesch lab [22]. We initially tested the hypothesis that this was due to

a requirement for PilT to bridge PilU with the platform protein PilC. Our BACTH results,

however, indicated that PilU may directly interact with PilC in the absence of PilT. While they

may interact, it is possible that PilU does not bind and engage the PilC platform in a

conformation that would facilitate retraction. Alternatively, it is possible that PilU does bind PilC

in a conformation that can facilitate retraction but additional interactions with PilT are required to

stimulate PilU ATPase activity. Another possibility is that PilT interacts with other components of

the pilus machine that are required for stimulating retraction, and that PilU lacks these

interactions. Future efforts focused on the interaction network of the retraction ATPases with the

pilus machine should shed light on this question.

Previous reports have shown that PilU is necessary for twitching motility in *P. aeruginosa* [13],

an activity that likely requires forceful retraction [30]. Consistent with this, our results indicate

that PilU is necessary for maximal retraction force for the competence pilus in *V. cholerae*.

Interestingly, because the Δ*pilU* mutant transforms similarly to the parent strain (**Fig 1B** and

2A), the forceful retraction mediated by PilU does not seem to be necessary for the function of

the *V. cholerae* competence pilus in DNA uptake under the conditions tested. However, forceful

PilU-dependent retraction may be necessary for DNA uptake when tDNA is absorbed onto

surfaces and/or in complex environments like biofilms.

Beyond altering the force of retraction, many mutants in PilT and PilU also reduced the speed of

pilus retraction (i.e. Δ*pilU* = ~1.3-fold, *pilT*^{K136A} = ~4.3-fold, *pilT*^{K136A} *pilU*^{L199C} = ~6.5-fold, *pilU*^{K134A}

= ~1.5-fold, *pilT*^{L201C} Δ*pilU* = ~1.7-fold). This revealed that altering the rates of these motor

ATPases results in a corresponding reduction in the rate of pilus retraction. Additionally, despite

this reduction in the speed of pilus retraction, these mutants still transformed at levels that were

comparable to the parent strain. Thus, these results suggest that the absolute speed of

competence pilus retraction is not critical for DNA uptake during natural transformation. Further

experiments will focus on defining the parameters that are essential for DNA uptake.

Mutants that completely inactivated the retraction machinery (e.g. Δ*pilT*, Δ*pilT* Δ*pilU*, *pilT*^{K136A}

Δ*pilU*, *pilT*^{K136A} *pilU*^{K134A}) resulted in a hyperpiliated phenotype, lower transformation frequency

and a marked reduction in the frequency of pilus retraction (as observed qualitatively by

epifluorescence time-lapse microscopy). These strains, however, did display rare pilus

retraction, which occurred at rates that were markedly slower than the parent (>40-fold), which

is independent of PilT and PilU. Recent work in *Neisseria* indicates that PilTU-independent

retraction is not unique to the *V. cholerae* competence pilus [38]. The mechanism underlying

PilTU-independent retraction is still unclear [5] and will be the focus of future work.

Materials and Methods

Bacterial strains and culture conditions

Vibrio cholerae and *A. baylyi* strains were routinely grown in LB Miller broth and on LB Miller

agar supplemented with erythromycin (10 µg/mL), kanamycin (50 µg/mL), and/or carbenicillin

(50 µg/mL) when appropriate.

Construction of mutant strains

All *V. cholerae* strains used throughout this study are derivatives of the El Tor isolate E7946. All

A. baylyi strains used were derivatives of strain ADP1 [39]. All mutant strains were constructed

via MuGENT and natural transformation using transforming DNA products made by splicing-by-

overlap extension (SOE) PCR exactly as previously described [40-43]. For a detailed list of all

mutant strains used throughout this study see **Table S1**. For a detailed list of all primers used to

construct mutant strains see **Table S2**.

Natural transformation assays

In order to induce competence in *V. cholerae* strains, the master competence regulator TfoX

was overexpressed using an IPTG-inducible or constitutive P_{tac} promoter and the cells were

genetically locked in a state of high cell density via deletion of *luxO* [41, 44-48]. Chitin-

independent transformation assays were performed exactly as previously described [42]. Briefly,

strains were grown overnight rolling at $30\,^{o}C$ with 100 µM IPTG then $\sim10^8$ colony forming units

(CFU) were subcultured into 3mL of LB + 100 µM IPTG + 20 mM $MgCl_2$ + 10 mM $CaCl_2$ and

grown to late log. Next, $\sim10^8$ CFU of this culture was diluted into instant ocean medium (7 g/L;

Aquarium Systems) supplemented with 100 µM IPTG and 500ng of transforming DNA was

added to each reaction and incubated statically at $30\,^{o}C$ overnight. The tDNA targeted VC1807,

a frame-shifted transposase, for deletion as previously described [40]. Negative control

reactions where no tDNA was added were also performed for each strain. After incubation with

tDNA, reactions were outgrown by adding 1mL of LB to each reaction and shaking (250 rpm) at

37 °C for ~2 hours. Reactions were then plated for quantitative culture onto medium selecting for

transformants (LB + 10 µg/mL Erythromycin) or onto plain LB for total viable counts. The

transformation frequency is defined as the number of transformants divided by the total viable

counts. For reactions where no transformants were obtained, a limit of detection was calculated

and plotted.

In order to sensitize natural transformation assays to differences in the efficiency of DNA

uptake, reactions were incubated with transforming DNA for shorter periods of time. These

assays were prepared exactly as described above, except that 10 units of DNase I (NEB) was

added to reactions 7 minutes after incubation with 500ng of transforming DNA. Reactions were

then incubated statically at 30°C overnight, outgrown, and plated exactly as described above.

For *A. baylyi*, transformations were performed exactly as previously described [49]. Briefly,

strains were grown overnight in LB medium. Then, ~10^8 cells were diluted into fresh LB

medium, and tDNA was added (~100 ng). Reactions were incubated at 30°C with agitation for 5

h and then plated for quantitative culture as described above to determine the transformation

frequency.

Pilin labelling, imaging and quantification

In order to label pili for observation with epifluorescence microscopy, strains were grown to the

late-log phase exactly as described above for natural transformation assays. Then, ~10^8 CFU

were spun down at 18,000 x g for 1 min and resuspended in instant ocean medium. Cells were

then incubated with 25 µg/mL AF488-mal for 15 minutes statically at room temperature in the

dark. Cells were washed twice using 100µL of instant ocean medium and resuspended to a final

concentration of ~10^7 CFU. Next, 2µL of these labeled cells were placed under an 0.2% gelzan

pad (made in instant ocean medium) on a coverslip and imaged on an inverted Nikon Ti-2

microscope with a Plan Apo ×60 objective, a green fluorescent protein filter cube, a Hamamatsu

ORCAFlash 4.0 camera and Nikon NIS Elements imaging software. In order to measure

retraction events, labelled cells were imaged by time-lapse microscopy in which a phase-

contrast (to image cell bodies) and fluorescent (to image labeled pili) image were taken every

second for 1-2 minutes or every 10 seconds for 10-15 minutes. Rates were calculated manually

using measurement tools in the NIS Elements analysis software. Only pili that were longer than

0.3 µm and completed retracting within the time-lapse window were analyzed.

In order to calculate the number of pili per cell, static images of cell bodies and labeled pili were

captured using phase-contrast and epifluorescence microscopy, respectively. Images from 3

independent biological replicates were sectioned into areas containing ~100-200 cells and the

total number of cells and the number of pili displayed on each cell was manually determined.

Representative images of the piliation state of strains were gathered and the lookup tables for

each phase or fluorescent image were adjusted to the same range.

To assess aggregation and pelleting of cultures, strains were grown to late log under the

conditions described above. Cultures were then allowed to sit statically at room temperature for

30-60 mins. Images of cultures were then taken against a white background.

Protein expression and purification

His tagged (6xHis) versions of PilT, PilT^{K136A}, PilU and PilU^{K134A} were cloned into a pHis

expression vector and verified by DNA sequencing (Eurofins). Vectors were transformed into $E.$

$coli$ BL21 DE3 for expression and purification.

For purification of His-PilU and His-PilU^{K134A}, cells were grown in a 1L flask at 37^{o}C with aeration

to an OD$_{600}$ = 0.6. The culture was then induced by adding IPTG (1 mM) and grown to a final

OD_{600} = 4.0 at 30°C with aeration. Cells were harvested by centrifugation at 8,000 x g for 15 min

at room temperature and then resuspended in Buffer A [100 mM Tris pH 8.5, 300 mM KCl, 10%

glycerol, 20 mM Imidazole] and stored at -80°C. Pellets were thawed on ice and then lysozyme

(1 mg/mL) and DNase I (2 mg/L) were added before lysing via sonication with a probe tip

sonicator. The soluble fraction was clarified by centrifugation at 12,000 x g for 1 hour at 4°C.

The supernatant was loaded onto a nickel-charged HisTrap HP column (1mL; GE) using an

ÄKTA FPLC at room temperature. The column was then washed with Buffer A and the protein

was eluted with a gradient of Buffer B [100 mM Tris pH 8.5, 300 mM KCl, 10% glycerol, 500 mM

imidazole] and fractions were collected on ice. Elution fractions were separated on 15% SDS-

PAGE gels and stained with Coomassie Brilliant Blue R-250 (BioRad). Peak fractions were

pooled and dialyzed (Fisher, MWCO 12 kDa) into 500 mL Buffer C [100 mM Tris pH 8.5, 300

mM KCl, 10% glycerol] overnight, on a stir plate at 4°C. Then a second round of dialysis was

performed using 500 mL Buffer C for 4-6 hours, on a stir plate at 4°C. Following dialysis, the

protein concentration was determined using a Bradford Assay and aliquots were stored at -80C.

For purification of His-PilT and His-PilT^{K136A}, cells were grown in a 1L flask at 37°C with aeration

to an OD_{600} = 0.6. The culture was then induced by adding IPTG (1 mM) and grown overnight at

22°C with aeration. Cells were harvested by centrifugation at 8,000 x g for 15min at room

temperature and then resuspended in Buffer A [25 mM Tris pH 8.5, 300 mM KCl, 10% glycerol,

1 mM EDTA, 1 mM $MgCl_2$, 5 mM β-mercaptoethanol, 20 mM Imidazole] and stored at -80°C.

Pellets were thawed on ice and lysozyme (1 mg/mL), DNase I (2 mg/L) and PMSF (1 mM) were

added before lysing via sonication with a probe tip sonicator. The soluble fraction was clarified

by centrifugation at 12,000 x g for 1 hour at 4°C. The supernatant was loaded onto a nickel-

charged HisTrap HP column (1 mL; GE) using an ÄKTA FPLC at room temperature. The

column was then washed with Buffer A and the protein was eluted with a gradient of Buffer B

[25 mM Tris pH 8.5, 300 mM KCl, 10% glycerol, 1 mM EDTA, 1 mM $MgCl_2$, 5 mM β-

mercaptoethanol, 500 mM Imidazole] and fractions were collected on ice. Elution fractions were

separated on 15% SDS-PAGE gels and stained with Coomassie Brilliant Blue R-250 (BioRad).

Peak fractions were pooled and dialyzed (Fisher, MWCO 12 kDa) into 500 mL Buffer C [25 mM

Tris pH 8.5, 50 mM KCl, 10% glycerol, 1 mM EDTA, 1 mM $MgCl_2$, 0.5mM TCEP] overnight, on a

stir plate at 4^oC. The next day, the eluate was concentrated with an Amicon conical filter

(MWCO 30 kDa) at 4,000 x g for 15 min at 4^oC and resuspended to the original volume in Buffer

C twice. The protein concentration was determined using a Bradford Assay and then aliquots of

protein were snap-frozen with liquid nitrogen and stored at -80C.

Bacterial Adenylate Cyclase Two-hybrid (BACTH) Assays

The Bacterial Adenylate Cyclase Two-Hybrid (BACTH) system was employed to study protein

interactions. Gene inserts were amplified by PCR and cloned into pUT18C (Carb[R]) and pKT25

(Kan[R]) vectors to generate N-terminal fusions to the T18 and T25 fragments of adenylate

cyclase, respectively. Miniprepped vectors (Qiagen plasmid miniprep kit) were then

cotransformed into *E. coli* BTH101 and outgrown with 500 µL LB for 1 hour at 37^oC with

shaking. Transformations were isolation streak plated on LB plates containing kanamycin 50

µg/mL and carbenicillin 100 µg/mL to select for transformants that had received both plasmids.

Strains were then grown statically at 30^oC from frozen stocks overnight in LB supplemented with

kanamycin 50 µg/mL and carbenicillin 100 µg/mL and 3.5µL of each strain was then spotted

onto LB plates containing kanamycin 50 µg/mL, carbenicillin 100µg/mL, 0.5 mM IPTG and 40

µg/mL X-gal and incubated at 30^oC for 48 hours.

BACTH Miller Assays

For BACTH Miller assays strains were prepared by growing the BTH101 strains with the

indicated pUT18C and pKT25 vectors statically overnight at 30^{o}C in LB supplemented with

kanamycin 50 μg/mL and carbenicillin 100 μg/mL. Then 5 μL of this overnight culture was

subcultured into 3mL of LB supplemented with kanamycin 50 μg/mL, carbenicillin 100μg/mL and

0.5 mM IPTG and grown rolling at 30^{o}C overnight. Cultures were then pelleted and resuspended

in 1 mL of Z buffer [60 mM Na_2HPO_4-$7H_2O$, 40 mM NaH_2PO_4-H_2O, 10 mM KCl, 2 mM $MgSO_4$-

$7H_2O$] to an OD_{600} = 0.5. 50μL of 1% SDS was added to each reaction and vortexed. 100μL of

chloroform was added to each reaction, vortexed again and then centrifuged for 1 min at 18,000

x g. The aqueous fraction was then mixed with ONPG (2.5 mg/mL) in a 96-well plate and the

OD_{420} and OD_{550} were measured every 5 minutes for 10 hours. Miller units were then calculated

at the time (in mins) when the A_{420} reached 0.3 or at 10 hours using the formula: [1000*(A_{420}-

(1.75*A_{550}))/ (mins*0.2 mL*A_{600})].

Western Blotting

Strains were grown to mid-log phase in LB with 100 μM IPTG if applicable and then

resuspended to an OD_{600} = 100 in TBS containing 1 mM PMSF. Cells were then lysed using a

FastPrep-24 (MP Bio). Lysates were centrifuged at 16,000 x g for 10 min at 4^{o}C, and the

supernatant was then mixed 1:1 with 2X SDS sample buffer [220 mM Tris pH 6.8, 25% glycerol,

1.8% SDS, 0.02% Bromophenol Blue, 5% β-mercaptoethanol] and then 2uL of each sample

was separated on a 15% SDS PAGE gel. Proteins were transferred to a PVDF membrane and

incubated with one of the following primary antibodies: α-FLAG polyclonal rabbit (Sigma) or α-

RpoA monoclonal mouse (Biolegend). Then, blots were washed and incubated with a α-rabbit

or α-mouse secondary antibodies conjugated to IRdye 800CW. Blots were then imaged using

an Odyssey classic LI-COR system.

Negative-stain electron microscopy

Negative stain specimens were prepared by applying 4 µL of protein solution (0.1 mg/mL + 1 mM AMP-PnP) onto a glow-discharged 300-mesh copper grid coated with continuous carbon film (EMS). After 30 s the protein solution was blotted with a piece of filter paper. The grid was washed with 4 µL of milli-Q water and stained with 4 µL of stain solution (1% (w/v) uranyl acetate, 0.5% (w/v) trehalose). The excess stain solution was removed by filter paper and the grid was left to air dry. Due to the availability of the instrument, we used two different transmission electron microscopes for data collection. For PilU, the data were collected on a 300-kV JEM-3200FS TEM equipped with FEG source and 8k x 8k Direct Electron DE-64 camera. A total of 32 micrograph images were taken at a nominal magnification of 60,000x, which is equivalent to 1.56 Å per pixel. For PilT, the data were collected on a 120-kV JEM-1400plus TEM equipped with LaB_6 source and a Gatan Oneview camera. A total of 59 images were taken at a nominal magnification of 60,000x and the pixel size is 1.94 Å. To avoid potential beam induced damage, minimum dose system was employed during data collection. Note that PilU data were collected on a direct electron detector; therefore, each image contains 50 frames and the frame alignment was performed using unblur program with exposure filter applied. Particle picking, defocus estimation and phase flipped particles were all done using EMAN2 software package [50]. These particles were then imported into RELION (v2.1) for 2D classification, Initial model building, 3D refinement, and 3D classification [51]. From 2D classification results, we observed clearly 6-fold symmetry; thus, the symmetry operation was applied to subsequent 3D refinement and 3D classification. The final 3D reconstructions were computed from 45204 particles to 13.1 Å for PilU and 17472 particles to 14.7 Å for PilT. The resolution estimation was calculated using gold-standard method at Fourier shell correlation of 0.143 as implemented in RELION. However, due to the nature of the negative stain, the effective resolution could be equal to or worse than 15 Å. The 3D reconstructions were rendered

525 using UCSF Chimera where the handedness of PilU was chosen to be consistent with PilT

526 [52].

Micropillar retraction assay

To ensure that all measurements recorded were due to type IV competence pili, all the

experiments were performed with *V. cholerae* strains lacking all external appendages other than

type IV competence pili as well as exopolysaccharide production (i.e. mutants lacking flagella,

MSHA pili, TCP pili and *Vibrio* polysaccharide (VPS) production). Strains were grown overnight

in LB either directly from frozen stocks or from a single colony from an LB agar plate. Of the

overnight liquid culture, 50 µl was subcultured into 3 ml LB + 100 µM IPTG + 20 mM $MgCl_2$ + 10

mM $CaCl_2$ and allowed to grow at 30°C for 4h. Then, 100 µl of this culture was centrifuged for 5

min at 20,000 x *g* and the bacteria were resuspended in instant ocean medium. At different

times, 23 µl of the resuspension were added to micropillars in an observation chamber as

previously reported [15]. Briefly, silica molds were inverted on activated coverslips with

polyacrylamide gels in between. The result is an array of flexible micropillars in a hexagonal

array of 3 µm × 3 µm.Once the bacteria were in contact with the micropillars, 10-Hz movies of

the top of the pillars were recorded. The motion of the tips of the pillars was tracked using a

cross-correlation algorithm in ImageJ. The amplitude and speeds of the pillars' motions were

then analyzed using MATLAB. Finally, we calibrated the pillars' stiffness constant using optical

tweezers as previously described [31]. The pillars used in this study have a stiffness constant of

$17\pm4 pN\mu m^{-1}$.

Statistics

548 Statistical differences were assessed by two-tailed Student's t-tests using GraphPad Prism

549 software.

Acknowledgements

552 We would like to thank CK Ellison for helpful discussions. We would like to acknowledge JD

553 Newman and CE Dann III for their advice during protein purification. We thank JC van Kessel

554 for providing BACTH strains and plasmids, and Lakshay Khosla and Vijay Deopersaud for

555 technical assistance with micropillar experiments. This work was supported by grants

556 R35GM128674 and AI118863 from the National Institutes of Health to ABD, and by grant

557 AI116566 from the National Institutes of Health to NB.

References

560 1. Burrows LL. Pseudomonas aeruginosa twitching motility: type IV pili in action. Annu Rev

561 Microbiol. 2012;66:493-520. doi: 10.1146/annurev-micro-092611-150055. PubMed PMID:

562 22746331.

563 2. Maier B, Wong GCL. How Bacteria Use Type IV Pili Machinery on Surfaces. Trends

564 Microbiol. 2015;23(12):775-88. doi: 10.1016/j.tim.2015.09.002. PubMed PMID: 26497940.

565 3. Hospenthal MK, Costa TRD, Waksman G. A comprehensive guide to pilus biogenesis in

566 Gram-negative bacteria. Nat Rev Microbiol. 2017;15(6):365-79. Epub 2017/05/13. doi:

567 10.1038/nrmicro.2017.40. PubMed PMID: 28496159.

568 4. Proft T, Baker EN. Pili in Gram-negative and Gram-positive bacteria - structure,

569 assembly and their role in disease. Cell Mol Life Sci. 2009;66(4):613-35. Epub 2008/10/28. doi:

570 10.1007/s00018-008-8477-4. PubMed PMID: 18953686.

5.   Craig L, Forest KT, Maier B. Type IV pili: dynamics, biophysics and functional

consequences. Nat Rev Microbiol. 2019. Epub 2019/04/17. doi: 10.1038/s41579-019-0195-4.

PubMed PMID: 30988511.

6.   Craig L, Pique ME, Tainer JA. Type IV pilus structure and bacterial pathogenicity. Nat

Rev Microbiol. 2004;2(5):363-78. doi: 10.1038/nrmicro885. PubMed PMID: 15100690.

7.   McCallum M, Tammam S, Khan A, Burrows LL, Howell PL. The molecular mechanism of

the type IVa pilus motors. Nat Commun. 2017;8:15091. doi: 10.1038/ncomms15091. PubMed

PMID: 28474682; PubMed Central PMCID: PMCPMC5424180.

8.   Bischof LF, Friedrich C, Harms A, Sogaard-Andersen L, van der Does C. The Type IV

Pilus Assembly ATPase PilB of Myxococcus xanthus Interacts with the Inner Membrane

Platform Protein PilC and the Nucleotide-binding Protein PilM. J Biol Chem. 2016;291(13):6946-

57. doi: 10.1074/jbc.M115.701284. PubMed PMID: 26851283; PubMed Central PMCID:

PMCPMC4807279.

9.   Chang YW, Rettberg LA, Treuner-Lange A, Iwasa J, Sogaard-Andersen L, Jensen GJ.

Architecture of the type IVa pilus machine. Science. 2016;351(6278):aad2001. doi:

10.1126/science.aad2001. PubMed PMID: 26965631.

10.   Chiang P, Habash M, Burrows LL. Disparate subcellular localization patterns of

Pseudomonas aeruginosa Type IV pilus ATPases involved in twitching motility. J Bacteriol.

2005;187(3):829-39. doi: 10.1128/JB.187.3.829-839.2005. PubMed PMID: 15659660; PubMed

Central PMCID: PMCPMC545728.

11.   Craig L, Volkmann N, Arvai AS, Pique ME, Yeager M, Egelman EH, et al. Type IV pilus

structure by cryo-electron microscopy and crystallography: implications for pilus assembly and

functions. Mol Cell. 2006;23(5):651-62. doi: 10.1016/j.molcel.2006.07.004. PubMed PMID:

16949362.

12.   Jakovljevic V, Leonardy S, Hoppert M, Sogaard-Andersen L. PilB and PilT are ATPases

acting antagonistically in type IV pilus function in Myxococcus xanthus. J Bacteriol.

2008;190(7):2411-21. doi: 10.1128/JB.01793-07. PubMed PMID: 18223089; PubMed Central

PMCID: PMCPMC2293208.

13.   Whitchurch CB, Mattick JS. Characterization of a gene, pilU, required for twitching

motility but not phage sensitivity in Pseudomonas aeruginosa. Mol Microbiol. 1994;13(6):1079-

91. Epub 1994/09/01. PubMed PMID: 7854122.

14.   Blair KM, Turner L, Winkelman JT, Berg HC, Kearns DB. A molecular clutch disables

flagella in the Bacillus subtilis biofilm. Science. 2008;320(5883):1636-8. doi:

10.1126/science.1157877. PubMed PMID: 18566286.

15.   Ellison CK, Kan J, Dillard RS, Kysela DT, Ducret A, Berne C, et al. Obstruction of pilus

retraction stimulates bacterial surface sensing. Science. 2017;358(6362):535-8. doi:

10.1126/science.aan5706. PubMed PMID: 29074778.

16.   Ellison CK, Dalia TN, Dalia AB, Brun YV. Real-time microscopy and physical

perturbation of bacterial pili using maleimide-conjugated molecules. Nat Protoc. 2019. doi:

10.1038/s41596-019-0162-6. PubMed PMID: 31028374.

17.   Ellison CK, Dalia TN, Vidal Ceballos A, Wang JC, Biais N, Brun YV, et al. Retraction of

DNA-bound type IV competence pili initiates DNA uptake during natural transformation in Vibrio

cholerae. Nat Microbiol. 2018;3(7):773-80. doi: 10.1038/s41564-018-0174-y. PubMed PMID:

29891864.

18.   Seitz P, Blokesch M. DNA-uptake machinery of naturally competent Vibrio cholerae.

Proc Natl Acad Sci U S A. 2013;110(44):17987-92. Epub 2013/10/16. doi: 1315647110 [pii]

10.1073/pnas.1315647110. PubMed PMID: 24127573.

19.   Bieber D, Ramer SW, Wu CY, Murray WJ, Tobe T, Fernandez R, et al. Type IV pili,

transient bacterial aggregates, and virulence of enteropathogenic Escherichia coli. Science.

1998;280(5372):2114-8. PubMed PMID: 9641917.

20.   Chamot-Rooke J, Mikaty G, Malosse C, Soyer M, Dumont A, Gault J, et al.

Posttranslational modification of pili upon cell contact triggers N. meningitidis dissemination.

Science. 2011;331(6018):778-82. doi: 10.1126/science.1200729. PubMed PMID: 21311024.

21.   Helaine S, Carbonnelle E, Prouvensier L, Beretti JL, Nassif X, Pelicic V. PilX, a pilus-

associated protein essential for bacterial aggregation, is a key to pilus-facilitated attachment of

Neisseria meningitidis to human cells. Mol Microbiol. 2005;55(1):65-77. doi: 10.1111/j.1365-

2958.2004.04372.x. PubMed PMID: 15612917.

22.   Adams DW, Stutzmann S, Stoudman C, Blokesch M. DNA-uptake pilus of Vibrio

cholerae capable of kin-discriminated auto-aggregation. bioRxiv. 2018:doi.org/10.1101/354878.

Epub June 26, 2018.

23.   Dietrich M, Mollenkopf H, So M, Friedrich A. Pilin regulation in the pilT mutant of

Neisseria gonorrhoeae strain MS11. FEMS Microbiol Lett. 2009;296(2):248-56. doi:

10.1111/j.1574-6968.2009.01647.x. PubMed PMID: 19486161; PubMed Central PMCID:

PMCPMC4428587.

24.   Merz AJ, So M, Sheetz MP. Pilus retraction powers bacterial twitching motility. Nature.

2000;407(6800):98-102. doi: 10.1038/35024105. PubMed PMID: 10993081.

25.   Pujol C, Eugene E, Marceau M, Nassif X. The meningococcal PilT protein is required for

induction of intimate attachment to epithelial cells following pilus-mediated adhesion. Proc Natl

Acad Sci U S A. 1999;96(7):4017-22. PubMed PMID: 10097155; PubMed Central PMCID:

PMCPMC22412.

26.   Hockenberry AM, Hutchens DM, Agellon A, So M. Attenuation of the Type IV Pilus

Retraction Motor Influences Neisseria gonorrhoeae Social and Infection Behavior. MBio.

2016;7(6). doi: 10.1128/mBio.01994-16. PubMed PMID: 27923924; PubMed Central PMCID:

PMCPMC5142622.

27.   Wolfgang M, Lauer P, Park HS, Brossay L, Hebert J, Koomey M. PilT mutations lead to

simultaneous defects in competence for natural transformation and twitching motility in piliated

Neisseria gonorrhoeae. Mol Microbiol. 1998;29(1):321-30. PubMed PMID: 9701824.

28.   Leighton TL, Dayalani N, Sampaleanu LM, Howell PL, Burrows LL. Novel Role for PilNO

in Type IV Pilus Retraction Revealed by Alignment Subcomplex Mutations. J Bacteriol.

2015;197(13):2229-38. doi: 10.1128/JB.00220-15. PubMed PMID: 25917913; PubMed Central

PMCID: PMCPMC4455263.

29.   Aukema KG, Kron EM, Herdendorf TJ, Forest KT. Functional dissection of a conserved

motif within the pilus retraction protein PilT. J Bacteriol. 2005;187(2):611-8. doi:

10.1128/JB.187.2.611-618.2005. PubMed PMID: 15629932; PubMed Central PMCID:

PMCPMC543540.

30.   Tala L, Fineberg A, Kukura P, Persat A. Pseudomonas aeruginosa orchestrates

twitching motility by sequential control of type IV pili movements. Nat Microbiol. 2019. Epub

2019/02/26. doi: 10.1038/s41564-019-0378-9. PubMed PMID: 30804544.

31.   Biais N, Higashi D, So M, Ladoux B. Techniques to measure pilus retraction forces.

Methods Mol Biol. 2012;799:197-216. doi: 10.1007/978-1-61779-346-2_13. PubMed PMID:

21993648; PubMed Central PMCID: PMCPMC5160128.

32.   Georgiadou M, Castagnini M, Karimova G, Ladant D, Pelicic V. Large-scale study of the

interactions between proteins involved in type IV pilus biology in Neisseria meningitidis:

characterization of a subcomplex involved in pilus assembly. Mol Microbiol. 2012;84(5):857-73.

doi: 10.1111/j.1365-2958.2012.08062.x. PubMed PMID: 22486968.

33.   Karimova G, Pidoux J, Ullmann A, Ladant D. A bacterial two-hybrid system based on a

reconstituted signal transduction pathway. Proc Natl Acad Sci U S A. 1998;95(10):5752-6. Epub

1998/05/20. PubMed PMID: 9576956; PubMed Central PMCID: PMCPMC20451.

34.   Forest KT, Satyshur KA, Worzalla GA, Hansen JK, Herdendorf TJ. The pilus-retraction

protein PilT: ultrastructure of the biological assembly. Acta Crystallogr D Biol Crystallogr.

2004;60(Pt 5):978-82. Epub 2004/04/23. doi: 10.1107/S0907444904006055. PubMed PMID:

15103158.

35.   Jones CJ, Utada A, Davis KR, Thongsomboon W, Zamorano Sanchez D, Banakar V, et

al. C-di-GMP Regulates Motile to Sessile Transition by Modulating MshA Pili Biogenesis and

Near-Surface Motility Behavior in Vibrio cholerae. PLoS Pathog. 2015;11(10):e1005068. doi:

10.1371/journal.ppat.1005068. PubMed PMID: 26505896; PubMed Central PMCID:

PMCPMC4624765.

36.   Takhar HK, Kemp K, Kim M, Howell PL, Burrows LL. The platform protein is essential for

type IV pilus biogenesis. J Biol Chem. 2013;288(14):9721-8. doi: 10.1074/jbc.M113.453506.

PubMed PMID: 23413032; PubMed Central PMCID: PMCPMC3617274.

37.   McCallum M, Tammam S, Little DJ, Robinson H, Koo J, Shah M, et al. PilN Binding

Modulates the Structure and Binding Partners of the Pseudomonas aeruginosa Type IVa Pilus

Protein PilM. J Biol Chem. 2016;291(21):11003-15. doi: 10.1074/jbc.M116.718353. PubMed

PMID: 27022027; PubMed Central PMCID: PMCPMC4900251.

38.   Zollner R, Cronenberg T, Maier B. Motor properties of PilT-independent type 4 pilus

retraction in gonococci. J Bacteriol. 2019. doi: 10.1128/JB.00778-18. PubMed PMID: 30692169.

39.   Juni E, Janik A. Transformation of Acinetobacter calco-aceticus (Bacterium anitratum). J

Bacteriol. 1969;98(1):281-8. PubMed PMID: 5781579; PubMed Central PMCID:

PMCPMC249934.

40.   Dalia AB. Natural Cotransformation and Multiplex Genome Editing by Natural

Transformation (MuGENT) of Vibrio cholerae. Methods Mol Biol. 2018;1839:53-64. doi:

10.1007/978-1-4939-8685-9_6. PubMed PMID: 30047054.

41.   Dalia AB, McDonough E, Camilli A. Multiplex genome editing by natural transformation.

Proc Natl Acad Sci U S A. 2014;111(24):8937-42. doi: 10.1073/pnas.1406478111. PubMed

PMID: 24889608; PubMed Central PMCID: PMC4066482.

42.   Dalia TN, Yoon SH, Galli E, Barre FX, Waters CM, Dalia AB. Enhancing multiplex

genome editing by natural transformation (MuGENT) via inactivation of ssDNA exonucleases.

Nucleic Acids Res. 2017;45(12):7527-37. doi: 10.1093/nar/gkx496. PubMed PMID: 28575400;

PubMed Central PMCID: PMCPMC5499599.

43.   Dalia AB, Lazinski DW, Camilli A. Characterization of undermethylated sites in Vibrio

cholerae. J Bacteriol. 2013;195(10):2389-99. doi: 10.1128/JB.02112-12. PubMed PMID:

23504020; PubMed Central PMCID: PMC3650525.

44.   Zhu J, Miller MB, Vance RE, Dziejman M, Bassler BL, Mekalanos JJ. Quorum-sensing

regulators control virulence gene expression in Vibrio cholerae. Proc Natl Acad Sci U S A.

2002;99(5):3129-34. Epub 2002/02/21. doi: 10.1073/pnas.052694299

052694299 [pii]. PubMed PMID: 11854465; PubMed Central PMCID: PMC122484.

45.   Lo Scrudato M, Blokesch M. The regulatory network of natural competence and

transformation of Vibrio cholerae. PLoS Genet. 2012;8(6):e1002778. Epub 2012/06/28. doi:

10.1371/journal.pgen.1002778

PGENETICS-D-12-00108 [pii]. PubMed PMID: 22737089; PubMed Central PMCID:

PMC3380833.

46.   Lo Scrudato M, Blokesch M. A transcriptional regulator linking quorum sensing and chitin

induction to render Vibrio cholerae naturally transformable. Nucleic Acids Res. 2013;41(6):3644-

58. Epub 2013/02/06. doi: gkt041 [pii]

10.1093/nar/gkt041. PubMed PMID: 23382174; PubMed Central PMCID: PMC3616704.

47.   Meibom KL, Blokesch M, Dolganov NA, Wu CY, Schoolnik GK. Chitin induces natural

competence in Vibrio cholerae. Science. 2005;310(5755):1824-7. Epub 2005/12/17. doi:

310/5755/1824 [pii]

10.1126/science.1120096. PubMed PMID: 16357262.

48.   Dalia AB, Lazinski DW, Camilli A. Identification of a membrane-bound transcriptional

regulator that links chitin and natural competence in Vibrio cholerae. MBio. 2014;5(1):e01028-

13. doi: 10.1128/mBio.01028-13. PubMed PMID: 24473132; PubMed Central PMCID:

PMC3903286.

49.   Nero TM, Dalia TN, Wang JC, Kysela DT, Bochman ML, Dalia AB. ComM is a

hexameric helicase that promotes branch migration during natural transformation in diverse

Gram-negative species. Nucleic Acids Res. 2018;46(12):6099-111. doi: 10.1093/nar/gky343.

PubMed PMID: 29722872.

50.   Ludtke SJ. Single-Particle Refinement and Variability Analysis in EMAN2.1. Methods

Enzymol. 2016;579:159-89. Epub 2016/08/31. doi: 10.1016/bs.mie.2016.05.001. PubMed

PMID: 27572727; PubMed Central PMCID: PMCPMC5101015.

51.   Scheres SH. Semi-automated selection of cryo-EM particles in RELION-1.3. J Struct

Biol. 2015;189(2):114-22. Epub 2014/12/09. doi: 10.1016/j.jsb.2014.11.010. PubMed PMID:

25486611; PubMed Central PMCID: PMCPMC4318617.

52.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al.

UCSF Chimera--a visualization system for exploratory research and analysis. J Comput Chem.

2004;25(13):1605-12. Epub 2004/07/21. doi: 10.1002/jcc.20084. PubMed PMID: 15264254.

Fig 1. PilU mediates type IVa competence pilus retraction in a PilT-dependent manner in *V. cholerae*. (A) Natural transformation assays of the indicated strains. Reactions were incubated with 500 ng of transforming DNA overnight. Parent, $n = 11$, all other strains, $n = 4$. **(B)** Retraction rates of the indicated strains measured by epifluorescence time-lapse microscopy of AF488-mal labeled cells. Data are from three independent biological replicates; Parent, $n = 68$. $\Delta pilU$, $n = 56$. $\Delta pilT$, $n = 9$. $\Delta pilT\Delta pilU$, $n = 18$. $pilT^{K136A}$, $n = 66$. $pilT^{K136A}\Delta pilU$, $n = 9$. $pilT^{E204A}$, $n = 5$. $pilT^{E204A/K136A}$, $n = 50$. $pilT^{E204A}\Delta pilU$, $n = 17$. $pilT^{E204A/K136A}\Delta pilU$, $n = 6$. **(C)** The percentage of cells with 0, <3 or ≥3 surface pili was determined from a static image. Data are from three independent biological replicates; Parent, $n = 453$. $\Delta pilU$, $n = 413$. $\Delta pilT$, $n = 381$. $\Delta pilT\Delta pilU$, $n = 425$. $pilT^{K136A}$, $n = 417$. $pilT^{K136A}\Delta pilU$, $n = 413$. $pilT^{E204A}$, $n = 579$. $pilT^{E204A/K136A}$, $n = 713$. $pilT^{E204A}\Delta pilU$, $n = 806$. $pilT^{E204A/K136A}\Delta pilU$, $n = 705$. **(D)** Representative images of surface piliation (top) and aggregation phenotypes (bottom) for the indicated strains. Scale bar, 1 µm. **(E)** Retraction force (Parent, $n = 101$. $\Delta pilU$, $n = 90$. $\Delta pilT$, $n = 372$. $\Delta pilT\Delta pilU$, $n = 43$) and **(F)** retraction rate (Parent, $n = 95$. $\Delta pilU$, $n = 83$. $\Delta pilT$, $n = 324$. $\Delta pilT\Delta pilU$, $n = 125$) was measured via micropillar assays. All bar graphs are shown as the mean ± SD. All Box plots represent the median and the upper and lower quartile, while the whiskers demarcate the range. Asterisk(s) directly above bars denote comparisons to the parent strain. All comparisons were made by Student's two-tailed t-test. LOD, limit of detection; NS, not significant; ** = $P < 0.01$, *** = $P < 0.001$.

Fig 2. The ATPase activity of PilT and PilU mediate pilus retraction. (A) Retraction rates of the indicated strains measured by epifluorescence time-lapse microscopy of AF488-mal labeled cells. Box plots represent the median and the upper and lower quartile, while the whiskers demarcate the range. Data are from three independent biological replicates; $pilU^{K134A}$, $n = 70$. $pilT^{K136A}\ pilU^{K134A}$, $n = 8$. $pilT^{K136A}\ pilU^{L199C}$, $n = 46$. $pilU^{L199C}$, $n = 49$. $pilT^{L201C}$, $n = 54$. $pilT^{L201C}\ \Delta pilU$, $n = 54$. $\Delta pilT\ pilU^{L199C}$, $n = 11$. Data for the retraction rates of the parent, $\Delta pilU$, and $pilT^{K136A}$ strains are identical to those shown in **Fig 1B** and plotted again here to facilitate comparisons to the mutant strains indicated. **(B)** Natural transformation assays of the indicated strains incubated with 500 ng of transforming DNA overnight. Data are shown as the mean ± SD. Parent, $n = 6$. All other strains, $n = 4$. **(C)** Representative images of surface piliation (top) and aggregation phenotypes (bottom) for the indicated strains. Scale bar, 1 μm. Asterisk(s) directly above bars denote comparisons to the parent strain. All comparisons were made by Student's two-tailed t-test. NS, not significant; ** = $P < 0.01$, *** = $P < 0.001$.

Fig 3. PilT and PilU interact with each other and the pilus platform protein PilC. Bacterial Adenylate Cyclase Two-Hybrid (BACTH) assays were performed to test pair-wise interactions between T18 and T25-tagged PilT, PilU, PilT^{K136A}, PilU^{K134A}, and PilC. Each construct was also tested for interactions with empty T18 and T25 vectors as a negative control. Leucine zipper fusions to T18 and T25 were used as the positive control (+) and empty T18 and T25 vectors served as an additional negative control (-). **(A)** Representative image of the BACTH assay plated on X-gal-containing media. **(B)** BACTH miller assays of the indicated pairwise interactions. The dotted line represents the β-galactosidase activity of the empty vector negative control. Data are shown as the mean ± SD and are from three independent experiments.

Fig 4. PilT and PilU form independent homomeric complexes *in vitro* and *in vivo* to facilitate pilus retraction. (A) Representative negative stain transmission electron micrographs of purified ^{6X}His-PilT and ^{6X}His-PilU (left; scale bar, 50 nm). 2D class averages revealing hexameric rings (middle; scale bar, 10 nm) and 3D reconstructions (right) of the ring complex imposing C6 symmetry. **(B)** Schematic of the experimental approach and expected outcomes for testing whether PilT and PilU form homomeric or heteromeric complexes to facilitate pilus retraction *in vivo*. **(C)** Natural transformation assays of indicated strains incubated with 500 ng of tDNA and after 7 minutes, DNAseI was added to prevent additional DNA uptake. The genotype of native *pilT* and *pilU* alleles are written below each bar graph and the presence or absence of a chromosomally-integrated IPTG-inducible ectopic allele of *pilT^{K136A}* or *pilU^{K134A}* is indicated by a + or -. Strains were grown with 100 μM IPTG if they contained an ectopic expression construct. Data are shown as the mean ± SD and are from four independent biological replicates. **(D)** Representative images of surface piliation (top) and aggregation phenotypes (bottom) for the indicated strains. Scale bar, 1 μm. Asterisk(s) directly above bars denote comparisons to the parent strain. All comparisons were made by Student's two-tailed t-test. NS, not significant; * = *P* < 0.05; ** = *P* < 0.01.

Fig 5. PilU supports pilus retraction in a PilT-dependent manner in *A. baylyi.* Natural transformation assays of the indicated strains incubated with 500ng of transforming DNA for 5 hours. Data are shown as the mean ± SD. Parent, *n* = 5. Δ*comP*, *n* = 6. All other strains, *n* = 4. Asterisk(s) directly above bars denote comparisons to parent strain. All comparisons were made by Student's two-tailed t-test. LOD, limit of detection; ** = *P* < 0.01; *** = *P* < 0.001.

Supporting Information for

PilT and PilU are homohexameric ATPases that coordinate to retract type IVa pili

Jennifer L. Chlebek[1], Hannah Q. Hughes[1], Aleksandra S. Ratkiewicz[2], Rasman Rayyan[2],

Joseph Che-Yen Wang[3], Brittany E. Herrin[1], Triana N. Dalia[1], Nicolas Biais[2], and Ankur B.

Dalia[1,*]

[1]Department of Biology, Indiana University, 1001 E. 3[rd] Street, Bloomington, IN 47405.

[2]Biology Department, CUNY Brooklyn College, 2900 Bedford Avenue, Brooklyn, NY 11210 and Graduate Center of CUNY, 365 5[th] Avenue, NY 10016.

[3]Electron Microscopy Center, Indiana University, Bloomington, IN 47405.

*Correspondence to: ankdalia@indiana.edu.

This PDF file includes:

     S1-S4 Figs.
     S1-S2 Tables

S1 Fig. Complementation analysis of *V. cholerae* mutant strains via ectopic expression of *pilT* or *pilU*.
Natural transformation assays of the indicated strains. All strains contain either a chromosomally integrated IPTG-inducible P_{tac}-*pilT* (white bars) or P_{tac}-*pilU* (gray bars) construct as indicated. Data show that ectopic overexpression of PilT and PilU does not affect the transformation frequency of the parent strain. Ectopic overexpression of PilT rescued the transformation of all strains that showed a substantial reduction in natural transformation in **Fig 1A** and **Fig 2B** (i.e. Δ*pilT*, Δ*pilT* Δ*pilU*, *pilT^{K136A}* Δ*pilU*, *pilT^{E204A}* Δ*pilU*, *pilT^{E204A/K136A}* Δ*pilU*, and *pilT^{K136A} pilU^{K134A}*). Ectopic overexpression of PilU rescued all strains except for strains that lacked a copy of *pilT* (i.e. Δ*pilT*, Δ*pilT* Δ*pilU*). All strains were grown with 100 µM IPTG to ectopically induce PilT or PilU. Data are shown as the mean ± SD. Parent, *n* = 8. All other strains, *n* = 4. Asterisk(s) directly above data denote comparisons to the parent strain. All comparisons were made by Student's two-tailed t-test. *** = *P* < 0.001.

S2 Fig. PilT and PilU interact with each other and the pilus platform protein PilC. BACTH Miller assays showing all data from **Fig 3B** with additional negative controls for every T25- or T18-fusion. Data are shown as the mean ± SD from three independent experiments. The dotted line represents the β-galactosidase activity of the empty vector negative control (T18-empty x T25-empty).

S3 Fig. PilT and PilU form homomeric complexes to facilitate pilus retraction. **(A)** Natural transformation assays showing that 6xHis and 3xFLAG N-terminal fusions to PilT and PilU are functional. The indicated strains were incubated with 500 ng of transforming DNA overnight. Parent, $n = 8$. All others, $n = 4$. **(B)** Natural transformation assays showing that overexpression of 3xFLAG tagged PilT/PilU/PilT^{K136A}/PilU^{K134A} behave the same as non-tagged proteins when ectopically overexpressed in the indicated backgrounds. Strains were incubated with 500 ng of transforming DNA and after 7 minutes, DNAse I was added to prevent additional DNA uptake. Parent, $n = 5$. $pilT^{K136A}$, $n = 5$. All others, $n = 4$. **(C)** Western blot of the indicated strains to detect FLAG tagged-proteins and RpoA as a loading control. Strains containing an IPTG regulated P$_{tac}$ constuct were grown in the presence of 100 μM IPTG. This blot indicates that ectopic induction of 3xFLAG tagged PilT/PilU/PilT^{K136A}/ PilU^{K134A} results in robust overexpression of proteins above native levels. Data is representative of three independent experiments. **(D)** Natural transformation assays where the indicated strains were incubated with 500 ng of transforming DNA for 7 minutes prior to the addition of DNAse I to prevent additional DNA uptake. These data indicate that ectopic overexpression of PilT or PilU can rescue strains with a substantial reduction in transformation frequency. All strains with P$_{tac}$ constructs were grown with 100 μM IPTG to overexpress PilT or PilU. All bar graphs are shown as the mean ± SD. Asterisk(s) directly above bars denote comparisons to the parent strain. All comparisons were made by Student's two-tailed t-test. LOD, limit of detection; * = $P < 0.05$, ** = $P < 0.01$. **(E)** Representative negative stain transmission electron micrographs showing that purified ^{6X}His-PilT^{K136A} and ^{6X}His-PilU^{K134A} form hexamers *in vitro*. Scale bar, 100 nm.

S4 Fig. Complementation analysis of *A. baylyi* mutant strains via ectopic expression of *pilT* or *pilU*. Natural transformation assays of the indicated *A. baylyi* strains. Strains harbored pMMB-*pilT* (white bars), pMMB-*pilU* (gray bars), or no vector (black bar). Data indicate that ectopic overexpression of PilT and PilU do not affect the transformation frequency of the parent strain. Ectopic overexpression of PilT rescued the transformation of all strains that showed a significant reduction in natural transformation in **Fig 5** (i.e. Δ*pilT*, Δ*pilT* Δ*pilU*, *pilT^{K136A}* Δ*pilU*, and *pilT^{K136A} pilU^{K137A}*). Ectopic overexpression of PilU rescued all strains except for strains that lacked a copy of pilT (i.e. Δ*pilT* and Δ*pilT* Δ*pilU*). All strains were grown with 100 µM IPTG to induce expression of *pilT* or *pilU*. Data are shown as the mean ± SD and are from three independent experiments.

S1 Table. Strains used in this study

Strain name in manuscript	Genotype and antibiotic resistances	Figure	Strain #
Parent	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C	Fig 1, Fig 2	TND0905 (SAD2468)
ΔpilU	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::SpecR, ΔVC1807::CmR, pilA S67C, comEA-mCherry, ΔpilU::TmR *	Fig 1, Fig 2	SAD2135
ΔpilT	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔpilT::TmR	Fig 1, Supp Fig 3	TND1035 (SAD2469)
ΔpilTU	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔpilTU::TmR	Fig 1	JLC227 (SAD2470)
pilT^{K136A}	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, pilT K136A	Fig 1, Fig 2	TND1077 (SAD2471)
pilT^{K136A} ΔpilU	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, pilT K136A, ΔpilU::TmR	Fig 1	JLC444 (SAD2472)
pilT^{E204A}	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A	Fig 1	TND1203 (SAD2473)
pilT$^{E204A/K136A}$	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A/K136A	Fig 1	JLC324 (SAD2474)
pilT$^{E204A/K136A}$ ΔpilU	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A/K136A, ΔpilU::TmR	Fig 1	JLC530 (SAD2475)
pilT^{E204A} ΔpilU	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A, ΔpilU::TmR	Fig 1	JLC531 (SAD2476)
ΔpilA	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::SpecR, ΔVC1807::KanR, ΔpilA::miniFRT	Fig 1	TND0855 (SAD2477)
Parent (micropillar assays)	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::SpecR, ΔVC1807::CmR, pilA S67C, ΔflaA::CarbR, ΔmshA::miniFRT, Δvps-rbmA::ZeoR, ΔtcpA::KanR	Fig 1	TND0752 (SAD2478)
ΔpilT (micropillar assays)	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::SpecR, ΔVC1807::CmR, pilA S67C, ΔflaA::CarbR, ΔmshA::miniFRT, Δvps-rbmA::ZeoR, ΔtcpA::KanR, ΔpilT::TmR	Fig 1	TND0767 (SAD2479)
ΔpilU (micropillar assays)	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::SpecR, ΔVC1807::CmR, pilA S67C, ΔflaA::CarbR, ΔmshA::miniFRT, Δvps-rbmA::ZeoR, ΔtcpA::KanR, ΔpilU::TmR	Fig 1	TND1168 (SAD2480)
P_{tac}-PilT	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔlacZ::P_{tac}-PilT SpecR	Supp Fig 1	JLC426 (SAD2481)

P_{tac}-PilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔlacZ::P_{tac}-PilU SpecR	Supp Fig 1	JLC427 (SAD2482)
P_{tac}-PilT ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔlacZ::P_{tac}-PilT SpecR, ΔpilTU::TmR	Supp Fig 1	JLC420 (SAD2483)
P_{tac}-PilU ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔlacZ::P_{tac}-PilU SpecR, ΔpilTU::TmR	Supp Fig 1	JLC421 (SAD2484)
P_{tac}-PilT pilT^{K136A} ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-PilT SpecR, pilT K136A, ΔpilU::TmR	Supp Fig 1	JLC450 (SAD2485)
P_{tac}-PilU pilT^{K136A} ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-PilU SpecR, pilT K136A, ΔpilU::TmR	Supp Fig 1	JLC451 (SAD2486)
P_{tac}-PilT pilT^{K136A} pilU^{K134A}	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-PilT SpecR, pilT K136A pilU K134A	Supp Fig 1	JLC424 (SAD2487)
P_{tac}-PilU pilT^{K136A} pilU^{K134A}	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-PilU SpecR, pilT K136A pilU K134A	Supp Fig 1	JLC425 (SAD2488)
P_{tac}-PilT ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔlacZ::P_{tac}-PilT SpecR, ΔpilTU::TmR	Supp Fig 1	JLC438 (SAD2489)
P_{tac}-PilU ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔlacZ::P_{tac}-PilU SpecR, ΔpilTU::TmR	Supp Fig 1	JLC439 (SAD2490)
P_{tac}-PilT pilT$^{E204A/K136A}$ ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A/K136A, ΔpilU::TmR, ΔlacZ::P_{tac}-PilT SpecR	Supp Fig 1	JLC540 (SAD2491)
P_{tac}-PilU pilT$^{E204A/K136A}$ ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A/K136A, ΔpilU::TmR, ΔlacZ::P_{tac}-PilU SpecR	Supp Fig 1	JLC541 (SAD2492)
P_{tac}-PilT pilT^{E204A} ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A, ΔpilU::TmR, ΔlacZ::P_{tac}-PilT SpecR	Supp Fig 1	JLC542 (SAD2493)
P_{tac}-PilU pilT^{E204A} ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT E204A, ΔpilU::TmR, , ΔlacZ::P_{tac}-PilU SpecR	Supp Fig 1	JLC543 (SAD2494)
PilU^{L199C}	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilU L199C	Fig 2	JLC409 (SAD2495)
ΔpilT PilU^{L199C}	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S67C, ΔpilT::TmR, pilU L199C	Fig 2	JLC411 (SAD2496)
pilT^{K136A} pilU^{L199C}	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilU L199C, pilT K136A	Fig 2	JLC414 (SAD2497)

pilT^{L201C}	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::SpecR, ΔVC1807::CmR, pilA S67C, comEA-mCherry, pilT L201C *	Fig 2	TND0989 (SAD2498)
pilT^{L201C} ΔpilU	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::SpecR, ΔVC1807::CmR, pilA S67C, comEA-mCherry, pilT L201C, ΔpilU::TmR *	Fig 2	TND1140 (SAD2499)
pilU^{K134A}	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, pilU K134A	Fig 2	TND1092 (SAD2500)
pilT^{K136A} pilU^{K134A}	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, pilT K136A, pilU K134A	Fig 2	TND1094 (SAD2501)
^{6X}His-pilU, pilT^{K136A}	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, ^{6X}His-pilU, pilT K136A	Supp Fig 3	JLC298 (SAD2502)
^{6X}His-pilT	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, ^{6X}His-pilT	Supp Fig 3	HKQ046 (SAD2503)
^{6X}His-pilU	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, ^{6X}His-pilU	Supp Fig 3	JLC398 (SAD2504)
pilT^{K136A} 3xFLAG-pilU	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT K136A, 3xFLAG-pilU	Supp Fig 3	JLC396 (SAD2505)
3xFLAG-pilU	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, 3xFLAG-pilU	Supp Fig 3	JLC326 (SAD2506)
3xFLAG-pilU^{K134A}	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, 3xFLAG-pilU^{K134A}	Supp Fig 3	JLC316 (SAD2507)
3xFLAG-pilT	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, 3xFLAG-pilT	Supp Fig 3	JLC310 (SAD2508)
3xFLAG-pilT^{K136A}	E7946 SmR, ΔlacZ::lacIq, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, 3xFLAG-pilT K136A	Supp Fig 3	JLC349 (SAD2509)
^{6X}His-pilU (purification)	BL21 DE3 pHisTev-6xHis-pilU	Fig 4	HKQ051 (SAD2510)
^{6X}His-pilU^{K134A} (purification)	BL21 DE3 pHisTev-6xHis-pilU K134A	Supp Fig 3	SAD2316
^{6X}His-pilT (purification)	BL21 DE3 pHisTev-6xHis-pilT	Fig 4	TND1277 (SAD2511)
^{6X}His-pilT^{K136A} (purification)	BL21 DE3 pHisTev-6xHis-pilT K136A	Supp Fig 3	TND1281 (SAD2512)
PilT x PilT	BTH101 pUT18C-PilT (CarbR), pKT25-PilT (KanR)	Fig 3, Supp Fig 2	JLC465
PilU x PilT	BTH101 pUT18C-PilU (CarbR), pKT25-PilT (KanR)	Fig 3, Supp Fig 2	JLC466

PilT^{K136A} x PilT	BTH101 pUT18C-PilT K136A (CarbR), pKT25-PilT (KanR)	Fig 3, Supp Fig 2	JLC467
PilU^{K134A} x PilT	BTH101 pUT18C-PilU K134A (CarbR), pKT25-PilT (KanR)	Fig 3, Supp Fig 2	JLC468
PilC x PilT	BTH101 pUT18C-PilC (CarbR), pKT25-PilT (KanR)	Fig 3, Supp Fig 2	JLC469
vector x PilT	BTH101 pUT18C-vector (CarbR), pKT25-PilT (KanR)	Fig 3, Supp Fig 2	JLC470
PilT x PilU	BTH101 pUT18C-PilT (CarbR), pKT25-PilU (KanR)	Fig 3, Supp Fig 2	JLC471
PilU x PilU	BTH101 pUT18C-PilU (CarbR), pKT25-PilU (KanR)	Fig 3, Supp Fig 2	JLC472
PilT^{K136A} x PilU	BTH101 pUT18C-PilT K136A (CarbR), pKT25-PilU (KanR)	Fig 3, Supp Fig 2	JLC473
PilU^{K134A} x PilU	BTH101 pUT18C-PilU K134A (CarbR), pKT25-PilU (KanR)	Fig 3, Supp Fig 2	JLC474
PilC x PilU	BTH101 pUT18C-PilC (CarbR), pKT25-PilU (KanR)	Fig 3, Supp Fig 2	JLC475
vector x PilU	BTH101 pUT18C-vector (CarbR), pKT25-PilU (KanR)	Fig 3, Supp Fig 2	JLC476
PilT x PilT^{K136A}	BTH101 pUT18C-PilT (CarbR), pKT25-PilT K136A (KanR)	Fig 3, Supp Fig 2	JLC477
PilU x PilT^{K136A}	BTH101 pUT18C-PilU (CarbR), pKT25-PilT K136A (KanR)	Fig 3, Supp Fig 2	JLC478
PilT^{K136A} x PilT^{K136A}	BTH101 pUT18C-PilT K136A (CarbR), pKT25-PilT K136A (KanR)	Fig 3, Supp Fig 2	JLC479
PilU^{K134A} x PilT^{K136A}	BTH101 pUT18C-PilU K134A (CarbR), pKT25-PilT K136A (KanR)	Fig 3, Supp Fig 2	JLC480
PilC x PilT^{K136A}	BTH101 pUT18C-PilC (CarbR), pKT25-PilT K136A (KanR)	Fig 3, Supp Fig 2	JLC481
vector x PilT^{K136A}	BTH101 pUT18C-vector (CarbR), pKT25-PilT K136A (KanR)	Fig 3, Supp Fig 2	JLC482

PilT x PilU^{K134A}	BTH101 pUT18C-PilT (CarbR), pKT25-PilU K134A (KanR)	Fig 3, Supp Fig 2	JLC483
PilU x PilU^{K134A}	BTH101 pUT18C-PilU (CarbR), pKT25-PilU K134A (KanR)	Fig 3, Supp Fig 2	JLC484
PilT^{K136A} x PilU^{K134A}	BTH101 pUT18C-PilT K136A (CarbR), pKT25-PilU K134A (KanR)	Fig 3, Supp Fig 2	JLC485
PilU^{K134A} x PilU^{K134A}	BTH101 pUT18C-PilU K134A (CarbR), pKT25-PilU K134A (KanR)	Fig 3, Supp Fig 2	JLC486
PilC x PilU^{K134A}	BTH101 pUT18C-PilC (CarbR), pKT25-PilU K134A (KanR)	Fig 3, Supp Fig 2	JLC487
vector x PilU^{K134A}	BTH101 pUT18C-vector (CarbR), pKT25-PilU K134A (KanR)	Fig 3, Supp Fig 2	JLC488
PilT x PilC	BTH101 pUT18C-PilT (CarbR), pKT25-PilC (KanR)	Fig 3, Supp Fig 2	JLC489
PilU x PilC	BTH101 pUT18C-PilU (CarbR), pKT25-PilC (KanR)	Fig 3, Supp Fig 2	JLC490
PilT^{K136A} x PilC	BTH101 pUT18C-PilT K136A (CarbR), pKT25-PilC (KanR)	Fig 3, Supp Fig 2	JLC491
PilU^{K134A} x PilC	BTH101 pUT18C-PilU K134A (CarbR), pKT25-PilC (KanR)	Fig 3, Supp Fig 2	JLC492
PilC x PilC	BTH101 pUT18C-PilC (CarbR), pKT25-PilC (KanR)	Fig 3, Supp Fig 2	JLC493
vector x PilC	BTH101 pUT18C-vector (CarbR), pKT25-PilC (KanR)	Fig 3, Supp Fig 2	JLC494
PilT x vector	BTH101 pUT18C-PilT (CarbR), pKT25-vector (KanR)	Fig 3, Supp Fig 2	JLC495
PilU x vector	BTH101 pUT18C-PilU (CarbR), pKT25-vector (KanR)	Fig 3, Supp Fig 2	JLC496
PilT^{K136A} x vector	BTH101 pUT18C-PilT K136A (CarbR), pKT25-vector (KanR)	Fig 3, Supp Fig 2	JLC497
PilU^{K134A} x vector	BTH101 pUT18C-PilU K134A (CarbR), pKT25-vector (KanR)	Fig 3, Supp Fig 2	JLC498

PilC x vector	BTH101 pUT18C-PilC (CarbR), pKT25-vector (KanR)	Fig 3, Supp Fig 2	JLC499
vector x vector	BTH101 pUT18C-vector (CarbR), pKT25-vector (KanR)	Fig 3, Supp Fig 2	JLC500
Zip x Zip	BTH101 pUT18C-Leucine Zipper (CarbR), pKT25-Leucine Zipper (KanR)	Fig 3, Supp Fig 2	JLC501
Parent (P_{const}-tfoX)	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C	Fig 4, Supp Fig 3	TND1076 (SAD2513)
ΔpilU	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔpilU::TmR	Fig 4, Supp Fig 3	JLC251 (SAD2514)
ΔpilT	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔpilT::TmR	Fig 4, Supp Fig 3	JLC250 (SAD2515)
pilT^{K136A}	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilT K136A	Fig 4	JLC299 (SAD2516)
P_{tac}-pilT^{K136A}	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac} -pilT K136A SpecR	Fig 4	JLC277 (SAD2517)
ΔpilU P_{tac}-pilT^{K136A}	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔpilU::TmR, ΔlacZ::Ptac-pilT K136A SpecR	Fig 4	SAD2243
P_{tac}-pilU^{K134A}	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-pilU K134A SpecR	Fig 4	JLC282 (SAD2518)
pilT^{K136A} P_{tac}-pilU^{K134A}	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, ΔlacZ::P_{tac}-pilU K134A SpecR, pilT K136A	Fig 4	JLC300 (SAD2519)
P_{tac}-pilT^{K136A} P_{tac}-pilU^{K134A}	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S67C, ΔlacZ::P_{tac}-pilU K134A SpecR, ΔVC1807::Ptac-pilT K136A CmR	Fig 4	SAD2242
P_{tac}-3xFLAG-PilU	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-3xFLAG-pilU SpecR	Supp Fig 3	JLC333 (SAD2520)
P_{tac}-3x FLAG-PilU K134A	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-3xFLAG-pilU K134A SpecR	Supp Fig 3	JLC336 (SAD2521)
P_{tac}-3x FLAG-PilT	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-3xFLAG-pilT SpecR	Supp Fig 3	JLC337 (SAD2522)
P_{tac}-3x FLAG-PilT^{K136A}	E7946 SmR, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P_{tac}-3xFLAG-pilT K136A SpecR	Supp Fig 3	JLC353 (SAD2523)
ΔpilA	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::TmR, ΔpilA::miniFRT	Supp Fig 3	TND1357 (SAD2524)

ΔpilTU	E7946 SmR, ΔlacZ::lacIq, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔpilTU::TmR	Supp Fig 3	JLC252 (SAD2525)
pilU^{K134A}	E7946 SmR, ΔlacZ::lacIq, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::CmR, pilA S67C, pilU K134A	Supp Fig 3	JLC308 (SAD2526)
pilT^{K136A} ΔpilU	E7946 SmR, ΔlacZ::lacIq, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, pilT K136A, ΔpilU::TmR	Supp Fig 3	JLC447 (SAD2527)
pilT^{K136A} pilU^{K134A}	E7946 SmR, ΔlacZ::lacIq, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, pilT K136A, pilU K134A	Supp Fig 3	JLC307 (SAD2528)
P$_{tac}$-pilT	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-pilT SpecR	Supp Fig 3	JLC279 (SAD2529)
ΔpilT P$_{tac}$-pilT	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-pilT SpecR, ΔpilT::TmR	Supp Fig 3	JLC280 (SAD2530)
P$_{tac}$-pilU	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-pilU SpecR	Supp Fig 3	JLC283 (SAD2531)
pilT^{K136A} ΔpilU P$_{tac}$-pilU	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-pilU SpecR, pilT K136A, ΔpilU::TmR	Supp Fig 3	JLC449 (SAD2532)
ΔpilT P$_{tac}$-3xFLAG-pilT	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-3x-FLAG-pilT SpecR, ΔpilT::TmR	Supp Fig 3	JLC373 (SAD2533)
pilT^{K136A} ΔpilU P$_{tac}$-3x FLAG-pilU	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-3x-FLAG-pilU SpecR, pilT K136A, ΔpilU::TmR	Supp Fig 3	JLC448 (SAD2534)
ΔpilU P$_{tac}$-3x FLAG-PilT^{K136A}	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-3xFLAG-pilT K136A SpecR, ΔpilU::TmR	Supp Fig 3	JLC379 (SAD2535)
pilT^{K136A} P$_{tac}$-3x FLAG-pilU^{K134A}	E7946 SmR, P$_{const}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S67C, ΔlacZ::P$_{tac}$-3x-FLAG-pilU K134A SpecR, pilT K136A	Supp Fig 3	JLC380 (SAD2536)
Parent (*A. baylyi*)	ADP1, comP T129C	Fig 5	TND0428 (SAD2537)
PilT^{K136A} ΔpilU	ADP1, comP T129C, PilT K136A ΔpilU::ZeoR	Fig 5	TND1135 (SAD2538)
PilT^{K136A} PilU^{K137A}	ADP1, , comP T129C, PilT K136A PilU K137A	Fig 5	TND1142 (SAD2539)
PilT^{K136A}	ADP1, comP T129C, PilT K136A	Fig 5	TND1122 (SAD2540)
PilU^{K137A}	ADP1, comP T129C, PilU K137A	Fig 5	TND1111 (SAD2541)
ΔpilT	ADP1, comP T129C, ΔpilT::SpecR	Fig 5	TND0779 (SAD2542)
ΔpilU	ADP1, comP T129C, ΔpilU::ZeoR	Fig 5	TND0798 (SAD2543)

ΔpilTU	ADP1, comP T129C, ΔpilTU::ZeoR	Fig 5	TND0799 (SAD2544)
ΔcomP	ADP1, ΔcomP::SpecR	Fig 5	TND0115 (SAD2545)
P_{tac}-pilT	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT	Supp Fig 4	TND1703 (SAD2546)
P_{tac}-pilT ΔpilTU	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT, ΔpilTU::ZeoR	Supp Fig 4	TND1704 (SAD2547)
P_{tac}-pilT ΔpilT	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT, ΔpilT::SpecR	Supp Fig 4	TND1705 (SAD2548)
P_{tac}-pilT ΔpilU	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT, ΔpilU::ZeoR	Supp Fig 4	TND1706 (SAD2549)
P_{tac}-pilT pilT^{K136A}	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT, pilT K136A	Supp Fig 4	TND1707 (SAD2550)
P_{tac}-pilT pilT^{K136A} ΔpilU	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT, pilT K136A, ΔpilU::ZeoR	Supp Fig 4	TND1708 (SAD2551)
P_{tac}-pilT pilU^{K137A}	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT, pilU K137A	Supp Fig 4	TND1709 (SAD2552)
P_{tac}-pilT pilT^{K136A} pilU^{K137A}	ADP1, comP T129C, pMMB67EH- P_{tac}-pilT, pilT K136A, pilU K137A	Supp Fig 4	TND1710 (SAD2553)
P_{tac}-pilU	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU	Supp Fig 4	TND1711 (SAD2554)
P_{tac}-pilU ΔpilTU	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU, ΔpilTU::ZeoR	Supp Fig 4	TND1712 (SAD2555)
P_{tac}-pilU ΔpilT	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU, ΔpilT::SpecR	Supp Fig 4	TND1713 (SAD2556)
P_{tac}-pilU ΔpilU	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU, ΔpilU::ZeoR	Supp Fig 4	TND1714 (SAD2557)
P_{tac}-pilU pilT^{K136A}	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU, pilT K136A	Supp Fig 4	TND1715 (SAD2558)
P_{tac}-pilU pilT^{K136A} ΔpilU	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU, pilT K136A, ΔpilU::ZeoR	Supp Fig 4	TND1716 (SAD2559)
P_{tac}-pilU pilU^{K137A}	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU, pilU K137A	Supp Fig 4	TND1717 (SAD2560)
P_{tac}-pilU pilT^{K136A} pilU^{K137A}	ADP1, comP T129C, pMMB67EH- P_{tac}-pilU, pilT K136A, pilU K137A	Supp Fig 4	TND1718 (SAD2561)

* comEA-mCherry does not affect natural transformation or retraction rates [17].

S2 Table. Primers used in this study

Primer name	Sequence (5' –> 3')	Description
BBC374	TGGCAAAAAGCGAGAGAAGAAG	ΔluxO F1
BBC375	gtcgacggatccccggaatCATGAGGACATATTTTGTTTTCTGC	ΔluxO R1
BBC376	gaagcagctccagcctacaTAAGCGATGAGAGAATGGATCAAC	ΔluxO F2
BBC377	TCACACCCGAATTTCCATCATGC	ΔluxO R2
ABD344	GATTAGCAACGATTCTAGCGCAGGAG	ΔVC1807 F1
ABD340	gtcgacggatccccggaatACGTTTCATTAGTCACCTCTATTGT TAACTTGTTC	ΔVC1807 R1
ABD341	gaagcagctccagcctacaTAGTCGAAAATAAAAAAAGAGG CTCGCCTC	ΔVC1807 F2
ABD345	CTTGCTAACCGTTGGTGTTACCAGC	ΔVC1807 R2
pilA S81C F	AAAACTAATATTGAAGATTATATTGCGACAGAAGGC TgTTTTCCTGCAACAACTGCAGG	pilA S67C F
pilA R2	CATTAATCGCGGTTTCAAAGTGCA	pilA S67C R2
DOG0400	ACTTCTGGCTGAAGGTCAATTTTC	ΔpilT/U F1
DOG0401	gtcgacggatccccggaatCATTTAAATTCCTTAATAAAGTCTG GC	ΔpilT R1
DOG0402	gaagcagctccagcctacaTAGGTAGGTAAAGACAGATGGAG	ΔpilT F2
DOG0403	TCACGTGTTCGGCCAAAATC	ΔpilT/U R2
HQP0001	GGGCTCGGGTgctTCGACGACCTTGGCG	PilT K136A R1
HQP0002	CGCCAAGGTCGTCGAagcACCCGAGCCC	PilT K136A F2
BBC2508	GTTTCTTGGTCACGCAGagCACCGACGAGGATCAC	PilT E204A R1
BBC2509	GTGATCCTCGTCGGTGctCTGCGTGACCAAGAAAC	PilT E204A F2
BBC401	ACCAGCAAAGCTAATAAAATCGAG	ΔpilA F1
BBC402	gtcgacggatccccggaatGAGCATATGCCTTGCTACACAAG	ΔpilA R1
BBC403	gaagcagctccagcctacaACTGCAGGTGCAACAATTAACTAA	ΔpilA F2
BBC404	CGCCATACTAACCCAATACACTC	ΔpilA R2
BBC1752	CTGAATGATTTCCATGAGACG	ΔflaA F1
BBC1753	gtcgacggatccccggaatCATAGTTTGCTCTCCTATCGAG	ΔflaA R1
BBC1754	gaagcagctccagcctacaTTGCAGTAGTTCACGGTACCTTC	ΔflaA F2
BBC1755	TTATACGCTCTTTTGCGTGATGG	ΔflaA R2
BBC1450	CACTCAACGAGCTCAATACG	ΔmshA F1
ABD653	gtcgacggatccccggaatCATCTCTCTTTCATGTGAATACGCA GC	ΔmshA R1
ABD654	gaagcagctccagcctacaGCGCAATAATTTAAATATGGCTCG TGC	ΔmshA F2

BBC1453	ATAGCCTTGCTGTTCATTTTGG	ΔmshA R2
BBC2022	TGCCGATCTCGTTTATGGACG	Δvps-rbmA locus F1
BBC2023	gtcgacggatccccggaatTGCCATTTTGATTGCCTCTGG	Δvps-rbmA locus R1
BBC2024	gaagcagctccagcctacaCAAAGAGAGCCTTATTAGGCTCTC	Δvps-rbmA locus F2
BBC2025	TGAAAGAGGTTGCTCTAGAACTCG	Δvps-rbmA locus R2
BBC1457	TTTCAAGACTTTGGGCAATAG	ΔtcpA F1
BBC1968	gtcgacggatccccggaatCTGTTTTAATAATTGCATATTTATA TAACTCCACC	ΔtcpA R1
BBC1969	gaagcagctccagcctacaTTTGGTAACAGTTAATCTACACCAT TATCTTG	ΔtcpA F2
BBC1463	TTGAATTCGTCCATCATCTAAG	ΔtcpA R2
BBC1833	GATATCCATGTCCATGTCCCAG	comEA-mCherry F1
BBC1834	cttgctcactccaccacttccacctgcTAACAAGATCCTTGCGGCATTG	comEA-mCherry R1
BBC1837	acgagctgtacaagtaaGCTGCAATCATGTGTCTCG	comEA-mCherry F2
BBC1838	GTCTTTCTATCAGTTCGGACTTAGTC	comEA-mCherry R2
BBC1835	gcaggtggaagtggtggaGTGAGCAAGGGCGAGGAGG	mCherry (for comEAmCherry) F
BBC1836	gattgcagcTTACTTGTACAGCTCGTCC	mCherry (for comEAmCherry) R
DOG0405	gtcgacggatccccggaatCATCTGTCTTTACCTACCTAAAATGC	ΔpilU R1
DOG0406	gaagcagctccagcctacaTAGCGATGGTTACTTTGTGCG	ΔpilU F2
JCP048	CAAGCCCCCGATATGATTTgcATTGGTGAAATCCGCAGC	PilU L199C F
JCP049	GCTGCGGATTTCACCAATgcAAATCATATCGGGGGCTTG	PilU L199C R
BBC2287	CATTGCGCTCGGCATTGCGTGAAGACCCAGATGTGATCtgtGTCGGTGAGCTGCGTGAC	PilT L201C F2
BBC2288	CGCAATGCCGAGCGCAATG	PilT L201C R1
BBC2369	GGCTGCCATAGTGGTCGAggcACCCGAACCGGTTGCGC	PilU K134A R1
BBC2347	GCGCAACCGGTTCGGGTgccTCGACCACTATGGCAGCC	PilU K134A F2
BBC2313	gcaggtggaagtggtggaCATCATCATCATCATCACgcaggtggagcaggtgga	6xHis (for 6xHis-PilU/T) F
BBC2314	tccacctgctccacctgcGTGATGATGATGATGATGtccaccacttccacctgc	6xHis (for 6xHis-PilU/T) R
BBC2272	gcaggtggaagtggtggagattataaagaccatgatggtgactacaaggatcacgacattgattataaggatgacgatgacaaagcaggtggagcaggtgga	3xFLAG (for 3xFLAG-PilU/T) F
BBC2273	tccacctgctccacctgctttgtcatcgtcatccttataatcaatgtcgtgatccttgtagtcaccatcatggtctttataatctccaccacttccacctgc	3xFLAG (for 3xFLAG-PilU/T) R
JCP028	tccaccacttccacctgcCATCTGTCTTTACCTACCTAAAATGC	X-PilU (N-term tags) R1

JCP029	gcaggtggagcaggtggaGAGTTGAATCAATATCTGGATGGC	X-PilU (N-term tags OR BACTH TXX-pilU) F2
JCP026	tccaccacttccacctgcCATTTAAATTCCTTAATAAAGTCTGGCTATG	X-PilT (N-term tags) R1
JCP027	gcaggtggagcaggtggaGATATCGCTGAGTTACTGGAATTTAG	X-PilT (N-term tags or BACTH TXX-pilT) F2
ABD332	GGCTGAACGTGGTTGTCGAAAATGAC	P_{tac}-PilT/U @lacZ F1
BBC244	CCCGGGATCCTGTGTGAAATTGTTATCCGC	P_{tac}-PilT/U @lacZ or @VC1807 R1
ABD255	TTAAGCTACTAAAGCGTAGTTTTCG	P_{tac}-PilT/U @lacZ F2
ABD256	CAATTTCACACAGGATCCCGGGAGGAGGTAACGTAATGGTGAGCAAAGGCGAAG	P_{tac}-PilT/U @lacZ R2
BBC1723	caatttcacacaggatcccgggAGGAGGTATTTAAATGGATATCGCTGAGTTAC	Amplify pilT (for P_{tac}-pilT) F
BBC1724	tgtaggctggagctgcttcCTAAAATGCTTTTAAATCCAACTCG	Amplify pilT (for P_{tac}-pilT OR BACTH TXX-pilT) R
BBC1725	caatttcacacaggatcccgggAGGAGGTAGACAGATGGAGTTGAATCAATATC	Amplify pilU (for P_{tac}-pilU) F
BBC1726	tgtaggctggagctgcttcCTACTCCATATCAATCTTCACATTC	Amplify pilU (for P_{tac}-pilU OR BACTH TXX-pilU) R
ABD342	ATTTTTCAGTTGGCCTACAATGCTTTCC	P_{tac}-PilT/U @VC1807 F1
BBC719	CACCATACCCACGCCGAAACAAGGATTTTGAATTAAACGTTTCATTAGTC	P_{tac}-PilT/U @VC1807 R1
BBC243	TTGTTTCGGCGTGGGTATGGTG	P_{tac}-PilT/U @VC1807 F2
BBC2667	tccaccacttccacctgcCATTACGTTACCTCCTCCCG	P_{tac}-X-PilT/U (X is 3xFLAG or 6xHis) R
BBC2513	gaagcagctccagcctacaGATCCCCGGGTACCTAAGTAAC	BACTH pKT25-X F
BBC2512	tccacctgctccacctgcCTCTAGAGTCGACCCTGCAG	BACTH pKT25-X R
BBC2515	gaagcagctccagcctacaCCGAGCTCGAATTCATCGATATAAC	BACTH pUT18C-X F
BBC2514	tccacctgctccacctgcCTCTAGAGTCGACCTGCAG	BACTH pUT18C-X R
BBC2518	gcaggtggagcaggtggaAAAGCGACCCAAACCTTACC	BACTH TXX-PilC F
BBC2524	tgtaggctggagctgcttcTTAACCCAATACACTCATTAAGTTAAAG	BACTH TXX-PilC R
BBC2358	GGTATATCTCCTTCTTAAAGTTAAACAAAATTATTTCTAGAGGGGAATTGTTATCC	pHisTev-X F
BBC2360	cggatccGGCTGCTAACAAAGCCCGAAAGGAAGCTGAGTTGGCTGCTGC	pHisTev-X R
BBC2478	TTTTGTTTAACTTTAAGAAGGAGATATACCATGgcaggtggaagtggtgga	Amplify 6xHis-pilT or 6xHis-PilU to clone into pHistev F
BBC2479	CCTTTCGGGCTTTGTTAGCAGCCggatccgCTAAAATGCTTTTAAATCCAACTCG	Amplify 6xHis-pilT to clone into pHistev R
BBC2480	CCTTTCGGGCTTTGTTAGCAGCCggatccgCTACTCCATATCAATCTTCACATTCTG	Amplify 6xHis-pilU to clone into pHistev R
BBC706	CCACACATTATACGAGCCGATG	P_{const}-tfoX R1
BBC2226	catcggctcgtataatgtgtggaGGGGAACGTGATTAAAGGA	P_{const}-tfoX F2

ABD332	GGCTGAACGTGGTTGTCGAAAATGAC	ΔlacZ::lacIq F1
ABD331	GAACATGGGGTGTACGGCAGTGCtATTTAACGATGTGCGGGTTTTGCCAATCTTG	ΔlacZ::lacIq R1
ABD255	gaagcagctccagcctacaCCACAATAAGCCAGAGAGCCTTAAG	ΔlacZ::lacIq F2
ABD256	CCCAAATACGGCAACTTGGCG	ΔlacZ::lacIq R2
ABD841	GGCACTGCCGTACACCCCATGTTCCGGTACCCGACACCATCGAATG	Amplify lacIq F
ABD842	GGCTCTCTGGCTTATTGTGGGGGGGAAACCTGTCGTGCCAGCTGC	Amplify lacIq R
DOG0175	AAACTATCCAGATAAGGGAAAGC	comP (*A. baylyi*) F1
CE87	GAAGTTGTCGTACATTTCCAacaTACAGCGCCACTAGCATATG	comP T129C (*A. baylyi*) R1
CE88	CATATGCTAGTGGCGCTGTATGTTGGAAATGTACGACAACTTC	comP T129C (*A. baylyi*) F2
DOG0178	AGGATCTGTAATGACGGGTTGAG	comP (*A. baylyi*) R2
DOG0176	gtcgacggatccccggaatCATAAAATTTCTCCACCAATGTTG	ΔcomP (*A. baylyi*) R1
DOG0177	gaagcagctccagcctacaTGATAGTAGTACTATATGGCTTTAAAAG	ΔcomP (*A. baylyi*) F2
BBC1934	GTTACAAAGTCAGGGACGTAAAG	pilT or pilU (*A. baylyi*) F1
BBC1935	gtcgacggatccccggaatATCCATATTTTCCCCGAAGATCG	ΔpilT (*A. baylyi*) R1
BBC1936	gaagcagctccagcctacaTAAGAATAACGCTACTCGATCTG	ΔpilT (*A. baylyi*) F2
BBC1939	GGTATTCAGATTGATCGTCAGTTAG	pilT or pilU (*A. baylyi*) R2
BBC1937	gtcgacggatccccggaatCATCGTATTGTTTTCCTCTCAGATC	ΔpilU (*A. baylyi*) R1
BBC1938	gaagcagctccagcctacaCGCCGTAATCTCAAATAAAAGAG	ΔpilU (*A. baylyi*) F2
BBC2403	GCTGCAAGGGTGGTTGAagcACCGGAACCAGTGG	pilT^{K136A} (*A. baylyi*) R1
BBC2404	CCACTGGTTCCGGTgctTCAACCACCCTTGCAGC	pilT^{K136A} (*A. baylyi*) F2
BBC2408	GGATGCCAAAGACGTGGAagcTCCTGTACCAGTTGC	pilU^{K137A} (*A. baylyi*) R1
BBC2409	GCAACTGGTACAGGAgctTCCACGTCTTTGGCATCC	pilU^{K137A} (*A. baylyi*) F2
BBC2807	caatttcacacaggatcccgggAGGAGGTGAAAATATGGATATAACAGAATTACTCG	pilT amplify for insertion into P$_{tac}$ construct (*A. baylyi*) F
BBC2808	tgtaggctggagctgcttcTTAGAACGACTCTGGAGATTTTG	pilT amplify for insertion into P$_{tac}$ construct or pMMB67EH (*A. baylyi*) R
BBC2809	caatttcacacaggatcccgggAGGAGGTAATACGATGGACTTTAATGATTTACTC	pilU amplify for insertion into P$_{tac}$ construct (*A. baylyi*) F
BBC2810	tgtaggctggagctgcttcTTATTTGAGATTACGGCGTTGAC	pilU amplify for insertion into P$_{tac}$ construct or pMMB67EH (*A. baylyi*) R

BBC2822	aatttcacacaggaaacagaattcgagctcTTGTTTCGGCGTGGGTATGGTG	Clone P$_{tac}$-pilT or P$_{tac}$-PilU into pMMB67EH (*A. baylyi*) F
BBC2824	gagctcgaattctgtttcctgtgtgaaattTTTGCCAGAACCGTTATGATGTC	Amplify pMMB67EH to clone P$_{tac}$-pilT or P$_{tac}$-pilU (*A. baylyi*) F
BBC2357	GGTACCCGGGGATCCTCTAGAGTCGACCTGCAGGCATGCAAGCTTGGC	Amplify pMMB67EH to clone P$_{tac}$-pilT or P$_{tac}$-pilU (*A. baylyi*) R