Ultrafast Structural Changes Decomposed from Serial Crystallographic Data

Zhong Ren

Department of Chemistry, University of Illinois at Chicago, Chicago, IL 60607, USA

Renz Research, Inc., Westmont, IL 60559, USA

zren@uic.edu

Summary

Direct imaging of ultrafast and subtle structural events during a biochemical reaction, such as a single electronic transition from one atomic or molecular orbital to another, is highly desirable but has been beyond the reach of protein crystallography. It entails the capability of observing changes in electronic distributions at both an ultrafast time scale and an ultrahigh spatial resolution (Itatani et al., Nature 432, 867, 2004). The recent developments in femtosecond serial crystallography at X-ray free electron lasers (XFELs) have brought the achievable temporal resolution within a striking distance. This paper presents the electron density map decomposed from the XFEL data that shows the remanence of several $3d$ atomic orbitals of the heme iron at an available spatial resolution although the map component is not an accurate image of the atomic orbitals. A key strategy that has enabled the findings here is a numerical deconvolution to resolve concurrent variations in a series of time-resolved electron density maps so that the electron densities influenced by an electron transfer event can be isolated as a partial change from the overwhelming presence of the bulk electrons that are not directly involved in bonding. Even at the limited spatial resolution, the subtle changes in electron distribution due to a spin crossover can be decoupled from far greater changes due to atomic displacements. Direct observations of electronic orbitals could offer unprecedented mechanistic insights into a myriad of chemical and biochemical

reactions such as electron transfer in redox reactions, and formation, rupture,

and isomerization of chemical bonds.

Ligand photodissociation in carbonmonoxy myoglobin (MbCO) has been a

benchmark for studying ultrafast protein dynamics in a biological system. A

number of studies in time-resolved crystallography have progressively improved

the time resolution (from Šrajer et al., Science 274, 1726, 1996 to Barends et al.,

Science 350, 445, 2015). This paper presents an in-depth analysis of the serial

crystallographic datasets of MbCO that Barends & Schlichting et al. (2015)

contributed to the Protein Data Bank. First, a component of electron density

distributions clearly shows the characteristic shape of the high-spin 3d orbitals

reappeared at the heme iron upon the photodissociation of the CO ligand despite

the limited accuracy of the orbital image due to the available spatial resolution.

Second, the dynamic behaviors of these newly regained 3d orbitals within

picoseconds after the photolysis provide long-awaited structural validation for

previous spectroscopic observations and computational simulations.

Specifically, the newly formed densities are oscillating with the heme iron at a

high frequency of a thousand wavenumbers and developing a positional

modulation during the first few picoseconds (Champion, Science 310, 980, 2005).

The iron pops out of the heme plane at a few picoseconds and recoils back and

pops out again afterwards. The dominant oscillation at a low frequency of

several tens wavenumbers previously detected by coherence spectroscopy can be

clearly resolved from the time series of electron density maps. The associated

changes in electron density during the first cycle of the oscillation are largely

located in the solvent rather than on the protein or heme, which suggests that the

low frequency oscillations in a number of heme proteins, including MbCO, likely

originate from a photolysis triggered pressure wave propagating in the solvated

protein. Finally, these findings of chemical signals are isolated from coexisting

thermal artifacts also by the numerical deconvolution. It is modeled in this study

that the ultrashort XFEL pulses cause a transient spike of the local temperature at

the heme site of hundreds of K.

Keywords

ultrafast dynamics, heme protein, X-ray free electron laser, serial crystallography,

singular value decomposition

Abstract

Direct visualization of electronic and molecular events during chemical and

biochemical reactions will offer unprecedented mechanistic insights. Ultrashort

pulses produced by X-ray free electron lasers (XFELs) offer an unprecedented

opportunity for direct observations of transient events as short-lived as tens of

femtoseconds. This paper presents an in-depth analysis of the serial

crystallographic datasets collected by Barends & Schlichting et al. (Science 350,

445, 2015) that probe the ligand photodissociation in carbonmonoxy myoglobin

(MbCO), a long-serving hallmark for observing ultrafast dynamics in a biological

system. This analysis reveals electron density changes that are caused directly

by the formation of high-spin $3d$ atomic orbitals of the heme iron upon the CO

departure and the dynamic behaviors of these newly formed orbitals in a time

series within the first few picoseconds. The heme iron is found vibrating at a

high frequency and developing a positional modulation that causes the iron to

pop out of and recoil back into the heme plane in succession. These findings

here provide long-awaited visual validations for previous works using ultrafast

spectroscopy and molecular dynamics simulations. This analysis also extracts

electron density variations largely in the solvent during the first period of a low

frequency oscillation previously detected by coherence spectroscopy. This work

demonstrates the power and importance of the analytical methods in detecting

and isolating transient, often very weak signals of electronic changes arising

from chemical reactions in proteins.

Introduction

One of the major applications of X-ray free electron lasers (XFELs) is to directly

observe ultrafast protein dynamics at atomic resolution during photochemical

reactions by serial femtosecond (fs) crystallography (SFX). Time-resolved

experiments (Barends et al., 2015; Coquelle et al., 2017; Nango et al., 2016; Nogly

et al., 2018; Pande et al., 2016; Tenboer et al., 2014) have been successfully

conducted by several groups at XFELs to follow the photodissociation of

carbonmonoxy myoglobin (MbCO; Fig. S1), photoisomerization of the *p*-

coumaric acid chromophore in photoactive yellow protein (PYP), and the

hydroxybenzylidene imidazolinone chromophore in a photoswitchable

fluorescent protein (Levantino et al., 2015a). While the time resolution, that is,

the ability to differentiate transient changes, is limited to tens of fs determined by

the duration of the XFEL pulses, the reported time resolution for MbCO and PYP

was around hundreds of fs due to some timing jitter between a short laser pump

and an XFEL probe.

It is well established that a ligand bound to the heme iron helps to split its

five $3d$ orbitals into two energy levels, the triply degenerate t_{2g} at a lower energy

and the doubly degenerate e_g at a higher energy, so that the six $3d$ electrons of

Fe(II) form three pairs occupying the lower energy orbitals, thus the net spin is

zero. Upon photodissociation, the CO ligand parks at a nearby docking site,

while the heme iron moves out of the heme plane towards the proximal side.

The absence of a bound ligand returns the five degenerate $3d$ orbitals of the iron

back to approximately identical energy levels so that they are all occupied

resulting in a high-spin, deoxy iron. As the porphyrin structure becomes

buckled, several side chains and helices of the globin move accordingly (Fig. 1).

These light-induced structural changes are highly reproducible and have been

observed by both static and time-resolved crystallography, repeatedly (Bourgeois

et al., 2003, 2006; Chu et al., 2000; Kachalova et al., 1999; Schotte et al., 2003; Šrajer

et al., 1996, 2001) . Recently, they have been captured once again at a sub-

picosecond (sub-ps) time scale in MbCO crystals using XFEL pulses (Barends et

al., 2015), a major achievement in SFX. However, no significantly new structural

finding is beyond those already obtained by time-resolved Laue diffraction at

synchrotrons (Bourgeois et al., 2007; Clifton et al., 1997; Ren et al., 1999), which is

one thousand times slower at a time resolution of 100 ps limited by the bunch

length of the synchrotrons (Aranda et al., 2006; Bourgeois et al., 2006; Jung et al.,

2013; Ren et al., 2012; Schotte et al., 2012).

The signals of ultrafast structural events could be too weak to be observed

clearly by the standard crystallographic analyses. In this work, I reexamine the

time-resolved XFEL datasets of MbCO deposited in the Protein Data Bank (PDB;

5cmv, 5cn4-9, and 5cnb-g) by Barends & Schlichting et al. (2015) with a

previously developed analytical method based on singular value decomposition

(SVD; Ren, 2013a; Ren et al., 2012, 2013; Schmidt et al., 2003). This SVD analysis

unravels several ultrafast events that have been identified by spectroscopy and

computational studies but have not been directly observed as images of electron

density distribution. The new findings along with several previously known

structural events upon photolysis of MbCO are summarized in a Gantt chart

(Fig. 1).

Heterogeneous signals in dynamic crystallography

Compared to the well-established structural events of MbCO photolysis such as

ligand docking and heme doming, the ultrafast signals are small in amplitude

and low in population. As a result, they do not stand out at a given time point,

thus went unnoticed in the conventional analyses. As demonstrated below, these

weak yet consistent signals can be manifested as major components resulting

from SVD when multiple datasets collected at different time points are jointly

examined. This necessity for decomposition presents both a challenge and a

unique opportunity for dynamic crystallography, broadly defined as

experimental and analytical approaches that tightly join crystallographic datasets

and their metadata (Ren et al., 2013). We have developed a numerical process of

deconvolution, which is able to identify and extract the dynamic signals from a

large collection of datasets acquired at varying experimental conditions such as

time points and temperatures (Bandara et al., 2017; Kim et al., 2017; Ren, 2016;

Ren et al., 2012; Yang et al., 2011; Zeng et al., 2015). In essence, this strategy takes

advantage of the structural heterogeneity, rather than preemptively avoiding

them as in the common practice of static protein crystallography. In a nutshell,

dynamic crystallography consists of two major steps: first, to experimentally

introduce a functionally-relevant structural heterogeneity; and second, to

numerically resolve the heterogeneous conformational species as long as their

compositions vary as a function of an experimental parameter (Ren et al., 2013).

Here, time-resolved serial crystallography, with the variable being the ultrashort

time delay, is simply a special case in the broader scope of dynamic

crystallography. Therefore, the previously established methodology is largely

applicable (Methods).

Ultrafast dynamics during photodissociation of MbCO

The departure of the CO ligand from the iron porphyrin system has been said

within the first 15-20 fs upon photolysis (Dunietz et al., 2003; Falahati et al., 2017,

2018). This signal is clear in the earliest time point of ~30 fs (nominally -0.1 ps;

Fig. S2). However, the photolyzed CO ligand is already observed in its docking

site 3.5-4.5 Å away from the heme iron at tens to hundreds of fs (Figs. 2c and

3agj) unlike recently suggested by quantum wavepacket dynamics that the

ligand departs only 0.4±0.1 Å (Falahati et al., 2018). A transient charge transfer

from the iron to porphyrin subsequently taking place during 50-300 fs has been

identified as a key event by resonance Raman spectra (Franzen et al., 2001). A

high-spin iron is resulting in the charge transfer from the porphyrin back to the

iron. The transient charge transfer state is not visible in the maps calculated from

the datasets of Barends & Schlichting et al. perhaps due to a broad distribution of

the transferred electron over the porphyrin. However, the larger, high-spin iron

is clearly manifested in the earliest maps (Figs. 2, 3, and S2). Vibrational signals

have been observed by ultrafast spectroscopy, which is attributed to a

discontinuity or "seam" at the intersection of two energy landscapes of the

photoexcited MbCO and the ligand dissociated product (Champion, 2005).

Dissipation of such vibrational energy in the protein matrix and its solvent

environment consists of at least two distinct components: localized, high

frequency vibration modes of the heme at approximately one thousand

wavenumbers; and global, low frequency oscillations of the protein at several

tens wavenumbers (Fujisaki and Straub, 2005). This analysis provides direct

visual evidences to validate both components and to delineate the low frequency

oscillations in greater details (Fig. 7).

The most observed high frequency modes range from hundreds to more than

a thousand wavenumbers (Armstrong et al., 2003; Kitagawa et al., 2002;

Mizutani, 1997). The corresponding vibrational periods of these high frequency

modes of 24-50 fs are comparable to the XFEL pulse duration. Hence an

individual cycle of such vibrations is beyond what can be traced out by the

current XFEL datasets. Nevertheless, the averaged effect of these high frequency

vibrations is detectable (Figs. 2 and 3) in the XFEL datasets of Barends &

Schlichting et al. Amplitude modulation and damping of the high frequency

mode ν_7 was simulated with a period of 350 fs and a time constant of 1.3 ps

(Barends et al., 2015). There are not sufficient time points in the existing datasets

to determine these amplitude modulation and damping. However, a slower

modulation of the high frequency modes is clearly present in these datasets (Figs.

2 and S4; Movie S1). These in-plane modes are thought to be coupled to out-of-

plane motions of the heme and its iron (Nagy et al., 2005), which take place

within a fraction of ps and continue to develop non-exponentially for another

hundreds of ps according to molecular dynamics simulations (Henry et al., 1985;

Kuczera et al., 1993). Recent simulations (Barends et al., 2015; Falahati et al.,

2017) suggested that many transient species, including those with iron in or out

of the plane, are mixed together before 1 ps. It was also suspected that the iron

may recoil back into the heme plane after its initial drop out of the plane (Gruia

et al., 2008). This analysis of the XFEL datasets finds that the out-of-plane

displacement of the iron takes place simultaneously with an in-plane shift of the

porphyrin (Movie S3). There is clear visual evidence to show that the first out-of-

plane move peaks at a few ps and that the recoil and the second drop take place

afterwards (Figs. 2cf and S4c; Movie S1). The signal associated with the iron

displacement lingers on for many μs before the ligand rebinding as captured by

the time-resolved Laue diffraction (Šrajer et al., 2001).

At a low frequency region, the dominant oscillations range between 25 and 45

cm^{-1}, corresponding to a period around 1 ps (Gruia et al., 2008; Rosca et al., 2002).

Such a low frequency oscillation is within the reach of the sub-ps time resolution

achieved with XFEL pulses. Therefore, each individual cycle of the low

frequency oscillation should be, and indeed is, visible in the datasets of Barends

& Schlichting et al. Here I demonstrate that such an oscillation occurs more in

the solvent channels than in the protein or the heme at a low frequency

comparable to those detected by spectroscopy (Fig. 7). This analysis supplies the

long missing visual evidence to support such a low-frequency oscillation in a

solvated protein.

Imaging the geometric shapes of atomic orbitals

The shape of a *d* orbital of copper was first experimentally imaged by Zuo et al.

in their difference electron density map (Humphreys, 1999; Zuo et al., 1999),

which is not, however, without controversy (Mulder, 2011; Scerri, 2000).

Nevertheless, experimenters continued to draw parallels between captured

images and atomic or molecular orbitals, sometimes with the analytical power of

digital reconstructions. Molecular orbital tomography constructs images of

highest occupied molecular orbital from many projections of aligned molecules

(Itatani et al., 2004; Ogilvie, 2011; Schwarz, 2006; Stapelfeldt, 2004; Vozzi et al.,

2011). Both the highest occupied and lowest unoccupied molecular orbitals of

flat molecules, such as pentacene, were scanned using scanning tunneling

microscopy (Chaika, 2014; Patera et al., 2019; Repp et al., 2005). Field-emission

electron microscopy and angle-resolved photoemission spectroscopy can directly

capture or indirectly reconstruct orbital images or transformation of orbital

configurations (Mikhailovskij et al., 2009; Puschnig et al., 2009). More recently, a

photoionization experiment showed the nodal structures of the atomic orbital of

excited hydrogen atoms (Stodolna et al., 2013). A charge-density analysis at an

ultrahigh resolution of 0.48 Å showed iron $3d$ and sulfur $3p$ densities (Hirano et

al., 2016). In all these achievements, the experimentally captured electron

densities with their unique shapes are directly influenced by electrons occupying

specific orbitals. Therefore, in this paper, the images and motion pictures of

difference electron densities with their characteristic nonspherical shape are

attributed to a superposition of several atomic orbitals and their dynamics.

However, this is not to say that the accurate image of an atomic orbital has been

captured, which is not a proper task for protein crystallography, given the limit

in spatial and temporal resolutions.

Results

Decomposition of difference maps and simulated annealing omit maps

To analyze the time-resolved serial crystallographic data of horse heart (hh)

MbCO, one dark and 12 light datasets are obtained from PDB (Fig. 1; Barends et

al., 2015), among which three light datasets collected at the shortest time delays

of 0 and ±0.1 ps span both sides of time 0 that is centered on the excitation laser

flash of 150 fs. These datasets are expected to capture ultrafast signals at delay

times as short as a few fs to tens of fs. However, these ultrafast signals are also

inevitably mixed with signals at longer delays as far as the X-ray pulses stretch

(Haldrup et al., 2011). These sub-ps time points were taken at the same
controlled time delay of 0.5 ps. But the diffraction images were binned according
to the timestamp records from an experimental timing tool (Bionta et al., 2011,
2014), which produced a number of short delays before 1 ps. Several longer time
points ranging from 3 to 150 ps are also included in this analysis (Fig. 1) along
with a deoxy hhMb dataset collected from a synchrotron beamline at 7°C, which
represents the end product of the MbCO photolysis, and two hhMb structures
(1dws/t) in the photolyzed state captured by cryo-trapping (Chu et al., 2000).
277

Two types of electron density maps, namely difference Fourier map and
simulated annealing omit map (SAOM), are produced and subjected to the SVD
analyses separately. Each type carries its own advantage (Methods). While
difference Fourier maps are sensitive for detecting weak signals, they are also
highly susceptible to systematic errors that often overwhelm the signal. It is
critically important to choose a proper reference dataset, which is preferably
obtained under the identical experimental condition, if not from the very same
crystal as the light dataset. An alternative is to use SAOMs, where a small
portion of a protein structure is set aside from structure refinement using the
simulated annealing algorithm. In such omit maps, the electron densities within
the omitted region shall not be biased by the protein model, thus are considered
as the most authentic representation of the experimentally observed electron
densities (Hodel et al., 1992). A collection of SAOMs within a common omitted
region, although less sensitive to subtle changes compared to difference maps,
permits a joint analysis of maps derived under very different experimental
conditions, which is usually difficult with difference maps.
294

Difference Fourier maps are synthesized using a Fourier coefficient set of
F_{light}-$F_{reference}$ combined with the best possible phase set. While $F_{reference}$ is typically
the structure factor amplitudes from a dark dataset (5cmv), it is important that a
light dataset at a given time point is also used as a reference (Table S1) to reveal
changes with respect to that specific time point (Methods). Therefore, a
collection of difference maps produced with the same reference dataset
constitutes a time series that show continuous changes from the reference time
point and on. For example, -F_0 series refers to the difference maps with the

reference dataset at the nominal 0 ps. For omit maps, the heme and several side

chains within its immediate vicinity are omitted to produce SAOMs of the ligand

binding site (Methods). A few static structures determined at room temperature

of sperm whale (sw) Mb in the aquomet (1bz6), deoxy (1bzp), and CO-bound

(1bzr) forms are also included (Kachalova et al., 1999), along with three hhMb

structures (1dwr/s/t) (Chu et al., 2000) as control data points in this joint analysis.

The resolutions of these static datasets are truncated to 1.75 Å to match the XFEL

datasets.

Altogether, a total of 93 difference maps (Table S1 and Fig. S2) and 20 omit

maps are obtained for two independent SVD analyses. The electron densities are

masked around the heme site unless stated otherwise (Methods). In this work,

the significant signals are identified by the top nine components from SVD (Fig.

2e). A numerical procedure called rotation in the SVD space plays an important

role to extract chemically sensible components from these (difference) electron

density maps (Henry and Hofrichter, 1992; Ren, 2013a, 2013b, 2016). Proper SVD

rotations reveal the correlation between the core data, i.e., electron densities, and

their metadata, i.e., delay time, sample temperature, and other experimental

parameters (Methods). Given their highly dynamic nature, the ultrafast signals

due to multiple causes are often entangled before suitable rotations are found in

the multi-dimensional SVD space (Methods). Using this decomposition and

rotation strategy, I identify signals that reveal the electronic redistributions

between high and low spin of the iron orbitals at the available spatial resolution.

Molecular image of low-spin to high-spin crossover

The ultrafast signals within several fs to a few tens of fs are manifested in the

fourth component, denoted as U_4, of 93 difference maps. The component U_4

exists only in the map series with the dark dataset as reference (red, $-F_{dark}$ series

in Fig. 2). That is to say, all signals in this component have occurred by the time

of the very first time point at ~30 fs (nominally -0.1 ps), therefore the map

series $-F_{-0.1}$ shows the cancellation of this component (orange series in Fig. 2).

Some of these signals start to decay slightly before 1 ps and diminish

significantly after 3 ps (Fig. 2cd). The corresponding map of U_4 reveals the

previously well-established events arising from the CO docking, out-of-plane

motion of the iron, and an uneven heme doming upon photolysis of MbCO (Fig.

3). The negative densities associated with the pyrrole rings and their propionic

acid side chains increase in intensity from ring A to B to C to D (Fig. 3c). Positive

densities are found at a cross section 1.1 Å above the heme plane towards the

distal side (Fig. 3b).

More remarkably, U_4 captures some hitherto unseen signals characterized by

a cubic-shaped network of positive densities that cage the iron atom. To

facilitate a more concise description, a cross section of $Z = 0$ is defined as the

heme plane and the cross section of $X = 0$ is the dividing plane between two

propionic acid side chains, from which the third orthogonal cross section of $Y = 0$

can be derived (Fig. S1). The eight corners of the positive cube are oriented along

the X and Y cross sections. Strong positive densities appear in the middle of four

six-membered rings encircled by adjacent pyrroles, the connecting methane

bridge, and the heme iron (marked 1-4 in Fig. 3c). These positive features reach

their four minima when they cross the heme plane. These positive densities

become more prominent above and below the heme plane (Fig. 3). The eight

corners of this cube are located at 1.7±0.2 Å away from the iron. This cubic-

shaped network is asymmetric with a notable skew towards CD rings (Fig. 3fh).

Meanwhile, the iron skews in the opposite direction towards the AB rings with

an out-of-plane displacement towards the proximal side (Fig. 3e). But the major

features of this cube seem more symmetric in the Y cross section.

Similar features are also evident in the seventh component U_7 from an

independent SVD analysis of 20 SAOMs. It is clear that the seventh dimension

after a proper rotation (Methods) represents the reaction coordinate of the ligand

dissociation. Mb structures along this reaction coordinate form a lineup on c_7

dimension regardless of sample temperatures: from the ligated structures of

MbCO, to the time-resolved structures, to several photoexcited species including

the static structures Mb*CO and **CO, eventually to deoxy Mb (Fig. 4e). U_7 of

the omit maps contains the usual signals of ligand dissociation, such as strong

negative densities on the ligand indicating dissociation and a positive layer on

the distal side of the heme indicating its doming (Fig. 5), except that the quality

of these signals are not as good as those from difference maps because the

SAOMs are derived from datasets in very different experiments. Despite the

systematic errors, the seventh component U_7 of the omit maps again clearly

shows the cubic network of positive densities around the iron. This cubic

network also skews towards the CD rings of the heme. This is a reoccurrence of

the same signals shown in U_4 of the difference maps (compare Figs. 3 and 5). The

eight corners of the positive cube are 1.9±0.2 Å away from the iron.

It is clear that the positive densities arranged in the cubic network around the

iron, either in the difference maps or in the omit maps, cannot be easily

attributed to accidental noise or systematic errors such as thermal effect or

Fourier truncation error (see Discussion). They originate from the intrinsic

signals induced by light and coexist with the previously seen signals of ligand

dissociation. While multiple interpretations of these distinct cube-shaped signals

may apply, this positive network clearly indicates a larger, high-spin iron as an

immediate overall effect upon photolysis. Here I further attribute the distinct

shape of the positive feature to the instantaneous regaining of the high-spin $3d$

orbitals of the heme iron. However, the observed cubic densities here are not an

accurate image of a probability distribution of a $3d$ electron. They are resulted

from a net gain of electron densities as a fully ligated iron transitions to the high-

spin deoxy state when several $3d$ orbitals are reoccupied by unpaired electrons.

Calculations of the combined probability distributions of the textbook $3d_{xz}$ and

$3d_{yz}$ with either $3d_{xy}$ or $3d_{x^2-y^2}$ orbitals show features highly similar to the

observed cubic-shaped map (Fig. 3lm). The strongest electron densities are

located at eight corners of the cube where the lobes of $3d$ orbitals intercept. It is

entirely possible that the observed cubic densities result from not only a

superposition of several $3d$ orbitals but also an average of multiple electronic

configurations. See Discussion below. Such orbital changes have not been

captured previously in protein crystals by X-ray crystallography.

However, the loss of the paired low-spin electrons in the ligated state is not

clear in the negative densities. Presumably, the loss of the CO ligand and the

drop of the iron out of the heme plane cause the displacement of many more

electrons than the low-spin electrons. Negative densities in the component U_4

are overwhelming. At the available spatial resolution, the loss of the low-spin

electrons is not separated from the main events upon photolysis.

Interestingly, the signals in U_4 of difference maps decay after a few ps leading

up to the static structures Mb*CO, **CO, and the deoxy Mb (Fig. 2cd). Given in-

plane vibration modes (Abe et al., 1978; Nagy et al., 2005) with time periods

comparable to the XFEL pulse duration, the averaged electron density between

vibrating atoms is expected to increase, as shown in the six-membered rings.

The observed decay of U_4 can be interpreted as the damping of these vibrations

(see below). These in-plane vibrations trigger the iron to pop out of the heme

plane (Franzen et al., 1995). However, it remains unclear from this observation

which, or a combination of which, vibration modes plays the dominant role.

Importantly, this decay does not reach zero even for the static deoxy Mb (Fig.

2cd), which suggests that the above interpretation of the cubic network of

positive densities as the consequence of the spin crossover largely holds, since

the cubic network cannot be fully explained by high frequency vibrations. This

decay of U_4 after a few ps is also an evidence to support the model of iron

recoiling back into the heme plane (Gruia et al., 2008), which has to be discussed

with the top component U_1 below.

Development, modulation, and damping after 1 ps

If the cubic network of positive densities is indeed influenced by the newly

formed 3*d* orbitals, the dynamic behaviors of these high-spin electrons should be

visible in the time points shortly after the initial formation. The observed

evolution of the cubic network provides the necessary validation. Along the

dimension c_1 in the SVD space of 93 difference maps, all time points after 1 ps,

including the deoxy Mb (5d5r, representing a time point of ∞), are well separated

from the sub-ps time points around $c_1 = 0$ (Fig. 2ac). Therefore, the first

component U_1 of the difference maps indicates a major development after 1 ps

and represents more permanent changes afterwards. The cubic positive densities

in U_4 have no sign in U_1. Among the major features captured by U_1 (Fig. S3), the

most noticeable is a rather skewed displacement of the iron towards the AB rings

of the heme. In the meanwhile, the motion of His93 corroborates the skewed

displacement of the iron (Fig. S3ae). It is also shown below that the entire heme

moves in its plane towards the AB rings as well. Such long-lived signals in U_1

from 1 ps and on are in good agreement with a wide distribution of individual

trajectories and interquartile range of iron displacement predicted by molecular

dynamics simulations (Barends et al., 2015; Henry et al., 1985; Kuczera et al.,

1993). However, the cryo-trapped structures, Mb*CO and **CO, are two

exceptions. They exhibit very little characteristics in U_1 (Fig. 2ac), even though

they presumably represent long time delays. It can be reasoned that the skewed

displacement of iron requires concerted motions in the proximal His93 and helix

F, which are evidently forbidden at cryogenic temperatures.

Both U_1 and U_4 feature the signal of the iron out-of-plane displacement (Figs.

3e and S3c). Therefore, the large positive coefficients c_1 and c_4 occurring

simultaneously around several ps indicate that the iron pops out of the heme

plane most at this time but only momentarily. The later decay of the fourth

component U_4 records a recoiling of the iron back into the heme plane around

100 ps (Movie S1). Another major signal revealed after 1 ps is a modulation of

the new formation of $3d$ orbitals indicated by the cubic-shaped positive densities

around the iron. This is a positional modulation manifested in the sixth and

eighth components of the difference maps. While the corresponding coefficients

c_6 and c_8 are nearly constant in the sub-ps range, they start to ramp up and vary

at a few ps. This variation can be most easily modeled as a two-dimensional

oscillation at a period of 220±20 ps given the limited time points (Fig. 2f).

Additional observations in the future may update this positional oscillation. The

U_6 component features a distorted network of positive densities around the heme

iron (Fig. S4) similar to U_4 but with an inclination towards ring B. In U_6, the

strong negative density on iron is skewed towards the proximal side in contrast

to U_4 (compare Figs. 3e and S4c). The positive peak on the proximal side of the

iron is also extremely skewed (Fig. S4c). In a linear combination sense, U_6 and U_8

can be interpreted as variations that are combined into U_4 resulting in an

oscillation of a period at a few hundreds of ps (Fig. 2f). Specifically, the cubic-

shaped densities lean towards ring D from 0.5 to 3 ps, return to the mean

position after 50 ps, then sway towards ring B in the opposite direction (Fig. 2f

and Movie S1). Near the end of the first oscillation, the iron again pops out of

the heme plane for the second time and severely skews towards AB rings (Movie

S1). These observations of the iron motions support the previous model of iron

recoiling (Gruia et al., 2008). Such oscillating behavior suggests modulation and

damping of the vibrational modes. In other words, the cubic network of positive

densities is not only observed in the earliest time points. The motions of this

network are captured in later time points as well. This 220-ps oscillation is

largely along BD cross section (Fig. S1), while damping of the vibrational

amplitude was only recorded up to 150 ps thus insufficient to depict a complete

picture. Two other more significant components U_2 and U_3 also show an

oscillation of a distinct type as discussed below (Fig. 2b).

Vibration induced by XFEL pulses and instantaneous local temperature jump

An independent SVD analysis of 20 simulated annealing omit maps (SAOMs)

masked around the heme site (Methods) offers a direct comparison of the XFEL

data with previous synchrotron data. Among the top eight components of the

SAOMs, the first component reveals a well-defined molecular image of the heme

and its surrounding (Fig. S5); and the second and fourth components apparently

account for differences between hhMb and swMb. It is the third component U_3

that clearly distinguishes the XFEL datasets from the synchrotron datasets in a

temperature-dependent manner (Fig. 4ace). In other words, the cryo structures

(1dwr-t), ambient temperature structures (1bz6/p/r) including the deoxy hhMb at

7°C (5d5r), and the room temperature XFEL structures line up in an ascending

order along the dimension c_3. Not incidentally, the photolyzed structures

Mb*CO and **CO have the most negative coefficient c_3 because they were

obtained at temperatures even lower than 100 K (Chu et al., 2000). The

corresponding decomposed map U_3 reveals the thermal effects in electron

densities of the heme as a nearly symmetric shell of spherical positive densities

around the iron with very little positive or negative density on the iron itself (Fig.

6), which transitions into a square shape with four corners centered at the six-

membered rings in the heme plane (marked 1-4 in Fig. 6e). This symmetric U_3 of

the omit maps shows no usual signals for ligand dissociation and heme doming

therefore contrasts with the asymmetric U_4 of the difference maps characterized

by the distinct cube-shaped positive density maxima at the eight corners. Two

sheets of positive densities are found at the cross sections parallel to, and 1.3 Å

away from, the heme plane on both sides (Fig. 6df). They are connected via

pillar-shaped positive densities that run through the six-membered rings

(marked 1-4 in Fig. 6e) and the pyrrole rings (marked 5-8 in Fig. 6e) along with

some other positive densities outside the heme (marked 9-20 in Fig. 6e). It is

noteworthy that such thermal effects do not stand out as a major component in

the SVD analysis of difference maps because they are largely self-canceled

during the subtraction of the reference.

The temperature dependency and the symmetry of the signals in U_3 strongly

suggest that these features are related to the local temperature at which the

MbCO molecules were probed, rather than artifacts arising from structural

refinement or difference in spatial resolutions among datasets. All datasets used

for the analysis are unified to a similar spatial resolution (Methods). The

positional parameters of the models within the omitted region and the thermal

parameters of the entire globin are intentionally "forgotten" in the calculation of

the omit maps. The large c_3 coefficients of the XFEL datasets are due to

instantaneous temperature elevation at the heme site upon exposure to each

ultrashort XFEL pulse, which introduces isotropic vibrations at high frequencies

to the entire tetrapyrrole cofactor as captured by U_3. This molecular image of U_3

shows an increasing intensity from cryo to ambient temperatures and to a very

high local temperature induced by XFEL pulses (Movie S2). Solely based on an

estimate by linear extrapolation without other validations, the instantaneous

temperature of the heme may exceed 500 K during an 80-fs XFEL pulse (Fig. 6i).

However, it remains to be seen whether this analysis of the omit maps restricted

within the heme site can be extended to show that the temperature of the globin

or the crystals has a significant elevation during an XFEL pulse.

Identification of such symmetric and temperature-dependent signals is

important for interpretation of the highly dynamic ultrafast signals. The fact that

the symmetric U_3 and the asymmetric U_7 of the omit maps are orthogonal to each

other mathematically guarantees that these coexisting components are not

interchangeable and not cross contaminated between them (Methods). The cubic

shaped U_7 of the omit maps and its counterpart U_4 of the difference maps are the

chemical consequence of ligand dissociation; the symmetrical and spherical

shaped U_3 originates from the thermal consequence. One cannot compensate for

the other and both coexist in all maps. The equivalents can be stated: The cubic

positive densities influenced by the $3d$ orbitals are not due to thermal vibrations;

the symmetrically enlarged iron and the thickened heme due to an instantaneous

temperature elevation have been isolated from the chemically relevant signals. It

is worth noting that the ultrafast onset of U_4 of the difference maps (Fig. 2cd) is

not in contradiction with U_7 of the omit maps contained not only in time-

resolved XFEL datasets but also in the static structures of deoxy Mb and

photolyzed Mb*CO and **CO at different temperatures (Fig. 4e). Instead,

regaining of the high-spin $3d$ orbitals of the iron occurs before the first time point

at tens of fs; this permanent change lasts throughout the dissociation reaction

into the deoxy product.

The asymmetric U_7 of the omit maps also describes an important but

unfamiliar structural event upon the ligand dissociation – the heme in-plane

movement. This type of motion is harder to capture in difference maps due to

little net gain or loss of electron density that would be caused by an in-plane

movement of the heme. With respect to the protein framework, the heme moves

towards its AB rings in addition to the other changes (Movie S3). Heme in-plane

sliding is consistent with the strong signals of the skewed iron displacement and

the proximal His93 (Fig. S3) that are closely related to structural responses in the

globin. Similar in-plane sliding is even more significant in tetrameric

hemoglobins (Baldwin and Chothia, 1979), which is the origin of the cooperative

oxygen affinity (Ren, 2013b). However, the functional role of the heme in-plane

sliding in Mb is not yet clear and this discussion branches away from the topic

here on ultrafast structural changes.

Global oscillation at low frequency

A recent solution scattering experiment using XFEL pulses showed that the

radius of gyration R_g first increases within 1 ps upon photodissociation of MbCO

and then oscillates with a period of 3.6 ps, equivalent to 9 cm^{-1}, a frequency lower

than those observed by coherence spectroscopy (Levantino et al., 2015b). The

molecular volume increases afterward with a phase shift. These observations

indicate a wavelike motion propagating outward in the protein. Recent

molecular dynamics simulations have confirmed that a pressure wave generated

by photolysis reaches the molecular surface at 300 fs along the direction

perpendicular to the heme plane (Brinkmann and Hub, 2016), which corresponds

to a frequency of 28 cm^{-1} if the wave is reflected by the surface and oscillates. It

was found by the simulations that the oscillation in solvent would be

overdamped thus dissipates quickly.

Some SVD components derived here display strict correlation suggesting

their dependency on each other. The U_2-U_3 pair of the difference maps (Fig. 2b)

is evidently equivalent to the U_5-U_8 pair of the omit maps (Fig. 4b). When the

difference maps of the entire globin are analyzed, the same signals emerge as the

top-ranked U_1-U_3 pair suggesting that the local signals at the heme site concur

with global signals over the entire globin (Fig. 7). Interestingly, this pair of

components displays a near-perfect circular correlation at sub-ps delays but is

almost absent at longer delays > 1 ps (Fig. 7afgh). Such an oscillation at a low

frequency requires at least two orthogonal components to depict. The oscillating

signals have spread over the entire globin at sub-ps delays while their

amplitudes decay to nearly zero after a few ps. Based on least-squares fittings of

the corresponding coefficients (Methods), the frequency of the oscillation is

determined to be 36±1 cm^{-1}, corresponding to a period of 0.93 ps. The equivalent

oscillation frequencies determined from the difference maps and omit maps at

the heme site are 42±2 cm^{-1} (a period of 0.8 ps) and 37±1 cm^{-1} (a period of 0.89 ps),

respectively.

The low frequency oscillation determined from the XFEL datasets of Barends

& Schlichting et al. is highly comparable to those measured by fs vibrational

coherence spectroscopy (Gruia et al., 2008; Rosca et al., 2002). I postulate that this

low frequency reflects the molecular size of the solvated Mb rather than the

property of the heme. First, the corresponding period of 0.93 ps is about the time

that a mechanical wave takes to traverse 14 Å, that is, an average distance from

the heme to the molecular surface of Mb, given a speed of sound at 1.5 km/s or

nm/ps (Fig. S1). Second, the molecular sizes of several heme proteins, such as

cytochrome c (12 kDa), Mb (17 kDa), and cystathionine β-synthase (63 kDa),

show good anticorrelation with their dominant low frequencies of 44, 40, and 25

cm^{-1}, respectively (Gruia et al., 2008; Karunakaran et al., 2010, 2014). Third, no

608 association between the oscillating signals and specific structural features can be

609 found, such as the heme, ligand, surrounding side chains, or helices (Fig. 7ik).

610 These oscillating signals sharply contrast with strong, non-oscillating signals

611 associated with the heme and various helices evidenced by the second and fourth

612 components of the difference maps (Fig. 7jl), where helices E, G, and H move

613 away from the heme in response to ligand dissociation (Barends et al., 2015).

614 These motion signals of various structural elements are much enhanced by SVD.

616   The oscillating signals at low frequency seem to propagate via solvent within

617 the protein matrix rather than the globin. Strong positive and negative peaks in

618 U_5, U_6, and U_8 of the omit maps are consistently found in the solvent region away

619 from the heme and protein. However, a specific description on how the solvent

620 responses to ligand dissociation is difficult, since the solvent structure is far less

621 well determined than the globin structure. An overall contribution of U_5, U_6, and

622 U_8 can be judged by a sum of the absolute values $|U_5|+|U_6|+|U_8|$ that enhances

623 the distribution of signals. The heme and globin are located in low intensities of

624 these oscillating components (Fig. 7b-e). These global oscillations occur only a

625 few times before dissipation (Gruia et al., 2008; Rosca et al., 2002). However,

626 there are not enough time points to determine the damping time constant, which

627 is roughly estimated to be 1-4 ps. Contrary to the common belief that low

628 frequency oscillations are due to the properties of the heme, here I attribute the

629 low frequency oscillation to the global response of protein and solvent as a whole

630 to a sudden molecular event – photodissociation of the ligand, where the

631 vibrational energy released from the photodissociation is quickly dissipated into

632 the environment.

634 **Discussion**

635 *Are atomic orbitals observed?*

636 It requires ultrahigh spatial resolution in a Fourier synthesis to accurately depict

637 the probability distribution of a single electron occupying its orbital. For

638 example, iron $3d$ and sulfur $3p$ densities in an Fe_4S_4 cluster are described with

639 crystallographic data at a resolution of 0.48 Å with an analytical technique called

640 charge-density analysis (Hirano et al., 2016), in which atoms are considered as

641 aspherical and modeled by spherical harmonic functions in contrast to spherical

Gaussian functions used in everyday protein crystallography (Coppens, 1998).

Since the non-bonding inner electrons remain largely constant in a spherical

distribution as a whole, the inner electrons in a majority can easily overwhelm

the presence of the bonding electrons in a minority, let alone subtle changes in

electron transfer. Therefore, it is very difficult, if not impossible, to detect small

orbital changes upon bond rupture using regular electron density maps such as a

2Fo-Fc map. Difference Fourier technique is required as it is far more sensitive to

small changes (Henderson and Moffat, 1971), in which the constant, spherical

distribution of the inner electrons is self-canceled and the remaining densities

would reveal electron transfer. However, the reality is usually far less ideal

because of conformational changes. Atomic displacements involved in a

conformational change carry all electrons associated with the moving atoms and

result in significant positive and negative densities in a difference Fourier map,

that is, a spatial distribution of any gain and loss of electrons. Atomic

displacements such as a side chain rotation or the heme iron movement by a few

tenths of an Å again conceal the subtle signal of an electron transfer from one

orbital to another.

The difference maps derived here as SVD components represent partial,

variable signals in addition to the constant electron density map regardless of the

bonding change. An ultrahigh spatial resolution is required to depict slightly

aspherical features when an overwhelming spherical background is present.

Here, a combined use of difference Fourier technique and decomposition eases

the demand for spatial resolution to reveal subtle changes. Furthermore, the

notion that an ultrahigh spatial resolution is required for observations of orbital

changes contradicts the fact that there is no discontinuity in the probability

distribution of any electronic orbital. A reasonably high spatial resolution shall

be sufficient to analytically describe the smooth function of a gain or loss of an

electron. Therefore, the small amplitude of desired changes over a much greater

constant background hinders a reliable observation more than the lack of high

spatial frequencies does. This work presents the effectiveness of the strategy that

enables numerical deconvolution to resolve concurrent events so that the image

of a partial change, such as a single electron transfer, can be isolated from other

changes. At an available spatial resolution, the partial change due to a specific

electronic event deconvoluted from much greater conformational changes could

be the remanece of orbital densities instead of an accurate image of these orbitals.

A limited experimental spatial resolution also raises a doubt whether some

unusual features are the artifacts due to a Fourier truncation error. A Fourier

truncation error may occur to several static datasets included in this analysis, in

which the spatial resolutions are truncated to 1.75 Å to match that of XFEL

datasets. Such errors may appear as minor components after SVD (Figs. 2e and

4f). When the average intensity of an experimental dataset naturally decays as

function of the spatial resolution, Fourier truncation errors are damped to a

minimum. Difference Fourier technique again self-cancels the remaining errors

because both datasets involved in the subtraction would carry similar Fourier

truncation errors if any. In addition, the applied weighting scheme (Ren et al.,

2001, 2013; Šrajer et al., 2001; Ursby and Bourgeois, 1997) severely reduces any

large differences in Fourier synthesis (Methods), which also prevents Fourier

truncation errors. After the decomposition, proper rotations in the SVD space

validate that the signals under discussion, such as the positive cubic network, is

not due to Fourier truncation errors because of its correlation with the metadata.

An artifact due to Fourier truncation errors, if so conclude, needs to be explained

why it is correlated with a physical or chemical parameter.

The theoretical radii of the $3d$ orbitals for an isolated Fe(II) cation are about

0.36 Å at the maximum probability of these orbitals. However, the orbital

densities spread out many folds beyond the maxima (Waber and Cromer, 1965).

The majority of the iron $3d$ orbital densities in an Fe_4S_4 cluster span about 1 Å

from the nucleus as measured in a protein (Hirano et al., 2016). The observed

distances between the iron and the corners of the observed cubic densities are

significantly larger. Therefore, the observed cubic densities here are not an

accurate image of a probability distribution of the $3d$ electrons. However, this

density distribution, given the available spatial resolution, is clearly influenced

by the unpaired $3d$ electrons. The net gain of electron densities could be further

away from the nucleus compared to the maximum probability of these orbital.

For a practical purpose in protein crystallography, an accurate depiction of a $3d$

orbital is not the task. Rather, direct observations of a specific electronic event
would provide unprecedented insight into the essence of biochemistry.
711
Temporal resolution vs. resolution of heterogeneity
Ultrashort XFEL pulses improve the time resolution to sub-ps or even tens of fs
compared to 100 ps at synchrotrons. This ultrafast time resolution, when
combined with the spatial resolution of crystallography, offers an unprecedented
opportunity for direct observation of transient structural changes in a
photochemical reaction as detail as individual electrons. However, due to their
highly dynamic and heterogeneous nature, it is difficult to experimentally isolate
short-lived structural species along the temporal dimension regardless of the
ultrafast time resolution. In other words, resolving the population difference
between two chemical species at one fs from the next does not mean that one can
resolve two distinct structures. As demonstrated by this work, analytical
methods such as SVD are needed to deconvolute them from mixtures of
intermediate species. The strategy we proposed for resolution of structural
heterogeneity (Ren et al., 2013) is highly effective for separating independent
electronic and atomic events that often coexist during a chemical reaction or a
biological process.
728
This work shows that it is possible to observe structural changes as small as a
gain or loss of individual electrons by protein crystallography. Small structural
signals can be correlated with their corresponding metadata by proper rotations,
that is, deconvolution after decomposition, thus allowing the origin of these
signals be revealed. I posit that all observations so far are the consequence,
rather than the cause, of the ligand dissociation. The proposed transient event of
metal-to-ligand charge transfer is not captured here, presumably because the
transferred electron is delocalized all over the porphyrin (Franzen et al., 2001).
Therefore, the fundamental questions remain for photolysis of MbCO: Upon the
absorption of a quantum of energy from a green photon, what initial electronic
events lead to a partitioning of the bonding electrons in two separate atoms, that
is, the bond rupture? What electronic events cause spin crossover? To image
these events directly, even higher time resolution is required to capture electron
density changes prior to and during bond rupture.

Future outlook of ultrafast dynamic crystallography

Despite the recent developments in SFX and its applications to time-resolved

studies of a few model systems, it is yet premature to declare that the current

SFX platform applies to other systems of more significant biological importance

such as visual and microbial rhodopsins, plant and bacterial phytochromes,

photosynthetic reaction centers, and light-dependent DNA repair enzymes.

There are at least three major challenges. First, it is not practical to produce and

inject an astronomical number of nano or microcrystals of these proteins into an

XFEL beam to support data collections of a time series. The current serial

protocol used to obtain the MbCO and other data has a diminishingly small yield

of useful diffraction data, based on the ratio between the number of diffracting

crystals and the total number of crystals produced (Ren et al., 2018). That is to

say, the vast majority of the crystals have no chance to diffract X-ray even when

they are injected into the X-ray beam. Such a yield, defined differently from the

hit rate, is not sustainable for other photosensitive protein crystals. Several

"fixed target" implementations of crystal delivery systems have been reported,

including one that we have recently proposed (Ren et al., 2018). However, it

remains to be seen how effective they are in achieving a higher yield while

maintaining the diffraction quality.

Second, since the XFEL pulses usually carry insufficient bandwidths to fully

integrate the Bragg reflections and in the meanwhile angular rotation of a crystal

during an ultrashort XFEL pulse is not attainable, tens to hundreds of thousands

of diffraction images are required to obtain integrated intensities for all Bragg

reflections in a complete sampling of the reciprocal space. A possibility to solve

this partiality problem is to use an X-ray beam with a significant convergence of

about a degree (Ho et al., 2002; Spence et al., 2014), which will produce

completely integrated intensities with a far fewer number of diffraction images,

thereby alleviating the demand for a large quantity of purified proteins.

Third, data merging from a large crystal pool is a major source of error in

producing difference signals for photosensitive crystals, particularly those with

much lower diffraction power than Mb crystals. To address this challenge, we

also proposed a protocol to obtain difference signals that bypasses the integrated

intensities of Bragg reflections. We reason that integrated intensities are not

necessary if the dark and light diffraction images can be obtained within a single

exposure of an XFEL pulse because the percentage change in integrated

intensities of a given reflection, that is, the time-resolved signal, is already

captured even if this reflection is partially measured in both dark and light

conditions. A plan has been laid to implement this protocol, based on a

dichromator instrument that achieves angular split and temporal delay at the

same time (Ren and Yang, 2016). When each of these technical hurdles is

overcome, direct observations of atomic and electronic movements during

ultrafast processes will be closer to broad applications for many light sensitive

crystals of great biological and medical significance.

Code availability

Computer software dynamiX™ is available from Renz Research, Inc.

Acknowledgements

The following database and software are used in this work: CCP4 (ccp4.ac.uk),

Coot (www2.mrc-lmb.cam.ac.uk/Personal/pemsley/coot), dynamiX™ (Renz

Research, Inc.), gnuplot (gnuplot.info), Mathematica (wolfram.com), PDB

(rcsb.org), PHENIX (phenix-online.org), PyMOL (pymol.org), Python

(python.org), and SciPy (scipy.org).

Competing interests

ZR is the founder of Renz Research, Inc. that currently holds the copyright of the

computer software dynamiX™.

References

Abe, M., Kitagawa, T., and Kyogoku, Y. (1978). Resonance Raman spectra of
octaethylporphyrinato-Ni(II) and meso-deuterated and 15N substituted derivatives. II. A normal
coordinate analysis. J. Chem. Phys. *69*, 4526–4534.

Aranda, R., Levin, E.J., Schotte, F., Anfinrud, P.A., and Phillips, G.N. (2006). Time-dependent
atomic coordinates for the dissociation of carbon monoxide from myoglobin. Acta Crystallogr. D
Biol. Crystallogr. *62*, 776–783.

Armstrong, M.R., Ogilvie, J.P., Cowan, M.L., Nagy, A.M., and Miller, R.J.D. (2003). Observation
of the cascaded atomic-to-global length scales driving protein motion. Proc. Natl. Acad. Sci. *100*,
4990–4994.

Baldwin, J., and Chothia, C. (1979). Haemoglobin: the structural changes related to ligand
binding and its allosteric mechanism. J. Mol. Biol. *129*, 175–200.

Bandara, S., Ren, Z., Lu, L., Xiaoli, Z., Shin, H., Zhao, K.-H., and Yang, X. (2017). Photoactivation
mechanism of a carotenoid-based photoreceptor. Proc. Natl. Acad. Sci. U. S. A. *114*, 6286–6291.

Barends, T.R.M., Foucar, L., Ardevol, A., Nass, K., Aquila, A., Botha, S., Doak, R.B., Falahati, K.,
Hartmann, E., Hilpert, M., et al. (2015). Direct observation of ultrafast collective motions in CO
myoglobin upon ligand dissociation. Science *350*, 445–450.

Bionta, M.R., Lemke, H.T., Cryan, J.P., Glownia, J.M., Bostedt, C., Cammarata, M., Castagna, J.-C.,
Ding, Y., Fritz, D.M., Fry, A.R., et al. (2011). Spectral encoding of x-ray/optical relative delay.
Opt. Express *19*, 21855.

Bionta, M.R., Hartmann, N., Weaver, M., French, D., Nicholson, D.J., Cryan, J.P., Glownia, J.M.,
Baker, K., Bostedt, C., Chollet, M., et al. (2014). Spectral encoding method for measuring the
relative arrival time between x-ray/optical pulses. Rev. Sci. Instrum. *85*, 083116.

Bourgeois, D., Vallone, B., Schotte, F., Arcovito, A., Miele, A.E., Sciara, G., Wulff, M., Anfinrud,
P., and Brunori, M. (2003). Complex landscape of protein structural dynamics unveiled by
nanosecond Laue crystallography. Proc. Natl. Acad. Sci. *100*, 8704–8709.

Bourgeois, D., Vallone, B., Arcovito, A., Sciara, G., Schotte, F., Anfinrud, P.A., and Brunori, M.
(2006). Extended subnanosecond structural dynamics of myoglobin revealed by Laue
crystallography. Proc. Natl. Acad. Sci. *103*, 4924–4929.

Bourgeois, D., Schotte, F., Brunori, M., and Vallone, B. (2007). Time-resolved methods in
biophysics. 6. Time-resolved Laue crystallography as a tool to investigate photo-activated protein
dynamics. Photochem. Photobiol. Sci. *6*, 1047.

Brinkmann, L.U.L., and Hub, J.S. (2016). Ultrafast anisotropic protein quake propagation after CO
photodissociation in myoglobin. Proc. Natl. Acad. Sci. *113*, 10565–10570.

Chaika, A.N. (2014). Visualization of electron orbitals in scanning tunneling microscopy. JETP
Lett. *99*, 731–741.

Champion, P.M. (2005). Chemistry: Following the flow of energy in biomolecules. Science *310*,
980–982.

Chu, K., Vojtchovský, J., McMahon, B.H., Sweet, R.M., Berendzen, J., and Schlichting, I. (2000).
Structure of a ligand-binding intermediate in wild-type carbonmonoxy myoglobin. Nature *403*,
921–923.

Clifton, I.J., Duke, E.M.H., Wakatsuki, S., and Ren, Z. (1997). Evaluation of Laue diffraction
patterns. Meth Enzym. *277*, 448–467.

Coppens, P. (1998). Charge-Density Analysis at the Turn of the Century. Acta Crystallogr. A *54*,
779–788.

Coquelle, N., Sliwa, M., Woodhouse, J., Schirò, G., Adam, V., Aquila, A., Barends, T.R.M., Boutet,
S., Byrdin, M., Carbajo, S., et al. (2017). Chromophore twisting in the excited state of a
photoswitchable fluorescent protein captured by time-resolved serial femtosecond
crystallography. Nat. Chem. *10*, 31–37.

Dunietz, B.D., Dreuw, A., and Head-Gordon, M. (2003). Initial steps of the photodissociation of
the CO ligated heme group. J. Phys. Chem. B *107*, 5623–5629.

Falahati, K., Tamura, H., Burghardt, I., and Huix-Rotllant, M. (2017). Quantum dynamics of
ultrafast heme-carbon monoxide photolysis and spin-crossover. ArXiv170903402 Cond-Mat
Physicsphysics.

Falahati, K., Tamura, H., Burghardt, I., and Huix-Rotllant, M. (2018). Ultrafast carbon monoxide
photolysis and heme spin-crossover in myoglobin via nonadiabatic quantum dynamics. Nat.
Commun. *9*, 4502.

Franzen, S., Bohn, B., Poyart, C., DePillis, G., Boxer, S.G., and Martin, J.-L. (1995). Functional
aspects of ultra-rapid heme doming in hemoglobin, myoglobin, and the myoglobin mutant
H93G. J. Biol. Chem. *270*, 1718–1720.

Franzen, S., Kiger, L., Poyart, C., and Martin, J.-L. (2001). Heme photolysis occurs by ultrafast
excited state metal-to-ring charge transfer. Biophys. J. *80*, 2372–2385.

Fujisaki, H., and Straub, J.E. (2005). Vibrational energy relaxation in proteins. Proc. Natl. Acad.
Sci. *102*, 6726–6731.

Gruia, F., Kubo, M., Ye, X., and Champion, P.M. (2008). Investigations of vibrational coherence in
the low-frequency region of ferric heme proteins. Biophys. J. *94*, 2252–2268.

Haldrup, K., Harlang, T., Christensen, M., Dohn, A., van Driel, T.B., Kjær, K.S., Harrit, N.,
Vibenholt, J., Guerin, L., Wulff, M., et al. (2011). Bond shortening (1.4 Å) in the singlet and triplet
excited states of [Ir2(dimen)4]2+ in solution determined by time-resolved X-ray scattering. Inorg
Chem *50*, 9329–9336.

Henderson, R., and Moffat, J.K. (1971). The difference Fourier technique in protein
crystallography: errors and their treatment. Acta Cryst *B27*, 1414–1420.

Henry, E.R., and Hofrichter, J. (1992). Singular value decomposition: Application to analysis of
experimental data. In Numerical Computer Methods, (Academic Press), pp. 129–192.

Henry, E.R., Levitt, M., and Eaton, W.A. (1985). Molecular dynamics simulation of
photodissociation of carbon monoxide from hemoglobin. Proc. Natl. Acad. Sci. U. S. A. *82*, 2034–
2038.

Hirano, Y., Takeda, K., and Miki, K. (2016). Charge-density analysis of an iron–sulfur protein at
an ultra-high resolution of 0.48 Å. Nature *534*, 281–284.

Ho, J.X., Ruble, J.R., McInnis, T.R., Carter, D.C., Huang, H., and Gibson, W.M. (2002).
Convergent-beam method in macromolecular crystallography. Acta Crystallogr. D Biol.
Crystallogr. *58*, 2087–2095.

Hodel, A., Kim, S.H., and Brünger, A.T. (1992). Model bias in macromolecular crystal structures.
Acta Crystallogr. A *48*, 851–858.

Humphreys, C.J. (1999). Electrons seen in orbit. Nature *401*, 21–22.

Itatani, J., Levesque, J., Zeidler, D., Niikura, H., Pépin, H., Kieffer, J.C., Corkum, P.B., and
Villeneuve, D.M. (2004). Tomographic imaging of molecular orbitals. Nature *432*, 867–871.

Jung, Y.O., Lee, J.H., Kim, J., Schmidt, M., Moffat, K., Šrajer, V., and Ihee, H. (2013). Volume-
conserving trans–cis isomerization pathways in photoactive yellow protein visualized by
picosecond X-ray crystallography. Nat. Chem. *5*, 212–220.

Kachalova, G.S., Popov, A.N., and Bartunik, H.D. (1999). A steric mechanism for inhibition of CO
binding to heme proteins. Science *284*, 473–476.

Karunakaran, V., Benabbas, A., Sun, Y., Zhang, Z., Singh, S., Banerjee, R., and Champion, P.M.
(2010). Investigations of low-frequency vibrational dynamics and ligand binding kinetics of
cystathionine β-synthase. J. Phys. Chem. B *114*, 3294–3306.

Karunakaran, V., Sun, Y., Benabbas, A., and Champion, P.M. (2014). Investigations of the low
frequency modes of ferric cytochrome c using vibrational coherence spectroscopy. J. Phys. Chem.
B *118*, 6062–6070.

Kim, T.H., Mehrabi, P., Ren, Z., Sljoka, A., Ing, C., Bezginov, A., Ye, L., Pomès, R., Prosser, R.S.,
and Pai, E.F. (2017). The role of dimer asymmetry and protomer dynamics in enzyme catalysis.
Science *355*, eaag2355.

Kitagawa, T., Haruta, N., and Mizutani, Y. (2002). Time-resolved resonance Raman study on
ultrafast structural relaxation and vibrational cooling of photodissociated carbonmonoxy
myoglobin. Biopolymers *67*, 207–213.

Kuczera, K., Lambry, J.-C., Martin, J.-L., and Karplus, M. (1993). Nonexponential relaxation after
ligand dissociation from myoglobin: A molecular dynamics simulation. Proc. Natl. Acad. Sci. U.
S. A. *90*, 5805–5807.

Levantino, M., Yorke, B.A., Monteiro, D.C., Cammarata, M., and Pearson, A.R. (2015a). Using
synchrotrons and XFELs for time-resolved X-ray crystallography and solution scattering
experiments on biomolecules. Curr. Opin. Struct. Biol. *35*, 41–48.

Levantino, M., Schirò, G., Lemke, H.T., Cottone, G., Glownia, J.M., Zhu, D., Chollet, M., Ihee, H.,
Cupane, A., and Cammarata, M. (2015b). Ultrafast myoglobin structural dynamics observed with
an X-ray free-electron laser. Nat. Commun. *6*, 6772.

Mikhailovskij, I.M., Sadanov, E.V., Mazilova, T.I., Ksenofontov, V.A., and Velicodnaja, O.A.
(2009). Imaging the atomic orbitals of carbon atomic chains with field-emission electron
microscopy. Phys. Rev. B *80*.

Mizutani, Y. (1997). Direct observation of cooling of heme upon photodissociation of
carbonmonoxy myoglobin. Science *278*, 443–446.

Mulder, P. (2011). Are orbitals observable? Int. J. Philos. Chem. *17*, 24–35.

Nagy, A.M., Raicu, V., and Miller, R.J.D. (2005). Nonlinear optical studies of heme protein
dynamics: Implications for proteins as hybrid states of matter. Biochim. Biophys. Acta BBA -
Proteins Proteomics *1749*, 148–172.

Nango, E., Royant, A., Kubo, M., Nakane, T., Wickstrand, C., Kimura, T., Tanaka, T., Tono, K.,
Song, C., Tanaka, R., et al. (2016). A three-dimensional movie of structural changes in
bacteriorhodopsin. Science *354*, 1552–1557.

Nogly, P., Weinert, T., James, D., Carbajo, S., Ozerov, D., Furrer, A., Gashi, D., Borin, V.,
Skopintsev, P., Jaeger, K., et al. (2018). Retinal isomerization in bacteriorhodopsin captured by a
femtosecond x-ray laser. Science eaat0094.

Ogilvie, J.F. (2011). Is a molecular orbital measurable by means of tomographic imaging? Found.
Chem. *13*, 87–91.

Pande, K., Hutchison, C.D.M., Groenhof, G., Aquila, A., Robinson, J.S., Tenboer, J., Basu, S.,
Boutet, S., DePonte, D.P., Liang, M., et al. (2016). Femtosecond structural dynamics drives the
trans/cis isomerization in photoactive yellow protein. Science *352*, 725–729.

Patera, L.L., Queck, F., Scheuerer, P., and Repp, J. (2019). Mapping orbital changes upon electron
transfer with tunnelling microscopy on insulators. Nature *566*, 245–248.

Puschnig, P., Berkebile, S., Fleming, A.J., Koller, G., Emtsev, K., Seyller, T., Riley, J.D., Ambrosch-
Draxl, C., Netzer, F.P., and Ramsey, M.G. (2009). Reconstruction of molecular orbital densities
from photoemission data. Science *326*, 702–706.

Ren, Z. (2013a). Reaction trajectory revealed by a joint analysis of Protein Data Bank. PLoS ONE
8, e77141.

Ren, Z. (2013b). Reverse engineering the cooperative machinery of human hemoglobin. PLoS
ONE *8*, e77363.

Ren, Z. (2016). Molecular events during translocation and proofreading extracted from 200 static
structures of DNA polymerase. Nucleic Acids Res. *6*, 1–13.

Ren, Z., and Yang, X. (2016). Angular-split/temporal-delay approach to ultrafast protein
dynamics at XFELs. Acta Crystallogr. Sect. Struct. Biol. *72*, 871–882.

Ren, Z., Bourgeois, D., Helliwell, J.R., Moffat, K., Srajer, V., and Stoddard, B.L. (1999). Laue
crystallography: coming of age. J Synchrotron Rad *6*, 891–917.

Ren, Z., Perman, B., Srajer, V., Teng, T.-Y., Pradervand, C., Bourgeois, D., Schotte, F., Ursby, T.,
Kort, R., Wulff, M., et al. (2001). A molecular movie at 1.8 Å resolution displays the photocycle of
photoactive yellow protein, a eubacterial blue-light receptor, from nanoseconds to seconds.
Biochemistry *40*, 13788–13801.

Ren, Z., Šrajer, V., Knapp, J.E., and Royer, W.E. (2012). Cooperative macromolecular device
revealed by meta-analysis of static and time-resolved structures. Proc. Natl. Acad. Sci. U. S. A.
109, 107–112.

Ren, Z., Chan, P.W.Y., Moffat, K., Pai, E.F., Royer, W.E., Šrajer, V., and Yang, X. (2013). Resolution
of structural heterogeneity in dynamic crystallography. Acta Cryst *D69*, 946–959.

Ren, Z., Ayhan, M., Bandara, S., Bowatte, K., Kumarapperuma, I., Gunawardana, S., Shin, H.,
Wang, C., Zeng, X., and Yang, X. (2018). Crystal-on-crystal chips for in situ serial diffraction at
room temperature. Lab. Chip 2246–2256.

Repp, J., Meyer, G., Stojković, S.M., Gourdon, A., and Joachim, C. (2005). Molecules on insulating
films: Scanning-tuneling microscopy imaging of individual molecular orbitals. Phys. Rev. Lett.
94.

Rosca, F., Kumar, A.T.N., Ionascu, D., Ye, X., Demidov, A.A., Sjodin, T., Wharton, D., Barrick, D.,
Sligar, S.G., Yonetani, T., et al. (2002). Investigations of anharmonic low-frequency oscillations in
heme proteins. J. Phys. Chem. A *106*, 3540–3552.

Scerri, E.R. (2000). Have orbitals really been observed? J. Chem. Educ. *77*, 1492.

Schmidt, M., Rajagopal, S., Ren, Z., and Moffat, K. (2003). Application of singular value
decomposition to the analysis of time-resolved macromolecular X-ray data. Biophys. J. *84*, 2112–
2129.

Schotte, F., Lim, M., Jackson, T.A., Smirnov, A.V., Soman, J., Olson, J.S., Phillips, G.N., Wulff, M.,
and Anfinrud, P.A. (2003). Watching a protein as it functions with 150-ps time-resolved X-ray
crystallography. Science *300*, 1944–1947.

Schotte, F., Cho, H.S., Kaila, V.R.I., Kamikubo, H., Dashdorj, N., Henry, E.R., Graber, T.J.,
Henning, R., Wulff, M., Hummer, G., et al. (2012). Watching a signaling protein function in real
time via 100-ps time-resolved Laue crystallography. Proc. Natl. Acad. Sci. *109*, 19256–19261.

Schwarz, W.H.E. (2006). Measuring Orbitals: Provocation or Reality? Angew. Chem. Int. Ed. *45*,
1508–1517.

Spence, J.C.H., Zatsepin, N.A., and Li, C. (2014). Coherent convergent-beam time-resolved X-ray
diffraction. Philos. Trans. R. Soc. B Biol. Sci. *369*, 20130325–20130325.

Šrajer, V., Teng, T. -y., Ursby, T., Pradervand, C., Ren, Z., Adachi, S. -i., Schildkamp, W.,
Bourgeois, D., Wulff, M., and Moffat, K. (1996). Photolysis of the carbon monoxide complex of
myoglobin: nanosecond time-resolved crystallography. Science *274*, 1726–1729.

Šrajer, V., Ren, Z., Teng, T.-Y., Schmidt, M., Ursby, T., Bourgeois, D., Pradervand, C., Schildkamp,
W., Wulff, M., and Moffat, K. (2001). Protein conformational relaxation and ligand migration in
myoglobin: A nanosecond to millisecond molecular movie from time-resolved Laue X-ray
diffraction. Biochemistry *40*, 13802–13815.

Stapelfeldt, H. (2004). Electrons frozen in motion. Nature *432*, 809–810.

Stodolna, A.S., Rouzée, A., Lépine, F., Cohen, S., Robicheaux, F., Gijsbertsen, A., Jungmann, J.H.,
Bordas, C., and Vrakking, M.J.J. (2013). Hydrogen atoms under magnification: Direct observation
of the nodal structure of Stark states. Phys. Rev. Lett. *110*.

Tenboer, J., Basu, S., Zatsepin, N., Pande, K., Milathianaki, D., Frank, M., Hunter, M., Boutet, S.,
Williams, G.J., Koglin, J.E., et al. (2014). Time-resolved serial crystallography captures high-
resolution intermediates of photoactive yellow protein. Science *346*, 1242–1246.

Ursby, T., and Bourgeois, D. (1997). Improved estimation of structure-factor difference
amplitudes from poorly accurate data. Acta Crystallogr. A *53*, 564–575.

Vozzi, C., Negro, M., Calegari, F., Sansone, G., Nisoli, M., De Silvestri, S., and Stagira, S. (2011).
Generalized molecular orbital tomography. Nat. Phys. *7*, 822–826.

Waber, J.T., and Cromer, D.T. (1965). Orbital radii of atoms and ions. J. Chem. Phys. *42*, 4116–
4123.

Yang, X., Ren, Z., Kuk, J., and Moffat, K. (2011). Temperature-scan cryocrystallography reveals
reaction intermediates in bacteriophytochrome. Nature *479*, 428–432.

Zeng, X., Ren, Z., Wu, Q., Fan, J., Peng, P.-P., Tang, K., Zhang, R., Zhao, K.-H., and Yang, X.
(2015). Dynamic crystallography reveals early signaling events in ultraviolet photoreceptor
UVR8. Nat Plants *1*, 14006.

Zuo, J.M., Kim, M., O'Keeffe, M., and Spence, J.C.H. (1999). Direct observation of d-orbital holes
and Cu–Cu bonding in Cu2O. Nature *401*, 49–52.

1011

1012 **Figures and Legends**

1013

1014

1015

1016 Figure 1. A Gantt chart summarizing structural events during photodissociation

1017 of MbCO. Previously known structural events are in pink. The excitation laser

1018 pulse and XFEL pulses are in yellow. The newly observed structural events from

1019 decomposition of the serial crystallographic datasets are in dark blue.

1020

1021

1022

Figure 2. SVD of 93 difference maps around the heme site. These difference

maps are divided into several series. Each series consists of the difference maps

produced from subtracting the same reference (Methods and Table S1). Each of

the 93 difference maps can be expressed as a linear combination of many

components $c_1 U_1 + c_2 U_2 + \ldots$ after the decomposition. Among a common set of

basis components U_k ($k = 1, 2, \ldots$) shared by all difference maps, also known as

left singular vectors in linear algebra, the first few components are the most

significant. Each of the difference maps requires a coefficient set c_k for the linear

combination. This coefficient set varies from one map to the other, that is, a

function of the delay time or other metadata associate with the datasets. Each

coefficient c_k indicates how much of the corresponding component U_k exists in

the linear combination. Each coefficient set is derived from a right singular

vector as traditionally known in linear algebra (Methods). These coefficients

produced by SVD should not be confused with the Fourier coefficients used to

synthesize each map. (a-d) Orthographic projections of coefficients in some of

the significant dimensions. Each difference map is represented by a dot at the

location that marks the coefficients associated with the major components U_1

through U_4. Difference maps in a series are connected by lines in a

corresponding color. The multi-dimensional space is projected to various planes

and tiled together so that any two adjacent panels from (a) to (d), horizontal or

vertical, can be imagined erecting a three-dimensional subspace. (e) Singular

values indicating the significance of the components are sorted in a descending

order of their values before a rotation. A multi-dimensional rotation alters their

values. (f) Coefficients c_6 and c_8 of the $-F_{0.1}$ series in yellow are least-squares

fitted with a two-dimensional oscillation (Methods).

1048

U_4 of difference maps at heme site

1050

Figure 3. Ultrafast formation of high-spin 3*d* orbitals of the heme iron. Each

component resulting from SVD, that is, a left singular vector, can also be

presented as a difference map. The fourth component U_4 of the difference maps

captures a cubic shaped positive density network. It is rendered as a difference

map in two styles, two-dimensional cross sections and three-dimensional

contours. The corresponding coefficient c_4 is plotted in Fig. 2cd. (a-h) Images of

two-dimensional cross sections through the component difference map U_4. Cross

sections defined in Fig. S1 are labeled in the top right corner of each panel. Non-

centric cross sections are marked to indicate small offsets from the centric cross

sections. The structure of the heme, CO ligand in both bound and docked

positions, and some side chains are also projected onto the images. C, N, O, and

Fe atoms are in black, blue, red, and rusty dots, respectively. The imidazole

rings of the distal His64 and proximal His93 are approximately coplanar with X

and AC cross sections, respectively. Two propionic acid side chains of AD rings

extend to the left in (bcd). But they are not coplanar with these cross sections. (i-

k) Three-dimensional mesh contours superimposed over a structure are

traditionally rendered to display electron density maps. Green and red meshes

are positive and negative contours at $\pm 7\sigma$, respectively. (l and m) Theoretical

contours of probability distributions of electrons in $3d$ orbitals of iron. The

combined probability distributions of $3d_{xz}$ and $3d_{yz}$ with $3d_{xy}$ (l) or with $3d_{x^2-y^2}$

(m) are both in the shape of a cube with the largest probability at the eight

corners. A 1/16 cutaway in each contour shows the cross sections. This

characteristic shape of electron distribution is observed in U_4 of the difference

maps.

1075

Figure 4. SVD of 20 simulated annealing omit maps (SAOMs) around the heme

site. The first component is the average of all omit maps (Fig. S5). (a-e) Each

omit map is represented by a dot at the location that marks the coefficients

associated with the major components. SAOMs derived from the structures at

cryo and ambient temperatures are in red and blue, respectively. SAOMs from

the XFEL datasets are in green. The multi-dimensional space is projected to

various planes and tiled together so that any two adjacent panels from (a) to (e),

horizontal or vertical, can be imagined erecting a three-dimensional subspace.

(b) Coefficients c_5 and c_8 show a damped two-dimensional oscillation as marked

by the curved arrow. (f) Singular values indicating the significance of the

components are sorted in a descending order of their values before a multi-

dimensional rotation.

1090

U_7 of SAOMs at heme site

1092

Figure 5. The seventh component of SAOMs correlated with photolysis at all

temperatures. The coefficient corresponding to U_7 of SAOMs increases as the

reaction of ligand dissociation proceeds from MbCO to deoxy Mb in static, cryo-

trapping, and time-resolved XFEL experiments (Fig. 4e). The hypothetical

transition from MbCO to deoxy Mb at 0 K is shown in Movie S3. (a-h) Cross

sections through the seventh component U_7. A cubic network of positive

densities is marked 1-4 in (c). (i-k) Three-dimensional meshes of this component

in green and red are contoured at $\pm15\sigma$, respectively. See Fig. 3 legend for more

detail.

1102

U_3 of SAOMs at heme site

1104

1105 Figure 6. Thermal vibrations isolated from the chemical signals of photolysis.

1106 The third component of the SAOMs U_3 is highly correlated with the sample

1107 temperature (Fig. 4ace). A linear extrapolation based on the datasets available at

1108 cryo and ambient temperatures predicts that the instantaneous local temperature

1109 jump of the heme may have exceeded 500 K in the XFEL datasets (i). (a and b)

1110 Linear combinations of $900U_1$ - $586U_3$ and $900U_1$ + $476U_3$ reconstitute the omit

1111 maps of the heme at 0 and 600 K, respectively, excluding all chemical signals

1112 related to photolysis, where the coefficient c_1 = 900 of U_1 is nearly constant for all

1113 omit maps. The transition from 0 to 600 K free of the chemical signals from

1114 photolysis is shown in Movie S2. (c-h) Cross sections through the third

1115 component U_3. Positive densities in 20 pillars connecting two sheets of positive

1116 densities above and below the heme in the cross sections of Z±1.3 Å are marked

1117 in (e). (j-l) Three-dimensional meshes of this component in green and red are

1118 contoured at ±15σ, respectively. See Fig. 3 legend for more detail.

1119

a

b

His93

$|U_5|+|U_6|+|U_8|$ of SAOMs at heme site

c

d

e

Fitted wavenumber = 36.0 cm^{-1}; period = 0.93 ps; damping time constant = 1.5 ps

f

g

h

Reference (ps)
dark
−0.1
0
0.1
0.2
0.3
0.4
0.5
0.6

Reference: dark −0.1 0 0.1 0.2 0.3 0.4 0.5 0.6

i Helix G Helix H U_1

j Helix E U_2

k Helix E U_3

l Helix G Helix H U_4

U_1 through U_4 of difference maps in the entire globin

Figure 7. Low frequency oscillations in the solvent. (a) SVD analysis of the

difference maps in the entire globin produces the first and third components

circularly oscillating before 1 ps and quickly diminishing during the first few ps.

Similar oscillations are also observed only around the heme site in both the

difference maps (Fig. 2b) and the omit maps (Fig. 4b). (b-e) Cross sections

through an averaged map of both the positive and negative halves of U_5, U_6, and

U_8 of the SAOMs. Low densities in white and light green are located consistently

on the heme and protein, which suggests that the oscillating signals are more

associated with the solvent rather than the globin. (f) The circularly oscillating

coefficients c_1 and c_3 are plotted as function of delay time in many series. The

offset sinusoidal curves indicate a circular oscillation. Each map series is derived

from subtracting a different reference dataset as indicated by the color bar below.

(g and h) c_1 and c_3 of the $-F_{dark}$ series are least-squares fitted with a two-

dimensional damped oscillation (Methods). (i-l) Comparison of several

components of difference maps in the entire globin. U_1 and U_3 contain

widespread signals, as they are ranked the most significant components. But no

signal is concentrated on the heme or the protein. Instead, U_2 and U_4 clearly

contain strong signals associated with the heme and secondary structures of the

protein.

1140

1141

Ultrafast Structural Changes Decomposed from Serial Crystallographic Data

Zhong Ren

Department of Chemistry, University of Illinois at Chicago, Chicago, IL 60607, USA

Renz Research, Inc., Westmont, IL 60559, USA

zren@uic.edu

Methods

Difference Fourier maps

Difference Fourier maps are commonly used in protein crystallography. A difference Fourier map is synthesized from a Fourier coefficient set of F_{light}-$F_{reference}$ with the best available phase set. Before Fourier synthesis, F_{light} and $F_{reference}$ must be properly scaled to the same level so that the distribution of difference values is centered at zero and not skewed either way. All available datasets from Barends & Schlichting et al. (2015) are carefully scaled and difference Fourier maps are calculated using a software package dynamiX™ according to previously published protocols including a weighting scheme under the assumption that a greater amplitude of a difference Fourier coefficient F_{light}-$F_{reference}$ is more likely caused by noise than by signal (Ren et al., 2001, 2013; Šrajer et al., 2001; Ursby and Bourgeois, 1997).

Both the dark and light datasets can serve as a reference in difference maps. If a light dataset at a certain delay is chosen as a reference, the difference maps show the changes since that delay time but not the changes prior to that delay. These 91 maps in a full matrix provide independent observations for all changes (Table S1). This strategy is important to the subsequent singular value decomposition (SVD) analysis. In addition, two more difference maps of the photolyzed Mb*CO and **CO (1dws/t) are also included in the -F_{dark} series for SVD analysis.

1175

Each difference map is masked around the heme site. All densities at those

grid points are set to zero, if they are greater than 5 Å away from the heme, the

bound or docked CO, the side chain of the proximal His93, and that of the distal

His64. Difference maps around the heme site include the most significant signals

and exclude noises in the protein and solvent. However, the unmasked

difference maps in the entire globin are also used a separate SVD analysis to

include the protein and solvent regions for observation of global changes (Fig. 7i-

l).

1184

Simulated annealing omit maps

The heme, the ligand, the side chains of proximal His93 and distal His64 are

removed from all structures listed in Table S2. The remaining structures are

refined using simulated annealing method with phenix.refine starting from a

simulated temperature of 5,000 K. Fourier coefficients are obtained from the

differences between the structure factor amplitudes $F_{observed}$ observed in the time-

resolved experiments and the structure factor amplitudes $F_{calculated}$ calculated from

the omitted structures. The resulting $F_{observed}$-$F_{calculated}$ maps, known as simulated

annealing omit maps (SAOMs), reveal the unbiased electron densities in the

omitted region. Since the sperm whale (sw) Mb structures are in a different unit

cell compared to the horse heart (hh) Mb structures, the SAOMs are aligned

together according to a least-squares fitting of the main chain atoms in residues

2-151. The heme and ligand are not used in the alignment procedure. Therefore,

the motion of heme found in these omit maps is with respect to the globin

framework (Fig. 5 and Movie S3). 20 aligned omit maps are subjected to the

subsequent SVD analysis.

1201

1202   *Singular value decomposition of (difference) electron density maps*

1203   The principle of SVD can be found in texts on linear algebra. SVD analysis of

1204   (difference) electron density maps was previously described (Ren et al., 2013;

1205   Schmidt et al., 2003). Here I briefly recap this numerical procedure with an

1206   emphasis on the physical meaning of the outcome.

1207

1208  An electron density map, sometimes a difference map, consists of density

1209 values on a collection of grid points within a mask (see above), also known as

1210 voxels that are equivalent to pixels in a digitized image. All M voxels in a three-

1211 dimensional map can be serialized into a one-dimensional sequence of density

1212 values according to a specific protocol. It is not important what the protocol is as

1213 long as a consistent protocol is used to serialize all maps of the same grid setting

1214 and size, and a reverse protocol is available to erect a three-dimensional map

1215 from a sequence of M densities. Therefore, a set of N serialized maps, also

1216 known as vectors in linear algebra, can fill the columns of a data matrix $\mathbf{A}$ with

1217 no specific order so that the width of $\mathbf{A}$ is N columns and the length is M rows.

1218 Often, $M \gg N$, thus $\mathbf{A}$ is an elongated matrix. In this work, $N = 93$ for the

1219 difference maps and $N = 20$ for the omit maps. If a consistent protocol of

1220 serialization is used, the corresponding voxel in all N maps occupies a single row

1221 of matrix $\mathbf{A}$. For example, the voxels at the heme iron in all maps could occupy

1222 the jth row of matrix $\mathbf{A}$. This strict correspondence in a row of matrix $\mathbf{A}$ is

1223 important. Changes of the density values in a row are due to either signals,

1224 systematic errors, or noises.

1225

1226  SVD of the data matrix $\mathbf{A}$ results in $\mathbf{A} = \mathbf{UWV}^{\mathrm{T}}$, also known as matrix

1227 factorization. Matrix $\mathbf{U}$ has the same shape as $\mathbf{A}$, that is, N columns and M rows.

1228 The N columns contain decomposed basis components $\boldsymbol{U}_k$, known as left singular

1229 vectors of length M, where $k = 1, 2, \ldots, N$. Therefore, each component $\boldsymbol{U}_k$ can be

1230 erected using the reverse protocol to form a three-dimensional map. This

1231 decomposed elemental map can be presented in the same way as the original

1232 maps, for example, rendered in molecular graphics software such as Coot and

1233 PyMol. The second matrix $\mathbf{W}$ is a square matrix that contains all zeros except for

1234 N positive values on its major diagonal, known as singular values w_k. The

1235 magnitude of w_k is considered as a weight or significance of its corresponding

1236 component $\boldsymbol{U}_k$. The third matrix $\mathbf{V}$ is also a square matrix of $N \times N$. Each column

1237 of $\mathbf{V}$ or row of its transpose $\mathbf{V}^{\mathrm{T}}$, known as a right singular vector, contains the

1238 relative compositions of $\boldsymbol{U}_k$. That is to say, the N values in a column V_k describe

1239 the relative compositions of the component $\boldsymbol{U}_k$ in the N original maps. A singular

1240 triplet denotes 1) a decomposed component $\boldsymbol{U}_k$, 2) its singular value w_k, and 3) the

1241 composition function V_k. Singular triplets are often sorted in a descending order

of their singular values w_k. Only a small number of n significant singular triplets

identified by the greatest singular values w_1 through w_n can be used in a linear

combination to reconstitute a set of composite maps that closely resemble the

original ones in matrix $\mathbf{A}$, where $n < N$. For example, the original map in the ith

column of matrix $\mathbf{A}$ can be closely represented by the ith composite map $w_1 v_{1i} \boldsymbol{U}_1 +$

$w_2 v_{2i} \boldsymbol{U}_2 + \dots + w_n v_{ni} \boldsymbol{U}_n$, where $(v_{1i}, v_{2i}, \dots)$ is the ith row of matrix $\mathbf{V}$. $c_k = w_k v_{ki}$ in

front of $\boldsymbol{U}_k$ is the coefficients for the linear combination. Excluding the

components after $\boldsymbol{U}_n$ in this linear combination is based on an assumption that

the singular values after w_n are very small relative to those from w_1 through w_n.

As a result, the structural information evenly distributed in all N original maps is

effectively concentrated into a far fewer number of n significant components,

known as information concentration or dimension reduction. In this study, the

difference maps seem to contain 15 major components; the top nine are the most

significant (Fig. 2e). The components 10 through 15 may very likely harbor less

and less signals or systematic differences. For example, Fourier truncation

errors, if significant enough, could occupy one or more of these components.

However, the components 10 through 15 have not been examined in detail.

1259

On the other hand, the trailing components in matrix $\mathbf{U}$ contain inconsistent

fluctuations and random noises. Excluding these components effectively rejects

noises. However, no clear boundary is guaranteed between signals, systematic

errors, and noises. Systematic errors could be more significant than the desired

signals, for example, the second and fourth components $\boldsymbol{U}_2$ and $\boldsymbol{U}_4$ of the omit

maps describe significant differences between hhMb and swMb, a valid scientific

point to investigate but outside the scope of this paper.

1267

The orthonormal property of SVD

The solution set of SVD must guarantee that the columns in $\mathbf{U}$ and $\mathbf{V}$, the left and

right singular vectors $\boldsymbol{U}_k$ and $\boldsymbol{V}_k$, are orthonormal, that is, $\boldsymbol{U}_h \bullet \boldsymbol{U}_k = \boldsymbol{V}_h \bullet \boldsymbol{V}_k = 0$

(ortho) and $\boldsymbol{U}_k \bullet \boldsymbol{U}_k = \boldsymbol{V}_k \bullet \boldsymbol{V}_k = 1$ (normal), where $h \neq k$ but both are from 1 to N. The

orthonormal property also holds for the row vectors. As a result, each

component $\boldsymbol{U}_k$ is independent of the other components. In other words, a

component cannot be represented by a linear combination of any other

components. For example, the symmetrical, spherical, and thermal component

U_3 (Fig. 6) and the asymmetrical, cubic component of U_7 (Fig. 5) of the omit maps

are orthonormal. Both components show positive densities around the iron. But

the cubic positive densities in U_7 are associated with the strong negative densities

on the ligand depicting dissociation, which is a direct link from ligand

dissociation to regaining of the high-spin $3d$ orbitals of the iron. On the other

hand, no sign of ligand dissociation is found in the component U_3 that features

the spherical densities, which strongly suggests that this component describes

the thermal effect. These two components are not interchangeable despite some

similarity in the arrangement of positive densities, that is, one of the physical

meanings of the orthonormal property of SVD. However, two physical or

chemical parameters, such as temperature and pH, may cause different changes

to a structure, but not necessarily orthogonal changes (see below).

1288

Due to the orthonormal property of SVD, an N-dimensional Euclidean space

is established, and the first n dimensions define its most significant subspace.

Each coefficient set $c_k = w_k v_{ki}$ of the ith composite map is located in this subspace.

All coefficient sets for $i = 1, 2, …, N$ in different linear combinations to

approximate the N original maps in a least-squares sense can be represented by

N points in the n-dimensional Euclidean subspace. In this work, many two-

dimensional orthographical projections of this n-dimensional subspace are

presented as scatter plots with each map represented as a dot located at a

position determined by the coefficient set of this map. These scatter plots are

highly informative to reveal the relationship between the (difference) electron

density maps and their metadata. For example, the difference maps at short time

delays < 1 ps are consistently different from those at long time delays > 1 ps in a

certain aspect (Fig. 2ac) and that two major components are circularly correlated

but only before 1 ps, that is, a damped oscillation (Fig. 2b).

1303

Rotation in SVD space

Dimension reduction is indeed effective in data analysis of dynamic

crystallography when many datasets are evaluated at the same time. However,

the interpretation of a basis component U_k, that is, "what-does-it-mean", requires

a clear demonstration of the relationship between the core data, here (difference)

electron density maps, and their metadata, such as time, temperature, and other

experimental conditions. The outcome of SVD does not guarantee any physical

meaning in a basis component. Therefore, SVD provides no direct answer to

"what-does-it-mean". The factorized set of matrices $\mathbf{U}$, $\mathbf{W}$, and $\mathbf{V}$ from SVD is

not a unique solution. That is to say, they are not the only solution to factorize

matrix $\mathbf{A}$. The default solution set from SVD may not carry an obvious physical

meaning to depict "what-does-it-mean", thus its usefulness is very limited to

merely a mathematical construction. A more straightforward analogy is that one

can walk along two perpendicular streets to reach the opposite corner of a street

block as if walking a diagonal block despite the fact that a diagonal street does

not physically exist. This is also the origin of the debate on whether atomic

orbitals as a basis set of mathematical constructions are physical and observable

(Scerri, 2000). Therefore, it is very important to find one or more alternative

solution sets that are physically meaningful to elucidate a structural

interpretation. The concept of rotation after SVD was introduced by Henry &

Hofrichter (1992). But they suggested a protocol that fails to preserve the

orthonormal and least-squares properties of SVD. The rotation process, also

known as Givens rotation, is a connection between the analytical tool SVD and

scientific findings. This rotation shall not be confused with a rotation in the

three-dimensional real space, in which a molecular structure resides.

1329

1330   A rotation in the n-dimensional Euclidean subspace is necessary to change

1331 the perspective before a clear relationship emerges to elucidate scientific

1332 findings. It can be shown that two linear combinations are identical before and

1333 after a rotation of an angle θ in a two-dimensional subspace of h and k. That is,

1334

$$c_h\mathbf{U}_h + c_k\mathbf{U}_k = f_h\mathbf{R}_h + f_k\mathbf{R}_k, \tag{1}$$

1335

1336 where c_h and c_k are the coefficients of the components $\mathbf{U}_h$ and $\mathbf{U}_k$ before the

1337 rotation as defined above; and f_h and f_k are the coefficients of the rotated basis

1338 components $\mathbf{R}_h$ and $\mathbf{R}_k$, respectively. The same Givens rotation is applied to both

1339 the bases and their coefficients:

1340

$$\begin{cases} \mathbf{R}_h = \mathbf{U}_h\cos\theta - \mathbf{U}_k\sin\theta; \\ \mathbf{R}_k = \mathbf{U}_h\sin\theta + \mathbf{U}_k\cos\theta. \end{cases} \tag{2}$$

1341

1342 Obviously, the rotated components R_h and R_k remain mutually orthonormal and

1343 orthonormal to other components. And

1344

$$\begin{cases} f_h = s_h t_h = c_h\cos\theta - c_k\sin\theta = w_h v_h\cos\theta - w_k v_k\sin\theta; \\ f_k = s_k t_k = c_h\sin\theta + c_k\cos\theta = w_h v_h\sin\theta + w_k v_k\cos\theta. \end{cases} \tag{3}$$

1345

Here $s_{h|k} = \sqrt{\sum f_{h|k}^2}$ are the singular values that replace w_h and w_k, respectively,

after the rotation. They may increase or decrease compared to the original

singular values so that the descending order of the singular values no longer

holds. $T_{h|k} = (t_{h|k1}, t_{h|k2}, \ldots, t_{h|kN}) = (f_{h|k1}, f_{h|k2}, \ldots, f_{h|kN})/s_{h|k}$ are the right singular

vectors that replace V_h and V_k, respectively. T_h and T_k remain mutually

orthonormal after the rotation and orthonormal to other right singular vectors

that are not involved in the rotation.

1353

A rotation in two-dimensional subspace of h and k has no effect in other

dimensions, as the orthonormal property of SVD guarantees. Multiple Givens

rotations can be carried out in many two-dimensional subspaces consecutively to

achieve a multi-dimensional rotation. Rotation in the SVD space converts one

solution set $\mathbf{A} = \mathbf{UWV}^T$ to other alternative solutions $\mathbf{A} = \mathbf{RST}^T$ so that an

appropriate perspective can be found to elucidate the relationship between the

core data and metadata clearly and concisely. A solution derived from a rotation

retains the orthonormal property of SVD. SVD analyses presented in this paper

employ rotations extensively except that no distinction is made in the symbols of

components and coefficients before and after a rotation.

1364

For example, a large gap between all difference maps < 1 ps and those > 1 ps

indicates major differences between these two distinct time scales (Fig. 2a). A

rotation does not change this fact except that this large gap used to orient along

some inclined direction before the rotation. Therefore, the original component

U_1 cannot describe the structural differences between the short delays < 1 ps and

the longer delays > 1 ps, because the original U_1 used to be a linear combination

of several factors. The best rotation is to separate these two groups of difference

maps along a major dimension, that is, a deterministic solution. The large gap

exists as soon as an SVD analysis is performed and can be easily revealed in an

interactive three-dimensional plotting on the computer screen, although it is not
obvious in any of the orthographical projections before a proper rotation is
found. One cannot create nor eliminate this large gap by any rotation. A
different rotation may reorient this large gap along another major dimension.
But the structural conclusion would be equivalent.
1379
All difference maps have their coefficients $c_4 \approx 0$ except those calculated with
the dark dataset as a reference (red trace in Fig. 2cd). The best rotation can be
deterministically obtained by fitting all difference maps into the plane of $c_4 = 0$
except the $-F_{dark}$ series standing outside the plane. The resulting component $\boldsymbol{U}_4$
depicts features only in the $-F_{dark}$ series (Fig. 3). Again, the fact cannot be
changed with any rotation that light-dark difference maps standout from the
other difference maps arranged on a plane that is not necessarily $c_4 = 0$ before a
proper rotation. Therefore, the original $\boldsymbol{U}_4$ are hard to interpret with several
physical meanings fixed in it. In this sense, it is the rotation after the
decomposition that completes the deconvolution of physical and chemical
causes. See also an example of rotation in an SVD analysis of atomic coordinates
of protein structures (Ren, 2016).
1392
Least-squares fittings of oscillating components
Oscillating components are repeatedly found in the difference maps and omit
maps. Usually, a pair of components are oscillating with the same frequency,
which does not exclude the possibility that a third weaker component is also
oscillating at the same frequency. A pair of coefficients in a two-dimensional
damped oscillation of a period p is least-squares fitted as x and y:
1399

$$\begin{cases} x = x_0 + ae^{-t/\tau}\sin(2\pi t/p + \varphi); \\ y = y_0 + be^{-t/\tau}\sin(2\pi t/p + \psi). \end{cases} \tag{4}$$

1400
1401 x_0 and y_0 are the center of the oscillation. a and b are the amplitudes of the
1402   elliptical oscillation in two dimensions. They are damped according to an
1403   exponential decay with a time constant τ. φ and ψ are the phase angles of the
1404   oscillation in two dimensions. They are not strictly $\pi/2$ apart.
1405

1406   The sixth and eighth components of the difference maps appear to gain a

1407   two-dimensional oscillation after 1 ps (Fig. 2f). Their coefficients are least-

1408   squares fitted as x and y with an exponential onset:

1409

$$\begin{cases} x = x_0 + a\left(1 - e^{-t/\tau}\right)\cos(2\pi t/p + \varphi)\,; \\ y = y_0 + a\left(1 - e^{-t/\tau}\right)\sin(2\pi t/p + \varphi)\,. \end{cases} \tag{5}$$

1410

1411   Due to the lack of sufficient time points, the oscillation is forced to take a circular

1412   form.

1413

1414   **Supplementary Table**

Table S1. The full matrix of difference Fourier maps

Reference (ps)	Dark	Time points (ps)												
[1]Dark (5cmv)	[2].	-0.1	0	0.1	0.2	0.3	0.4	0.5	0.6	3	10	50	150	deoxy
-0.1 (5cn4)	[3]-	.	0	0.1	0.2	0.3	0.4	0.5	0.6	3	10	50	150	deoxy
0 (5cn5)	-	-	.	0.1	0.2	0.3	0.4	0.5	0.6	3	10	50	150	deoxy
0.1 (5cn6)	-	-	-	.	0.2	0.3	0.4	0.5	0.6	3	10	50	150	deoxy
0.2 (5cn7)	-	-	-	-	.	0.3	0.4	0.5	0.6	3	10	50	150	deoxy
0.3 (5cn8)	-	-	-	-	-	.	0.4	0.5	0.6	3	10	50	150	deoxy
0.4 (5cn9)	-	-	-	-	-	-	.	0.5	0.6	3	10	50	150	deoxy
0.5 (5cnb)	-	-	-	-	-	-	-	.	0.6	3	10	50	150	deoxy
0.6 (5cnc)	-	-	-	-	-	-	-	-	.	3	10	50	150	deoxy
3 (5cnd)	-	-	-	-	-	-	-	-	-	.	10	50	150	deoxy
10 (5cne)	-	-	-	-	-	-	-	-	-	-	.	50	150	deoxy
50 (5cnf)	-	-	-	-	-	-	-	-	-	-	-	.	150	deoxy
150 (5cng)	-	-	-	-	-	-	-	-	-	-	-	-	.	deoxy
Deoxy (5d5r)	-	-	-	-	-	-	-	-	-	-	-	-	-	.
PDB entry	5cmv	5cn4	5cn5	5cn6	5cn7	5cn8	5cn9	5cnb	5cnc	5cnd	5cne	5cnf	5cng	5d5r

[1]Traditionally, the dark dataset is chosen as the reference in calculation of difference Fourier maps.

[2]Dot (.) indicates that these are empty maps.

[3]Dash (-) indicates that these are negated maps.

1415

1416

Table S2. Dataset labeling		
PDB entry	Label	Description
1bz6	aquomet	Aquomet Mb from sperm whale at room temperature
1bzp	swMb	Deoxy Mb from sperm whale at room temperature
1bzr	swMbCO	MbCO from sperm whale at room temperature
1dwr	MbCO	MbCO from horse heart at 100 K
1dws	Mb*CO	Photolyzed MbCO from horse heart at 22 K
1dwt	Mb**CO	Photorelaxed Mb from horse heart at 88 K
5cmv	dark	XFEL dark dataset of MbCO from horse heart
5cn4	-0.1	
5cn5	0	
5cn6	0.1	
5cn7	0.2	
5cn8	0.3	XFEL light datasets binned by delay time < 1 ps
5cn9	0.4	
5cnb	0.5	
5cnc	0.6	
5cnd	3	
5cne	10	
5cnf	50	XFEL light datasets with delay time > 1 ps
5cng	150	
5d5r	Mb	Deoxy Mb from horse heart at 7°C

1417

1418

1419 **Supplementary Figures and Legends**

1420

1422

Figure S1. Orientations in myoglobin. Myoglobin (Mb) is a 17 kDa protein in
muscle that carries molecular oxygen O_2. The protein structure is rendered in
ribbon and surface models in (a) and (b). The non-protein prosthetic group heme
in stick model is anchored to helix F of the protein through His93 also in sticks.
Two propionic acid side chains of the heme are facing the molecular surface. The
sixth coordination site of the heme iron in sphere could bind O_2, CO, and NO
ligands also in spheres. Both a bound CO and a dissociated CO docked nearby
are shown. Traditionally, the anchor side of the heme is called proximal with
respect to the opposite distal side, where the ligands bind. The most remote part
of the protein is 30 Å away from the heme iron. The proximal and distal
directions extend 19 and 20 Å in the protein from the heme iron, respectively.
The tetrapyrrole rings are traditionally named ABCD as marked in (c). In this
paper, the heme plane is called Z cross section. Other cross sections X, Y, AC,
and BD are defined in (c).

1437

Figure S2. The difference Fourier maps for SVD analysis. Difference maps are

calculated according to the protocol described in Methods. Only the series

from -0.1 to 150 ps with the dark dataset (5cmv) as the reference is displayed

here. Cross sections are marked on top. See the legends of Figs. S1 and 3 for

definition of cross sections. See Table S1 for the full matrix of difference maps. A

visual inspection of these difference maps at different delay times may conclude

that they are largely identical. Very subtle variations could be seen by a careful

comparison of these maps in a time series. A numerical deconvolution is

necessary to extract the inherent components.

1448

1449

1450
1451
Figure S3. The most significant component U_1 of the difference maps. No doubt
the most significant event shown in the difference maps is the out-of-plane
displacement of the iron. This component shows the severely skewed iron
displacement. The corresponding coefficient c_1 is plotted in Fig. 2ac. This
component clearly separates the delays > 1 ps from those < 1 ps. (a-d) Images of
two-dimensional cross sections through the first component. (e) and (f) Three-
dimensional mesh contours of the component difference map. Green and red
meshes are positive and negative contours at $\pm 7\sigma$, respectively.

1460
1461
Figure S4. Modification to the cubic positive densities around the iron in the
sixth component U_6 of the difference maps. The corresponding coefficient c_6 is
plotted in Fig. 2f. This component and the eighth component U_8 form a two-
dimensional oscillation that ramps up in the delays > 1 ps (Fig. 2f). The effect of
modulation and oscillation is shown in Movie S1. (a-d) Images of two-
dimensional cross sections through the sixth component. (e) and (f) Three-
dimensional mesh contours of the component difference map. Green and red
meshes are positive and negative contours at $\pm 7\sigma$, respectively.

1470

1471

1472 Figure S5. The most significant component of SAOMs. Since an omit map is the

1473 least biased experimental electron density map, the most significant component

1474 of the omit maps, by far (Fig. 4f), is an averaged experimental electron density

1475 map of 20 omit maps. Only the ligand is not clear due to its variation. All other

1476 components will be added onto or subtracted from this average.

1477

1479

Movie S1. Reconstituted difference maps showing modulation and oscillation.

Linear combination of components U_1, U_4, U_6, and U_8 of the difference maps span

from 0.02 to 500 ps with some extrapolation on both ends. Other major

components carrying systematic difference from the deoxy Mb or featuring

oscillations in the solvent are not included in the linear combination to produce

this mivie. The four panels are cross sections of Z+1.1 Å, Z, Z-1.1 Å, and Y,

respectively. See Figs. S1 and 3 for the definition of cross sections. The cubic

positive densities first lean towards ring D from 0.5 to 3 ps. Then the position

returns back to the mean position after 50 ps, sways in the opposite direction

towards ring B, and ramps up after 1 ps into an oscillation of a period of 220 ps

(Fig. 2f). It is clear from this movie that the iron reaches its most out-of-plane

displacement at a few ps and recoils back during 100-200 ps and pops out again

after 200 ps.

1493

1494

1495

1496

Movie S2. Temperature effect on electron density map of the heme. The first and

third components of the SAOMs are recombined to show the temperature effect

only. The components related to the chemical signals of the photolysis are not

included. Therefore, this movie shows the pure thermal effect. Slight

extrapolation at both ends covers from 0 to 600 K. The two panels are cross

sections Z and Y, respectively. See also Fig. 6ab for electron density maps at 0

and 600 K.

1504

1505

1507

Movie S3. Comparison of electron density of the heme in ligated and unligated

deoxy states. The first, third, and seventh components of the SAOMs are

recombined to directly contrast the electron density maps of the heme in ligated

and deoxy states. The coefficients for the first and third components are constant

in the reconstituted maps at 0 K, that is to say, the thermal effect is excluded.

The coefficient c_7 for the seventh components is set to -400 for the ligated state

and +300 for the deoxy state (Fig. 4e). The panels are cross sections of Z+1.1 Å, Z,

Z-1.1 Å, and Y, respectively. The ligation state and the iron out-of-plane motion

are shown in cross section Y. Cross section Z+1.1 Å on the distal side clearly

shows greater densities of the heme in deoxy state, signaling the heme doming.

Most importantly, the heme is sliding towards the AB rings in deoxy state. This

is consistent with the skewed displacement of the iron and the motion of the

proximal His93.

1521

Ultrafast Structural Changes Decomposed from Serial Crystallographic Data

Zhong Ren

Department of Chemistry, University of Illinois at Chicago, Chicago, IL 60607, USA
Renz Research, Inc., Westmont, IL 60559, USA

zren@uic.edu

Point-by-point responses to the reviewers' comments

This manuscript has been extensively reviewed by two reviewers, both of whom evaluated the potential significance of the work and constructively suggested possible improvements. The reviewer 1 identified himself as Vladimir Y. Lunin of Russian Academy of Sciences. The reviewers' comments are copied below verbatim in *italics* followed by my response. The topics are arranged by relevance to one another. The complete comments are appended at the end.

Reviewer 1: The author challenges an ambitious task to extract tiny structural information possessing X-ray scattering data sets of a resolution only about 1.8 Å while usually this requires data of much higher resolution, atomic 1.0-1.2 Å or even ultrahigh, about 0.8 Å.

Reviewer 1: In particular, this leads to the fact that individual electrons and hydrogen atoms become distinguishable in Fourier difference syntheses only when working with data sets at resolution of 1.0-1.2 Å (atomic resolution), while partial charges (density of valence electrons) become visible only at about 0.8 Å (ultra-high resolution) and only under certain restrictions on the structure refinement protocol (Afonine et al., 2004, Acta Cryst. D60, 260-274).

Reviewer 2: C. Finally, some aspects of the central results could be presented with greater caution, considering that they are somewhat speculative, and that they depend on interpretation.
1) The author explains that the initial report of the observation of 3d orbitals (Zuo el al 1999) was somewhat controversial. In particular Scerri (2000) states that orbitals of single electrons are fundamentally not observable, and recommends that observations be described

in terms of electron density. This would be the most cautious language. However, haven't the philosophical arguments now run their course after 20 years, so it should now be safe to speak of the U4 component as revealing the high-spin orbitals? Unfortunately, the author's own description shows that the electron density is only marginally interpretable as a 3d orbital. Line 368 states that the 3d orbital for an isolated Fe(II) has 0.36 Angstrom radius, and line 369 that the orbital density in an Fe4S4 cluster has a maximum radius of 1 Angstrom. Meanwhile, the data shown here place the electron density at 1.7 to 1.9 Angstrom from the iron, depending on which map is considered, so it appears that this analysis is "first of its kind" result, rather than the observation of a 3d orbital by established methods. It is somewhat difficult to understand the author's point of view in the following text (lines 372-394). It is stated that the cubic density is not the probability distribution of a 3d electron, yet the paper states in many places that we observe a high-spin 3d orbital.

Can crystallographic data at the resolution of 1.8 Å reveal the electron density map of a single orbital? Clearly, this becomes one of the central questions to be addressed. I have consolidated previously scattered discussion into one new section in Discussion with a subtitle of "Are atomic orbitals observed?" (lines 635-710). This section should clarify my answer. A simple answer is No. However, the reasons behind this negative answer must be examined. More importantly, an accurate orbital density map of a single electron is not a goal in protein crystallography. These are my main points: First, the reason why an ultrahigh spatial resolution is required in charge-density analysis is because of the overwhelming spherical distribution of the inner electrons. Many techniques other than crystallography can capture images of atomic and molecular orbitals because these techniques are selectively sensitive to certain orbitals but not because they can achieve ultrahigh spatial resolution. If SVD is applied to charge-density analysis, the requirement for spatial resolution will reduce. Sadly, my conversation with Philip Coppens cannot continue after his pass. Second, if there is no discontinuity in any probability distribution functions of electrons, why ultrahigh frequency terms are required in Fourier synthesis? There is probably no discontinuity in the first derivatives of these distributions either, which means no kinks. Ultrahigh spatial resolution is to describe discontinuities and kinks. Therefore, ultrahigh spatial resolution appears necessary more because of the demand to high accuracy of measurements and less because of the high spatial frequencies. Third, although difference Fourier technique has been used for decades, it did not ease the demand for

ultrahigh spatial resolution because conformational changes brought even more complexity into the picture, which has prevented aspherical orbital densities from showing up. What makes the difference in revealing the remanence of individual orbitals at a given median resolution is the decomposition and deconvolution (continued below).

Reviewer 1: Creation of such tools would be a great step in development of X-ray research methods. Unfortunately, the results presented do not allow one to agree unconditionally with the author's conclusions about the progress achieved, as well as with the interpretation of the results obtained when processing previously published experimental data. In particular, the annotated author's conclusions on the visualization of the 3d orbitals and structural rearrangements in the spin crossover process raise significant doubts. The suggestions and findings of the author could be a basis for further discussions, but the categorical style of interpretation of the results does not correspond to the level of evidence of these conclusions and may give rise to false illusions for non-experts in X-ray data processing. All this requires a substantial revision of the manuscript and of the conception.

Reviewer 2: The author is completely correct (lines 237-239) to describe the literature-reported electron densities with unique shapes as being directly influenced by electrons occupying specific orbitals. However, there is a very large difference between the current diffraction dataset that diffracts to 1.8 Angstrom resolution, and the literature report of Hirano et al. (2016), showing iron 3d and sulfur 3p density at 0.48 Angstrom resolution. It is important to specifically compare the numerical resolution limits for the Hirano and the present work at both lines 236 and 370; in order to fully communicate that the present work is not simply a routine report of orbital density. Rather, the current work is lower-resolution and more speculative in nature.

It has been stated multiple times in the revised manuscript that the captured image is NOT an accurate depiction of the 3*d* orbitals (lines 387-388 and 703-704), unlike the iron 3*d* and sulfur 3*p* orbitals at 0.48 Å. The difference in spatial resolution has been mentioned everywhere it can (lines 250 and 639). It is not a proper task for protein crystallography to provide accurate depictions of orbital densities. Instead, any direct evidence, at a good spatial resolution, to indicate a specific electronic event would be a desirable probe into the essence of biochemistry. This is one of the contributions of this work (continued below).

Reviewer 2: The analysis leads to several astonishing claims highly relevant to myoglobin/CO binding, including that 1) the high-spin 3d orbitals can be visualized throughout the time series, 2) the XFEL pulse heats up the specimen to ~500K, and 3) there is an oscillation with a 220 ps period.

In addition to the visual evidence to indicate the reappearance of high-spin $3d$ electrons, other conclusions such as the positional modulation, recoiling of the heme iron, the XFEL pulse induced temperature jump, and the low frequency oscillation in the solvent are as novel as the indication of the high-pin orbitals. These results are long awaited visual evidences to validate and to interpret decades of data in spectroscopy and computation. This work represents the highest achievement in serial femtosecond crystallography using XFELs. Each of the conclusions comes from a solid mathematical deduction of the XFEL datasets and corresponds to spectroscopic experiments and computational simulations in the field. However, making a reader "unconditionally" agree with my interpretations is beyond my paygrade.

Reviewer 1: Similarly, the author ignores that, when an electron transfers from one orbital to another, the image of the orbital in a difference map should appear twice: as a negative density of the orbital that has lost an electron, and a positive density of the orbitals that have gained an electron. He does not mention either the question on a principal possibility to see such a subtle change in 1.8 Å resolution maps. The absence of the U4 component in the light-light series of difference maps is presented as evidence of the speed of this transition.

This is a very constructive suggestion. Indeed, it would be ideal to see some negative densities showing the loss of the low-spin electrons in the ligated state. However, the loss of the CO ligand and the drop of the iron cause the displacement of many more electrons than the low-spin electrons. Negative densities in the component U_4 are overwhelming. At this resolution, the loss of the low-spin electrons cannot be separated from the main events upon photolysis. This note has been added (lines 400-405).

Reviewer 2: Investigating the light-triggered release of CO from myoglobin, the author collectively re-examines electron density maps from room temperature and cryo-trapped

structures, as well as the time-dependent XFEL paper [Barends et al. (2015)] covering the -0.1 ps to 150 ps time evolution. Mutual difference maps and single-dataset omit maps are analyzed with the Schmidt et al. (2003) method of singular value decomposition (SVD), to identify features in the electron density that change with either time, sample, or temperature. This reviewer does not know if SVD has ever been applied over so many independent variables, and if so it is worth mentioning.

Reviewer 2: 3) It is further stated (lines 390-393) that normal 2Fo-Fc maps cannot reveal small orbital changes, but that SVD analysis is highly sensitive. If true, this would be a critical factor, and everyone should be using SVD to analyze structure comparisons. It would be highly beneficial to lay out the evidence and controls, or provide literature references.

Applications of SVD in protein crystallography have had limited success in the past 20 years until Givens rotations are applied, mainly by me (Ren PLoS One 8, e77141 & e77363, 2013; Ren Nucleic Acids Res 6, 1, 2016). This is due to the fact that the default solution of SVD guarantees no physical meaning in the basis components. Instead, each component is a linear combination or weighted sum of several meaningful images. SVD guarantees all basis components are orthonormal. However, physically meaningful images are not necessarily orthonormal. Therefore, a satisfactory interpretation of an SVD result is based on luck, which is the "drawback" of SVD.

Reviewer 1: An essential drawback of the suggested approach is the stage of the "manual" rotation of the found set of basic density distributions with unclear criteria of the best choice. This makes it difficult to reproduce the presented results.

Reviewer 1: The author's approach to the study of a series of experimental electron density distributions close to each other is based on a representation of each of these distributions as a weighted sum of a fixed set of density distributions pairwise uncorrelated, considered as a set of basis vectors mutually orthogonal. A visual study of the evolution of electron density under changing of external conditions is substituted by a numeric analysis of the weights with which these basis functions enter into one or another experimental density distribution. Sets of basic vectors can be selected in many ways and the basic distributions themselves may have no structural meaning. The recommendation of the author is to choose the basis in which "... an appropriate perspective can be found to elucidate the relationship between the core data and metadata clearly and concisely". This recommendation is extremely vague,

The concept of rotation was introduced much earlier by Henry & Hofrichter (in Num Comp Meth 210, 129, 1992). But their protocol specializes in the field of spectroscopy. More importantly, it fails to preserve the orthonormal and least-squares properties of SVD after rotation. This manuscript is the first time that I explicitly describe a rotation procedure with the orthonormal and least-squares properties preserved. That is to say, the rotation procedure finds an alternative solution for SVD. The rotation does not change the internal relationship among datasets except that it finds a different perspective to view the relationship. This rotation procedure may require a user decision. But this is not a drawback; this is a major advance in data analysis of dynamic crystallography. It is this rotation that finally connects the core crystallographic data to the metadata of experimental conditions. This rotation finally accomplishes the deconvolution of physical or chemical factors after decomposition. It is absolutely true that "everyone should be using SVD to analyze structure comparisons." I also believe that this advance is not limited to crystallography. It is going to impact other fields using SVD.

For example in Fig. 2a, there is a large gap between all datasets < 1 ps and those > 1 ps, where "large" is with respect to the spacings among the datasets in each group. A rotation does not change this fact except that this gap was not originally along the first dimension c_1. Therefore, the original $\boldsymbol{U}_1$ cannot describe the structural meaning of this gap, that is, the difference between the short delays < 1 ps and the longer delays > 1 ps, because the original $\boldsymbol{U}_1$ used to be a linear combination of several factors. The criteria of

the best choice are clear, that is, to separate these two groups along a major dimension, a deterministic solution. However, did I know this large gap before SVD? Of course not. This is where a user input is necessary. Nevertheless, one cannot create this large gap by any rotation. It always exists even I don't see it before a proper rotation is found. In fact, an interactive 3D plotting on the computer screen easily shows this large gap. What if another person decides to rotate in a different way? It is entirely possible that this large gap ends up orienting along another dimension. But the conclusion would not change. What if one decides not to apply any rotation? A linear combination of several components can still explain the inclined large gap with an identical conclusion.

For another example in Fig. 2cd, all difference maps have $c_4 \approx 0$ except those calculated with the dark dataset as a reference, the red trace. The criteria of the best rotation are very clear and deterministic so that U_4 depicts features only in the difference maps with the dark dataset as a reference. Again, the fact cannot be changed with any rotation that light-dark difference maps standout from the other maps except that the other maps were not originally on the plane of $c_4 = 0$. Therefore, the original U_4 was puzzling with several physical meanings fixed in it. In this sense, it can be stated that the Ren rotation protocol completes a deconvolution of physical and chemical factors after a singular value decomposition. A more concise version of these discussions has been incorporated into Methods (lines 1365-1391).

Reviewer 2: If I were to perform an SVD analysis on a new protein system, how would I go about applying the Ren methodology to this new system? The core issue here is reproducibility—it would be good if more details were given about how the rotations were derived, for this particular work. Would the conclusions have been robust if slightly different U-vectors were chosen?

A satisfactory interpretation in an SVD analysis, the rotation after SVD in particular, highly depends on the scientific question. A major basis component may very likely describe an uninteresting fact that has been known for many years, the so-called trivial component. A major basis may also describe a systematic error, interesting but off-topic. The key to a proper rotation is the correlation between the core data and metadata, the reason why the name meta-analysis (Ren et al. PNAS 109, 107, 2012).

With a clear scientific question in mind, a rotation is deterministic. However, before the data are evaluated by SVD, it is indeed difficult to predict what SVD may turn out. Since two different physical parameters, say temperature and pH, may not cause orthogonal changes to a structure, a single rotation is not sufficient to satisfy both dimensions. That is to say, multiple rotations, thus multiple perspectives, are needed to answer all the scientific questions. If two parameters are largely orthogonal to each other, the solutions are deterministic and stable. However, if two parameters are highly colinear, that is, they cause very similar structural responses, the solutions could be less stable. Discussion and Methods have been expanded. However, this is not a paper on applications of SVD.

Reviewer 1: A key problem in identifying small changes in the electron density of an object in a series of experiments under variable conditions is that one never has the true distribution of electron density. What a researcher has instead is a Fourier synthesis of electron density, always of a limited resolution. Saying nothing about experimental errors, these syntheses contain larger or smaller distortion, caused by lack of the necessary data. The magnitude of such distortions, called Fourier series truncation errors, can be significant and these distortions can mask the subtle changes in the electron density.

Reviewer 1: Even 0.6 Å-resolution syntheses can contain Fourier series truncation errors exceeding the signal level and mimicking structural details. The results presented by the author have been obtained at resolution of 1.8 Å, which is intermediate between "medium" and "high" resolutions according to the existing crystallographic classification. At such conditions, sufficiently large distortions of Fourier syntheses are expected. In this regard, the questions arise: "What are the changes studied in the manuscript? Are they those in the electronic structure of the sample or just a game of the Fourier series truncation errors?" The author does not pay any attention to these circumstances. Moreover, the author regularly speaks of "ultrahigh resolution", which can confuse the reader.

The topic of Fourier truncation error is not new. Its impact to electron density maps has been well understood. Errors from this source are not an unpredictable ghost. Therefore, we cannot attribute everything that we don't understand to this error. First, a Fourier truncation error may most likely occur to several static datasets included in this analysis, in which the spatial resolutions are truncated to 1.75 Å to match that of XFEL datasets. However, such errors are not clearly present in these datasets, thanks to

the difference Fourier technique and the simulated annealing protocol. Second, when the average intensity of an experimental dataset naturally decays as function of the spatial resolution, Fourier truncation errors are damped to a minimum. A low resolution dataset not necessarily contains more severe Fourier truncation errors; it simply lacks more details. Third, difference Fourier technique again self-cancels the remaining errors because both datasets would carry similar Fourier truncation errors. Fourth, the applied weighting scheme severely reduces any large differences in Fourier synthesis (Methods), which also prevents Fourier truncation errors. Fifth, even when Fourier truncation errors occur, such errors may appear as minor components after SVD (Figs. 2e and 4f), therefore isolated from the signals under discussion. Sixth, the SVD rotations validate that the signals under discussion, such as the positive cubic network, is not due to Fourier truncation errors because of its correlation with the metadata. An artifact due to Fourier truncation errors, if so conclude, needs to be explained why it is correlated with a physical or chemical parameter. As a conclusion, Fourier truncation errors are well guarded from. New findings cannot be dismissed under the fear of such errors. Discussion on Fourier truncation errors has been added (lines 679-695).

Reviewer 1: The start data is a set of several series of difference maps. The first series is based on the differences between results of the "dark" (without the use of exciting light pulses) experiment and ones in which the crystals were irradiated both light X-ray pulses ("light" sets), with different time delay

$F_i^{light} - k_{(i,0)} \cdot F^{dark}$,i=1,2,3,…

The remaining difference maps were calculated using only "light" data sets

$F_i^{light} - k_{(i,j)} \cdot F_j^{light}$,i=j,j+1,…

Here $k_{(i,j)}$ are scaling factors, individual for each map.

I start with a naive crystallographic explanation of the results. The first series of maps is shown in Fig. S2. These maps do not reproduce the typical pattern of difference maps, but rather reproduce a flipped (the one with the sign changed) image of the electron density distribution in the vicinity of the heme (the initial distribution is shown in Fig. S5). This is apparently caused by a imbalance in the difference syntheses of the first series, i.e. by too large values of the coefficients $k_{(i,0)}$, that results in the $-F^{dark}$ component dominating the maps.

It remains very puzzling to me after many readings of the above comments that Lunin, the first reviewer, calls the difference maps displayed in Fig. S2 "not typical" and "flipped sign". There are two ways to interpret Lunin's comments: The positive and negative densities in the difference maps are mistakenly flipped or the difference maps look like a negation of an electron density map in Fig. S5. Neither makes sense to me. The convention in producing these difference maps is light-dark so that positive densities show the product of light and negative densities show the loss of the dark state. The first column of Fig. S2 Z+1Å shows a cross section parallel to the heme plane and 1 Å towards to distal side. The second column Z shows the heme plane. It is obvious that Z+1Å cross section has more positive densities and Z cross section has more negative densities. This is a typical difference map of light-dark at the heme site and cannot have a flipped sign because the heme buckles towards the distal side as a result of the light illumination. Also, the bound CO ligand is in an intense negative peak indicating the loss of the ligand and an intense positive peak skewed on the proximal side of the iron (right column in Fig. S2). Again, this is typical and without sign flipping. These maps are identical to those originally produced by Barends & Schlichting (Science 350, 445, 2015) except in an arguably better quality. The scaling of the light and dark datasets is perfectly balanced as judged by equal amounts of positive and negative densities. Alternatively, a sign flipped image from Fig. S5 in Fig. S2? Of course, this is light-dark, the part of -dark should be a sign flipped image (see below). It seems that there exist some fundamental misunderstandings I have not figured out.

Reviewer 1: After decomposition, a balanced differential density distribution appears as the basic U1 distribution (Fig. S3). This map is close to the difference map of 1.8 Å resolution. The excessive presence of the term $-F^{dark}$ in the initial difference syntheses $F^{light}-$ ⟦$k \cdot F$⟧ ^{dark} is compensated by U4 component (Fig. 3c), which resembles again the flipped image of density in the vicinity of the heme (Fig. S5). As noted by the author, in light-to-light difference syntheses $F_i^{light}-k \cdot F_j^{light}$ the U4 component is absent. This can be explained by a better balance of syntheses coefficients when working with data sets obtained under the same experimental conditions (in the presence of both "light" and "dark" X-ray pulses) and, as a result, the absence of a "ghost" corresponding to the initial heme density distribution.

Essentially, Lunin questions a possibility that the light-dark series is not scaled as well as the other light-light series, because the light and dark datasets were obtained

under slightly different conditions, such as no laser pulse for the dark dataset. As a result, a "ghost" component exists in the light-dark series but not in the light-light series that has been decomposed by SVD as U_4. The principle in meta-analysis is the correlation between core data and metadata. If the component U_4 is an artifact due to scaling errors, it must be explained why these errors remain constant for short delays < 1 ps and decay as function of time > 1 ps. The most significant systematic error should occur to the deoxy dataset collected at a synchrotron instead of an XFEL (marked as Mb in Fig. 2). However, this dataset fits smoothly into the trace. Again, an unusual result cannot be simply dismissed by an unexplained "error", here a scaling error.

Reviewer 1: The author suggests a totally different explanation declaring U4 component as visualization of 3d orbital. An attention is not paid to the similarity of this distribution with the density distributions in the vicinity of the heme, shown in Fig. S2 and S5, as well as to the discrepancy between the size of the orbitals and the size of the density fragments in the U4 component noted by the author.

Again, this comment is very puzzling to me. There is a negative image of Fig. S5 in Fig. S2. This is correct, since Fig. S2 shows light-dark difference maps and Fig. S5 shows an average map of all datasets, which is very close to the dark map. The negative part of Fig. S2 is supposed to be similar to Fig. S5. As the size of the density features, there is not enough spatial resolution to depict an accurate map of the 3d orbitals (see above). However, this is not to say that the net gain of densities further away from the iron is completely meaningless. The shape of the net gain coincides with the calculated orbital densities, which strongly suggests the influence of the high-spin 3d orbitals.

Reviewer 2: 3) It is difficult to understand why the time evolution in Fig. 2f is described as an "oscillation". Not even one full cycle is observed, so it should not be assumed to be a circular progression. If there were more time points available they could prove or disprove the model.

Here an oscillation is only a model that requires the least number of parameters to describe the variation. This can be updated when more observations become available (line 460).

Reviewer 2: A. The author does a good job at describing how the analysis was done, and providing literature references to place the work in context. However, I am concerned that the data presentation and logical arguments are quite lengthy and detailed, and are challenging to keep straight. To assist the wider audience, a few things could readily be done to enhance the text's accessibility:

This reviewer offered a number of excellent suggestions to make the manuscript better.

Reviewer 2: 2) The claim that the XFEL beam pumps the sample to 500 K relies on evidence that is quite indirect—Fig 6i assumes that the synchrotron datasets at various thermometer temperatures can be plotted on a straight line with respect to the c3 coefficient, and that the XFEL data can then be placed directly on this line. It would be more cautious if the text admitted that there is no other physical evidence for this claim. In other words, this is an interesting but unique claim for which other evidence should be gathered.

This is a good suggestion. A note has been added to state that the temperature jump is solely based on a linear extrapolation (lines 525-527).

Reviewer 2: B. The subject matter and techniques are extremely specialized, and I believe that more must be done to help the audience place the methods and claims into a larger context. Four specific concerns are:
1) Even for an expert protein crystallographer, the spin states of CO-myoglobin may not be common knowledge. It is important to give enough basic chemistry background in the Introduction so the reader does not erroneously conclude that the data actually distinguish between high spin and low spin electrons. Rather, we know fundamentally that in MbFe(II)CO, Fe(II) has 6 electrons, all are paired so the total spin is S=0. This is because the 5 d-orbitals are split by the ligand field into 3 lower t2g + 2 upper eg; so the lower orbitals are all full. On photolysis of CO, the ligand field and orbital splitting decrease, so there are four unpaired electrons with total spin S=2. Including this textbook material would help to capture a larger audience.

This is an excellent point. I have incorporated a little more into the brief background of spin crossover in Introduction (lines 108-116).

Reviewer 2: 1) There are exactly 11 novel claims made in Fig. 1, represented by blue lines in the time chart with corresponding text description. It would be good for each of these blue lines to indicate which Figures contain the supporting evidence. For example, the "heme heating up to 500-600K" is clearly a consequence of Fig. 6i. Cross-referencing in this manner would help the reader keep track of the logical arguments.
2) A similar comment applies to the claims in the Introduction. Clearly line 172 "high spin iron" follows from Fig. 3; however, this reviewer cannot locate the evidence for "high frequency vibrational mode" and "low frequency oscillations" in lines 178-179.

I have implemented these cross referencing.

Reviewer 2: 3) Part of the problem is that the reader does not necessarily have the information about what is "high frequency" and "low frequency" in a particular context; after all, the paper discusses 18-orders of magnitude of time in Fig. 1. It seems safer to avoid the use of "high" and "low" and possibly give specific numerical ranges or functional descriptions.

I have added the corresponding wavenumbers where high or low frequency is mentioned.

Reviewer 2: 4) Also the line 213-214 claim that an oscillation occurs more in the solvent channels than in the protein or heme is similarly difficult to find in the visual evidence. Is it within Figure 6 somewhere? One possible strategy would be to take specific claims out of the Introduction since they are repeated anyway in the sequence of Results.

Fig. 7 is about the oscillation largely in the solvent. Cross referencing has been added.

Reviewer 2: Various typographical and language corrections were found:
Line 56 "of hundreds of K"
Line 80 "first few"
Line 151 "preemptively avoiding"
Line 192 "not sufficient time points"
Line 204 please specify what kind of shift

Line 577 "features can be found"

Line 608 "structural"

Line 630 "conceals"

Line 929 "first few"

All fixed.

Reviewers' comments:

Reviewer #1 (Remarks to the Author):

The author challenges an ambitious task to extract tiny structural information possessing X-ray scattering data sets of a resolution only about 1.8 Å while usually this requires data of much higher resolution, atomic 1.0-1.2 Å or even ultrahigh, about 0.8 Å. Creation of such tools would be a great step in development of X-ray research methods. Unfortunately, the results presented do not allow one to agree unconditionally with the author's conclusions about the progress achieved, as well as with the interpretation of the results obtained when processing previously published experimental data. In particular, the annotated author's conclusions on the visualization of the 3d orbitals and structural rearrangements in the spin crossover process raise significant doubts. An essential drawback of the suggested approach is the stage of the "manual" rotation of the found set of basic density distributions with unclear criteria of the best choice. This makes it difficult to reproduce the presented results. The suggestions and findings of the author could be a basis for further discussions, but the categorical style of interpretation of the results does not correspond to the level of evidence of these conclusions and may give rise to false illusions for non-experts in X-ray data processing. All this requires a substantial revision of the manuscript and of the conception.

A key problem in identifying small changes in the electron density of an object in a series of experiments under variable conditions is that one never has the true distribution of electron density. What a researcher has instead is a Fourier synthesis of electron density, always of a limited resolution. Saying nothing about experimental errors, these syntheses contain larger or smaller distortion, caused by lack of the necessary data. The magnitude of such distortions, called Fourier series truncation errors, can be significant and these distortions can mask the subtle changes in the electron density. In particular, this leads to the fact that individual electrons and hydrogen atoms become distinguishable in Fourier difference syntheses only when working with data sets at resolution of 1.0-1.2 Å (atomic resolution), while partial charges (density of valence electrons) become visible only at about 0.8 Å (ultra-high resolution) and only under certain restrictions on the structure refinement protocol (Afonine et al., 2004, Acta Cryst. D60, 260-274). Even 0.6 Å-resolution syntheses can contain Fourier series truncation errors exceeding the signal level and mimicking structural details. The results presented by the author have been obtained at resolution of 1.8 Å, which is intermediate between "medium" and "high" resolutions according to the existing crystallographic classification. At such conditions, sufficiently large distortions of Fourier syntheses are expected. In this regard, the questions arise: "What are the changes studied in the manuscript? Are they those in the electronic structure of the sample or just a game of the Fourier series truncation errors?" The author does not pay any attention to these circumstances. Moreover, the author regularly speaks of "ultrahigh resolution", which

can confuse the reader.

The author's approach to the study of a series of experimental electron density distributions close to each other is based on a representation of each of these distributions as a weighted sum of a fixed set of density distributions pairwise uncorrelated, considered as a set of basis vectors mutually orthogonal. A visual study of the evolution of electron density under changing of external conditions is substituted by a numeric analysis of the weights with which these basis functions enter into one or another experimental density distribution. Sets of basic vectors can be selected in many ways and the basic distributions themselves may have no structural meaning. The recommendation of the author is to choose the basis in which "... an appropriate perspective can be found to elucidate the relationship between the core data and metadata clearly and concisely". This recommendation is extremely vague, opening up the scope for the selection of the "perspective" that confirms most the hypothesis of the researcher, and makes it difficult to reproduce and validate the results obtained.

The start data is a set of several series of difference maps. The first series is based on the differences between results of the "dark" (without the use of exciting light pulses) experiment and ones in which the crystals were irradiated both light X-ray pulses ("light" sets), with different time delay
$F_i^{light} - k_{(i,0)} \cdot F^{dark}$, $i = 1,2,3,\dots$
The remaining difference maps were calculated using only "light" data sets
$F_i^{light} - k_{(i,j)} \cdot F_j^{light}$, $i = j, j+1, \dots$
Here $k_{(i,j)}$ are scaling factors, individual for each map.

I start with a naive crystallographic explanation of the results. The first series of maps is shown in Fig. S2. These maps do not reproduce the typical pattern of difference maps, but rather reproduce a flipped (the one with the sign changed) image of the electron density distribution in the vicinity of the heme (the initial distribution is shown in Fig. S5). This is apparently caused by a imbalance in the difference syntheses of the first series, i.e. by too large values of the coefficients $k_{(i,0)}$, that results in the $-F^{dark}$ component dominating the maps. After decomposition, a balanced differential density distribution appears as the basic U1 distribution (Fig. S3). This map is close to the difference map of 1.8 Å resolution. The excessive presence of the term $-F^{dark}$ in the initial difference syntheses $F^{light} - ⟦k \cdot F⟧^{dark}$ is compensated by U4 component (Fig. 3c), which resembles again the flipped image of density in the vicinity of the heme (Fig. S5). As noted by the author, in light-to-light difference syntheses $F_i^{light} - k \cdot F_j^{light}$ the U4 component is absent. This can be explained by a better balance of syntheses coefficients when working with data sets obtained under the same experimental conditions (in the presence of both "light" and "dark" X-ray pulses) and, as a result, the absence of a "ghost" corresponding to the initial heme density distribution.

The author suggests a totally different explanation declaring U4 component as visualization of 3d orbital. An attention is not paid to the similarity of this distribution with the density distributions in the vicinity of the heme, shown in Fig. S2 and S5, as well as to the discrepancy between the size of the orbitals and the size of the density fragments in the U4 component noted by the author. Similarly, the author ignores that, when an electron transfers from one orbital to another, the image of the orbital in a difference map should appear twice: as a negative density of the orbital that has lost an electron, and a positive density of the orbitals that have gained an electron. He does not mention either the question on a principal possibility to see such a subtle change in 1.8 Å resolution maps. The absence of the U4 component in the light-light series of difference maps is presented as evidence of the speed of this transition.

By this particular example, I would like to illustrate that the results obtained allow different interpretations that could be a subject of discussion, and that the author's interpretation is rather vague and should not be presented as a proof.

Vladimir Y. Lunin

Reviewer #2 (Remarks to the Author):

Investigating the light-triggered release of CO from myoglobin, the author collectively re-examines electron density maps from room temperature and cryo-trapped structures, as well as the time-dependent XFEL paper [Barends et al. (2015)] covering the -0.1 ps to 150 ps time evolution. Mutual difference maps and single-dataset omit maps are analyzed with the Schmidt et al. (2003) method of singular value decomposition (SVD), to identify features in the electron density that change with either time, sample, or temperature. This reviewer does not know if SVD has ever been applied over so many independent variables, and if so it is worth mentioning.

The analysis leads to several astonishing claims highly relevant to myoglobin/CO binding, including that 1) the high-spin 3d orbitals can be visualized throughout the time series, 2) the XFEL pulse heats up the specimen to ~500K, and 3) there is an oscillation with a 220 ps period.

A. The author does a good job at describing how the analysis was done, and providing literature references to place the work in context. However, I am concerned that the data presentation and logical arguments are quite lengthy and detailed, and are challenging to keep straight. To assist the wider audience, a few things could readily be done to enhance the text's accessibility:

1) There are exactly 11 novel claims made in Fig. 1, represented by blue lines in the time chart with corresponding text description. It would be good for each of these blue lines to indicate which Figures

contain the supporting evidence. For example, the "heme heating up to 500-600K" is clearly a consequence of Fig. 6i. Cross-referencing in this manner would help the reader keep track of the logical arguments.

2) A similar comment applies to the claims in the Introduction. Clearly line 172 "high spin iron" follows from Fig. 3; however, this reviewer cannot locate the evidence for "high frequency vibrational mode" and "low frequency oscillations" in lines 178-179.

3) Part of the problem is that the reader does not necessarily have the information about what is "high frequency" and "low frequency" in a particular context; after all, the paper discusses 18-orders of magnitude of time in Fig. 1. It seems safer to avoid the use of "high" and "low" and possibly give specific numerical ranges or functional descriptions.

4) Also the line 213-214 claim that an oscillation occurs more in the solvent channels than in the protein or heme is similarly difficult to find in the visual evidence. Is it within Figure 6 somewhere? One possible strategy would be to take specific claims out of the Introduction since they are repeated anyway in the sequence of Results.

B. The subject matter and techniques are extremely specialized, and I believe that more must be done to help the audience place the methods and claims into a larger context. Four specific concerns are:

1) Even for an expert protein crystallographer, the spin states of CO-myoglobin may not be common knowledge. It is important to give enough basic chemistry background in the Introduction so the reader does not erroneously conclude that the data actually distinguish between high spin and low spin electrons. Rather, we know fundamentally that in MbFe(II)CO, Fe(II) has 6 electrons, all are paired so the total spin is S=0. This is because the 5 d-orbitals are split by the ligand field into 3 lower t2g + 2 upper eg; so the lower orbitals are all full. On photolysis of CO, the ligand field and orbital splitting decrease, so there are four unpaired electrons with total spin S=2. Including this textbook material would help to capture a larger audience.

2) The author is completely correct (lines 237-239) to describe the literature-reported electron densities with unique shapes as being directly influenced by electrons occupying specific orbitals. However, there is a very large difference between the current diffraction dataset that diffracts to 1.8 Angstrom resolution, and the literature report of Hirano et al. (2016), showing iron 3d and sulfur 3p density at 0.48 Angstrom resolution. It is important to specifically compare the numerical resolution limits for the Hirano and the present work at both lines 236 and 370; in order to fully communicate that the present work is not simply a routine report of orbital density. Rather, the current work is lower-resolution and more speculative in nature.

3) It is further stated (lines 390-393) that normal 2Fo-Fc maps cannot reveal small orbital changes, but that SVD analysis is highly

sensitive. If true, this would be a critical factor, and everyone should be using SVD to analyze structure comparisons. It would be highly beneficial to lay out the evidence and controls, or provide literature references.

4) The left-singular vectors are not unique—this is correctly described in the supplementary material starting at line 1201. At 1249 it is stated that a Givens rotation is applied so that an appropriate perspective can be found to elucidate the relationship between the core data and metadata. However, this leaves us with a situation where the main results of the paper (e.g., the U4 difference map showing high-spin density) are not based on a deterministic procedure! To paraphrase what I understand from this description, U4 was chosen by manually finding a Givens rotation that most clearly correlates with time-dependence. However, the author does not give a protocol to find that rotation, nor a metric to judge when the rotation is optimized. If I were to perform an SVD analysis on a new protein system, how would I go about applying the Ren methodology to this new system? The core issue here is reproducibility—it would be good if more details were given about how the rotations were derived, for this particular work. Would the conclusions have been robust if slightly different U-vectors were chosen?

C. Finally, some aspects of the central results could be presented with greater caution, considering that they are somewhat speculative, and that they depend on interpretation.

1) The author explains that the initial report of the observation of 3d orbitals (Zuo el al 1999) was somewhat controversial. In particular Scerri (2000) states that orbitals of single electrons are fundamentally not observable, and recommends that observations be described in terms of electron density. This would be the most cautious language. However, haven't the philosophical arguments now run their course after 20 years, so it should now be safe to speak of the U4 component as revealing the high-spin orbitals? Unfortunately, the author's own description shows that the electron density is only marginally interpretable as a 3d orbital. Line 368 states that the 3d orbital for an isolated Fe(II) has 0.36 Angstrom radius, and line 369 that the orbital density in an Fe4S4 cluster has a maximum radius of 1 Angstrom. Meanwhile, the data shown here place the electron density at 1.7 to 1.9 Angstrom from the iron, depending on which map is considered, so it appears that this analysis is "first of its kind" result, rather than the observation of a 3d orbital by established methods. It is somewhat difficult to understand the author's point of view in the following text (lines 372-394). It is stated that the cubic density is not the probability distribution of a 3d electron, yet the paper states in many places that we observe a high-spin 3d orbital.

2) The claim that the XFEL beam pumps the sample to 500 K relies on evidence that is quite indirect—Fig 6i assumes that the synchrotron datasets at various thermometer temperatures can be plotted on a straight line with respect to the c3 coefficient, and that the XFEL

data can then be placed directly on this line. It would be more cautious if the text admitted that there is no other physical evidence for this claim. In other words, this is an interesting but unique claim for which other evidence should be gathered.

3) It is difficult to understand why the time evolution in Fig. 2f is described as an "oscillation". Not even one full cycle is observed, so it should not be assumed to be a circular progression. If there were more time points available they could prove or disprove the model.

Various typographical and language corrections were found:
Line 56 "of hundreds of K"
Line 80 "first few"
Line 151 "preemptively avoiding"
Line 192 "not sufficient time points"
Line 204 please specify what kind of shift
Line 577 "features can be found"
Line 608 "structural"
Line 630 "conceals"
Line 929 "first few"