Title: BIOTIN ATTACHMENT DOMAIN-CONTAINING proteins, inhibitors of

ACCase, are regulated by WRINKLED1

Hui Liu[1], Zhiyang Zhai[1], Kate Kuczynski, Jantana Keereetaweep, Jorg Schwender, John Shanklin[#]

Hui Liu

ORCID: 0000-0001-8524-1759

Dept. of Biology

BNL 463,

50 Bell Ave,

Upton NY 11973, USA

631 344 5360

huiliu1@bnl.gov

Zhiyang Zhai

ORCID: 0000-0003-3181-1773

Dept of Biology

BNL 463,

50 Bell Ave,

Upton NY 11973, USA

631 344 5360

zzhai@bnl.gov

Jantana Keereetaweep

ORCID: 0000-0001-8314-9289

Dept of Biology

BNL 463,

50 Bell Ave,

Upton NY 11973, USA

631 344 5360

keereetaweep@bnl.gov

Kate Kuczynski

ORCID: 0000-0002-0625-1017

Dept of Biology

BNL 463,

50 Bell Ave,

Upton NY 11973, USA

kkuczynski@bnl.gov

Tel: 631 344 5068

Jorg Schwender

ORCID: 0000-0003-1350-4171

Dept of Biology

BNL 463,

50 Bell Ave,

Upton NY 11973, USA

schwend@bnl.gov

Tel: 631 344 3797

[#]John Shanklin (corresponding author)

ORCID: 0000-0002-6774-8043

Dept of Biology

BNL 463,

50 Bell Ave,

Upton NY 11973, USA

shanklin@bnl.gov

Tel: 631 344 3414

One sentence summary:

WRI1 regulates genes encoding both fatty acid synthesis and inhibitors of FA synthesis (BADCs),
creating a lipid homeostatic mechanism in which the transcription of FA synthetic capacity is coordinated
with the capacity to biochemically downregulate FA synthesis.

Running title: WRI1 regulates BADC proteins

Key words: Metabolic regulation, homeostatic mechanism, lipid metabolism, triacylglycerol

[1]Liu and Zhai contributed equally to this work and are both considered first authors.

Footnotes: This work was supported by the Division of Chemical Sciences, Geosciences, and
Biosciences, Office of Basic Energy Sciences, US Department of Energy (grant DOE KC0304000 JSh)
(and DE-SC0012704 JSc), and by a National Science Foundation under which initial discussions took
place and several reagents were made (Grant DBI 1117680 JSh and JSc).

The authors declare no conflict of interest.

70   **List of author contributions**

71   H.L, Z.Z. J.Sc. J.K., and J.Sh. conceived the original research plans and designed the

72   experiments; H.L, K.K., Z.Z. and J.K. performed the experiments; H.L, Z.Z. J.Sc. J.K., and J.Sh.

73   analyzed the data and wrote the article with contributions of all the authors; J.Sh. agrees to serve

74   as the author responsible for contact and ensures communication.

Corresponding author:
John Shanklin
Biology Department
Brookhaven National Laboratory
Upton, NY 11973, USA

Phone: (631) 344-3414

Fax: (631) 344-3407

shanklin@bnl.gov

ACCESSION NUMBERS:

Sequence data from this article can be found in The Arabidopsis Information Resource under the following

accession numbers: *WRI1* (AT3G54320), *BADC1* (AT3G56130), *BADC2* (AT1G52670), *BADC3*

(AT3G15690), *BCCP1* (AT5G16390), *BCCP2* (AT5G15530), *Fbox* (AT5G15710) and *UBQ10*

(AT4G05320).

Abstract

WRINKLED1 (WRI1) is a transcriptional activator that binds to AW boxes in the promoters of many genes

from central metabolism and FA synthesis, resulting in their transcription. BIOTIN ATTACHMENT

DOMAIN-CONTAINING (BADC) proteins are homologs of BIOTIN CARBOXYL CARRIER PROTEIN

(BCCP) that lack a biotin-attachment domain and are therefore inactive. In the presence of excess FA,

BADC1 and BADC3 are primarily responsible for the observed long-term irreversible inhibition of

ACETYL-COA CARBOXYLASE (ACCase), and consequently FA synthesis. Purified WRI1 bound with

high affinity (*Kd*s in the low nanomolar range) to canonical AW-boxes from the promoters of all three

BADC genes. Consistent with this observation, the expression of *BADC1, BADC2 and BADC3* genes and

BADC1 protein levels were reduced in *wri1-1* relative to wild type (WT), and *BADCs* gene expression and

BADC1 protein levels also were elevated upon *WRI1* overexpression. The double mutant *badc1badc2*

phenocopied *wri1-1* with respect to both reduction in root length, and elevation of indole-3-acetic acid-Asp

(IAA-Asp) levels relative to WT. Overexpression of *BADC1* in *wri1-1* decreased its IAA-Asp and partially

rescued its short-root phenotype demonstrating a role for BADCs in seedling establishment. That WRI1

positively regulates genes encoding both FA synthesis and BADCs i.e., conditional inhibitors of FA

synthesis, represents a coordinated mechanism to achieve lipid homeostasis in which plants couple the

transcription their FA synthetic capacity with their capacity to biochemically downregulate it.

Introduction

Lipids are primary metabolites in cells, acting as structural components of cell membranes, energy-dense

storage compounds and cell signaling molecules. Fatty acids (FAs) are major components of lipids and

triacylglycerols (TAGs) which are storage lipids that accumulate mostly in oil bodies in plant seeds (Li-

Beisson et al., 2013). De novo synthesis of FAs occurs in the plastid via a well-established pathway

(Ohlrogge and Browse, 1995; Rawsthorne, 2002), In Arabidopsis, transcription factors (TF): LEAFY

COTYLEDON1 (LEC1) and LEC2, FUSCA3 (FUS3), and ABSCISIC ACID INSENSITIVE3 (ABI3),

serve as the master regulators of embryogenesis and seed maturation (Parcy et al., 1997; Lotan et al., 1998;

To et al., 2006) and also regulate accumulation of TAG (Kagaya et al., 2005; Mendoza et al., 2005;

Braybrook et al., 2006). WRINKlED1 (WRI1), an APETALA2 (AP2)-type transcription factor, functions

downstream of LEC1 and LEC2 (Baud et al., 2007; Mu et al., 2008) and is a master regulator of FAs

synthesis because seeds of the *wri1* mutant show an 80% reduction in TAGs compared to wild type (WT)

(Focks and Benning, 1998). More than 20 *WRI1* target genes have been identified by comparing gene

expression in WT with that of *wri1* and *WRI1* overexpression-lines, and the use electrophoretic mobility

shift assay (EMSA) confirmed WRI1-gene promoter binding. Based on promoter sequence comparisons

[CnTnG](n)$_7$[CG] was identified as the consensus AW-binding site. WRI1 target sequences are found

upstream of genes coding for enzymes involved in glycolysis (sucrose synthase (SUS2), pyruvate kinase,

and pyruvate dehydrogenase), Glu6P and PEP plastidial transporters (Glu6P/phosphate translocator and

PEP/phosphate translocator), subunits of acetyl-CoA carboxylase (ACCase) (biotin carboxyl carrier protein

2 (BCCP2), biotin carboxylase (BC) and carboxyltransferase (CTα)), fatty acid synthesis (malonyl-

CoA:ACP malonytransferase, ketoacyl-ACP synthase, hydroxyacyl-ACP dehydrase, enoyl-ACP reductase,

acyl-carrier proteins, oleoyl-ACP thioesterase (FatA)) and genes involved in lipoic acid synthesis, a

cofactor of pyruvate dehydrogenase (Ruuska et al., 2002; Baud et al., 2007; Maeo et al., 2009; Fukuda et

al., 2013; Li et al., 2015). Interestingly, genes involved in TAG synthesis and oil body assembly such as

diacylglycerol acyltransferase 1 (DGAT1) that catalyzes the final step of TAG synthesis and oleosins and

caleosins appear not to be regulated by WRI1, though some species-specific variation has been reported

(Maeo et al., 2009; Grimberg et al., 2015). By characterizing gene expression of subunits of ACCase, it

was shown that the distance between the AW site and the translational initiation site (TIS) strongly

influences the function of the AW box (Fukuda et al., 2013) in that the majority of AW sites fall within

200bp of the TIS in *bona fide* WRI1 target genes (Maeo et al., 2009; Fukuda et al., 2013).

A class of proteins annotated as biotin attachment domain-containing (BADC) (Olinares et al., 2010) have

been recently identified to be inhibitors of FAs synthesis (Salie et al., 2016; Keereetaweep et al., 2018).

BADC proteins are homologs of BCCP, sharing 25-30% sequence similarity, but they lack a biotin

attachment site, rendering them inactive. All three Arabidopsis BADC isoforms interact with BCCP.

Displacement of functional BCCP subunits in ACCase by inactive BADC subunits was believed to reduce

the ACCase catalytic activity (Salie et al., 2016). ACCase is subject to feedback-regulation upon exposure

to exogenously supplied fatty acids in the form of Tween esters (Tween-80 containing predominantly oleic

acid; 18:1). Short-term exposure results in reversible inhibition of ACCase, whereas longer-term exposure

results in irreversible inhibition (Andre et al., 2012). Oleoyl-ACP was shown to mediate the reversible

phase of inhibition (Andre et al., 2012), whereas BADC1 and BADC3 are primarily responsible for the

phase of irreversible inhibition (Keereetaweep et al., 2018).

In this work we present evidence that the expression of all three *BADC* genes are under the control of the

WRI1 transcription factor. The short-root phenotype and elevated conjugated IAA levels common to the

wri1-1 and the *badc1badc2* double mutant led us to hypothesize that BADC-deficiency is responsible for

the observed short-root phenotype. Restoration of near-WT root length upon overexpression of BADC1 in

both *wri1* and *badc1badc2* is consistent with a yet-to-be identified role for BADCs in seedling development.

Results

badc1 badc2 *double mutant shows reduced primary root growth and fewer lateral roots than WT.*

Seeds of the *badc1 badc2* double mutant generated in our previous study, contained 17% and 23% increases

in total fatty acid and TAG accumulation, respectively compared to WT (Keereetaweep et al., 2018). When

grown on vertical MS plates, *badc1 badc2* exhibited a reduction in primary root length by 62% and a

decrease in lateral root number by 39% compared with WT (Fig. 1A, 1B and 1C). In contrast, root growth

of *badc1* or *badc2* single mutants showed no differences from that of WT. A complementation experiment

was performed in which either *BADC1* or *BADC2* was expressed under the control of the constitutive 35S

promoter in the *badc1 badc2* mutant. More than 10 independent lines were generated for each of *BADC1*

and *BADC2* constructs into *badc1 badc2* (SFig. 1). All transgenic lines demonstrated restored root growth

to varying degrees, indicating the involvement of *BADC1* and *BADC2* in seedling development (Fig. 1D

and 1E). Because the *wri1* mutant (*wri1-1*) was recently shown to have short-root phenotype during

seedling establishment (Kong et al., 2017), we compared the growth of *badc1 badc2* and *wri1-1* under the

same conditions and show that their phenotypes are visually indistinguishable (SFig. 2).

Levels of conjugated auxin were elevated in badc1 badc2 *plants compared to those of wild-type.*

In *wri1-1* the levels of conjugated auxin, i.e., indole-3-acetic acid (IAA)-Asp, were reported to be

significantly elevated compared to WT (Kong et al., 2017). The similarity between the root phenotypes of

badc1 badc2 and *wri1-1* during seedling establishment prompted us to measure the levels of several growth

hormones. As shown in Fig. 2, IAA-Asp levels in 7-day-old seedlings showed a highly significant

approximately 50% increase relative to levels found in WT. In contrast, the levels of other plant hormones

i.e., ABA, JA and SA were not significantly different between *badc1 badc2* and WT (Table S1).

WRI1 interacts with AW boxes from the promoters of BADCs.

That *badc1 badc2* mimics *wri1-1* in both its short-root phenotype and IAA-Asp content led us to

hypothesize that *BADCs* are under transcriptional control of WRI1. Sequence analysis identified a canonical

AW-box consensus sequence within 200bp upstream of the translational initiation site (TIS) in the

promoters of all three *BADC* isoforms (Fig. 3A and Table S2). To test whether WRI1 binds directly to these

AW boxes, microscale thermophoresis (MST) assays were performed, in which a 28bp dsDNA fragment

(Table S2), containing each of AW boxes from *BADCs* was titrated against purified WRI1 DNA binding

domain (AA58-240) fused with GFP (SFig. 3). As shown in Fig. 3B and 3C, WRI1 exhibited low

equilibrium dissociation constants (*Kd*s) i.e., tight binding affinities for all three putative AW boxes with

Kd values between 0.27 and 5.24 nM, comparable with the affinity of WRI1 for the bona fide AW box1

from the promoter of the WRI1 target gene *BCCP2* of 0.65 nM (Maeo et al., 2009). Among the three

BADC genes, WRI1 showed the highest affinity for the BADC1 AW box at 0.27 ± 0.27 nM. The AW box

from the *BCCP1* promoter was also tested because although *BCCP1* contains an AW box (-60/-47 relative

to TIS) the gene was shown not to be regulated by WRI1 (Fukuda et al., 2013). Consistent with this, the *Kd*

value for dsDNA that includes the *BCCP1* AW box and WRI1 showed an approximately 37-fold higher *Kd*

value than those of for *BCCP2* and WRI1 (SFig. 4). To test whether the consensus sequence

[CnTnG](n)$_7$[CG] in BADCs AW boxes is critical for WRI1 binding, 3 different mutant sequences were

generated for each *BADC* AW box (Table S2) and tested in MST. As shown in Table S3, mutations in

conserved nucleotides dramatically decreased their affinities to WRI1. Mutations in the 7 nucleotides

between also decreased the affinity but to a less degree (Table S3).

BADCs *gene expression and BADC1 protein levels are lower in developing siliques of* wri1-1 *than those*

of WT.

Gene expression quantification of *WRI1* and *BADCs* in different tissues (flower, silique, root and mature

leaf) showed generally similar tissue-specific expression pattern to that of *WRI1* (Fig. 4A), i.e., highly

expressed in siliques (developing seeds) and low expression in mature leaves. Because of the highest

expression of *WRI1* and *BADCs* in developing seeds among analyzed tissues (Fig. 4A), siliques were

selected for investigating the relationship between *WRI1* and *BADCs*. Gene expression of *BADCs* in

developing siliques (7 days after flowering) of *wri1-1* were only 10-20% of that in WT (Fig. 4B). We next

used a BADC1 antibody that showed high specificity towards BADC1 protein as (Sfig.5) to probe BADC1

levels in *wri1-1*. Consistent with the observed decreased expression of *BADC1* in *wri1-1*, relative to WT,

the level of BADC1 protein in *wri1-1* observed by immunoblotting was only 23% of that in WT (Fig. 4C).

Since relative high gene expressions of *WRI1* and *BADCs* were also observed in seedling roots (Fig. 4A),

BADC1 protein level was quantified in seedling roots and found to be significantly lower in *wri1-1*

seedlings roots (12 day old) compared with wild type (Fig. S6).

BADCs *gene expression and BADC1 protein levels are significantly higher in developing siliques of* WRI1

overexpression transgenic plant than those of WT.

To test whether overexpression of *WRI1* would increase gene expression of *BADCs* and the abundance of

BADC1 polypeptide, two ethanol-inducible *WRI1* transgenic lines that were generated in our previous study

(Zhai et al., 2017) were used. Both lines demonstrated elevated WRI1 accumulation after induction with

2% ethanol treatment (Zhai et al., 2017). As shown in figure 5A, *BADCs*, *WRI1* and *BCCP2* gene expression

in developing siliques of *WRI1* transgenic plants that were induced by irrigation with 2% ethanol for 4 days

were significantly higher than either WT or the corresponding *WRI1* transgenic plant lines that were not

exposed to ethanol treatment (Fig. 5A). Consistent with the elevated gene expressions of *BADC1*, BADC1

protein levels were also significantly higher upon WRI1 overexpression (Fig. 5B and 5C). To test whether

global regulation of WRI1 on BADCs, gene expression and protein level were also quantified in the

seedling roots of *WRI1* transgenics, as shown in Fig. S7, both gene expression and protein level were higher

in *WRI1* transgenics induced by ethanol treatment than either non-ethanol treatment control or WT (Fig.

S7).

Overexpression of BADC1 *in* wri1-1 *partially rescues wri1-1 short-root phenotype*

The data presented above supports the hypothesis that the expression of *BADC* genes are under the control

of WRI1, and that the short-root phenotype of *wri1-1* primarily results from a deficiency of *BADC1* and

BADC2. To test this hypothesis, the CDS of *BADC1* was placed under the control of the 35S promoter and

transformed into *wri1-1*. 15 out of 17 independent transgenics demonstrated significantly longer primary

root length than that of *wri1-1* when germinated and grown vertically on ½ strength MS medium plates for

16 days (Fig. 6A and 6B). Immunoblot showed that BADC1 protein levels were significantly increased in

those *BADC1/wri1-1* transgenics (Fig. S8). To test whether IAA-Asp is somehow associated with the short-

root phenotype of *wri1-1*, IAA contents were measured in 7-day-old seedlings of *BADC1/wri1-1*, *wri1-1*

and WT and showed decreasing in *BADC1/wri1-1* when compared with *wri1-1*(Fig. 6C), supporting

BADC1 being a downstream component of WRI1-dependent root development.

Discussion

In this work we establish that expression of *BADC1, BADC2* and *BADC3* genes are under transcriptional

control of WRI1. Evidence to support this comes from: 1) the identification of AW boxes within 200bp

upstream of the TIS; 2) that microscale thermophoresis measurement demonstrated tight binding,i.e., *Kd*s

in the low nanomolar range between WRI1 and the AW boxes from all three *BADC* genes; 3) that

expression of each of the three *BADC* transcripts are reduced (by approximately 85%) in the *wri1-1* mutant;

4) the protein levels of BADC1 are reduced similarly to reductions in the level of its transcript in the *wri1-*

1 mutant. 5) Both gene expressions of *BADCs* and protein Levels of BADC1 are increased upon

overexpression of *WRI1*.

That WRI1, the master transcriptional activator of genes involved in FA synthesis, simultaneously activate

the transcription of BADCs which have been shown to act as negative regulators of ACCase might at first

appear paradoxical. However, FA synthesis is metabolically expensive in terms of energy and reducing

power, creating an imperative for tight metabolic control. For example, ACCase, the rate limiting step in

FA synthesis is under tight genetic and biochemical control (Salie et al., 2016). When free fatty acid levels

in the cell were increased beyond demand by supplementation with Tween-esters, two phases of feedback

inhibition were reported, one short term, the second longer term (Andre et al., 2012). Short term inhibition

occurs when the levels of oleoyl-ACP, the terminal product of the ACP-track of FA synthesis in the plastid,

increase. Oleoyl-ACP inhibits ACCase by binding to it in a reversible manner (Andre et al., 2012). If,

however, excess fatty acid levels persist for two or more days, an irreversible phase of inhibition of ACCase

is initiated that is primarily mediated by BADC1 and BADD3 (Keereetaweep et al., 2018). In this context,

BADC subunits should not be regarded as a simple inhibitor, but rather as a conditional inhibitor of

ACCase. Thus, coexpression of the approximately 20 genes that promote FA synthesis along with the three

BADC subunits makes biological sense because as WRI1 increases cellular metabolic demand by activating

FA synthesis, it concomitantly increases the capacity to downregulate ACCase, and thereby FA synthesis,

by increasing abundance of the conditionally inhibitory BADC subunits. Co-regulation of FAS and the

BADCs thus allows the cell to match synthetic capacity with its regulatory capacity. The biochemical

regulation of ACCase described herein is complementary to the transcriptional regulation of FAS we

recently reported in which levels of the WRI1 polypeptide are tightly coupled to the availability cellular

carbon in the form of sugars. When the carbon/energy status of the cell decreases, the SnRK1 carbon-

sensing kinase becomes activated and it phosphorylates WRI1, leading to its selective proteasomal

degradation thereby reducing activation of FAS genes (Zhai et al., 2017). Conversely, when carbon/energy

abundance increases, levels of the SnRK1 kinase inhibitor trehalose 6-phosphatae increase, reducing WRI1

phosphorylation/degradation, thereby increasing the amount of WRI1 available to activate the transcription

of FAS genes (Zhai et al., 2018).

In this study we also observed the *badc1 badc2* double mutant exhibits reduced primary root growth, a

phenotype that resembled that of the *wri1-1* mutant (Kong et al., 2017), which in combination of our finding

that WRI1 controls BADC transcription led us to ask whether the short-root phenotype characteristic of

both *wri1-1* and *badc1badc2* mutants result from a deficit of BADC. If correct, overexpression of BADC

in *wri1-1* would mitigate the *wri1-1* short-root phenotype. This view is supported by the observations that

overexpression of *BADC1* in *wri1-1* partially rescued its short-root phenotype in a manner similar to its

overexpression in the *badc1badc2* mutant background (and as overexpression of *BADC2* had in the

badc1badc2 mutant background) and reduced IAA-Asp content in *BADC1/wri1-1*. Taken together this

work suggests that deficiency of BADC levels, resulting either directly via mutation of *badc* genes, or

indirectly through the mutation of their upstream transcriptional activator, *wri1*, result in a short-root

phenotype *via* an as yet-to-be determined mechanism. Further work will be required to identify details of

the mechanism, that is likely related to increased levels of conjugated IAA that were observed for both

wri1-1 and *badc1badc2* mutants.

Taken together, we show all three BADC genes are direct target genes of WRI1 and can play a role in root

development. The observation that WRI1 not only upregulates genes contributing to FA synthesis, but also

BADCs which are conditional regulators of FA synthesis identifies an additional layer of regulation with

respect to FAs synthesis.

Materials and Methods

Plant Materials and Growth Conditions

Wild-type (WT) *Arabidopsis thaliana* (ecotype Columbia-0) and *wri1-1*(CS69538), *badc1* (Salk 000817C),

badc2 (Salk 021108C) and *badc3* (CS2103834) were used in this study. Double mutant *badc1 badc2* was

generated and described previously (Keereetaweep et al., 2018). For growth on plates, seeds were surface-

sterilized with 70% ethanol, followed by 30% bleach containing 0.01% Tween20, and rinsed three times

with sterile water. Seeds were stratified for 3 days at 4°C in the dark and germinated on half Murashige and

Skoog (MS) medium supplemented with 1% w/v sucrose plates in an incubator with a light/dark cycle of

18h/6h at 23°C, photosynthetic photon flux density of 250 μmol m^{-2} s^{-1}, and 75% relative humidity.

Genetic Constructs

CDS of *BADC1* and *BADC2* were amplified by PCR from cDNAs using primers listed in Table S4. The

PCR products were then cloned into the Invitrogen GATEWAY^M pDONR/Zeo vector (Thermo Fisher

Scientific, Waltham, MA; www.thermofisher.com) using the BP reaction and sub-cloned (LR reaction) into

the plant GATEWAY^{TM} binary vector: pGWB414 (Nakagawa et al., 2007). *BADC1*/pGWB414 and

BADC2/pGWB414 were introduced into *badc1 badc2* double mutant by agrobacterium (AgL0)-mediated

transformation respectively. For rescuing short-root phenotype of *wri1-1*, *BADC1*/pGWB414 was

transformed into *wri1-1* mutant.

Quantification of IAA, IAA-Asp in badc1 badc2 double mutant Seedlings

7-day-old Arabidopsis WT and *badc1 badc2* seedlings grown vertically on half MS medium

supplemented with 1% w/v sucrose were harvested (20-30 seedlings per replicate, ~100 mg FW) and

frozen in liquid Nitrogen. Frozen samples were sent to the Proteomics & Mass Spectrometry Facility at

the Donald Danforth Plant Science Center for acidic hormone analysis where frozen plant material is

extracted twice with ice-cold acetonitrile/methanol (1:1 v:v) using a bead beater and resulting samples

were dried with the use of a speedvac. Samples were reconstituted in 30% MeOH and separated via high

performance liquid chromatography using a Waters (Milford, MA) Acquity UPLC® BEH C18 1.0×100

mm, 1.7 μm column kept at 50 °C with a flow rate of 15 μL/min coupled to a a Sciex 6500 QTrap® mass

spectrometer (Framingham, MA) operated in multiple reaction monitoring (MRM) mode employing

polarity-switching for simultaneous detection of positive and negative ions. Data analysis was completed

using MultiQuant 3.0.2 (AB Sciex) with unlabeled compound peak areas normalized to their respective

labeled internal standard peak areas. Labeled standards for indole-3-acetic acid and N-(3-indolylacetyl)-

DL-aspartic acid were obtained from Sigma-Aldrich. Calibration curves were linear (r values = >0.99)

within the ranges provided above and applying 1/x weighting.

RNA Isolation and RT-qPCR

To quantify gene expression in mutants and transgenic plants, total RNAs were isolated using the RNeasy

Plant Mini Kit (Qiaqen, Gaithersburg, MD). cDNA was prepared using SuperScript™ IV VILO™ Master

Mix with ezDNase™ Enzyme (Invitrogen, Carlsbad, CA). RT-qPCR reactions were setup with

SsoAdvanced™ Universal SYBR® Green Supermix (Bio-Rad) and gene specific primers (primer sequences

are listed in Table S4) and performed with the CFX96 qPCR Detection System (Bio-Rad). Statistical

analysis of qRT-PCR data was carried out with REST2009.

Expression of purification of recombinant GFP-WRI1$_{58-240}$

DNA sequence of full length of WRI1 fused with GFP on its N-terminus (*GFP-WRI1*) was amplified from

a OWD5 vector previously described in Zhai et al. (Zhai et al., 2017). *GFP-WRI1* then was inserted into

pet28b between *XhoI* and *NcoI* by in-fusion cloning. Full length of *WRI1* in the resulted *GFP-WRI1*/pet28b

was replaced with DNA sequence corresponding to WRI1 DNA binding domain (WRI1$_{58-240}$) between *XhoI*

and *BsrGI* (Kong et al., 2017). The primer pairs used in building this construct is listed in Table S4.

Recombinant GFP-WRI1$_{58-240}$ was expressed in *E.coli* BL21(DE3). Protein purification was performed as

reported by Zhai (Zhai et al., 2017).

MicroScale Thermophoresis (MST)

Thermophoretic assays were conducted using a Monolith NT.115 apparatus (NanoTemperTechnologies,

South San Francisco, CA; nanotempertech.com). For determining Kds, 8 nM of GFP-WRI1$_{58-240}$ was

incubated with a serial dilution of the ligand (dsDNA) from 1.25 µM to 38.81 pM. Samples of

approximately 10 µl were loaded into capillaries and inserted into the MST instrument loading tray

(Monolith NT.115). The thermophoresis experiments were carried out using 40% MST power and 80%

LED power at 25 °C.

Antibody and Immunoblotting

Anti-BADC1 and anti-histone H3 antibodies were used in this study. Anti-BADC1 antibodies were

requested from Dr. Jay Thelen (Salie et al., 2016). Histone H3 polyclonal antibodies were purchased from

Agrisera (Catalog No. AS10710, Vännäs, Sweden; www.agrisera.com). A total of 50 mg of siliques (7 days

after flowering, DAF) of WT and *wri1-1* mutant plants were ground in liquid nitrogen and then mixed with

200 mL of preheated protein extraction buffer (8 M urea, 2% SDS, 0.1 M DTT, 20% glycerol, 0.1 M Tris-

HCl, pH (6.8), and 0.004% Bromophenol Blue). Samples were centrifuged at 17,000 g before transferring

the supernatant to a new microcentrifuge tube and then loading into SDS-PAGE. 20 ul of supernatant was

loaded each and resolved in SDS-PAGE and transferred onto PVDC membrane for immunoblot analysis.

Primary antibodies anti-BADC1 were used at a 1:3000 dilution, and the anti-H3 antibodies were used at a

1: 5000 dilutions. Immunoblots of targeted proteins were visualized using HRP-conjugated secondary

antibodies (Catalog No. AP187P, Millipore) with SuperSignal^{TM} West Femto Maximum Sensitivity

Substrate (Catalog No. 34095, ThermoFisher). Immunoblot signals were detected and digitalized with

Image Quant LAS4000 and quantified with GelAnalyzer2010a.

Supplemental Figure Legends

Supplemental Figure 1. *BADC1* expression in transgenic *badc1badc2* double mutant overexpressing

BADC1 lines.

Supplemental Figure 2. *badc1badc2* shows similar short-root phenotype as *wri1-1*.

Supplemental Figure 3 Purified recombinant GFP-WRI1$_{58-240}$.

Supplemental Figure 4. Putative AW boxes show varied binding affinity with WRI1.

Supplemental Figure 5. The BADC1 antibody used in this study specifically recognizes BADC1.

Supplemental Figure 6. BADC1 protein levels are lower in the roots of seedlings of *wri1-1* than WT.

Supplemental Figure 7. Both *BADC1* expression and protein levels are elevated in the roots of *WRI1*

inducible expression transgenic plants.

Supplemental Figure 8. BADC1 protein levels in the *BADC1/wri1-1* transgenic line.

Acknowledgements

We thank Dr. Jay Thelen (University of Missouri) for the gift of BADC1 antibody.

Table S1. Quantification of plant hormones in *badc1 badc2* double mutant seedlings.

	ABA	JA	SA	OPDA	JA-Ile	t-Z riboside
WT	2.80±0.45	12.8±4.15	21.06±1.30	238.82±109.75	0.5±0.31	2.63±0.93
badc1 badc2	3.33±0.53	12.2±7.35	26.36±7.43	285.13±140.5	0.35±0.31	1.96±1.90

378   Plant hormones concentration are shown as ng per g DW in this table.

380   **Table S2**. **Nucleotide sequences of AW box in thermophoretic experiment.**

Gene	AW box sequences (5'-3')
BCCP2(-36/-8)	TGGGTTT**C**T**T**CG**T**AAGCAT**CG**AAATCAGA
BCCP2(-143/-115)	CCTACTT**CC**T**CG**GTTTCAT**CG**TCCACTG
BADC1(-169/-142)	CCTTCTT**C**T**TT**GGTAACTC**CG**TCCGCCG
BADC1m1	CCTTCTT**C**T**TT**GGTAACTC**A**TT**CCGCCG
BADC1m2	CCTTCTT**T**T**CT**A**GTAACTC**CG**TCCGCCG
BADC1m3	CCTTCTT**C**T**TT**GGT**GCC**TC**CG**TCCGCCG
BADC2(-102/-75)	TCGTCTT**CC**T**CG**TTCGTAT**CG**TTCACTC
BADC2m1	TCGTCTT**CC**T**CG**TTCGTAT**A**TT**CACTC
BADC2m2	TCGTCTT**T**CC**C**A**TTCGTAT**CG**TTCACTC
BADC2m3	TCGTCTT**CC**T**CG**TTC**AC**AT**CG**TTCACTC
BADC3(-94/-67)	GTTGTGT**CC**T**CG**TTCATAT**CG**TTATCGT

BADC3m1	GTTGTGT**CCT**C**G**TTCATATAT̲TTATCGT
BADC3m2	GTTGTGTT̲CC̲CA̲TTCATAT**CG**TTATCGT
BADC3m3	GTTGTGT**CCT**C**G**TTA̲CATC**G**TTATCGT
BCCP1(-67/-40)	TTGTGCT**C**T**T**GGTTTCAT**CG**AGAGAGA
Neg. C	TGTTACAATCTAGTTACTATTCAGGGTAA
Neg. C1	CATGTTATGGTACTAACAAAAATTTGTTG
Neg. C2	TACTAAATGCTACTTTTTGTTTTGTTTCT

381 Nucleotides in bold are consensus sequence in AW box. Nucleotides underlined are mutated in BADCs

382 AW mutants.

384 **Table S3. Binding affinity of *BADCs* AW boxes and their mutated ones to WRI1 were quantified as**

385 **equilibrium dissociation constants (*kd*) using microscale thermophoresis.**

AW boxex	*Kd* (nM)
BADC1	0.27±0.27
BADC2	2.73±0.72
BADC3	5.24±2.59
BADC1m1	> 1000
BADC2m1	> 1000
BADC3m1	> 1000
BADC1m2	> 1000
BADC2m2	> 1000
BADC3m2	> 1000
BADC1m3	70.67 ± 27.80
BADC2m3	149.18 ± 138.63
BADC3m3	22.88 ±2.94

388 **Table S4**. **Oligonucleotide sequences of primers used in this study.**

Gene	Applications	Primer Pair Sequences (forward and reverse primer, 5'-3')
BADC1	cloning	GTGGGGACAAGTTTGTACAAAAAAGCAGGCTT CATGGCGTCTTCTGCAGCTC and GTGGGGACCACTTTGTACAAGAAAGCTGGGTC TCACTGGATGTTGATGTCGTGG
BADC2	cloning	GTGGGGACAAGTTTGTACAAAAAAGCAGGCTT CATGAATTCCTGTAGCTTAGGAGC and GTGGGGACCACTTTGTACAAGAAAGCTGGGTC TTACTGAAGCTTCTTGATGCCAG
BADC1	qPCR	CAGTTGGGAACAGAACTTCCA and GCAACCAGAGGATCTCCATAAC
BADC2	qPCR	CGACTCTGCGATCTGTGAAA and TGCTACATACAGGCGGAAAC

BADC3	qPCR	CGAATGGATCAGCTTCCTCTAC and GGATCTCCTGAAGAACCCTACT	389
F-box	qPCR	GGCTGAGAGGTTCGAGTGTT and GGCTGTTGCATGACTGAAGA	390
UBQ10	qPCR	ACCATCACTTTGGAGGTGGA and GTCAATGGTGTCGGAGCTTT	391
GFP-WRI1	Cloning	GGTGGTGGTGCTCGAGTTCAGAACCAACGAAGCC and	392
		CAAACAAGTTTGTACACTAGTATGGCT TCTACCCGACG	393
WRI1$_{58}$ $_{-240}$	Cloning	GGTGGTGGTGCTCGAGCGGGAAAACACCTTTCTTTA AC and	394
		CAAACAAGTTTGTACACTAGTATGGCTTCTACCCGACG	395
			396

References

Andre C, Haslam RP, Shanklin J (2012) Feedback regulation of plastidic acetyl-CoA carboxylase by 18: 1-
acyl carrier protein in Brassica napus. Proceedings of the National Academy of Sciences **109:**
10107-10112
Baud S, Mendoza MS, To A, Harscoët E, Lepiniec L, Dubreucq B (2007) WRINKLED1 specifies the
regulatory action of LEAFY COTYLEDON2 towards fatty acid metabolism during seed maturation
in Arabidopsis. The Plant Journal **50:** 825-838
Braybrook SA, Stone SL, Park S, Bui AQ, Le BH, Fischer RL, Goldberg RB, Harada JJ (2006) Genes directly
regulated by LEAFY COTYLEDON2 provide insight into the control of embryo maturation and
somatic embryogenesis. Proceedings of the National Academy of Sciences **103:** 3468-3473
Focks N, Benning C (1998) wrinkled1: a novel, low-seed-oil mutant of Arabidopsis with a deficiency in the
seed-specific regulation of carbohydrate metabolism. Plant physiology **118:** 91-101

Fukuda N, Ikawa Y, Aoyagi T, Kozaki A (2013) Expression of the genes coding for plastidic acetyl-CoA
carboxylase subunits is regulated by a location-sensitive transcription factor binding site. Plant
molecular biology **82:** 473-483
Grimberg Å, Carlsson AS, Marttila S, Bhalerao R, Hofvander P (2015) Transcriptional transitions in
Nicotiana benthamiana leaves upon induction of oil synthesis by WRINKLED1 homologs from
diverse species and tissues. BMC plant biology **15:** 192
Kagaya Y, Okuda R, Ban A, Toyoshima R, Tsutsumida K, Usui H, Yamamoto A, Hattori T (2005) Indirect
ABA-dependent regulation of seed storage protein genes by FUSCA3 transcription factor in
Arabidopsis. Plant and Cell Physiology **46:** 300-311
Keereetaweep J, Liu H, Zhai Z, Shanklin J (2018) Biotin Attachment Domain-Containing Proteins
Irreversibly Inhibit Acetyl CoA Carboxylase. Plant physiology**:** pp. 00216.02018
Kong Q, Ma W, Yang H, Ma G, Mantyla JJ, Benning C (2017) The Arabidopsis WRINKLED1 transcription
factor affects auxin homeostasis in roots. Journal of experimental botany **68:** 4627-4634
**Li-Beisson Y, Shorrosh B, Beisson F, Andersson MX, Arondel V, Bates PD, Baud S, Bird D, DeBono A,
Durrett TP** (2013) Acyl-lipid metabolism. The arabidopsis book**:** e0161
Li Q, Shao J, Tang S, Shen Q, Wang T, Chen W, Hong Y (2015) Wrinkled1 accelerates flowering and
regulates lipid homeostasis between oil accumulation and membrane lipid anabolism in Brassica
napus. Frontiers in plant science **6:** 1015
**Lotan T, Ohto M-a, Yee KM, West MA, Lo R, Kwong RW, Yamagishi K, Fischer RL, Goldberg RB, Harada
JJ** (1998) Arabidopsis LEAFY COTYLEDON1 is sufficient to induce embryo development in
vegetative cells. Cell **93:** 1195-1205
Maeo K, Tokuda T, Ayame A, Mitsui N, Kawai T, Tsukagoshi H, Ishiguro S, Nakamura K (2009) An AP2-
type transcription factor, WRINKLED1, of Arabidopsis thaliana binds to the AW-box sequence
conserved among proximal upstream regions of genes involved in fatty acid synthesis. The Plant
Journal **60:** 476-487
Mendoza MS, Dubreucq B, Miquel M, Caboche M, Lepiniec L (2005) LEAFY COTYLEDON 2 activation is
sufficient to trigger the accumulation of oil and seed specific mRNAs in Arabidopsis leaves. FEBS
letters **579:** 4666-4670
Mu J, Tan H, Zheng Q, Fu F, Liang Y, Zhang J, Yang X, Wang T, Chong K, Wang X-J (2008) LEAFY
COTYLEDON1 is a key regulator of fatty acid biosynthesis in Arabidopsis. Plant physiology **148:**
1042-1054
Ohlrogge J, Browse J (1995) Lipid biosynthesis. The Plant Cell **7:** 957
Olinares PDB, Ponnala L, van Wijk KJ (2010) Megadalton complexes in the chloroplast stroma of
Arabidopsis thaliana characterized by size exclusion chromatography, mass spectrometry, and
hierarchical clustering. Molecular & Cellular Proteomics **9:** 1594-1615
Parcy F, Valon C, Kohara A, Miséra S, Giraudat J (1997) The ABSCISIC ACID-INSENSITIVE3, FUSCA3, and
LEAFY COTYLEDON1 loci act in concert to control multiple aspects of Arabidopsis seed
development. The Plant Cell **9:** 1265-1277
Rawsthorne S (2002) Carbon flux and fatty acid synthesis in plants. Progress in lipid research **41:** 182-196
Ruuska SA, Girke T, Benning C, Ohlrogge JB (2002) Contrapuntal networks of gene expression during
Arabidopsis seed filling. The Plant Cell **14:** 1191-1206
Salie MJ, Zhang N, Lancikova V, Xu D, Thelen JJ (2016) A family of negative regulators targets the
committed step of de novo fatty acid biosynthesis. The Plant Cell**:** tpc. 00317.02016
To A, Valon C, Savino G, Guilleminot J, Devic M, Giraudat J, Parcy F (2006) A network of local and
redundant gene regulation governs Arabidopsis seed maturation. The Plant Cell **18:** 1642-1651
Zhai Z, Keereetaweep J, Liu H, Feil R, Lunn JE, Shanklin J (2018) Trehalose 6-phosphate positively
regulates fatty acid biosynthesis by stabilizing WRINKLED1. The Plant Cell**:** tpc. 00521.02018

Zhai Z, Liu H, Shanklin J (2017) Phosphorylation of WRINKLED1 by KIN10 results in its proteasomal
degradation, providing a link between energy homeostasis and Lipid biosynthesis. The Plant Cell
29: 871-889

Figure Legends

Figure 1. *badc1 badc2* double mutant shows reduced primary root growth and less lateral roots number

than wild type (WT). A) 9-day-old WT, *badc1*, *badc2*, and *badc1 badc2* seedlings grown vertically on

1/2MS media supplemented with 1% of sucrose. B) Primary root length measurement of each genotype in

A. C) Lateral root numbers for each genotype in A. Values in B and C represent means±SD from 10

individual plants for each genotype. D) Roots growth and E) primary roots length of 20-day-old plants of

WT, *badc1 badc2*, and representative transgenic *badc1 badc2* overexpressing *BADC1* or *BADC2* lines.

Values in E are means±SD from 10 individual plants for each indicated genotype. In the figure, Levels

indicated with different letters above histogram bars are significantly different (Student's T test for all pairs

of genotypes, P<0.01). Data in this figure is a representative of 3 independent repetitions.

Figure 2. IAA-Asp content in *badc1badc2* double mutant is significantly higher than that in WT. A)

Quantification of IAA in 7-day-old seedling of WT and *badc1badc2*. B) Quantification of IAA-Asp in WT

and *badc1badc2*. Values in this figure are means ±SD (n=4) for each sample of 30 seedlings. Asterisks

denote statistically significant difference from WT (Student's T test, **, P<0.01). Data in this figure is a

representative of 2 independent repetitions.

Figure 3. Purified WRI1 shows specific binding with AW boxes in the promoters of BADCs. In the

thermophoresis experiment, A 28bp-oligonucleotide containing putative AW box for each gene (A) was

titrated against purified WRI1 DNA binding domain (AA58-240). Dissociation curves for each pair are

shown (B) and fitted to the data for calculation of the *Kd* values (C). An AW box in the promoter of *BCCP2*

was used as a positive control. A random sequence is used as a negative control (Neg.C). AW box consensus

sequences are marked in bold. Data shown are the mean±SD, n=3 independent repetitions.

Figure 4. *BADCs* expression level and BADC1 protein level are significantly lower in *wri1-1* mutant

developing siliques than in WT. (A) Tissue expression pattern of *WRI1* and *BADCs* in flowers, siliques (7

days after flowering), seedling roots (SR, 10 days old) and mature leaves (ML). (B) RT-qPCR results of

BADC1, *BADC2*, and *BADC3* expression levels in siliques (7 days after flowering) of WT and *wri1-1*

mutant. Values are means±SD from three independent experiments. Expression of each gene in WT sets to

1. For each experiment, total RNA was isolated from pooled siliques from WT and *wri1-1* plants. *Fbox*

and *UBQ10* were used as reference genes. Asterisk denotes statistically significant difference from WT

(using mean crossing point deviation analysis computed by the relative expression (REST) software

algorithm, **, $P<0.01$). (C) BADC1 protein levels in siliques (7 days after flowering) of WT or *wri1-1*

mutant are shown by immunoblot with BADC1 specific antibody. Siliques were collected from 3

independent plants of WT or *wri1-1* mutant. Protein loading is shown by histone H3 in the same protein

samples. (D) The histogram shows relative BADC1 protein levels in (C) quantified with GelAnalyzer2010

and normalized against corresponding protein levels of histone H3. Asterisk denotes statistically significant

difference from the WT (Student's T test, *, $P<0.05$). Data in this figure is a representative of 3 independent

repetitions.

Figure 5. *BADCs* gene expression and BADC1 protein levels are elevated in inducible *WRI1* transgenic

plants. (A) Relative gene expression of *BADCs, BCCP2* and *WRI1* in siliques (7 days after flowering) of

WT or 2 independent ethanol inducible *WRI1* transgenic lines (AlcA:*WRI1* line1 and line4) treated (+) or

not treated (−) with 2% ethanol for 4 days. (B) BADC1 protein levels. (C) The histogram shows relative

BADC1 protein levels in (B) quantified with GelAnalyzer2010 and normalized against corresponding

protein levels of histone H3. Asterisk denotes statistically significant difference from the WT (Student's T

test, *, $P<0.05$). Data in this figure is a representative of 3 independent repetitions.

Figure 6. Overexpression of *BADC1* in *wri1-1* partially rescues *wri1-1* short-root phenotype. (A) Roots

growth and B) Primary roots length measurement of 16-day-old WT, *wri1-1* and representative transgenic

wri1-1 overexpressing *BADC1* lines. Values in (B) are means±SD from 10 individual plants for each

indicated genotype. Asterisk denotes statistically significant difference from *wri1-1* mutant (Student's T

test, *, $P<0.05$). (C) IAA-Asp was quantified in the 7-day-old seedlings of WT, *wri1-1* and *BADC1/wri1-1*

transgenic line 1 and 2. Data in this figure is a representative of 3 independent repetitions.

Figure 1. *badc1 badc2* double mutant shows reduced primary root growth and less lateral roots number than wild type (WT). (A) 9-day-old WT, *badc1*, *badc2*, and *badc1 badc2* seedlings grown vertically on 1/2MS media supplemented with 1% of sucrose. (B) Primary root length measurement of each genotype in (A). (C) Lateral root numbers for each genotype in (A). Values in (B) and (C) represent means±SD from 10 individual plants for each genotype. (D) Roots growth and (E) primary roots length of 20-day-old plants of WT, *badc1 badc2*, and representative transgenic *badc1 badc2* overexpressing *BADC1* or *BADC2* lines. Values in (E) are means±SD from 10 individual plants for each indicated genotype. In the figure, Levels indicated with different letters above histogram bars are significantly different (Student's T test for all pairs of genotypes, P<0.01). Data in this figure is a representative of 3 independent repetitions.

Figure 2. IAA-Asp content in *badc1badc2* double mutant is significantly higher than that in WT. (A) Quantification of IAA in 7-day-old seedling of WT and *badc1badc2*. (B) Quantification of IAA-Asp in WT and *badc1badc2*. Values in this figure are means ±SD (n=4) for each sample of 30 seedlings. Asterisks denote statistically significant difference from WT (Student's T test, **, P<0.01). Data in this figure is a representative of 2 independent repetitions.

A	BCCP2$_{(-143/-115)}$	TGGGTTT**CTTCG**TAAGCAT**CG**AAATCAG
	BADC1$_{(-169/-142)}$	CCTTCTT**CTTTG**GTAACTC**CG**TCCGCCG
	BADC2$_{(-102/-75)}$	TCGTCTT**CCTCG**TTCGTAT**CG**TTCACTC
	BADC3$_{(-94/-67)}$	GTTGTGT**CCTCG**TTCATAT**CG**TTATCGT
	Neg. C	TGTTACAATCTAGTTACTATTCAGGGTA

C

Target	Ligand	Kd (nM)
	BCCP2	0.65±0.39
	BADC1	0.27±0.27
GFP-WRI1$_{58-240}$	***BADC2***	2.73±0.72
	BADC3	5.24±2.59
	Neg.C	—

Figure 3. Purified WRI1 shows specific binding with AW boxes in the promoters of BADCs. In the thermophoresis experiment, A 28bp-oligonucleotide containing putative AW box for each gene (A) was titrated against purified WRI1 DNA binding domain (AA58-240). Dissociation curves for each pair are shown (B) and fitted to the data for calculation of the Kd values (C). An AW box in the promoter of *BCCP2* was used as a positive control. A random sequence is used as a negative control (Neg.C). AW box consensus sequences are marked in bold. Data shown are the mean±SD, n=3 independent repetitions.

Figure 4. *BADCs* expression level and BADC1 protein level are significantly lower in *wri1-1* mutant developing siliques than in WT. (A) Tissue expression pattern of *WRI1* and *BADCs* in flowers, siliques (7 days after flowering), seedling roots (SR, 10 days old) and mature leaves (ML). (B) RT-qPCR results of *BADC1*, *BADC2*, and *BADC3* expression levels in siliques (7 days after flowering) of WT and *wri1-1* mutant. Values are means±SD from three independent experiments. Expression of each gene in WT sets to 1. For each experiment, total RNA was isolated from pooled siliques from WT and *wri1-1* plants. *Fbox* and *UBQ10* were used as reference genes. Asterisk denotes statistically significant difference from WT (using mean crossing point deviation analysis computed by the relative expression (REST) software algorithm, **, P<0.01). (C) BADC1 protein levels in siliques (7 days after flowering) of WT or *wri1-1* mutant are shown by immunoblot with BADC1 specific antibody. Siliques were collected from 3 independent plants of WT or *wri1-1* mutant. Protein loading is shown by histone H3 in the same protein samples. (D) The histogram shows relative BADC1 protein levels in (C) quantified with GelAnalyzer2010 and normalized against corresponding protein levels of histone H3. Asterisk denotes statistically significant difference from the WT (Student's T test, *, P<0.05). Data in this figure is a representative of 3 independent repetitions.

Figure 5. *BADCs* gene expression and BADC1 protein levels are elevated in inducible *WRI1* transgenic plants. (A) Relative gene expression of *BADCs, BCCP2* and *WRI1* in siliques (7 days after flowering) of WT or 2 independent ethanol inducible *WRI1* transgenic lines (AlcA:*WRI1* line1 and line4) treated (+) or not treated (−) with 2% ethanol for 4 days. (B) BADC1 protein levels. (C) The histogram shows relative BADC1 protein levels in (B) quantified with GelAnalyzer2010 and normalized against corresponding protein levels of histone H3. Asterisk denotes statistically significant difference from the WT (Student's T test, *, P<0.05). Data in this figure is a representative of 3 independent repetitions.

Figure 6. Overexpression of *BADC1* in *wri1-1* partially rescues *wri1-1* short-root phenotype. (A) Roots growth and B) Primary roots length measurement of 16-day-old WT, *wri1-1* and representative transgenic *wri1-1* overexpressing *BADC1* lines. Values in (B) are means±SD from 10 individual plants for each indicated genotype. Asterisk denotes statistically significant difference from *wri1-1* mutant (Student's T test, *, P<0.05). (C) IAA-Asp was quantified in the 7-day-old seedlings of WT, *wri1-1* and *BADC1/wri1-1* transgenic line 1 and 2. Asterisk denotes statistically significant difference from *wri1-1* mutant (Student's T test, *, P<0.05). Data in this figure is a representative of 3 independent repetitions.