Complex community-wide consequences of consumer sexual dimorphism

Stephen P. De Lisle [1,2] *

Sebastian J. Schrieber [3]

Daniel I. Bolnick [1]

[1] Department of Ecology & Evolutionary Biology, University of Connecticut, Storrs, CT

06269

[2] Present Address: Evolutionary Ecology Unit, Department of Biology, Lund University,

Sölvegatan 37, 22362 Lund Sweden

[3] Department of Evolution and Ecology and Center for Population Biology, University of

California, Davis, CA 95616

* email: stephen.de_lisle@biol.lu.se phone: +46 70 892 8731

Running Title: Community effects of sexual dimorphism

Keywords: apparent competition, community assembly, competitive exclusion, ecological

sexual dimorphism, resource competition

Type of Article: Letter

Word Count: 148 (Abstract) 4739 (Main text)

References: 58

Figures: 5 **Tables:** 1

Statement of authorship: DB SD and SJS designed the study, SJS performed the

mathematical analysis, SD performed the simulations and drafted the manuscript. All authors

revised the manuscript.

Data accessibility statement: No data is used

Abstract

Sexual dimorphism is a ubiquitous source of within-species variation, yet the community-level consequences of sex differences remain poorly understood. Here, we analyze a bitrophic model of two competing resource species and a sexually-reproducing consumer species. We show that consumer sex differences in resource acquisition can have striking consequences for consumer-resource coexistence, abundance, and dynamics. Under both direct interspecific competition and apparent competition between two resource species, sexual dimorphism in consumers' attack rates can mediate coexistence of the resource species, while in other cases can lead to exclusion when stable coexistence is typically expected. Slight sex differences in total resource acquisition also can reverse competitive outcomes and lead to density cycles. These effects are expected whenever both consumer sexes require different amounts or types of resources to reproduce. Our results suggest that consumer sexual dimorphism, which is common, has wide-reaching implications for the assembly and dynamics of natural communities.

Introduction

Within-species variation is one key feature of natural populations that has emerged as a critical contributor to community ecology and species interactions (Hughes et al. 2008 , Bolnick et al. 2011, Des Roches et al. 2018). Even in the simplest scenario of a single-species community with a single optimum phenotype, within-population variation is expected to reduce population growth rate of a well-adapted population via effects on mean fitness (Haldane 1937). In more complex multi-species communities, phenotypic variation in consumers or their resources can either promote or constrain coexistence between competing species, under many circumstances yielding different conclusions than would be reached from models ignoring within species variation (Bolnick et al. 2011, Schreiber et al. 2011, Patel and Schreiber 2015, Cortez and Patel 2017). One emerging question (Bolnick et al. 2011) from this work is the degree to which understanding the specific source of phenotypic variation matters in community ecology.

Here, we explore the ecological consequences of sexual dimorphism, a central feature of metazoans and many plant populations that has not been fully incorporated into ecological theory. The gamete differences that define the sexes are expected to lead to divergence between males and females in a suite of life history traits (Trivers 1972, Shärer et al. 2012), and phenotypic traits. This divergence is iconically manifest as striking sexual dimorphism in sexually-selected traits such as body size or courtship displays (Darwin 1871, Andersson 1994). Although sexual selection and associated mate choice behaviors themselves may have relevance to interspecific ecological interactions (Gomez-Llano et al. 2018), the different life histories that define the sexes are often expected lead to different nutritional and resource requirements for males and females (Maklakov et al. 2008). As a specific example, in insects, differential contributions of longevity and fecundity to male and female lifetime reproductive success result in different combinations of macronutrients that maximize male

and female fitness (Maklakov et al. 2008, Reddiex et al. 2013, Garlapow et al. 2015, Jensen

et al. 2015, Camus et al. 2017). Similarly, nutritional models of optimal foraging have been

proposed to explain sex differences in diets of moose (Belovksy 1978). Consequently, males

and females of many species have evolved divergent resource use, either as an indirect

outcome of divergent reproductive roles or through other forms of sex-specific natural

selection, resulting in sex-differences in diet composition and often in corresponding trophic

morphology (Slatkin 1984, Temeles 1985, Shine 1989, Temeles et al. 2000, De Lisle and

Rowe 2015a, De Lisle 2019).

Data suggest that these 'ecological' sexual dimorphisms can have substantial

consequences for community structure (Fryxell et al. 2015, Pincheira-Donoso et al. 2018,

Start and De Lisle 2018, Tsuji and Fukami 2018, 2020). More generally, sex differences may

play an important role in the relationship between ecological and evolutionary dynamics

(Giery and Layman 2019, Svensson 2019, Fryxell et al. 2019). Sexual dimorphism thus

represents a key source of ecologically-relevant variation within species. Within-species

variation in resource specialization is commonplace (Bolnick et al. 2003), with important

consequences for species interactions and community assembly (Hughes et al. 2008 , Bolnick

et al. 2011, Des Roches et al. 2018). Critically, as a source of intraspecific variation, sexual

dimorphism may have different consequences from other types of variation. Unlike most

phenotypes, whose relative abundances can evolve to reflect local ratios of alternative

resources, the ratio of males to females (at birth) is expected to be maintained at 1:1 (Darwin

1871, Fisher 1930), though their phenotypes and demography may diverge to have

differential ecological impacts. Thus, the rate at which males are born (which determines

predation pressure on males' prey), is strongly dependent on females' foraging success.

Perhaps to a lesser extent (depending on mating system), the rate of female offspring

production can depend on male foraging success. Importantly, persistence of a sexually

reproducing consumer depends on the persistence of both sexes. This coupling of the

dynamics of two consumer phenotypes has unknown consequences for consumer-resource

dynamics, consumer mediated coexistence of the resource species, and apparent competition

between the resource species. Using a simple model of consumer-resource dynamics, we

show that consumer sexual dimorphism can influence (both positively and negatively)

coexistence between resource species (competing directly or apparently), and species'

abundance.

109   Materials and Methods

Our model represents a simple extension of a classic model of consumer-resource dynamics,

in which a consumer species exploits two resources that themselves undergo density-

dependent growth (Figure 1). Consumer growth rate is limited only by resource abundance.

The original version of this model did not consider within-species phenotypic variation, and

can lead to exclusion of one resource ('apparent competition'; (e.g. Holt 1977). More

recently, (Schreiber et al. 2011) showed that quantitative trait variation in the predator,

affecting attack rates on the two resources, can facilitate coexistence between the prey. Here,

we instead allow for the possibility that male and female consumers differ in resource-

specific attack rates. We describe the population dynamics of two resources (with densities

R_i, R_j) and consumer males (with density M) and females (with density F) using the system

of ordinary differential equations:

$$\frac{dR_i}{dt} = r_i R_i\big(1 - \alpha_{ii} R_i - \alpha_{ij} R_j\big) - a_{M,i} R_i M - a_{F,i} R_i F \text{ for } j \neq i \qquad (1a)$$

$$\frac{dM}{dt} = \frac{1}{2} B(M, F, R_1, R_2) - d_M M \qquad (1b)$$

$$\frac{dF}{dt} = \frac{1}{2} B(M, F, R_1, R_2) - d_F F. \tag{1c}$$

Each resource undergoes density-dependent population growth with intrinsic per-capita

growth rates r, with competition represented by competition coefficients α. Resources are

also regulated by consumers depending on sex-specific attack rates by males a_M and by

females a_F. Consumer dynamics are governed by females' birth rate B (half of the offspring

being female), and limited by sex-specific intrinsic mortality rates d_M and d_F.

An appropriate function describing birth rates has been a point of debate for

demographers, with the general conclusion being that any function describing birth rates

should capture the negative effects of extreme sex ratio skew on birth rates, with the extreme

being that birth rates should be zero when one sex is absent (Caswell and Weeks 1986), or

unable to breed. Generally, treating birth rates proportional to the harmonic mean density of

males and females is agreed as the best approach (Caswell and Weeks 1986, Lindstöm and

Kokko 1998), and is also empirically supported (Miller and Inouye 2011). Here, we extend

this harmonic mean birth function to account for each sex's foraging success, on the logic

that each sex must both be present and sufficiently well-fed to reproduce:

$$B(M, F, R_i, R_j) = 2 \frac{MI_M * FI_F}{MI_M + FI_F} \tag{2}$$

139   where I is the energy intake of a given sex:

$$I_F = b_F(a_{F,1}R_1 + a_{F,2}R_2) \text{ and } I_M = b_M(a_{M,1}R_1 + a_{M,2}R_2) \tag{3}$$

which couples the dynamics of consumers and resources. In this model the rate of consumer

births depends on not just the number of males and females but also on the abundance of both

prey and the prey preference of each sex, where MI_M is the number of males weighted by

their energy to reproduce (similarly with females), b is a scaling constant reflecting the

degree to which sex-specific resource acquisition influences birth rate, and $a_{M,i}, a_{F,i}$ are the

attack rates on the resource i. Although we make no genetic assumptions, we note that our

birth function, and thus our model, applies to sexually-reproducing diploid consumers. As an

alternative to the inclusion of constants b_M and b_F, equation 2 can be equivalently expressed

with the inclusion of a "harem size" parameter, h (Caswell and Weeks 1986, Lindstöm and

Kokko 1998; see supplemental material); changes in b_M relative to b_F alter the degree to

which each sex contributes to birth rates in an equivalent manner to the effects of deviations

of h from unity. We focus on the parameterization in equation 2 because it illustrates that the

mathematical/demographic effects of "harem size" can in fact be brought about by any factor

that alters the relative contribution of sex-specific density to birth rates.

Because we are specifically interested in understanding effects of sexual dimorphism

in prey preference, we relate male and female resource-specific attack rates via the degree of

sexual dimorphism, β, such that

$$a_{M,1} = a_{M,max}\left(\frac{1}{2} + \frac{\beta}{2}\right) \text{ and } a_{F,1} = a_{F,max}\left(\frac{1}{2} - \frac{\beta}{2}\right) \tag{4a}$$

$$a_{M,2} = a_{M,max}\left(\frac{1}{2} - \frac{\beta}{2}\right) \text{ and } a_{F,2} = a_{F,max}\left(\frac{1}{2} + \frac{\beta}{2}\right) \tag{4b}$$

where β ranges from zero to one, with $\beta = 0$ representing complete sexual monomorphism if

maximum attack rates are equal across the sexes (both sexes are generalists, attacking each

prey at the rates, $a_{F,max}/2$ and $a_{M,max}/2$), and $\beta = 1$ representing complete sexual

dimorphism such that males only attack R_1 and females only attack R_2. Sex differences in

a_{max} reflect a situation where males and females intake different total amounts of resources,

independent of any difference in resource preference. In our study we explore both types of

sex differences in resource acquisition; sexual dimorphism in resource preference captured by

β, as well as sex differences in total resource acquisition captured by sex-specific a_{max}.

By assuming that resource acquisition of males and females contributes to the

consumer birth rate, we are not necessarily making any assumption about parental care or

condition-dependent fecundity. Rather, this is an assumption that the fraction of each sex

available for reproduction depends in part on resource acquisition, a realistic feature of many

organisms. For instance, males' energetic costs of finding a mate, defending a territory, care

for offspring, or expressing a sexually-selected trait are all expected to depend on the

resource pool available to males (Rowe and Houle 1996). Females' ability to produce eggs,

gestate, and provision young (e.g., lactation in mammals) similarly depends on their energy

intake. Parental care and other factors can also lead to associations between sex-specific

resource abundance and consumer birth rates, beyond what would be expected beyond the

costs of simply being available to mate.

We analyzed this model to find equilibria when both one or two resources are present,

as well as invasion criteria for a resource into a two species community at equilibrium. Using

the mathematical theory of permanence (Schreiber 2000, Patel & Schreiber 2018), we can use

these invasion criteria to determine whether all three species coexist (in the sense of

permanence), exhibit a bistability, or competitive exclusion occurs. In addition to presenting

our analytical results below, we provide further mathematical details in Supplement A. We

also numerically explored the behavior of the model under a variety of scenarios using the

packages deSolve v. 1.21 (Soetgart et al. 2010) and caTools v. 1.17.1.2 (Tuszynski 2019) in R

v. 3.5.0 (R Core Team 2018). Importantly, using simulations allowed us to explore the

behavior of our model for cases where 2-species consumer-resource equilibria do not exist.

Because there are a large number of potential combinations of parameters that could be

explored, we focused our simulations on three different biological scenarios: 1) completely

symmetric male and female total attack rates and contributions to birth rate, 2) asymmetric

total attack rates across the sexes, and 3) asymmetric contribution of male and female

resource acquisition to consumer birth rates. The first scenario applies most readily to

organisms with biparental care and similar total caloric requirements across the sexes. The

second scenario likely applies to many organisms where total lifetime resource acquisition is

higher for one sex, which is the case for many organisms where the sexes differ, for example,

in body size. The third scenario is another realistic departure from the first, relevant for

organisms where resource acquisition in one sex (e.g., females) has a greater influence on

that sexes mating propensity, or cases with biased operational sex ratios due to polyandry or

polygyny. For each scenario we explore the consequences of consumer ecological sexual

dimorphism (β) for consumer persistence and coexistence. For each set of parameter values,

we simulated the model 1,000 time steps (to ensure the simulation reaches its equilibrium).

For each run, we determined the equilibrium abundances of M, F, R_1 and R_2 at t = 1,000. We

also calculated the standard deviation of species' abundances in the last 50 time-steps.

Although we present solutions simulated on the scale of α_{ii} = 0.1, we obtained qualitatively

equivalent conclusions rescaling under a wide range values of competition coefficients (see

Table S1B, Supplement B). The R script to generate all figures and results presented is

available in the supplementary material.

Results

Analytical results

Our analysis begins by considering a subsystem of a single resource species, say species i,

and the consumer species. That is, we explore the conditions under which a sexually

dimorphic consumer can persist on a single resource species. These two species can coexist

if

$$H_i \frac{1}{\alpha_{ii}} > 2 \tag{5}$$

where $H_i := 2\dfrac{\frac{b_M a_{i,M}}{d_M}\frac{b_F a_{i,F}}{d_F}}{\frac{b_M a_{i,M}}{d_M}+\frac{b_F a_{i,F}}{d_F}}$ is the harmonic mean of the lifetime per-capita resource

contributions of each sex to reproduction, and $\dfrac{1}{\alpha_{ii}}$ corresponds to the carrying capacity of

resource species i. Intuitively, $H_i\dfrac{1}{\alpha_{ii}}$ corresponds to the average number of offspring produced

by a mating pair during their lifetime when the resource species is at its carrying capacity.

These harmonic means decrease to zero with the degree of sexual dimorphism (i.e. H_i is a

decreasing function of β and $H_i=0$ when $\beta = 1$), there always is a critical degree of sexual

dimorphism above which the consumer cannot persist on a single resource species.

Intuitively, this arises when one of the sexes specializes too much on the other (absent)

resource species and, consequently, contributes too little to reproduction. When the

coexistence condition holds, the consumer-resource species pair coexist at the following

equilibrium densities:

227
$$R_i^* = 2/H_i$$

228
$$F_i^* = \frac{r_i\left(1-\frac{2\alpha_{ii}}{H_i}\right)}{a_{i,F}+\frac{a_{i,M}d_F}{d_M}} \tag{6}$$

229
$$M_i^* := \frac{d_F}{d_M}F_i^*$$

230 At these densities of the consumer and resource i, the second (rare) resource species j can

231 invade if its per capita growth rate

232
$$I_j = r_j\left(1 - \frac{2\alpha_{ji}}{H_i}\right) - A_jF_i^* \tag{7}$$

is positive. Here, $A_j = a_{F,j} + a_{M,j}d_F/d_M$ is the average attack rate on resource j when the

males and females are at an equilibrium. Provided the consumer species can persist on each

of the resource species individually, the invasion growth rates I_1 and I_2 determine the

ecological outcomes. If both invasion growth rates are positive (mutual invasibility), all three

species coexist. If both invasion rates are negative, there is a priority effect in which both

single resource-consumer equilibria are stable. If one invasion growth rate is positive and the

other is negative, this suggests (as confirmed by numerical simulations) that the resource

species with the positive invasion growth rate excludes the other resource species. In the

supplementary material (Supplement A), we provide more details about this analysis and

describe the invasion conditions for these outcomes or when the consumer only persists in the

presence of both resource species.

As a step toward understanding the general conditions for coexistence and exclusion,

we first examine two special cases corresponding to (i) resource competitive symmetry

whereby $\alpha_{11} = \alpha_{12} = \alpha_{21} = \alpha_{22}$ and (ii) pure apparent competition whereby $\alpha_{12} = \alpha_{21} =$

0 . When there is competitive symmetry, the resource species with the larger value of r_i/A_i

excludes the other resource species. The quantity r_i/A_i corresponds to the consumer density

supported by resource species i when intraspecific competition is very weak i.e. $\alpha_{ii} = 0$.

Hence, the resource species that supports the highest consumer equilibrium density (in the

absence of self-limitation) excludes the other. This is an analog of the P^* rule (Holt 1977,

Holt and Lawton 1993, Schreiber 2021). In this case, sexual dimorphism (as measured by β)

influences outcomes only if there is an asymmetry in the maximal attack rates of the female

and male (e.g., female masked boobies dive for prey more often than males; Weimerskirch et

al. 2009). For example, if the female-preferred resource (R_1) has the higher intrinsic rate of

growth ($r_1 > r_2$) and the female consumer has the higher maximal attack rate of the two

sexes, then the female-preferred resource species excludes the other resource species when

sexual dimorphism is low (i.e. $A_1=A_2$ when $\beta = 0$). However, as the average attack rate A_1

on the female-preferred resource increases with the magnitude β of the sexual dimorphism,

sexual dimorphism reverses the outcome of apparent competition whenever $r_1/a_{F,max} <$

$r_2/a_{M,max}$ and the self-limitation in the female-preferred resource is weak.

Next, we consider the case of pure apparent competition in which there is no direct

competition between the resources, i.e. $\alpha_{12} = \alpha_{21} = 0$. Then, coexistence occurs when

$$\frac{1}{1-2\alpha_{ii}/H_i} > \frac{r_i/A_i}{r_j/A_j} \text{ for } i \neq j \qquad (8)$$

The right-hand side of this condition determines the outcome of apparent competition in the

case of competitive symmetry i.e. the species with larger r_i/A_i wins. The left-hand increases

to infinity as the sexual dimorphism increases such that H_i approaches $2\alpha_{ii}$. This has two

implications: (i) when r_1/A_1 and r_2/A_2 are equal, the resource species coexist at any level of

sexual dimorphism, and (ii) when r_1/A_1 and r_2/A_2 are unequal, sufficiently strong sexual

dimorphisms (that still allow the consumer to persist, per equation 5) will ensure coexistence

of the two resource species by diluting the strength of apparent competition.

Finally, the general coexistence condition is

$$\frac{1-2\alpha_{ji}/H_i}{1-2\alpha_{ii}/H_i} > \frac{r_i/A_i}{r_j/A_j} \text{ for } i \neq j \qquad (9)$$

The effect of sexual dimorphism on the left-hand side term depends on the relative strengths

of intra- and interspecific competition. When intraspecific competition is greater than

interspecific competition (i.e. $\alpha_{ii} > \alpha_{ji}$), the left-hand side term increases to infinity with

increasing sexual dimorphism. In contrast, when interspecific competition between resources

is greater than intraspecific competition (i.e. $\alpha_{ji} > \alpha_{ii}$), the left-hand side term of the general

coexistence condition decreases to zero with increasing levels of sexual dimorphism. The

effect of sexual dimorphism on the right-hand side term of the general coexistence criterion is

as discussed for the case of competitive symmetry. Together these observations imply that if

resource 1 is the better direct competitor i.e. $\alpha_{21} > \alpha_{11}$ and $\alpha_{21} < \alpha_{22}$, then coexistence

requires that resource 2 is the better apparent competitor i.e. $r_2/A_2 > r_1/A_1$. In which case, a

sexual dimorphism can help consumer-mediated coexistence.

Numerical results: Symmetric resource acquisition across the sexes

Our numerical results generally matched conclusion from our analytical solutions and are

summarized in Table 1. The effects of consumer sexual dimorphism on resource population

dynamics are illustrated in Figure 2. Sexual dimorphism results in increased resource density

(Fig 2B vs. 2D; Fig 3), reduced consumer density (Figure 3), and can facilitate persistence of

a competitively inferior resource species (Fig 2A vs. 2B; note that this contrast is determined

by persistence of the consumer, which is possible under the conditions in 2B). When

competition is purely apparent and resource acquisition by the consumer is symmetric across

the sexes, increasing consumer sexual dimorphism increases the parameter space under

which resources can coexist (Fig. 2C vs 2D), consistent with our analytical results (Fig. 4A).

In particular, as sexual dimorphism approaches the extreme (each sex uses a different

resource exclusively), competitively inferior resource species can coexist with a superior

competitor that is more susceptible to apparent competition (Figure 4). This effect is in part

due to the fact that consumer persistence depends on the density of both resources at this

extreme.

When resources compete directly with symmetric intraspecific competition, resource

acquisition is symmetric across the sexes and intrinsic per-capita growth rates are equal

between resources, consumer sexual dimorphism has little effect on invasion of a resource

into a consumer-single-resource community at equilibrium (S1A-C, S4A). Yet even in the

absence of effects on coexistence consumer sexual dimorphism has strong effects on

equilibrium resource abundance, both under analytical solutions to a two species community

and under simulations of three species communities (Fig 3). As predicted by our analytical

conditions, when resource growth rates are unequal, sexual dimorphism changes expected

regions of coexistence of both resources and the consumer, leading to invasion and

persistence of competitively inferior resource that is a superior apparent competitor i.e. has a

larger value of r_i/A_i (Figure 5B, Figure S4B, Figure S1).

Numerical results: Asymmetric total resource acquisition

Introducing asymmetric total resource acquisition, represented as sex differences in the

maximal attack rates $a_{F,max}$ and $a_{M,max}$, has complex effects on both population dynamics and

coexistence between competing resources. Introducing asymmetric attack rates can promote

coexistence (as predicted by the analytical results) while simultaneously creating cyclical

resource dynamics (Fig 2B). Under pure apparent competition and asymmetric attack rates,

consumer sexual dimorphism has some similar effects to the case of symmetry, mediating

coexistence of resources that differ in intrinsic per-capita growth rates (Fig 4B).

Under direct interspecific competition, sexual dimorphism in resource specialization

has striking effects on resource persistence when the sexes differ in total resource acquisition,

expanding regions of coexistence (Figs 5A, B) and in some cases leading to persistence of a

competitively inferior resource (Figs 5C, D; Supplement B Figs S4B, C). Moreover, sex

differences in total resource acquisition can reverse competitive outcomes when resources

differ in their intrinsic per-capita growth rates and the degree of resource specialization is

held constant (Figs 5C, D). The effect of sex differences in total resource acquisition on

patterns of prey persistence can be as striking as the effects of resource specialization

(Supplement B Figs S2, S1).

Asymmetric contribution of resource acquisition to birth rate

Introducing asymmetries in the contribution of sex-specific resource acquisition to birth

rate (e.g., polygyny or polyandry, manifest as changes of the constant b_M), had little

qualitative effect. This similarity is illustrated in Supplement B figure S3 (compare to S1).

Although changing these constants results in shifts of absolute regions of coexistence, the

influence of consumer sex differences was similar. Lack of sensitivity in asymmetries in the

contribution of resource acquisition to birth rates is consistent with our analytical results

(Supplement A) that apply to all first-order homogenous mating functions including

geometric and arithmetic contributions to reproduction.

Discussion

Using a general model of consumer-resource dynamics we show that consumer sexual

dimorphism has substantial consequences for community assembly. Competitive exclusion

via apparent competition is expected and observed when males and females are

monomorphic generalists and resources differ substantially in their intrinsic per-capita

growth rates. However, when male and female consumers differ in their resource-specific

attack rates, resource species that differ substantially in their intrinsic per-capita growth rates

can coexist.   Similar effects of consumer sexual dimorphism are observed when resources

compete directly, with sexual dimorphism in some cases permitting coexistence or

persistence of a competitively inferior resource (summarized in Table 1). However, for both

direct and apparent competition, consumer sexual dimorphism can also lead to competitive

exclusion between resources that would typically be expected to coexist. Thus, consumer sex

differences result in fundamental changes in the types of competing resources that can

establish during community assembly. These results also support conclusions from other

food web models suggesting trophic position may impact the observed ecological effects of

sexual reproduction (Kawatsu 2018). Moreover, equilibrium resource abundances and

temporal dynamics are altered by consumer sexual dimorphism even when long-term

ecological outcomes are unaffected. Our results in many ways echo recent work

demonstrating that ontogenetic differences in resource acquisition in a species can have

similar complex consequences for community assembly (de Roos 2020). However, a key

difference is that in sexually-reproducing species we expect to observe these types of effects

whenever consumer mating propensity depends, in part, on resource acquisition, a biological

reality for many sexually-reproducing consumers (e.g. widespread evidence of developmental

thresholds for transitions to reproduction, reveiwed in Wilbur and Collins 1973, Day and

Rowe 2002, and resource-dependent sexual displays, Bonduriansky 2007), and whenever

male abundance matters for population growth rates.

Although our results suggest that the effects of consumer sexual dimorphism on

community assembly can be complex, our analysis reveals some key predictions from our

model. First, consumer sex differences in resource-specific attack rates result in an increase

in the equilibrium density of each resource and a decrease in the abundance of the consumer

species. Intuitively, sexual dimorphism can frequently lead to slightly suboptimal resource

use of the species due to a mismatch between demand (restricted by a 50:50 sex ratio) and

resource availability which can be more variable and dynamic. Second, consumer sexual

dimorphism can promote coexistence when the more aggressively-feeding sex (higher total

attack rates) specializes on and suppresses what would otherwise be the competitively

superior resource. This later situation may be commonplace if sex-specific natural selection

favors specialization, by the sex with higher resource requirements, on the most abundant

resource. Finally, and related to the previous observation, the effects of consumer sexual

dimorphism are most pronounced when resources either differ in intrinsic per-capita growth

rates, or when consumer sexes differ in their total resource acquisition.

Although the assumption of complete female demographic dominance is common in

theoretical models in ecology and evolutionary biology, there is little empirical support for

this demographic extreme. Our simulation models instead focused on a harmonic mean birth

function, which is biologically realistic in that it captures severely reduced birth rates when

any one sex is rare, which is likely to be the case even if sperm limitation is unimportant at

sex ratios near 1:1. Empirically evaluating alternative birth functions in real systems is

difficult, requiring extreme variation in adult sex ratios rarely seen in many species.

However, some empirical data exists and supports the harmonic mean as the best-fitting

description of sexual birth rates (Miller and Inouye 2011a). Nonetheless, our analytical

results (see Supplement A for details) that our conclusions apply to any birth function that is

first order homogenous i.e. $B(aMI_M, aFI_F) = aB(MI_M, FI_F)$ for any positive constant $a>0$.

These functions include all functions typically considered candidate birth functions by

demographers (Caswell and Weeks 1986, Miller and Inouye 2011a). Our conclusions do

depend on the assumption that male abundance makes some contribution to birth rate.

Although even moderate sex differences in attack rates can change the conditions

under which resources coexist, the strongest effects are observed when the consumer

population approaches complete sexual dimorphism in resource specific attack rates. Cases

of such extreme sex differences in diet preference exist (Temeles et al. 2000), although

whether they occur with any regularity is unclear. Diet divergence between the sexes, and

sexual dimorphisms in trophic morphology, are common (Shine 1989), although overlap in

male and female diets can be substantial even in the presence of ecological sex differences

(e.g., Stamps et al. 1997, De Lisle and Rowe 2015a). Yet overlap in diet content may be

expected even in the case of large differences in attack rates, as diet content is the product of

both attack rate and resource abundance. Further, males and females may fail to diverge in

diet preference even when optimum diet nutritional content differs across the sexes (Reddiex

et al. 2013). These challenges to understanding sex-specific resource acquisition highlight

the need for further empirical studies disentangling expressed diet preference, male and

female nutritional optima, and the evolution of ecological sexual dimorphism.

We have assumed a Fisherian sex ratio, where the primary sex ratio is maintained at a

stable 1:1 ratio (Fisher 1930), although our model allows for deviations from a 1:1

operational sex ratio via sex-specific intrinsic mortality. In fact, our results show that even

slight sex differences in mortality rates can have dramatic consequences for community

assembly under both pure apparent and direct competition between resource species. Sex-

specific mortality is commonplace in natural populations and can occur for a variety of

reasons, such as sex-specific costs of reproduction, predation, or sexual conflict. In our

model, sex-specific mortality, as well as variation in consumer mating system (see

Supplemental material) had strong but predictable consequences for resource abundance,

leading to increases in the abundance of the resource favored by the sex with higher intrinsic

mortality.

Our analysis also assumes that the male and female trophic traits are constant. In

nature, however, the degree and even direction of sexual selection can vary dramatically

among closely related populations (e.g., Reimchen et al. 2016). Dimorphism clearly evolves

rapidly in response to spatially varying sexual selection and resource availability. Future

extensions to our strictly ecological model could add in eco-evolutionary dynamics of sexual

dimorphism.

Male and female densities were reduced with increasing ecological sexual

dimorphism, because consumer births are limited by resource acquisition in both sexes in our

model. In the extreme case, where the consumer cannot persist on a single resource (for

example, under complete sexual dimorphism), coexistence of the consumer depends on stable

coexistence of the resource species (Supplement A). Our simulations suggest robust

persistence of extremely dimorphic consumers in two-consumer communities, albeit at

reduced density. Both of these results – reduced consumer density and dependence on the

presence of both resource species in the community – suggest that sexually-dimorphic

populations may face a higher risk of extinction due to demographic stochasticity. However,

it is difficult from our purely deterministic ecological model to fully interpret the

consequences of sexual dimorphism on extinction probability. Evolution of sexual

dimorphism from a monomorphic ancestor that specializes on a single resource could lead to

increased population mean fitness, by increasing the total resource pool available across both

sexes (Rand 1952, Selander 1966, Slatkin 1984, Li and Kokko 2021); such a process

represents a form of within-species, between-sex ecological character displacement, and may

be particularly likely to occur when the sexes interact in small demes (Li and Kokko 2021).

More generally, evolution of sexual dimorphism is expected to be critical to population mean

fitness whenever optima differ for males and females (Lande 1980), and empirical data

suggest sexual dimorphism can be associated with reduced extinction probability at both the

macroevolutionary scale and in extant populations (De Lisle and Rowe 2015b). Nonetheless,

our results suggest a full understanding of sexual dimorphism's role in population persistence

could require integrating theory and data from population/community ecology and

evolutionary genetics. This conclusion is complemented by recent theory (de Vries and

Caswell 2019) indicating sexual dimorphism in demographic parameters can have important

consequences for maintaining genetic diversity.

Our model generally predicts that ecological sexual dimorphism may in some cases

promote, and in other cases reduce, diversity at lower trophic levels during community

assembly. Testing this prediction with correlative data (reviewed in Tsuji and Fukami 2020)

would be possible but challenging. An alternative and non-exclusive hypothesis, that

dimorphic predators are more likely to establish in communities with diverse prey

assemblages, would also generate patterns consistent with our results. Alternatively,

experiments that manipulate the expressed dimorphism of predators and track community

dynamics at lower trophic levels are tractable in some systems. Similar experiments have

been performed (Fryxell et al. 2015, Start and De Lisle 2018), in which effects of predator

sex ratio manipulation on prey communities are assessed in mesocosm designs. Although the

results of these experiments do suggest sex differences can have important community

consequences, no studies (to our knowledge) have compared communities in which the

magnitude of morphological sexual dimorphism is manipulated under a stable sex ratio. Such

designs are possible when distributions of male and female phenotypes exhibit substantial

variation, and would represent an ideal empirical test of the theoretical results presented here.

Ecological sex differences are commonplace, although the details of their

evolutionary drivers and ecological consequences are unclear. Emerging theory and data

indicate ecological sex differences may have important consequences for the evolutionary

genetics of adaptation (Zajitschek and Connallon 2017), the dynamics of diversification

(Bolnick and Doebeli 2003, De Lisle and Rowe 2015b) and community assembly (Fryxell et

al. 2015, Pincheira-Donoso et al. 2018, Start and De Lisle 2018). Our results add to this body

of work, indicating that sexual dimorphism can have substantial effects on the structure,

abundance, and dynamics of ecological communities, including changing conditions for

coexistence between competing resource species.

Acknowledgements. We thank Tim Connallon and Miguel Gomez for comments on an

early draft of this manuscript, as well as Gonzalo Hernando for assistance in the early stages

of the project. This work was supported by funds from the University of Connecticut to DIB,

an establishment grant from the Swedish Research Council to S.P.D. (2019-03706), and an

U.S. National Science Foundation Grant DMS176803 to SJS.

References

484
Andersson, M. B. 1994. Sexual selection. Princeton University Press, Princeton, N.J.

Belovksy, G. E. 1978. Diet optimization in a generalist herbivore: the

moose. Theoretical Population Biology **14**:105-134.

Bolnick, D. I., P. Amarasekare, M. S. Araujo, R. Burger, J. M. Levine, M. Novak, V. H. W.

Rudolf, S. Schreiber, M. C. Urban, and D. A. Vasseur. 2011. Why intraspecific trait

variation matters in community ecology. Trends in Ecology & Evolution **26**:183-192

Bolnick, D. I., and M. Doebeli. 2003. Sexual dimorphism and adaptive speciation: two sides

of the same ecological coin. Evolution **57**:2433-2449.

Bolnick, D. I., R. Svanbäck, J. A. Fordyce, L. H. Yang, J. M. Davis, C. D. Hulsey, and M. L.

Forister. 2003 The Ecology of Individuals: Incidence and implications of individual

specialization The American Naturalist **161:**1-28.

Bonduriansky, R. 2007. The evolution of condition-dependent sexual dimorphism The

American Naturalist **169**:9-19.

Camus, M. F., K. Fowler, M. W. D. Piper, and M. Reuter. 2017. Sex and genotype effects on

nutrient-dependent fitness landscapes in Drosophila melanogaster. Proceedings of the

Royal Society of London Series B. **284**:20172237.

Caswell, H., and D. E. Weeks. 1986. Two-sex models: chaos, extinction, and other dynamic

consequences of sex. American Naturalist **128**:707-735.

Cortez, M. H., and S. Patel. 2017. The effects of predator evolution and genetic variationon

predator-prey population-level dynamics. Bulletin of Mathematical Biology **79**:1510-

1538.

Darwin, C. 1871. The descent of man and selection in relation to sex. J. Murray, London.

Day, T., and L. Rowe. 2002. Developmental thresholds and the evolution of reaction norms

for age and size at life-history transitions. The American Naturalist **159**:338-350.

De Lisle, S. P. 2019. Understanding the evolution of ecological sex differences: Integrating

character displacement and the Darwin-Bateman paradigm. Evolution Letters **3**:444-

447.

De Lisle, S. P., and L. Rowe. 2015a. Ecological character displacement between the sexes.

The American Naturalist **186**:693-707.

De Lisle, S. P., and L. Rowe. 2015b. Independent evolution of the sexes promotes amphibian

diversification. Proceedings of the Royal Society B: Biological Sciences

282:20142213.

de Roos, A.M. 2020. Effects of life history and individual development on community

dynamics: A review of counterintuitive consequences. Ecological Research 35: 930–

946.

de Vries, C., and H. Caswell. 2019. Selection in two-sex stage structured populations:

Genetics, demography, and ploymorphism Theoretical Population Biology **130**:160-

169.

Des Roches, S., D. M. Post, N. E. Turley, J. K. Bailey, A. P. Hendry, M. T. Kinnison, J. A.

Schweitzer, and E. P. Palkovacs. 2018. The ecological importance of intraspecific

variation. Nature Ecology and Evolution **2**:57-64.

Fisher, R. 1930. The genetical theory of natural selection. Oxford University Press, Oxford,

UK.

Fryxell, D. C., H. A. Arnett, T. M. Apgar, M. T. Kinnison, and E. P. Palkovacs. 2015. Sex

ratio variation shapes the ecological effects of a globally introduced freshwater fish

Proceedings of the Royal Society of London Series B. **282**:20151970.

Fryxell, D. C., D. Weiler, M. Kinnison, and E. P. Palkovacs. 2019 Eco-evolutionary

dynamics of sexual dimorphism. Trends in Ecology and Evolution **34**:591-594.

Garlapow, M. E., W. Huang, M. T. Yarboro, K. R. Peterson, and T. F. C. Mackay. 2015.

Quantitative genetics of food intake in Drosophila melanogaster. PloS OnE

10:e0138129.

Giery, S. T., and C. G. Layman. 2019. Ecological consequences of sexually selected traits:

An eco-evolutionary perspective The Quarterly Review of Biology **94**:29-74.

Gomez-Llano, M. A., H. M. Bensch, and E. I. Svensson. 2018. Sexual conflict and ecology:

species composition and male density interact to reduce male mating harassment and

increase female survival Evolution **72**:906-915.

Haldane, J. B. S. 1937. The effect of variation on fitness. The American Naturalist **71**:337-

349.

Holt, R. D. 1977. Predation, apparent competition, and the structure of prey communities.

Theoretical Population Biology **12**:197-229.

Holt, R. D., and J. H. Lawton. 1993. Apparent competition and enemy-free space in insect

host-parasitoid communities. The American Naturalist **142**:623-645.

Hughes, A. R., B. D. Inouye, M. T. J. Johnson, N. Underwood, and M. Vellend. 2008

Ecological consequences of genetic diversity Ecology Letters **11**:609-623.

Jensen, K., C. McClure, N. K. Priest, and J. Hunt. 2015. Sex-specific effects of protein and

carbohyfrate intake on reproduction but not lifespan in Drosophila melanogaster.

Aging Cell **14**:605-615.

Kawatsu, K. 2018. Ecological effects of sex differ with trophic positions in a simple food

web. Ecology and Evolution **8**:1239-1246.

Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.

Evolution **34**:292-305.

Li, X.-Y., and H. Kokko. 2021. Sexual Dimorphism driven by intersexual resource

competition: Why it is rare, and where to look for it? Journal of Animal Ecology

90:1831-1843.

Lindstöm, J., and H. Kokko. 1998. Sexual reproduction and population dynamics: the role of

polygony and demographic sex differences Proceedings of the Royal Society of

London Series B. **265**:483-488.

Maklakov, A., S. J. Simpson, F. Zajitchek, M. D. Hall, J. Dessmann, F. Clissold, D.

Raubenheimer, R. Bondurianksy, and R. C. Brooks. 2008. Sex-specific effects of

nutrient intake on reproduciton and lifespan Current Biology **18**:1062-1066.

Miller, T. E. X., and B. D. Inouye. 2011a. Confronting two-sex demographic models with

data. Ecology **92**:2141-2151.

Miller, T. E. X., and B. D. Inouye. 2011b. Confronting two-sex demographic models with

data. Ecology **92**:2141-2151.

Patel, S., and S. J. Schreiber. 2015. Evolutionarily driven shifts in communities with

intraguild predation. The American Naturalist **186**:E98-E110.

Pincheira-Donoso, D., T. Tregenza, R. K. Butlin, and D. J. Hodgson. 2018. Sexes and species

as rival units of niche saturation during community assembly. Global Ecology and

Biogeography **27**:593-603.

R Core Team. 2018. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria.

Rand, A. L. 1952. Secondary sexual characters and ecological competition. Fieldiana-

Zoology **34**:65-70.

Reddiex, A. J., T. P. Gosden, R. Bonduriansky, and S. F. Chenoweth. 2013. Sex-specific

fitness consequences of nutrient intake and the evolvability of diet preferences. The

American Naturalist **182**:91-102.

Reimchen, T., D. Steeves, and C. A. Bergstrom. 2016. Sex matters for defence and trophic

traits of threespine stickleback. Evolutionary Ecology Research **17**:459-485.

Rowe, L., and D. Houle. 1996. The lek paradox and the capture of genetic variance by

condition dependent traits. Proceedings of the Royal Society of London Series B-

Biological Sciences **263**:1415-1421.

Schreiber, S. J. 2021. The P* rule in the stochastic Holt-Lawton model of apparent

competition. Discrete and Continuous Dynamical Systems-B **26**:633.

Schreiber, S. J., R. Burger, and D. I. Bolnick. 2011. The community effects of phenotypic

and genetic variation within a predator population Ecology **92**:1582-1593.

Selander, R. K. 1966. Sexual dimorphism and differential niche utilization in birds. The

Condor **68**:113-151.

Shine, R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the

evidence. Quarterly Review of Biology **64**:419-461.

Shärer, L., L. Rowe, and G. Arnqvist. 2012 Anisogamy, chance and the evolution of the sex

roles. Trends in Ecology & Evolution **27**:260-264.

Slatkin, M. 1984. Ecological causes of sexual dimorphism. Evolution **38**:622-630.

Soetgart, K., T. Petzoldt, and R. W. Setzer. 2010. Solving Differential Equations in R:

Package deSolve. Journal of Statisitcal Software **33**:1-25.

Stamps, J. A., J. B. Losos, and R. M. Andrews. 1997. A comparative study of population

density and sexual size dimorphism in lizards. American Naturalist **149**:64-90.

Start, D., and S. P. De Lisle. 2018. Sexual dimorphism in a top predator drives aquatic prey

community assembly. Proceedings of the Royal Society of London Series B.

2018:1717.

Svensson, E. I. 2019. Eco-evolutionary dynamics of sexual selection and sexual conflict

Functional Ecology **33**:60-72.

Temeles, E. J. 1985. Sexual size dimorphism of bird-eating hawks: the effect of prey

vulnerability. The American Naturalist **125**:485-499.

Temeles, E. J., I. L. Pan, J. L. Brennan, and J. N. Horwitt. 2000. Evidence for ecological

causation of sexual dimorphism in a hummingbird. Science **289**:441-443.

Trivers, R. L. 1972. Parental investment and sexual selection. Pages 136-179 *in* B. Campbell,

editor. Sexual selection and the descent of man. Aladine, Chicago.

Tsuji, K., and T. Fukami. 2018. Community-wide consequences of sexual dimorphism:

evidence from nectar microbes in dioecious plants. Ecology **99**:2476-2484.

Tsuji, K., and T. Fukami. 2020. Sexual dimorphism and species diversity: from clades to

sites. Trends in Ecology and Evolution **35**:105-114.

Tuszynski, J. 2019. caTools: Tools: moving window statistics, GIF, Base64, ROC AUC, etc..

R package version 1.17.1.2. https://CRAN.R-project.org/package=caTools.

Weimerskirch, H., M. Le Corre, H. Gadenne, D. Pinaud, A. Kato, Y. Ropert-Coudert, and C.-

A. Bost. 2009. Relationship between reversed sexual dimorphism, breeding

investment and forgaing ecology in a pelagic seabird, the masked booby. Oecologia

161:637-649.

Wilbur, H. M., and J. P. Collins. 1973. Ecological aspects of amphbian metamorphosis

Science **182**:1305-1314.

Zajitschek, F., and T. Connallon. 2017. Partitioning of resources: the evolutionary genetics of

sexual conlflict over resource acquisition and allocation. Journal of Evolutionary

Biology **30**:826-838.

Table 1. Summary of effects of sex differences in resource use. Arrows indicate effects on mean species density and the range of parameter space under which resource coexistence is observed.

Density	Coexistence	
Consumer ↓	Apparent competition	↑
Resource ↑	Direct competition, equal resource growth rates, symmetric acquisition	0
	Direct competition, different resource growth rates, symmetric acquisition	↑
	Direct competition, asymmetric acquisition	↑↓

Figure 1. Illustration of the model structure. Males (M) and females (F) of a consumer

species exploit two resources (R_1 and R_2) that may or may not also compete directly (double

arrow). Sex-specific attack rates (black arrows) generate sex differences in ecological niche

such that males and females preferentially attack R_1 and R_2, respectively, with varying

degrees of overlap (grey arrows). Although males and females may consume different

resources, fitness is equal across the sexes and so the dynamics of males, females, resource 1

and resource 2 are coupled.

Figure 2. Coexistence mediated by sexual dimorphic consumers. In panel A, resource 2

(red) is a superior competitor to resource 1 (blue), leading to exclusion of resource 1 and the

sexually-monomorphic consumer that cannot persist on a single resource species. However,

adding consumer sex differences in total resource acquisition, under otherwise identical

parameter values, leads to consumer persistence and coexistence of the resource species with

cyclical dynamics (B). When competition is completely apparent (C, D), the resource with

superior growth rate (red) excludes the inferior resource when consumers are sexually

monomorphic (C). Under the same parameterization but with sexually dimorphic consumers

(D), density of both resources is increased and they coexist. Parameter values: **A**, $\alpha_{11} = .1$,

$\alpha_{12} = .12$, $\alpha_{22} = .1$, $\alpha_{21} = .09$, $\beta = 1$, $d_M = d_F = 0.02$, $b_M = b_F = .1$, $a_{M, max} = a_{F, max} = 1$, $r_1 = r_2$

$= 1$; **B** $\alpha_{11} = .1$, $\alpha_{12} = .12$, $\alpha_{22} = .1$, $\alpha_{21} = .09$, $\beta = 1$, $d_M = d_F = 0.02$, $b_M = b_F = .1$, $a_{M, max} = .8$,

$a_{F, max} = 1.2$, $r_1 = 1$, $r_2 = 1$; **C** $\alpha_{11} = .1$, $\alpha_{12} = 0$, $\alpha = .1$, $\alpha_{21} = 0$, $\beta = 0$, $d_M = d_F = 0.02$, $b_M = b_F$

673      $= .1, a_{M, max} = 1, a_{F, max} = 1,\ r_1 = 1, r_2 = 1.1;$ **D** $\alpha_{11} = .1, \alpha_{12} = 0, \alpha_{22} = .1, \alpha_{21} = 0, \beta = 0.9, d_M$

674      $= d_F = 0.02, b_M = b_F = .1, a_{M, max} = 1, a_{F, max} = 1,\ r_1 = 1, r_2 = 1.1;$

Figure 3. Consumer sexual dimorphism results in increased resource density and reduced consumer density in two and three species communities. Panel A: Equilibrium consumer female density decreases and resource 1 density increases with increased sexual dimorphism in a two species (consumer- 1 resource) community (equation 6). Panel B: these results hold in simulations in three species communities, where resource 1 density is plotted against competition coefficients in both resource species under three levels of sexual dimorphism, increasing from left to right. Assuming symmetric total attack rates across the sexes, symmetric contribution of resource acquisition to consumer birth rates, and equal growth rates across resources.

Figure 4. Sexual dimorphism in resource acquisition expands regions of coexistence

between resources under apparent competition. Panels show relative density of resource

1 ($R_1/(R_1+R_2)$) from numerical simulations with starting conditions of $R_1 = R_2 = 1$ and $M = F$

$= 1$. Panel A shows the outcomes under symmetric maximal attack rates across the sexes

($a_{M, max} = a_{F, max} = 1$), and equal death rates $d_M = d_F = 0.02$. Panel B shows the outcomes

assuming sex differences in maximal attack rate ($a_{M, max} = .8$, $a_{F, max} = 1.2$). Panel C shows

outcomes under identical conditions to A, but with a sex difference in death rate of

approximately 11% (reduced male mortality, $d_M=0.018$, $d_f=0.02$). Panel D shows the

opposite sex difference (reduced female mortality, $d_M=0.02$, $d_f=0.018$). White dashed lines

indicate analytical solutions (equation 8) limited to the range of consumer coexistence

$H_i \frac{1}{\alpha_{ii}} > 2$, which corresponds to $\beta < .96$.

Figure 5. Sexual dimorphism alters regions of coexistence of resource species. Panels

show relative density of resource 1 ($R_1/(R_1 + R_2)$) from numerical simulations with starting

conditions of $R_1 = R_2 = 1$ and $M = F = 1$. Panel A shows outcomes under sexual

monomorphism ($\beta = 0$), unequal growth rates ($r_1 = 2$, $r_2 = 2.1$) across resources, and no sex

differences in total attack rate ($a_{M, max} = a_{F, max} = 1$). Panel B shows the outcome under the

same parameter values but with strong consumer sexual dimorphism ($\beta = 0.7$). Panels C and

D show the effect of sexual niche divergence when the sexes also differ in total attack rates

(total resource acquisition), with C showing the outcomes under unequal total attack rates

($a_{M, max} = .7$, $a_{F, max} = 1.3$), sexual monomorphism in prey preference ($\beta = 0$), and

differential resource growth rates ($r_1 = 1.6$, $r_2 = 1.8$), and D showing the outcome under the

same parameter values but with moderate sexual dimorphism in prey preference ($\beta = 0.4$).

Panels E and F show the effects of sex differences in mortality; in Panel E, $d_M = 0.018$, d_f

$= 0.02$. In Panel F, $d_M = 0.02$, $d_f = 0.018$, while $\beta = 0.5$ in both panels. Note that in the absence

of sex differences in mortality coexistence would be restricted to the bottom left quadrant

represented by black dashed lines, which demark equal inter and intraspecific competition

coefficients. White dashed box demarks coexistence determined by analytical invasion

criteria into a two-species community under equation 9.