Complex community-wide consequences of consumer sexual dimorphism

Stephen P. De Lisle [1,2] *

Sebastian J. Schrieber [3]

Daniel I. Bolnick [1]

[1] Department of Ecology & Evolutionary Biology, University of Connecticut, Storrs, CT 06269

[2] Present Address: Evolutionary Ecology Unit, Department of Biology, Lund University, Lund

Sweden

[3] Department of Evolution and Ecology and Center for Population Biology, University of

California, Davis, CA 95616

* email: stephen.de_lisle@biol.lu.se

Keywords: apparent competition, community assembly, competitive exclusion, ecological sexual

dimorphism, resource competition

Abstract

Sexual dimorphism is a ubiquitous source of within-species variation, yet the community-level

consequences of sex differences remain poorly understood. Here, we analyze a bitrophic model

of two competing resource species and a sexually-reproducing consumer species. We show that

consumer sex differences in resource acquisition can have striking consequences for consumer-

resource coexistence, abundance, and dynamics. Under both direct interspecific competition and

apparent competition between two resource species, sexual dimorphism in consumers' attack

rates can mediate coexistence of the resource species, while in other cases can lead to exclusion

when stable coexistence is typically expected. Slight sex differences in total resource acquisition

also can reverse competitive outcomes and lead to density cycles. These effects are expected

whenever both consumer sexes require different amounts or types of resources to reproduce. Our

results suggest that consumer sexual dimorphism, which is commonplace, has wide-reaching

implications for the assembly and dynamics of natural communities.

Introduction

Within-species variation is one key feature of natural populations that has emerged as a critical contributor to community ecology and species interactions (Hughes et al. 2008 , Bolnick et al. 2011, Des Roches et al. 2018). Even in the simplest scenario of a single-species community with a single optimum phenotype, within-population variation is expected to reduce population growth rate of a well-adapted population via effects on mean fitness (Haldane 1937). In more complex multi-species communities, phenotypic variation in consumers or their resources can either promote or constrain coexistence between competing species, under many circumstances yielding different conclusions than would be reached from models ignoring within species variation (Bolnick et al. 2011, Schreiber et al. 2011, Patel and Schreiber 2015, Cortez and Patel 2017). One emerging question (Bolnick et al. 2011) from this work is the degree to which understanding the specific source of phenotypic variation matters in community ecology.

Here, we explore the ecological consequences of sexual dimorphism, a central feature of metazoans and many plant populations that has not been fully incorporated into ecological theory. The gamete differences that define the sexes are expected to lead to divergence between males and females in a suite of life history traits (Trivers 1972, Shärer et al. 2012), and phenotypic traits. This divergence is iconically manifest as striking sexual dimorphism in sexually-selected traits such as body size or courtship displays (Darwin 1871, Andersson 1994). Although sexual selection and associated mate choice behaviors themselves may have relevance to interspecific ecological interactions (Gomez-Llano et al. 2018), the different life histories that define the sexes are often expected lead to different nutritional and resource requirements for males and females (Maklakov et al. 2008). As a specific example, in insects, differential contributions of longevity and fecundity to male and female lifetime reproductive success result

in different combinations of macronutrients that maximize male and female fitness (Maklakov et

al. 2008, Reddiex et al. 2013, Garlapow et al. 2015, Jensen et al. 2015, Camus et al. 2017).

Similarly, nutritional models of optimal foraging have been proposed to explain sex differences

in diets of moose (Belovksy 1978). Consequently, males and females of many species have

evolved divergent resource use, either as an indirect outcome of divergent reproductive roles or

through other forms of sex-specific natural selection, resulting in sex-differences in diet

composition and often in corresponding trophic morphology (Slatkin 1984, Temeles 1985, Shine

1989, Temeles et al. 2000, De Lisle and Rowe 2015a, De Lisle 2019).

Data suggest that these 'ecological' sexual dimorphisms can have substantial

consequences for community structure (Fryxell et al. 2015, Pincheira-Donoso et al. 2018, Start

and De Lisle 2018, Tsuji and Fukami 2018, 2020). More generally, sex differences may play an

important role in the relationship between ecological and evolutionary dynamics (Giery and

Layman 2019, Svensson 2019, Fryxell et al. 2019). Sexual dimorphism thus represents a key

source of ecologically-relevant variation within species. Within-species variation in resource

specialization is commonplace (Bolnick et al. 2003), with important consequences for species

interactions and community assembly (Hughes et al. 2008 , Bolnick et al. 2011, Des Roches et

al. 2018). Critically, as a source of intraspecific variation, sexual dimorphism may have different

consequences from other types of variation. Unlike most phenotypes, whose relative abundances

can evolve to reflect local ratios of alternative resources, the ratio of males to females (at birth) is

expected to be maintained at 1:1 (Darwin 1871, Fisher 1930), though their phenotypes and

demography may diverge to have differential ecological impacts. Thus, the rate at which males

are born (which determines predation pressure on males' prey), is strongly dependent on

females' foraging success. Perhaps to a lesser extent (depending on mating system), the rate of

female offspring production can depend on male foraging success. This coupling of the dynamics

of two consumer phenotypes has unknown consequences for consumer-resource dynamics,

consumer mediated coexistence of the resource species, and apparent competition between the

resource species. Using a simple model of consumer-resource dynamics, we show that consumer

sexual dimorphism can affect coexistence between resource species (competing directly or

apparently), and species' abundance.

Materials and Methods

Our model represents a simple extension of a classic model of consumer-resource dynamics, in

which a consumer species exploits two resources that themselves undergo density-dependent

growth (Figure 1). Consumer growth rate is limited only by resource abundance. The original

version of this model did not consider within-species phenotypic variation, and can lead to

exclusion of one resource ('apparent competition'; (e.g. Holt 1977). More recently, (Schreiber et

al. 2011) showed that quantitative trait variation in the predator, affecting attack rates on the two

resources, can facilitate coexistence between the prey. Here, we instead allow for the possibility

that male and female consumers differ in resource-specific attack rates. We can describe the

population dynamics of two resources (with densities R_i, R_j) and consumer males (with density

M) and females (with density F) using the system of ordinary differential equations:

$$\frac{dR_i}{dt} = r_i R_i \left(1 - \alpha_{ii} R_i - \alpha_{ij} R_j \right) - a_{M,i} R_i M - a_{F,i} R_i F \text{ for } j \neq i \qquad (1a)$$

$$\frac{dM}{dt} = \frac{1}{2} B(M, F, R_1, R_2) - d_M M \qquad (1b)$$

$$\frac{dF}{dt} = \frac{1}{2} B(M, F, R_1, R_2) - d_F F \qquad (1c)$$

Each resource undergoes density-dependent population growth with intrinsic per-capita growth

rates r, with competition represented by competition coefficients α. Resources are also regulated

by consumers depending on sex-specific attack rates by males a_M and by females a_F. Consumer

dynamics are governed by females' birth rate B (half of the offspring being female), and limited

by sex-specific intrinsic mortality rates d_M and d_F.

An appropriate function describing birth rates has been a point of debate for

demographers, with the general conclusion being that any function describing birth rates should

capture the negative effects of extreme sex ratio skew on birth rates, with the extreme being that

birth rates should be zero when one sex is absent (Caswell and Weeks 1986), or unable to breed.

Generally, treating birth rates proportional to the harmonic mean density of males and females is

agreed as the best approach (Caswell and Weeks 1986, Lindstöm and Kokko 1998). Here, we

extend this harmonic mean birth function to account for each sex's foraging success, on the logic

that each sex must both be present and sufficiently well-fed to reproduce:

$$B(M, F, R_i, R_j) = 2\frac{MI_M * FI_F}{MI_M + FI_F} \qquad (2)$$

128   where I is the energy intake of a given sex:

$$I_F = b_F(a_{F,1}R_1 + a_{F,2}R_2) \text{ and } I_M = b_M(a_{M,1}R_1 + a_{M,2}R_2) \qquad (3)$$

which couples the dynamics of consumers and resources. In this model the rate of consumer

births depends on not just the number of males and females but also on the abundance of both

prey and the prey preference of each sex, where MI_M is the number of males weighted by their

energy to reproduce (similarly with females), b is a scaling constant reflecting the number of

predator offspring created per prey eaten, and $a_{M,i}, a_{F,i}$ are the attack rates on the resource i.

Because we are specifically interested in understanding effects of sexual dimorphism in

prey preference, we relate male and female resource-specific attack rates via the degree of sexual

dimorphism, β, such that

$$a_{M,1} = a_{M,max}\left(\frac{1}{2} + \frac{\beta}{2}\right) \text{ and } a_{F,1} = a_{F,max}\left(\frac{1}{2} - \frac{\beta}{2}\right) \qquad (4a)$$

$$a_{M,2} = a_{M,max}\left(\frac{1}{2} - \frac{\beta}{2}\right) \text{ and } a_{F,2} = a_{F,max}\left(\frac{1}{2} + \frac{\beta}{2}\right) \qquad (4b)$$

where β ranges from zero to one, with $\beta = 0$ representing complete sexual monomorphism if

maximum attack rates are equal across the sexes (both sexes are generalists, attacking each prey

at the rate, $a_{max}/2$; we explore both $a_{M, max} = a_{F, max} = a_{max}$ as well as $a_{M, max} \neq a_{F, max}$), and $\beta = 1$

representing complete sexual dimorphism such that males only attack R_1 and females only

attack R_2.

By assuming that resource acquisition of males and females contributes to the consumer

birth rate, we are not necessarily making any assumption about parental care or condition-

dependent fecundity. Rather, this is an assumption that the fraction of each sex available for

reproduction depends in part on resource acquisition, a realistic feature of many organisms. For

instance, males' energetic costs of finding a mate, defending a territory, care for offspring, or

expressing a sexually-selected trait are all expected to depend on the resource pool available to

males (Rowe and Houle 1996). Females' ability to produce eggs, gestate, and provision young

(e.g., lactation in mammals) similarly depends on their energy intake. Parental care and other

factors can also lead to associations between sex-specific resource abundance and consumer birth

rates, beyond what would be expected beyond the costs of simply being available to mate.

We analyzed this model to find equilibria when both one or two resources are present, as

well as invasion criteria for a resource into a two species community at equilibrium. Using the

mathematical theory of permanence (Schreiber 2000, Patel & Schreiber 2018), we can use these

invasion criteria to determine whether all three species coexist (in the sense of permanence),

exhibit a bistability, or competitive exclusion occurs. In addition to presenting our analytical

results below, we provide further mathematical detail in Supplement A. We also numerically

explored the behavior of the model under a variety of scenarios using the packages deSolve v.

1.21 (Soetgart et al. 2010) and caTools v. 1.17.1.2 (Tuszynski 2019) in R v. 3.5.0 (R Core Team

2018). Importantly, using simulations allowed us to explore the behavior of our model for cases

where 2-species consumer-resource equilibria do not exist. Because there are a large number of

potential combinations of parameters that could be explored, we focused our simulations on three

different biological scenarios: 1) completely symmetric male and female total attack rates and

contributions to birth rate, 2) asymmetric total attack rates across the sexes, and 3) asymmetric

contribution of male and female resource acquisition to consumer birth rates. The first scenario

applies most readily to organisms with biparental care and similar total caloric requirements

across the sexes. The second scenario likely applies to many organisms where total lifetime

resource acquisition is higher for one sex, which is the case for many organisms where the sexes

differ, for example, in body size. The third scenario is another realistic departure from the first,

relevant for organisms where resource acquisition in one sex (e.g., females) has a greater

influence on that sexes mating propensity, or cases with biased operational sex ratios due to

polyandry or polygyny. For each scenario we explore the consequences of consumer ecological

sexual dimorphism (β) for consumer persistence and coexistence. For each set of parameter

values, we simulated the model 1,000 time steps (to ensure the simulation reaches its

equilibrium). For each run, we determined the equilibrium abundances of M, F, R_1 and R_2 at t =

1,000. We also calculated the standard deviation of species' abundances in the last 50 time-steps.

Although we present solutions simulated on the scale of $\alpha_{ii} = 0.1$, we obtained qualitatively

equivalent conclusions rescaling under a wide range values of competition coefficients (see Table

S1B, Supplement B). The R script to generate all figures and results presented is available in the

supplementary material.

Results

Analytical results

Our analysis begins by considering a subsystem of a single resource species, say species i, and

the consumer species. These two species can coexist if

$$H_i \frac{1}{\alpha_{ii}} > 2 \qquad\qquad (5)$$

where $H_i := 2 \frac{\frac{b_M a_{i,M} b_F a_{i,F}}{d_M}\frac{}{d_F}}{\frac{b_M a_{i,M}}{d_M}+\frac{b_F a_{i,F}}{d_F}}$ is the harmonic mean of the lifetime per-capita resource contributions

of each sex to reproduction, and $\frac{1}{\alpha_{ii}}$ corresponds to the carrying capacity of resource species i.

Intuitively, $H_i \frac{1}{\alpha_{ii}}$ corresponds to the average number of offspring produced by a mating pair

during their lifetime when the resource species is at its carrying capacity. These harmonic means

decrease to zero with the degree of sexual dimorphism (i.e. H_i is a decreasing function of β and

H_i=0 when $\beta = 1$), there always is critical degree of sexual dimorphism above which the

consumer cannot persist on a single resource species. Intuitively, this arises when one of the

sexes specializes too much on the other (absent) resource species and, consequently, contributes

too little to reproduction. When the coexistence condition holds, the consumer-resource species

pair coexist at the following equilibrium densities:

199
$$R_i^* = 2/H_i$$

200
$$F_i^* = \frac{r_i\left(1 - \frac{2\alpha_{ii}}{H_i}\right)}{a_{i,F} + \frac{a_{i,M}d_F}{d_M}} \tag{6}$$

201
$$M_i^* = \frac{d_F}{d_M} F_i^*$$

202   At these densities of the consumer and resource i, the second (rare) resource species j can invade

203   if its per capita growth rate

204
$$I_j = r_j\left(1 - \frac{2\alpha_{ji}}{H_i}\right) - A_j F_i^* \tag{7}$$

is positive. Here, $A_j = a_{F,j} + a_{M,j}d_F/d_M$ is the average attack rate on resource j when the males

and females are at an equilibrium. Provided the consumer species can persist on each of the

resource species individually, the invasion growth rates I_1 and I_2 determine the ecological

outcomes. If both invasion growth rates are positive (mutual invasibility), all three species

coexist. If both invasion rates are negative, there is a priority effect in which both single

resource-consumer equilibria are stable. If one invasion growth rate is positive and the other is

negative, this suggests (as confirmed by numerical simulations) that the resource species with the

positive invasion growth rate excludes the other resource species. In the supplementary material

(Supplement A), we provide more details about this analysis and describe the invasion conditions

for these outcomes or when the consumer only persists in the presence of both resource species.

As a step toward understanding the general conditions for coexistence and exclusion, we

first examine two special cases corresponding to (i) resource competitive symmetry whereby

$\alpha_{11} = \alpha_{12} = \alpha_{21} = \alpha_{22}$ and (ii) pure apparent competition whereby $\alpha_{12} = \alpha_{21} = 0$. When there

is competitive symmetry, the resource species with the larger value of r_i/A_i excludes the other

resource species. The quantity r_i/A_i corresponds to the consumer density supported by resource

species i when intraspecific competition is very weak i.e. $\alpha_{ii} = 0$. Hence, the resource species

that supports the highest consumer equilibrium density (in the absence of self-limitation)

excludes the other. This is an analog of the *P** rule (Holt 1977, Holt and Lawton 1993, Schreiber

2021). In this case, sexual dimorphism (as measured by β) influences outcomes only if there is

an asymmetry in the maximal attack rates of the female and male (e.g., female masked boobies

dive for prey more often than males; Weimerskirch et al. 2009). For example, if the female-

preferred resource (R_I) has the higher intrinsic rate of growth ($r_1 > r_2$) and the female consumer

has the higher maximal attack rate of the two sexes, then the female-preferred resource species

excludes the other resource species when sexual dimorphism is low (i.e. $A_1{=}A_2$ when $\beta = 0$).

However, as the average attack rate A_I on the female-preferred resource increases with the

magnitude β of the sexual dimorphism, sexual dimorphism reverses the outcome of apparent

competition whenever $r_1/a_{F,max} < r_2/a_{M,max}$ and the self-limitation in the female-preferred

resource is weak.

Next, we consider the case of pure apparent competition in which there is no direct

competition between the resources, i.e. $\alpha_{12} = \alpha_{21} = 0$. Then, coexistence occurs when

235
$$\frac{1}{1-2\alpha_{ii}/H_i} > \frac{r_i/A_i}{r_j/A_j} \quad \text{for } i \neq j \tag{8}$$

The right-hand side of this condition determines the outcome of apparent competition in the case

of competitive symmetry i.e. the species with larger r_i/A_i wins. The left-hand is always greater

than one and increases to infinity as the sexual dimorphism increases. This has two implications:

(i) when r_1/A_1 and r_2/A_2 are equal, the resource species coexist at any level of sexual

dimorphism, and (ii) when r_1/A_1 and r_2/A_2 are unequal, sufficiently strong sexual dimorphisms

(allowing the consumer to persist) will ensure coexistence of the two resource species by diluting

the strength of apparent competition.

Finally, the general coexistence condition is

$$\frac{1-2\alpha_{ji}/H_i}{1-2\alpha_{ii}/H_i} > \frac{r_i/A_i}{r_j/A_j} \text{ for } i \neq j \tag{9}$$

The effect of sexual dimorphism on the left-hand side term depends on the relative strengths of

intra- and interspecific competition. When intraspecific competition is greater than interspecific

competition (i.e. $\alpha_{ii} > \alpha_{ji}$), the left-hand side term increases to infinity with increasing sexual

dimorphism. In contrast, when interspecific competition between resources is greater than

intraspecific competition (i.e. $\alpha_{ji} > \alpha_{ii}$), the left-hand side term of the general coexistence

condition decreases to zero with increasing levels of sexual dimorphism. The effect of sexual

dimorphism on the right-hand side term of the general coexistence criterion is as discussed for

the case of competitive symmetry. Together these observations imply that if resource 1 is the

better direct competitor i.e. $\alpha_{21} > \alpha_{11}$ and $\alpha_{21} < \alpha_{22}$, then coexistence requires that resource 2

is the better apparent competitor i.e. $r_2/A_2 > r_1/A_1$. In which case, a sexual dimorphism can

help consumer-mediated coexistence.

Numerical results: Symmetric resource acquisition across the sexes

Our numerical results generally matched conclusion from our analytical solutions and are

summarized in Table 1. The effects of consumer sexual dimorphism on resource population

dynamics are illustrated in Figure 2. Sexual dimorphism results in increased resource density

(Fig 2B vs. 2D; Fig 3) and can facilitate persistence of a competitively inferior resource species

(Fig 2A vs. 2B). When competition is purely apparent and resource acquisition by the consumer

is symmetric across the sexes, increasing consumer sexual dimorphism increases the parameter

space under which resources can coexist (Fig. 2C vs 2D), consistent with our analytical results

(Fig. 4A). In particular, as sexual dimorphism approaches the extreme (each sex uses a different

resource exclusively), competitively inferior resource species can coexist with a superior

competitor that is more susceptible to apparent competition (Figure 4).

When resources compete directly with symmetric intraspecific competition, resource

acquisition is symmetric across the sexes and intrinsic per-capita growth rates are equal between

resources, consumer sexual dimorphism has little effect on invasion of a resource into a

consumer-single-resource community at equilibrium (S1A-C, S4A). Yet even in the absence of

effects on coexistence consumer sexual dimorphism has strong effects equilibrium resource

abundance, both under analytical solutions to a two species community and under simulations of

three species communities (Fig 3). As predicted by our analytical conditions, when resource

growth rates are unequal, sexual dimorphism changes expected regions of coexistence of both

resources and the consumer, leading to invasion and persistence of competitively inferior

resource that is a superior apparent competitor i.e. has a larger value of r_i/A_i (Figure 5B, Figure

S4B, Figure S1).

Numerical results: Asymmetric total resource acquisition

Introducing asymmetric total resource acquisition, represented as sex differences in the

maximal attack rates $a_{F,max}$ and $a_{M,max}$, has complex effects on both population dynamics and

coexistence between competing resources. Introducing asymmetric attack rates can promote

coexistence (as predicted by the analytical results) while simultaneously creating cyclical

resource dynamics (Fig 2B). Under pure apparent competition and asymmetric attack rates,

consumer sexual dimorphism has some similar effects to the case of symmetry, mediating

coexistence of resources that differ in intrinsic per-capita growth rates (Fig 4B).

Under direct interspecific competition, sexual dimorphism in resource specialization has

striking effects on resource persistence when the sexes differ in total resource acquisition,

expanding regions of coexistence (Figs 5A, B) and in some cases leading to persistence of a

competitively inferior resource (Figs 5C, D; Supplement B Figs S4B, C). Moreover, sex

differences in total resource acquisition can reverse competitive outcomes when resources differ

in their intrinsic per-capita growth rates and the degree of resource specialization is held constant

(Figs 5C, D). The effect of sex differences in total resource acquisition on patterns of prey

persistence can be as striking as the effects of resource specialization (Supplement B Figs S2,

S1).

Asymmetric contribution of resource acquisition to birth rate

Introducing asymmetries in the contribution of sex-specific resource acquisition to birth rate

(e.g., polygyny or polyandry, manifest as changes of the constant b_f), had little qualitative effect.

This similarity is illustrated in Supplement B figure S3 (compare to S1). Although changing

these constants results in shifts of absolute regions of coexistence, the influence of consumer sex

differences was similar. Lack of sensitivity in asymmetries in the contribution of resource

acquisition to birth rates is consistent with our analytical results (Supplement A) that apply to all

first-order homogenous mating functions including geometric and arithmetic contributions to

reproduction.

Discussion

Using a general model of consumer-resource dynamics we show that consumer sexual

dimorphism has substantial consequences for community assembly. Competitive exclusion via

apparent competition is expected and observed when males and females are monomorphic

generalists and resources differ substantially in their intrinsic per-capita growth rates. However,

when male and female consumers differ in their resource-specific attack rates, resource species

that differ substantially in their intrinsic per-capita growth rates can coexist.   Similar effects of

consumer sexual dimorphism are observed when resources compete directly, with sexual

dimorphism in some cases permitting coexistence or persistence of a competitively inferior

resource (summarized in Table 1). However, for both direct and apparent competition, consumer

sexual dimorphism can also lead to competitive exclusion between resources that would typically

be expected to coexist. Thus, consumer sex differences result in fundamental changes in the

types of competing resources that can establish during community assembly. These results also

support conclusions from other food web models suggesting trophic position may impact the

observed ecological effects of sexual reproduction (Kawatsu 2018). Moreover, equilibrium

resource abundances and temporal dynamics are altered by consumer sexual dimorphism even

when long-term ecological outcomes are unaffected. We expect to observe these types of effects

whenever consumer mating propensity depends, in part, on resource acquisition, a biological

reality for many sexually-reproducing consumers (e.g. widespread evidence of developmental

thresholds for transitions to reproduction, reveiwed in Wilbur and Collins 1973, Day and Rowe

2002, and resource-dependent sexual displays, Bonduriansky 2007).

Although our results suggest that the effects of consumer sexual dimorphism on

community assembly can be complex, our analysis reveals some key predictions from our model.

First, consumer sex differences in resource-specific attack rates result in an increase in the

equilibrium density of each resource and a decrease in the abundance of the consumer species.

Intuitively, sexual dimorphism can frequently lead to slightly suboptimal resource use of the

species due to a mismatch between demand (restricted by a 50:50 sex ratio) and resource

availability which can be more variable and dynamic. Second, consumer sexual dimorphism can

promote coexistence when the more aggressively-feeding sex (higher total attack rates)

specializes on and suppresses what would otherwise be the competitively superior resource. This

later situation may be commonplace if sex-specific natural selection favors specialization, by the

sex with higher resource requirements, on the most abundant resource. Finally, and related to the

previous observation, the effects of consumer sexual dimorphism are most pronounced when

resources either differ in intrinsic per-capita growth rates, or when consumer sexes differ in their

total resource acquisition.

Our simulation models focused on a harmonic mean birth function, which is biologically

realistic in that it captures severely reduced birth rates when any one sex is rare. Empirically

evaluating alternative birth functions in real systems is difficult, requiring extreme variation in

adult sex ratios rarely seen in many species. However, some empirical data exists and supports

the harmonic mean as the best-fitting description of sexual birth rates (Miller and Inouye 2011).

Nonetheless, our analytical results (see Supplement A for details) that our conclusions apply to

any birth function that is first order homogenous i.e. $B(aMI_M, aFI_F) = aB(MI_M, FI_F)$ for any

positive constant $a>0$. These functions include all functions typically considered candidate birth

functions by demographers (Caswell and Weeks 1986, Miller and Inouye 2011). Our

conclusions do depend on the assumption that male abundance makes some contribution to birth

rate.

Although even moderate sex differences in attack rates can change the conditions under

which resources coexist, the strongest effects are observed when the consumer population

approaches complete sexual dimorphism in resource specific attack rates. Cases of such extreme

sex differences in diet preference exist (Temeles et al. 2000), although whether they occur with

any regularity is unclear. Diet divergence between the sexes, and sexual dimorphisms in trophic

morphology, are common (Shine 1989), although overlap in male and female diets can be

substantial even in the presence of ecological sex differences (e.g., Stamps et al. 1997, De Lisle

and Rowe 2015a). Yet overlap in diet content may be expected even in the case of large

differences in attack rates, as diet content is the product of both attack rate and resource

abundance. Further, males and females may fail to diverge in diet preference even when

optimum diet nutritional content differs across the sexes (Reddiex et al. 2013). These challenges

to understanding sex-specific resource acquisition highlight the need for further empirical studies

disentangling expressed diet preference, male and female nutritional optima, and the evolution of

ecological sexual dimorphism.

We have assumed a Fisherian sex ratio, where the primary sex ratio is maintained at a

stable 1:1 ratio (Fisher 1930), although our model allows for deviations from a 1:1 operational

sex ratio via sex-specific intrinsic mortality. Sex-specific mortality is commonplace in natural

populations and can occur for a variety of reasons, such as sex-specific costs of reproduction,

predation, or sexual conflict. In our model, sex-specific mortality, as well as variation in

consumer mating system (see Supplemental material) had predictable consequences for resource

abundance, leading to increases in the abundance of the resource favored by the sex with higher

intrinsic mortality. Nonetheless, the general effects of sexual dimorphism hold under sex-biased

mortality and departures from monogamy, indicating our general conclusions may also apply to

systems with diverse mating systems or where the primary sex ratio, or the operational sex ratio,

deviates from 1:1.

Our analysis also assumes that the male and female trophic traits are constant. In nature,

however, the degree and even direction of sexual selection can vary dramatically among closely

related populations (e.g., Reimchen et al. 2016). Dimorphism clearly evolves rapidly in response

to spatially varying sexual selection and resource availability. Future extensions to our strictly

ecological model could add in eco-evolutionary dynamics of sexual dimorphism.

Male and female densities were reduced with increasing ecological sexual dimorphism,

because consumer births are limited by resource acquisition in both sexes in our model. In the

extreme case, where the consumer cannot persist on a single resource (for example, under

complete sexual dimorphism), coexistence of the consumer depends on stable coexistence of the

resource species (Supplement A). Our simulations suggest robust persistence of extremely

dimorphic consumers in two-consumer communities, albeit at reduced density. Both of these

results – reduced consumer density and dependence on the presence of both resource species in

the community – suggest that sexually-dimorphic populations may face a higher risk of

extinction due to demographic stochasticity. However, it is difficult from our purely

deterministic ecological model to fully interpret the consequences of sexual dimorphism on

extinction probability. Evolution of sexual dimorphism from a monomorphic ancestor that

specializes on a single resource could lead to increased population mean fitness, by increasing

the total resource pool available across both sexes (Rand 1952, Selander 1966, Slatkin 1984);

such a process represents a form of within-species, between-sex ecological character

displacement. More generally, evolution of sexual dimorphism is expected to be critical to

population mean fitness whenever optima differ for males and females (Lande 1980), and

empirical data suggest sexual dimorphism can be associated with reduced extinction probability

at both the macroevolutionary scale and in extant populations (De Lisle and Rowe 2015b).

Nonetheless, our results suggest a full understanding of sexual dimorphism's role in population

persistence could require integrating theory and data from population/community ecology and

evolutionary genetics. This conclusion is complemented by recent theory (de Vries and Caswell

2019) indicating sexual dimorphism in demographic parameters can have important

consequences for maintaining genetic diversity.

Our model generally predicts that ecological sexual dimorphism may in some cases

promote, and in other cases reduce, diversity at lower trophic levels during community assembly.

Testing this prediction with correlative data (reviewed in Tsuji and Fukami 2020) would be

possible but challenging. An alternative and non-exclusive hypothesis, that dimorphic predators

are more likely to establish in communities with diverse prey assemblages, would also generate

patterns consistent with our results. Alternatively, experiments that manipulate the expressed

dimorphism of predators and track community dynamics at lower trophic levels are tractable in

some systems. Similar experiments have been performed (Fryxell et al. 2015, Start and De Lisle

2018), in which effects of predator sex ratio manipulation on prey communities are assessed in

mesocosm designs. Although the results of these experiments do suggest sex differences can

have important community consequences, no studies (to our knowledge) have compared

communities in which the magnitude of morphological sexual dimorphism is manipulated under

a stable sex ratio. Such designs are possible when distributions of male and female phenotypes

exhibit substantial variation, and would represent an ideal empirical test of the theoretical results

presented here.

Ecological sex differences are commonplace, although the details of their evolutionary

drivers and ecological consequences are unclear. Emerging theory and data indicate ecological

sex differences may have important consequences for the evolutionary genetics of adaptation

(Zajitschek and Connallon 2017), the dynamics of diversification (Bolnick and Doebeli 2003, De

Lisle and Rowe 2015b) and community assembly (Fryxell et al. 2015, Pincheira-Donoso et al.

2018, Start and De Lisle 2018). Our results add to this body of work, indicating that sexual

dimorphism can have substantial effects on the structure, abundance, and dynamics of ecological

communities, including changing conditions for coexistence between competing resource

species.

Acknowledgements. We thank Tim Connallon and Miguel Gomez for comments on an early

draft of this manuscript, as well as Gonzalo Hernando for assistance in the early stages of the

project. This work was supported by funds from the University of Connecticut to DIB, an

establishment grant from the Swedish Research Council to S.P.D. (2019-03706), and an U.S.

National Science Foundation Grant DMS176803 to SJS.

References

Andersson, M. B. 1994. Sexual selection. Princeton University Press, Princeton, N.J.

Belovksy, G. E. 1978. Diet optimization in a generalist herbivore: the

moose. Theoretical Population Biology **14**:105-134.

Bolnick, D. I., P. Amarasekare, M. S. Araujo, R. Burger, J. M. Levine, M. Novak, V. H. W.

Rudolf, S. Schreiber, M. C. Urban, and D. A. Vasseur. 2011. Why intraspecific trait

variation matters in community ecology. Trends in Ecology & Evolution **26**:183-192

Bolnick, D. I., and M. Doebeli. 2003. Sexual dimorphism and adaptive speciation: two sides of

the same ecological coin. Evolution **57**:2433-2449.

Bolnick, D. I., R. Svanbäck, J. A. Fordyce, L. H. Yang, J. M. Davis, C. D. Hulsey, and M. L.

Forister. 2003 The Ecology of Individuals: Incidence and implications of individual

specialization. The American Naturalist **161** 1-28.

Bonduriansky, R. 2007. The evolution of condition-dependent sexual dimorphism. The

American Naturalist **169**:9-19.

Camus, M. F., K. Fowler, M. W. D. Piper, and M. Reuter. 2017 Sex and genotype effects on

nutrient-dependent fitness landscapes in Drosophila melanogaster. Proceedings of the

Royal Society of London Series B. **284**:20172237.

Caswell, H., and D. E. Weeks. 1986. Two-sex models: chaos, extinction, and other dynamic

consequences of sex. American Naturalist **128**:707-735.

Cortez, M. H., and S. Patel. 2017. The effects of predator evolution and genetic variationon

predator-prey population-level dynamics. Bulletin of Mathematical Biology **79**:1510-

1538.

Darwin, C. 1871. The descent of man and selection in relation to sex. J. Murray, London.

Day, T., and L. Rowe. 2002. Developmental thresholds and the evolution of reaction norms for

age and size at life-history transitions. The American Naturalist **159**:338-350.

De Lisle, S. P. 2019. Understanding the evolution of ecological sex differences: Integrating

character displacement and the Darwin-Bateman paradigm. Evolution Letters **3**:444-447.

De Lisle, S. P., and L. Rowe. 2015a. Ecological character displacement between the sexes. The

American Naturalist **186**:693-707.

De Lisle, S. P., and L. Rowe. 2015b. Independent evolution of the sexes promotes amphibian

diversification. Proceedings of the Royal Society B: Biological Sciences **282**:20142213.

de Vries, C., and H. Caswell. 2019. Selection in two-sex stage structured populations: Genetics,

demography, and ploymorphism. Theoretical Population Biology **130**:160-169.

Des Roches, S., D. M. Post, N. E. Turley, J. K. Bailey, A. P. Hendry, M. T. Kinnison, J. A.

Schweitzer, and E. P. Palkovacs. 2018. The ecological importance of intraspecific

variation. Nature Ecology and Evolution **2**:57-64.

Fisher, R. 1930. The genetical theory of natural selection. Oxford University Press, Oxford, UK.

Fryxell, D. C., H. A. Arnett, T. M. Apgar, M. T. Kinnison, and E. P. Palkovacs. 2015. Sex ratio

variation shapes the ecological effects of a globally introduced freshwater fish.

Proceedings of the Royal Society of London Series B. **282**:20151970.

Fryxell, D. C., D. Weiler, M. Kinnison, and E. P. Palkovacs. 2019 Eco-evolutionary dynamics of

sexual dimorphism. Trends in Ecology and Evolution **in press**.

Garlapow, M. E., W. Huang, M. T. Yarboro, K. R. Peterson, and T. F. C. Mackay. 2015.

Quantitative genetics of food intake in Drosophila melanogaster. PloS OnE **10**:e0138129.

Giery, S. T., and C. G. Layman. 2019. Ecological consequences of sexually selected traits: An

eco-evolutionary perspective. The Quarterly Review of Biology **94**:29-74.

Gomez-Llano, M. A., H. M. Bensch, and E. I. Svensson. 2018. Sexual conflict and ecology:

species composition and male density interact to reduce male mating harassment and

increase female survival. Evolution **72**:906-915.

Haldane, J. B. S. 1937. The effect of variation on fitness. The American Naturalist **71**:337-349.

Holt, R. D. 1977. Predation, apparent competition, and the structure of prey communities.

Theoretical Population Biology **12**:197-229.

Holt, R. D., and J. H. Lawton. 1993. Apparent competition and enemy-free space in insect host-

parasitoid communities. The American Naturalist **142**:623-645.

Hughes, A. R., B. D. Inouye, M. T. J. Johnson, N. Underwood, and M. Vellend. 2008 Ecological

consequences of genetic diversity. Ecology Letters **11**:609-623.

Jensen, K., C. McClure, N. K. Priest, and J. Hunt. 2015. Sex-specific effects of protein and

carbohyfrate intake on reproduction but not lifespan in Drosophila melanogaster. Aging

Cell **14**:605-615.

Kawatsu, K. 2018. Ecological effects of sex differ with trophic positions in a simple food web.

Ecology and Evolution **8**:1239-1246.

Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.

Evolution **34**:292-305.

Lindstöm, J., and H. Kokko. 1998. Sexual reproduction and population dynamics: the role of

polygony and demographic sex differences. Proceedings of the Royal Society of London

Series B. **265**:483-488.

Maklakov, A., S. J. Simpson, F. Zajitchek, M. D. Hall, J. Dessmann, F. Clissold, D.

Raubenheimer, R. Bonduriansky, and R. C. Brooks. 2008. Sex-specific effects of nutrient

intake on reproduciton and lifespan. Current Biology **18**:1062-1066.

Miller, T. E. X., and B. D. Inouye. 2011. Confronting two-sex demographic models with data.

Ecology **92**:2141-2151.

Patel, S., and S. J. Schreiber. 2015. Evolutionarily driven shifts in communities with intraguild

predation. The American Naturalist **186**:E98-E110.

Pincheira-Donoso, D., T. Tregenza, R. K. Butlin, and D. J. Hodgson. 2018. Sexes and species as

rival units of niche saturation during community assembly. Global Ecology and

Biogeography **27**:593-603.

R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Rand, A. L. 1952. Secondary sexual characters and ecological competition. Fieldiana-Zoology

34:65-70.

Reddiex, A. J., T. P. Gosden, R. Bonduriansky, and S. F. Chenoweth. 2013. Sex-specific fitness

consequences of nutrient intake and the evolvability of diet preferences. The American

Naturalist **182**:91-102.

Reimchen, T., D. Steeves, and C. A. Bergstrom. 2016. Sex matters for defence and trophic traits

of threespine stickleback. Evolutionary Ecology Research **17**:459-485.

Rowe, L., and D. Houle. 1996. The lek paradox and the capture of genetic variance by condition

dependent traits. Proceedings of the Royal Society of London Series B-Biological

Sciences **263**:1415-1421.

Schreiber, S. J. 2021. The P* rule in the stochastic Holt-Lawton model of apparent competition.

Discrete and Continuous Dynamical Systems-B **26**:633.

Schreiber, S. J., R. Burger, and D. I. Bolnick. 2011. The community effects of phenotypic and

genetic variation within a predator population. Ecology **92**:1582-1593.

Selander, R. K. 1966. Sexual dimorphism and differential niche utilization in birds. The Condor

68:113-151.

Shine, R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the

evidence. Quarterly Review of Biology **64**:419-461.

Shärer, L., L. Rowe, and G. Arnqvist. 2012 Anisogamy, chance and the evolution of the sex

roles. Trends in Ecology & Evolution **27**:260-264.

Slatkin, M. 1984. Ecological causes of sexual dimorphism. Evolution **38**:622-630.

Soetgart, K., T. Petzoldt, and R. W. Setzer. 2010. Solving Differential Equations in R: Package

deSolve. Journal of Statisitcal Software **33**:1-25.

Stamps, J. A., J. B. Losos, and R. M. Andrews. 1997. A comparative study of population density

and sexual size dimorphism in lizards. American Naturalist **149**:64-90.

Start, D., and S. P. De Lisle. 2018. Sexual dimorphism in a top predator drives aquatic prey

community assembly. Proceedings of the Royal Society of London Series B. **2018**:1717.

Svensson, E. I. 2019. Eco-evolutionary dynamics of sexual selection and sexual conflict.

Functional Ecology **33**:60-72.

Temeles, E. J. 1985. Sexual size dimorphism of bird-eating hawks: the effect of prey

vulnerability. The American Naturalist **125**:485-499.

Temeles, E. J., I. L. Pan, J. L. Brennan, and J. N. Horwitt. 2000. Evidence for ecological

causation of sexual dimorphism in a hummingbird. Science **289**:441-443.

Trivers, R. L. 1972. Parental investment and sexual selection. Pages 136-179 *in* B. Campbell,

editor. Sexual selection and the descent of man. Aladine, Chicago.

Tsuji, K., and T. Fukami. 2018. Community-wide consequences of sexual dimorphism: evidence

from nectar microbes in dioecious plants. Ecology **99**:2476-2484.

Tsuji, K., and T. Fukami. 2020. Sexual dimorphism and species diversity: from clades to sites.

Trends in Ecology and Evolution **35**:105-114.

Tuszynski, J. 2019. caTools: Tools: moving window statistics, GIF, Base64, ROC AUC, etc.. R

package version 1.17.1.2. https://CRAN.R-project.org/package=caTools.

Weimerskirch, H., M. Le Corre, H. Gadenne, D. Pinaud, A. Kato, Y. Ropert-Coudert, and C.-A.

Bost. 2009. Relationship between reversed sexual dimorphism, breeding investment and

forgaing ecology in a pelagic seabird, the masked booby. Oecologia **161**:637-649.

Wilbur, H. M., and J. P. Collins. 1973. Ecological aspects of amphbian metamorphosis. Science

182:1305-1314.

Zajitschek, F., and T. Connallon. 2017. Partitioning of resources: the evolutionary genetics of

sexual conlflict over resource acquisition and allocation. Journal of Evolutionary Biology

30:826-838.

Table 1. Summary of effects of sex differences in resource use. Arrows indicate effects on mean species density and the range of parameter space under which resource coexistence is observed.

Density	Coexistence	
Consumer ↓	Apparent competition	↑
Resource ↑	Direct competition, equal resource growth rates, symmetric acquisition	0
	Direct competition, different resource growth rates, symmetric acquisition	↑
	Direct competition, asymmetric acquisition	↑↓

Figure 1. Illustration of the model structure. Males (M) and females (F) of a consumer

species exploit two resources (R_1 and R_2) that may or may not also compete directly (double

arrow). Sex-specific attack rates (black arrows) generate sex differences in ecological niche such

that males and females preferentially attack R_1 and R_2, respectively, with varying degrees of

overlap (grey arrows). Although males and females may consume different resources, fitness is

equal across the sexes and so the dynamics of males, females, resource 1 and resource 2 are

coupled.

Figure 2. Coexistence mediated by sexual dimorphic consumers. In panel A, resource 2 (red)

is a superior competitor to resource 1 (blue), leading to exclusion of resource 1 despite the

presence of the consumer species. However, adding sex differences in total resource acquisition,

under otherwise identical parameter values, leads to coexistence with cyclical dynamics (B).

When competition is completely apparent (C, D), the resource with superior growth rate (red)

excludes the inferior resource when consumers are sexually monomorphic (C). Under the same

parameterization but with sexually dimorphic consumers (D), density of both resources is

increased and they coexist. Parameter values: **A**, $\alpha_{11} = .1$, $\alpha_{12} = .12$, $\alpha_{22} = .1$, $\alpha_{21} = .09$, $\beta = 1$, d_M

$= d_F = 0.02$, $b_M = b_F = .1$, $a_{M, max} = a_{F, max} = 1$, $r_1 = r_2 = 1$; **B** $\alpha_{11} = .1$, $\alpha_{12} = .12$, $\alpha_{22} = .1$, $\alpha_{21} =$

$.09$, $\beta = 1$, $d_M = d_F = 0.02$, $b_M = b_F = .1$, $a_{M, max} = .8$, $a_{F, max} = 1.2$, $r_1 = 1$, $r_2 = 1$; **C** $\alpha_{11} = .1$, $\alpha_{12} =$

611   0, $\alpha = .1$, $\alpha_{21} = 0$, $\beta = 0$, $d_M = d_F = 0.02$, $b_M = b_F = .1$, $a_{M,\,max} = 1$, $a_{F,\,max} = 1$, $r_1 = 1$, $r_2 = 1.1$; **D**

612     $\alpha_{11} = .1$, $\alpha_{12} = 0$, $\alpha_{22} = .1$, $\alpha_{21} = 0$, $\beta = 0.9$, $d_M = d_F = 0.02$, $b_M = b_F =$, $a_{M,\,max} = 1$, $a_{F,\,max} = 1$, r_1

613     $= 1$, $r_2 = 1.1$;

Figure 3. Consumer sexual dimorphism results in increased resource density and reduced consumer density in two and three species communities. Panel A: Equilibrium consumer female density decreases and resource 1 density increases with increased sexual dimorphism in a two species (consumer- 1 resource) community (equation 6). Panel B: these results hold in simulations in three species communities, where resource 1 density is plotted against competition coefficients in both resource species under three levels of sexual dimorphism, increasing from left to right. Assuming symmetric total attack rates across the sexes, symmetric contribution of resource acquisition to consumer birth rates, and equal growth rates across resources.

A

B

Figure 4. Sexual dimorphism expands regions of coexistence between resources under

apparent competition. Panels show relative density of resource 1 ($R_1/(R_1+R_2)$) from numerical

simulations with starting conditions of $R_1 = R_2 = 1$ and $M = F = 1$. The top panel A shows the

outcomes under symmetric maximal attack rates across the sexes ($a_{M,\,max} = a_{F,\,max} = 1$). Panel

B shows the outcomes assuming sex differences in maximal attack rate ($a_{M,\,max} = .8$, $a_{F,\,max} =$

1.2). White dashed lines indicate analytical solutions (equation 8).

Figure 5. Sexual dimorphism alters regions of coexistence of resource species. Panels show

relative density of resource 1 ($R_1/(R_1 + R_2)$) from numerical simulations with starting conditions

of $R_1 = R_2 = 1$ and $M = F = 1$. Panel A shows outcomes under sexual monomorphism ($\beta = 0$),

unequal growth rates ($r_1 = 2$, $r_2 = 2.1$) across resources, and no sex differences in total attack rate

($a_{M, max} = a_{F, max} = 1$). Panel B shows the outcome under the same parameter values but with

strong consumer sexual dimorphism ($\beta = 0.7$). Panels C and D show the effect of sexual niche

divergence when the sexes also differ in total attack rates (total resource acquisition), with C

showing the outcomes under unequal total attack rates ($a_{M, max} = .7$, $a_{F, max} = 1.3$), sexual

monomorphism in prey preference ($\beta = 0$), and differential resource growth rates ($r_1 = 1.6$, $r_2 = $

1.8), and D showing the outcome under the same parameter values but with moderate sexual

665   dimorphism in prey preference ($\beta = 0.4$). Black dashed lines demark equal inter and intraspecific

666   competition coefficients. White dashed box demarks coexistence determined by analytical

667   invasion criteria into a two-species community under equation 9.