Consumer sexual dimorphism promotes coexistence among competing resources

Stephen P. De Lisle [1] *

Gonzalo Hernando [2]

Daniel I. Bolnick [1]

[1] Department of Ecology & Evolutionary Biology, University of Connecticut, Storrs, CT 06269

[2] Healthway International Communications Inc., 4030 Braker Lane West, Suite 175, Austin, TX 78759

* email: stephen.de_lisle@uconn.edu

Keywords: apparent competition, community assembly, competitive exclusion, ecological sexual dimorphism, resource competition

Statement of authorship: DB and GH initially designed the study, GH and SD performed the simulations, SD wrote the manuscript with DB.

Data accessibility statement: Complete R script and numerical solution results are provided in the supplementary material

Abstract

Within-species variation is a salient feature of natural populations, of substantial importance for

species interactions. However, the community consequences of sexual dimorphism, one of the

most ubiquitous sources of within-species variance, remains poorly understood. Here, we extend

classical models of consumer-resource dynamics to explore the ecological consequences of

consumer sexual dimorphism. We show that sexual dimorphism in consumer attack rates on two

different resource species promotes coexistence between those resources, mitigating the effects

of both apparent competition and direct interspecific competition. Consumer sexual dimorphism

can prevent exclusion of a resource with inferior growth rates because reduction in any of the

two resources reduces consumer density, generating negative frequency dependence that

stabilizes coexistence between resources. Our work highlights ecological sex differences as a

potentially key factor governing the assembly of ecological communities, illustrating that the

specific source of within-species variance can have important implications for community

ecology.

Introduction

Species composition varies substantially across ecological communities, inspiring a rich body of theory and data aimed at identifying the processes that govern community assembly. Resulting models of consumer-resource dynamics yield general conclusions for the circumstances under which competing species can or cannot coexist. This modern coexistence theory tells us that competitive exclusion is expected when competing, or apparently-competing (via shared predation; Holt 1977), species differ substantially in per-capita growth rates and/or when interspecific competition is stronger than intraspecific competition (Chesson 2000). Mechanisms promoting coexistence thus act either to equalize per-capita growth rates or stabilize interactions between species. Many extensions of simple consumer-resource models (summarized in Ellner et al. 2018) have identified features typical of natural populations that affect these two pathways to coexistence.

Within-species variation is one key feature of natural populations that has emerged as a critical contributor to community ecology and species interactions (Bolnick et al. 2011, Start and Gilbert 2017, Des Roches et al. 2018, Start 2018). Phenotypic variation in consumers and/or their resources can either promote or constrain coexistence, under many circumstances yielding different conclusions than would be reached from models ignoring within species variation (Bolnick et al. 2011, Schreiber et al. 2011, Patel and Schreiber 2015, Cortez and Patel 2017). Such variation can arise from evolutionary processes, such as mutation, the maintenance of standing genetic variance (including differences between the sexes), and ontogenetic change, as well as environmental inputs and developmental plasticity. With the possible exception of ontogenetic change, the source of phenotypic variance is rarely considered in ecological models, to the extent that variation is considered at all.

Here, we explore the ecological consequences of sexual dimorphism, a central feature of

metazoans and many plant populations that has not been fully incorporated into ecological

theory. The gamete differences that define the sexes are expected to lead to divergence between

males and females in a suite of life history traits (Trivers 1972, Shärer et al. 2012) iconically

manifest as striking sexual dimorphism in sexually-selected traits (Darwin 1871, Andersson

1994). Although sexual selection and associated mate choice behaviors themselves may have

little obvious relevance to interspecific ecological interactions (but see Gomez-Llano et al.

2018), the different life histories that define the sexes are often expected lead to different

nutritional and resource requirements for males and females (Belovksy 1978, Maklakov et al.

2008). As a specific example, in insects, differential contributions of longevity and fecundity to

male and female lifetime reproductive success result in different combinations of macronutrients

that maximize male and female fitness (Maklakov et al. 2008, Reddiex et al. 2013, Garlapow et

al. 2015, Jensen et al. 2015, Camus et al. 2017). Consequently, males and females of many

species have evolved divergent resource use, either as an indirect outcome of divergent

reproductive roles or through other forms of sex-specific natural selection, resulting in sex-

differences in diet composition and often in corresponding trophic morphology (Slatkin 1984,

Temeles 1985, Shine 1989, Temeles et al. 2000, De Lisle and Rowe 2015a). Data suggest that

these 'ecological' sexual dimorphisms can have substantial consequences for community

structure (Fryxell et al. 2015, Pincheira-Donoso et al. 2018, Start and De Lisle 2018). More

generally, sex differences may play an important role in the relationship between ecological and

evolutionary dynamics (Giery and Layman 2019, Svensson 2019, Fryxell et al. 2019). Sexual

dimorphism thus represents a key source of ecologically-relevant variation within species.

Critically, this source of intraspecific variation may have different consequences from other

types of variation. Unlike most phenotypes, whose relative abundances can evolve, the ratio of

males to females (at birth) is usually genetically fixed at 50:50 (Fisher 1930), though their

phenotypes and demography may diverge to have differential ecological impacts. Using a

simple model of consumer-resource dynamics, we show that consumer sexual dimorphism has

important consequences for coexistence between resources that compete directly or apparently.

Model

Our model represents a simple extension of the classic model of consumer-resource dynamics, in

which a consumer species exploits two resources that themselves undergo density-dependent

growth (Figure 1). Consumer growth rate is limited only by resource abundance. Our model

mainly differs from existing models that do (Schreiber et al. 2011) and do not (e.g. Holt 1977)

incorporate within-species variance in that we explicitly model male and female demography,

allowing for the possibility that males and females differ in resource-specific attack rates. We

can describe the population dynamics of two resources (R_1 and R_2) and consumer males (M) and

females (F) using the system of ordinary differential equations:

$$\frac{dR_n}{dt} = r_n R_n \left(1 - \frac{R_n + a_R R_j}{K_n}\right) - a'_{M,n} R_n M - a'_{F,n} R_n F \tag{1a}$$

$$\frac{dM}{dt} = \frac{1}{2} B(M, F, R_1, R_2) F - d_M M \tag{2b}$$

$$\frac{dF}{dt} = \frac{1}{2} B(M, F, R_1, R_2) F - d_F F \tag{3c}$$

In which M, F, R_n (where $n = 1, 2$) are the population densities for consumer males, consumer

females, the resources, respectively, $M + F$ = Total Predator population (N) such that $\frac{dN}{dt} =$

$\frac{dM}{dt} + \frac{dF}{dt}$. Each resource n undergoes density-dependent population growth with birth rate r_n

towards its carrying capacity K_n, with interspecific competition represented by a_R. For

simplicity we assume symmetric competition between the resources. Resources are also

regulated by consumers depending on sex-specific attack rates by males $a'_{M,n}$ and by females

$a'_{F,n}$. Consumer dynamics are governed by the product of birth rates B and female density,

limited by intrinsic mortality d_M and d_F. An appropriate function describing birth rates has been

a point of debate for demographers, with the general conclusion being that any function

describing birth rates should capture Allee effects likely to limit birth rates in the case of extreme

sex ratio skew, with the extreme being that birth rates should be zero when one sex is absent

(Caswell and Weeks 1986). Generally, treating birth rates proportional to the harmonic mean

density of males and females is agreed as the best approach (Caswell and Weeks 1986, Lindstöm

and Kokko 1998). Here, we extend this harmonic mean birth function:

$$B(M, F, R_1, R_2) = 2k \frac{MI_M * FI_F}{MI_M + FI_F} \tag{2}$$

128 Where I is the energy intake of a given sex:

$$I_i = \sum_{j=1}^{2} a'_{ij} R_j \qquad for\ i = M, F \tag{3}$$

which couples the dynamics of consumers and resources. In this model the rate of consumer

births depends on not just the number of males and females but also on the abundance of both

prey and the prey preference of each sex, where MI_M is the number of males who have the

energy to reproduce (similarly with females), k is a scaling constant reflecting the number of

predator offspring created per prey eaten, and a_{ij} as the attack rate by the i predator to the j

resource. We define the degree of sexual dimorphism, α, such that

$$a'_{M,1} = a'_{F,2} = a'_{max}\left(1 - \frac{\alpha}{2}\right) \tag{4a}$$

$$a'_{M,2} = a'_{F,1} = a'_{max}\left(\frac{\alpha}{2}\right) \tag{4b}$$

where α ranges from zero to one, with α = 1 representing complete sexual monomorphism (both

sexes are generalists, attacking each prey at the same rate, $a'_{max}/2$), and α = 0 representing

complete sexual dimorphism such that males only attack R_1 and females only attack R_2. For

simplicity we assume a symmetric dimorphism in attack rates.

We explored deviations from the model in equation 1, including making consumer

growth rates a function of only B and d (removing the product with F), relaxing the implicit

assumption of monogamy, as well as including a term further penalizing population growth rates

under male-biased sex ratio skew (e.g., due to interlocus sexual conflict), which had nearly

equivalent effects to introducing simple sex differences in death rate. In all cases qualitative

conclusions remain unchanged. In the appendix, we explore the consequences of consumer

mating system variation, although report only the simplified cases in the main text as conclusions

appear robust to variation in consumer mating system.

We were unable to solve the system of equations 1 analytically, and so numerically

solved the differential equations using the packages deSolve v 1.21 (Soetgart et al. 2010) and

caTools v. 1.17.1.2 (Tuszynski 2019) in R v. 3.5.0 (R Core Team 2018). We iterated through

different values of dimorphism (α from 0 to 1 by 0.1), r_1 and r_2 (from 1 to 1.4 by 0.05), d_M and

d_F (0.02, 0.024), under three different scenarios of interspecific competition between resource 1

and 2: apparent competition, where $a_R = 0$ (see equation 1A), moderate interspecific competition

where $a_R = 0.5$, and strong interspecific competition where $a_R = 1$. For each set of parameter

values, we ran the numerical solution to equations 1a-c through 10,000 time steps (to ensure the

simulation reaches its equilibrium) by increments of 0.1 (to create an accurate discrete model).

For each run, we determined the equilibrium abundances of M, F, R_1 and R_2 at t = 10,000. We

also calculated the standard deviation of species' abundances in the last 50 time-steps (500 sub

time steps); all but 9 (all with sex-biased mortality and direct interspecific competition) of the

2244 parameter combinations reached stable equilibrium. Parameters held constant in our

numerical solutions were K = 10000, k = 0.0001, and $a'_{max} = 0.0009$. These values were chosen

to generate realistic values for the resulting per capita growth rates, and conclusions are not

qualitatively changed under deviations from them. The R script to generate all figures and

results presented, as well as a file containing numerical solutions to the model, is available in the

supplementary material.

Results

Exploring the effects of sexual dimorphism on consumer birth rates under the

assumptions of equation 2 reveals some important properties of the model. First, because

consumer birth rates depend on resource abundance, consumer birth rates decline when density

of any single resource declines (Figure 2), even under complete sexual monomorphism.

However, introducing sexual dimorphism generates a non-linear relationship between resource

abundance and consumer birth rates such that birth rates decline rapidly as a single resource

declines in abundance. When dimorphism is complete, birth rates approach zero as abundance of

a single resource approaches zero, even if the second resource is abundant (Figure 2). Thus,

sexual dimorphism effectively generates negative frequency dependence such that declining

abundance of a single resource is expected to reduce predation pressure via the feedback between

resource abundance and consumer population growth rate. Although mating system variation

may influence the sex ratio under which birth rates are maximized, these general effects are none

the less expected to hold even when the assumption of monogamy is relaxed (see Appendix).

Because resource population growth rates are assumed to be dependent in part on consumer

density (equation 1A), we may expect such a feedback to generally expand the parameter space

under which resources may coexist.

We next explore the temporal dynamics of R_1 and R_2 under three scenarios of apparent

competition. When growth rates differ across resources and consumers are sexually

monomorphic, the inferior resource rapidly declines to extinction (Figure 3, top panel). This

result reflects the classic insight that a resource cannot exist in a community dominated by other

resources with superior growth rates in the face of predation (Holt 1977, Holt and Bonsall 2017).

When sexes differ in their resource preference completely, such that each sex specializes on a

single resource, the inferior resource is maintained and each population in the community cycles

to a stable equilibrium (Figure 3, middle panel). This cycling appears to be the result of

nonlinearities introduced from both sexual dimorphism and differences in resource growth rates;

temporal stability increases in the case of equal growth rates across both resources (Figure 3,

bottom panel). These results suggest consumer sexual dimorphism and resource growth rates

together determine equilibrium densities of each resource, and we next explore this further by

seeking equilibrium solutions to equations 1.

When consumer male and female death rates are equal and resource competition is

apparent, increasing consumer sexual dimorphism has the effect of increasing the parameter

space under which resources can coexist. As sexual dimorphism approaches the extreme (each

sex uses a different resource), a resource with inferior growth rates can coexist with a superior

growing resource under apparent competition (Figure 4A). This effect of dimorphism on

coexistence holds under moderate to strong interspecific competition (Figure 4B-C), as well as

when the assumption of equal death rates across consumer sexes is relaxed. Introducing sex

differences in death rates has predicable effects on resource density, leading to an increase in

density of the resource preferred by the sex with higher mortality and an asymmetric response of

each resource to varying levels of sexual dimorphism (Figure 5). Allowing for deviations from

monogamy (Appendix Figure 1), where consumer birth rates are maximized under sex ratios

other than 1:1, and/or incorporating sexual conflict (Appendix Figure 2, 3), where female

survival is differentially impacted by sex ratio skew, had similar effects to simply incorporating

sex-specific intrinsic mortality (see Appendix).

Sexual dimorphism also effects equilibrium consumer density, leading to a reduction in

total density of males and females with increasing sexual dimorphism. However, these changes

(Appendix Figure 4A, B) were not as extreme as that seen in equilibrium densities of the two

resources, and generally consumer coexistence is expected. When interspecific resource

competition equals intraspecific competition ($a_R = 1$), sexually dimorphic consumers cannot exist

with resources that differ in growth rate (Appendix Figure 4C), as the superior resource drives

extinction of both the inferior resource and, as a result, the consumer. However, the biological

relevance of such a scenario, with resource ecological equivalence yet complete consumer

dimorphism in attack rates, seems limited.

Discussion

Using a simple model of consumer-resource dynamics we show that consumer sexual

dimorphism has substantial consequences for the coexistence of prey that compete directly or

apparently. When male and female consumers differ in their resource-specific attack rates,

resources that differ in per-capita growth rates can coexist, while competitive exclusion is

expected and observed when males and females are monomorphic generalists and resources

differ substantially in per capita growth rates. Coexistence occurs when the sexes differ in attack

rates because sexual reproduction results in a phenotypic subsidy (Bolnick et al. 2011) in which

male and female birth rates are coupled; decreasing abundance of the slower growing resource

decreases both consumer male and female birth rates, generating negative frequency dependence

that prevents competitive exclusion of the inferior resource. Thus, in this model consumer sexual

dimorphism results in a phenotypic subsidy that effectively generates a stabilizing mechanism of

coexistence between the resources (Chesson 2000). These results hold whether competition

between the resources is completely apparent, occurring only via shared predation (Holt 1977,

Holt and Bonsall 2017), or in part direct, indicating that sexual dimorphism may have general

stabilizing effects on coexistence between competing species at lower trophic levels whenever

consumer birth rates are affected by resource aquistion, and may in part explain the observation

that competitive exclusion under apparent competition has only rarely been observed empirically

(Schreiber et al. 2011). These results also support conclusions from other food web models

suggesting trophic position may impact the observed ecological effects of sexual reproduction

(Kawatsu 2018). Importantly, our work also indicates that the specific source of variation may

often need to be considered to fully understand the role of within-species variation in community

assembly.

Although even moderate ecological sexual dimorphism increases the parameter space

under which resources coexist, the strongest effects are observed as the consumer population

approaches complete sexual dimorphism in attack rates. Cases of such extreme sex differences

in diet preference exist (Temeles et al. 2000), although whether they occur with any regularity is

unclear. Diet divergence between the sexes, and sexual dimorphisms in trophic morphology, are

common (Shine 1989), although overlap in male and female diets can be substantial even in the

presence of ecological sex differences (e.g., Stamps et al. 1997, De Lisle and Rowe 2015a). Yet

overlap in diet content may be expected even in the case of large differences in attack rates, as

diet content is the product of both attack rate and resource abundance. Further, males and

females may fail to diverge in diet preference even when optimum diet nutritional content differs

across the sexes (Reddiex et al. 2013). These challenges to understanding sex-specific resource

acquisition highlight the need for further empirical studies disentangling expressed diet

preference, male and female nutritional optima, and the evolution of ecological sexual

dimorphism.

We have assumed a Fisherian sex ratio, where the primary sex ratio is maintained at a

stable 1:1 ratio (Fisher 1930), although our model allows for deviations from a 1:1 operational

sex ratio via sex-specific intrinsic mortality. Sex-specific mortality is commonplace in natural

populations and can occur for a variety of reasons, such as sex-specific costs of reproduction,

predation, or sexual conflict. In our model, sex-specific mortality, as well as variation in

consumer mating system (see Appendix) had predictable consequences for resource abundance,

leading to increases in the abundance of the resource favored by the sex with higher intrinsic

mortality. Nonetheless, the general effects of sexual dimorphism on coexistence hold under sex-

biased mortality and departures from monogamy, indicating our general conclusions may also

apply to systems with diverse mating systems or where the primary sex ratio deviates from 1:1.

Male and female densities were reduced with increasing ecological sexual dimorphism,

because consumer births are limited by resource acquisition in both sexes in our model. This

suggests that sexually-dimorphic populations may face a higher risk of extinction due to

demographic stochasticity. However, it is difficult from our ecological model, that always

assumes two resource populations, to fully interpret the consequences of sexual dimorphism on

extinction probability. Evolution of sexual dimorphism from a monomorphic ancestor that

specializes on a single resource could lead to increased population mean fitness, by increasing

the total resource pool available across both sexes (Rand 1952, Selander 1966, Slatkin 1984);

such a process represents a form of within-species, between-sex ecological character

displacement. More generally, evolution of sexual dimorphism is expected to be critical to

population mean fitness whenever optima differ for males and females (Lande 1980), and

empirical data suggest sexual dimorphism can be associated with reduced extinction probability

at both the macroevolutionary scale and in extant populations (De Lisle and Rowe 2015b).

Nonetheless, our results suggest a full understanding of sexual dimorphism's role in population

persistence could require integrating theory and data from population/community ecology and

evolutionary genetics.

Our model generally predicts that ecological sexual dimorphism may promote diversity at

lower trophic levels during community assembly. Testing this prediction with correlative data

would be possible but challenging. An alternative and non-exclusive hypothesis, that dimorphic

predators are more likely to establish in communities with diverse prey assemblages, would also

generate such a pattern. Alternatively, experiments that manipulate the expressed dimorphism of

predators and track community dynamics at lower trophic levels are tractable in some systems.

Similar experiments have been performed (Fryxell et al. 2015, Start and De Lisle 2018), in which

effects of predator sex ratio manipulation on prey communities are assessed in mesocosm

designs. Although the results of these experiments do suggest sex differences can have important

community consequences, no studies (to our knowledge) have compared communities in which

the magnitude of morphological sexual dimorphism is manipulated under a stable sex ratio. Such

designs are possible when distributions of male and female phenotypes exhibit substantial

variation, and would represent an ideal empirical test of the theoretical results presented here.

Ecological sex differences are commonplace, although the details of their evolutionary

drivers and ecological consequences are unclear. Emerging theory and data indicate ecological

sex differences may have important consequences for the evolutionary genetics of adaptation

(Zajitschek and Connallon 2017), the dynamics of diversification (Bolnick and Doebeli 2003, De

Lisle and Rowe 2015b) and community assembly (Fryxell et al. 2015, Pincheira-Donoso et al.

2018, Start and De Lisle 2018). Our work adds to this body of work, indicating that sexual

dimorphism can have predictable effects on the structure of ecological communities, by

introducing frequency-dependence that stabilizes coexistence between resource species.

Acknowledgements We thank Miguel Gomez for comments on the manuscript. This work was

supported by funds from the University of Connecticut to DIB.

Figure 1. Illustration of the model structure. Males (M) and females (F) of a consumer

species exploit two resources (R1 and R2) that may or may not also compete directly (double

arrow). Sex-specific attack rates (black arrows) generate sex differences in ecological niche such

that males and females preferentially attack R1 and R2, respectively, with varying degrees of

overlap (grey arrows). Although males and females may consume different resources, fitness is

equal across the sexes and so the dynamics of males, females, resource 1 and resource 2 are

coupled.

Figure 2. Consumer birth rate as a function of Resource 1 density, holding R2 and

consumer density constant. Birth rates are reduced with reduced R1 density even when

consumers are sexually monomorphic generalists (alpha = 1, blue line) because total resources

(R1 plus R2) are reduced. Adding in sexual dimorphism in attack rates generates nonlinearity

that leads to rapid reduction in consumer birth rates with decreasing R1 density (alpha = 0.5,

purple line, partial overlap between the sexes; alpha = 0, red line, complete sexual dimorphism).

Consumer and R2 density were fixed at their equilibrium values obtained as the numerical

solutions to equations 1.

Figure 3. Temporal resource dynamics under three scenarios of apparent competition. In

the top panel, differential growth rates across the two resources lead to rapid extinction of

resource 1 when consumers are sexually monomorphic. When consumers are completely

dimorphic in attack rates, resource 1 can coexist with the superior resource 2, undergoing cyclic

dynamics towards equilibrium; middle panel. The bottom panel illustrates dynamics when

consumers are completely dimorphic and resources have equal growth rates.

Figure 4. Effects of differential growth rates and consumer sexual dimorphism on resource

1 abundance when consumer mortality is equal across the sexes. Values on z axis are

numerical solutions to equation 1 for resource 1 density, with resource 1 growth rate fixed at 1.

Competitive exclusion of resource 1 is observed when growth rates differ substantially and

consumers are sexually monomorphic (niche overlap = 1). Increasing sexual dimorphism

increases the difference in growth rates under which resource 1 can coexist with resource 2. This

effect is observed when A) competition between the resources is due only to shared predation, as

well as when direct interspecific competition between resource 1 and 2 occurs (B and C).

Figure 5. Effects of differential growth rates and consumer sexual dimorphism on resource abundance when mortality is sex-specific (female-biased). Top panels (A-C) are Resource 1 abundance with fixed $r_1 = 1$, bottom panels (D-F) are Resource 2 abundance with fixed $r_2 = 1$. Values on z axis are numerical solutions to equation 1. Introducing sex-biased mortality effectively makes it more difficult for the resource favored by the more abundant sex (in this case, resource 1, preferred by males) to coexist with the resource favored by the rarer sex, as illustrated for resource 1 abundance under varying degrees of direct competition in panels A-B. Differential growth rates have less effect on existence of the resource favored by the rare sex, as illustrated for resource 2 in D-F. In all cases, consumer sexual dimorphism has the effect of promoting coexistence between the two resources. Note differences in z axis scale between top and bottom panels.

388 **References**
389
Andersson, M. B. 1994. Sexual selection. Princeton University Press, Princeton, N.J.

Arnqvist, G., and L. Rowe. 2005. Sexual Conflict. Princeton University Press

, Princeton NJ.

Belovksy, G. E. 1978. Diet optimization in a generalist herbivore: the

moose. Theoretical Population Biology **14**:105-134.

Bolnick, D. I., P. Amarasekare, M. S. Araujo, R. Burger, J. M. Levine, M. Novak, V. H. W.

Rudolf, S. Schreiber, M. C. Urban, and D. A. Vasseur. 2011. Why intraspecific trait

variation matters in community ecology. Trends in Ecology & Evolution **26**:183-192

.

Bolnick, D. I., and M. Doebeli. 2003. Sexual dimorphism and adaptive speciation: two sides of

the same ecological coin. Evolution **57**:2433-2449.

Camus, M. F., K. Fowler, M. W. D. Piper, and M. Reuter. 2017 Sex and genotype effects on

nutrient-dependent fitness landscapes in Drosophila melanogaster. Proceedings of the

Royal Society of London Series B. **284**:20172237.

Caswell, H., and D. E. Weeks. 1986. Two-sex models: chaos, extinction, and other dynamic

consequences of sex. American Naturalist **128**:707-735.

Chesson, P. 2000. Mechanisms of Maintenance of Species Diversity Annual Review of Ecology

and Systematics **31**:343-366.

Cortez, M. H., and S. Patel. 2017. The effects of predator evolution and genetic variationon

predator-prey population-level dynamics. Bulletin of Mathematical Biology **79**:1510-

1538.

Darwin, C. 1871. The descent of man and selection in relation to sex. J. Murray, London.

De Lisle, S. P., and L. Rowe. 2015a. Ecological character displacement between the sexes. The

413   American Naturalist **186**:693-707.

414   De Lisle, S. P., and L. Rowe. 2015b. Independent evolution of the sexes promotes amphibian

415   diversification. Proceedings of the Royal Society B: Biological Sciences **282**:20142213.

416   Des Roches, S., D. M. Post, N. E. Turley, J. K. Bailey, A. P. Hendry, M. T. Kinnison, J. A.

417   Schweitzer, and E. P. Palkovacs. 2018. The ecological importance of intraspecific

418   variation. Nature Ecology and Evolution **2**:57-64.

419   Ellner, S. P., R. E. Snyder, and P. B. Adler. 2018. An expanded modern coexistence theory for

420   empirical applications. Ecology Letters **22**:3-18.

421   Fisher, R. 1930. The genetical theory of natural selection. Oxford University Press, Oxford, UK.

422   Fryxell, D. C., H. A. Arnett, T. M. Apgar, M. T. Kinnison, and E. P. Palkovacs. 2015. Sex ratio

423   variation shapes the ecological effects of a globally introduced freshwater fish

424   Proceedings of the Royal Society of London Series B. **282**:20151970.

425   Fryxell, D. C., D. Weiler, M. Kinnison, and E. P. Palkovacs. 2019 Eco-evolutionary dynamics of

426   sexual dimorphism Trends in Ecology and Evolution **in press**.

427   Garlapow, M. E., W. Huang, M. T. Yarboro, K. R. Peterson, and T. F. C. Mackay. 2015.

428   Quantitative genetics of food intake in Drosophila melanogaster. PloS OnE **10**:e0138129.

429   Giery, S. T., and C. G. Layman. 2019. Ecological consequences of sexually selected traits: An

430   eco-evolutionary perspective The Quarterly Review of Biology **94**:29-74.

431   Gomez-Llano, M. A., H. M. Bensch, and E. I. Svensson. 2018. Sexual conflict and ecology:

432   species composition and male density interact to reduce male mating harassment and

433   increase female survival Evolution **72**:906-915.

Holt, R. D. 1977. Predation, apparent competition, and the structure of prey communities.

Theoretical Population Biology **12**:197-229.

Holt, R. D., and M. B. Bonsall. 2017. Apparent Competition. Annual Review of Ecology and

Systematics **48**:447-471.

Jensen, K., C. McClure, N. K. Priest, and J. Hunt. 2015. Sex-specific effects of protein and

carbohyfrate intake on reproduction but not lifespan in Drosophila melanogaster. Aging

Cell **14**:605-615.

Kawatsu, K. 2018. Ecological effects of sex differ with trophic positions in a simple food web.

Ecology and Evolution **8**:1239-1246.

Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.

Evolution **34**:292-305.

Lindstöm, J., and H. Kokko. 1998. Sexual reproduction and population dynamics: the role of

polygony and demographic sex differences Proceedings of the Royal Society of London

Series B. **265**:483-488.

Maklakov, A., S. J. Simpson, F. Zajitchek, M. D. Hall, J. Dessmann, F. Clissold, D.

Raubenheimer, R. Bonduriansky, and R. C. Brooks. 2008. Sex-specific effects of nutrient

intake on reproduciton and lifespan Current Biology **18**:1062-1066.

Patel, S., and S. J. Schreiber. 2015. Evolutionarily driven shifts in communities with intraguild

predation. The American Naturalist **186**:E98-E110.

Pincheira-Donoso, D., T. Tregenza, R. K. Butlin, and D. J. Hodgson. 2018. Sexes and species as

rival units of niche saturation during community assembly. Global Ecology and

Biogeography **27**:593-603.

R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Rand, A. L. 1952. Secondary sexual characters and ecological competition. Fieldiana-Zoology

34:65-70.

Rankin, D. J., and H. Kokko. 2007. Do males matter? The role of males in population dynamics.

Oikos **116**:335-348.

Reddiex, A. J., T. P. Gosden, R. Bonduriansky, and S. F. Chenoweth. 2013. Sex-specific fitness

consequences of nutrient intake and the evolvability of diet preferences. The American

Naturalist **182**:91-102.

Schreiber, S. J., R. Burger, and D. I. Bolnick. 2011. The community effects of phenotypic and

genetic variation within a predator population Ecology **92**:1582-1593.

Selander, R. K. 1966. Sexual dimorphism and differential niche utilization in birds. The Condor

68:113-151.

Shärer, L., L. Rowe, and G. Arnqvist. 2012 Anisogamy, chance and the evolution of the sex

roles. Trends in Ecology & Evolution **27**:260-264.

Shine, R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the

evidence. Quarterly Review of Biology **64**:419-461.

Slatkin, M. 1984. Ecological causes of sexual dimorphism. Evolution **38**:622-630.

Soetgart, K., T. Petzoldt, and R. W. Setzer. 2010. Solving Differential Equations in R: Package

deSolve. Journal of Statisitcal Software **33**:1-25.

Stamps, J. A., J. B. Losos, and R. M. Andrews. 1997. A comparative study of population density

and sexual size dimorphism in lizards. American Naturalist **149**:64-90.

Start, D. 2018. Keystone individuals alter ecological and evolutionary consumer-resource

dynamics. The American Naturalist **191**:277-286.

Start, D., and S. P. De Lisle. 2018. Sexual dimorphism in a top predator drives aquatic prey

community assembly. Proceedings of the Royal Society of London Series B. **2018**:1717.

Start, D., and B. Gilbert. 2017. Predator personality structures prey communities and trophic

cascades. Ecology Letters **20**:366-374.

Svensson, E. I. 2019. Eco-evolutionary dynamics of sexual selection and sexual conflict

Functional Ecology **33**:60-72.

Temeles, E. J. 1985. Sexual size dimorphism of bird-eating hawks: the effect of prey

vulnerability. The American Naturalist **125**:485-499.

Temeles, E. J., I. L. Pan, J. L. Brennan, and J. N. Horwitt. 2000. Evidence for ecological

causation of sexual dimorphism in a hummingbird. Science **289**:441-443.

Trivers, R. L. 1972. Parental investment and sexual selection. Pages 136-179 *in* B. Campbell,

editor. Sexual selection and the descent of man. Aladine, Chicago.

Tuszynski, J. 2019. caTools: Tools: moving window statistics, GIF, Base64, ROC AUC, etc.. R

package version 1.17.1.2. https://CRAN.R-project.org/package=caTools.

Zajitschek, F., and T. Connallon. 2017. Partitioning of resources: the evolutionary genetics of

sexual conlflict over resource acquisition and allocation. Journal of Evolutionary Biology

30:826-838.

Appendix

Here we relax two assumptions of equations 1 and 2 to explore how various

aspects of consumer mating system affect our conclusions. The birth function in

equation 2 implies monogamous mating, where each male and female has a single

mating partner. We can relax this assumption by including a parameter h, termed

'harem size' by demographers (Caswell and Weeks 1986, Lindstöm and Kokko

1998), that reflects departures from monogamy:

$$B(M,F,R_1,R_2) = 2k\frac{MI_M * FI_F}{MI_M + (FI_F) * h^{-1}} \tag{5}$$

In this modified birth function, a value of h = 1 reflects monogamy, while values of

h < 1 reflect polyandry and values h >1 reflect polygyny. Whenever h deviates

from unity, birth rates, and thus consumer population growth rates, are maximized

under a sex ratio skewed towards the sex that is monopolized and limits birth rates

more (Caswell and Weeks 1986). Thus, in our model, departures from monogamy

have the effect of reducing abundance of the resource favored by the sex that is

monopolized (Figure A1), because this resource places a greater limit on consumer

growth rates. None the less, qualitative conclusions on the effects of consumer

sexual dimorphism remain unchanged in the face of departures from monogamy.

In many taxa, sexual conflict over optimal mating rate can create scenarios

where selection can favor traits in one sex (typically males) that reduce absolute

fitness in the other (Arnqvist and Rowe 2005). These sexual conflicts can result in

fitness costs borne by females, that are often elevated as competition for mating

opportunities increase under sex ratio skew. Equation 1 assumes that equal

demographic effects of sex-ratio skew across the sexes, an assumption we can

relax by incorporating a term (simplified from a similar term in Rankin and Kokko

2007) into equation 1c:

$$\frac{dF}{dt} = \frac{1}{2} B(M, F, R_1, R_2) F - d_F F - c(\frac{M}{F}) \qquad (6)$$

that penalizes female survival under male-biased sex ratios. In this term, the

scaling constant c reflects the strength of sexual conflict. Incorporating this sexual

conflict term has similar effects to that of sex-biased demography, decreasing the

abundance of resource 1 (favored by males) relative to resource 2 (favored by

females) (Figure A2, A3); this is unsurprising as the addition of this term generated

female biased mortality in the consumer. Thus, variation in consumer mating

system has predictable consequences for the relative abundances of each resource.

Figure A1. Effects of altering harem size (polyandry vs. polygyny) for resource abundance

under apparent competition. Values on z axis are numerical solutions for resource 1 density

with resource 1 growth rate fixed at 1 (top panels) or resource 2 density with resource 2 growth

rate fixed at 1 (bottom panels). Three different departures from monogamy are illustrated: weak

polyandry (left column), and weak and strong polygyny. Increasing sexual dimorphism

increases the difference in growth rates under which resources can coexist. a_r was fixed at 0; all

other parameter values where as described in the Methods.

Figure A2. Effects of sexual antagonism under monogamy and polygyny for resource 1

abundance. Values on z axis are numerical solutions for resource 1 density with resource 1

growth rate fixed at 1. Two levels of sexual conflict are illustrated in the left and right column.

Under monogamy (top panels) introduction of sexual conflict has little effect, although the effect

of polygyny (bottom panels) on resource 1 abundance is amplified with the addition of sexual

conflict. a_r was fixed at 0; all other parameter values where as described in the Methods.

Figure A3. Effects of sexual antagonism under monogamy and polygyny for resource 2

abundance. Values on z axis are numerical solutions for resource 2 density with resource 2

growth rate fixed at 1. Two levels of sexual conflict are illustrated in the left and right column.

Sexual conflict has the effect on increasing resource 2 abundance (contrast with Figure A1) as

consumer female survival is reduced. a_r was fixed at 0; all other parameter values where as

described in the Methods.

Figure A4. Effects of differential growth rates and consumer sexual dimorphism on

consumer abundance. Values on z axis are numerical solutions to equation 1 for female

density, with resource 1 growth rate fixed at 1. Consumer density is reduced with increasing

sexual dimorphism (reduced niche overlap) under both apparent (A) and interspecific (B)

competition between the resources. However, consumer populations are maintained except when

interspecific competition between resources is equal to intraspecific competition (C). In this case,

sexually-dimorphic consumers cannot exist when resource growth rates differ; competitive

exclusion of the inferior resource leads to extinction of both the resource and the consumer.

Note differences in z axis scale.