Relationship Between Motor Capacity of the Contralesional and Ipsilesional Hand Depends on the Side of Stroke

Rini Varghese, MS[1]

Carolee J. Winstein, PhD[1,2]

1. Division of Biokinesiology and Physical Therapy, University of Southern California

2. Department of Neurology, Keck School of Medicine, University of Southern California

Corresponding Author:

Rini Varghese

1540 E Alcazar St, CHP 155

Los Angeles, CA 90089

rinivarg@pt.usc.edu

(323) 442-1196

Abstract

There is considerable evidence that after a stroke, ipsilesional deficits increase as contralesional impairment increases. However, it is unclear whether this relationship differs based on the side of stroke. Here, we tested the hypothesis that the ipsilesional hand motor capacity co-varies with contralesional hand impairment only in individuals with left hemisphere damage. Forty-two pre-morbidly right-handed chronic stroke survivors (left hemisphere damage, LHD = 21) with mild-to-moderate paresis (Upper Extremity Fugl-Meyer-UEFM range: 19-59) performed distal items of the Wolf Motor Function Test (dWMFT). We used univariate co-efficient of determination (R^2) and multiple linear regression to assess the relationship between motor capacity of ipsilesional and contralesional hands. Contralesional UEFM, and dWMFT were found to be significant predictors of ipsilesional hand motor capacity ($p < 0.0001$ for both). Importantly, the relationship between contralesional and ipsilesional hands was significantly modified by side of stroke (model adjusted $R^2 = 0.26$ and 0.42, respectively, $p < 0.01$). For individuals with LHD, contralesional impairment explained 42% and contralesional hand motor capacity explained 65% of the variance in ipsilesional hand motor capacity. However, this relationship was not statistically significant for individuals with right hemisphere damage (RHD, unadjusted $R^2 < 1\%$ for UEFM and 9% for dWMFT, $p > 0.05$). In chronic stroke survivors with mild-to-moderate impairment, our findings demonstrate that the relationship between contralesional and ipsilesional motor deficits depends on the side of stroke. Specifically, deficits co-vary between the limbs of stroke survivors with left hemisphere damage but not right hemisphere damage.

Keywords

Ipsilesional deficits, stroke, hemispheric differences

Introduction

It is now well known that unilateral stroke not only results in contralesional arm deficits, but also significant albeit more subtle motor deficits in the ipsilesional limb compared to age-matched non-disabled adults.[1-4] Previous work that examined the relationship between motor capability or capacity of the contralesional and ipsilesional hands are in agreement that the presence of motor deficits in the ipsilesional arm and hand are related to the severity of motor deficits in the contralesional upper limb, especially in the chronic phase after stroke.[5-9] For example, Boyd and colleagues demonstrated that the ability to learn an implicit motor task with the ipsilesional hand was inversely correlated with the degree of motor impairment in the contralesional upper extremity.[5] Rinne and colleagues[6] used a force-tracking task and showed that grip strength and tracking accuracy co-varied between the ipsilesional and contralesional hands.[6] Similarly, two recent studies reported that deficits in the ipsilesional hand were most impaired for a group classified with relatively more "severe" contralesional motor impairments, defined as an Upper Extremity Fugl-Meyer score (UEFM) of less than 28,[7] or moderate impairment, 32 to 57.[10]

In addition to the studies aimed at characterizing the relationship between contralesional and ipsilesional hands, there is mounting evidence that the unilateral motor deficits observed for contralesional and ipsilesional are hemisphere-specific and thus depend on side of stroke lesion.[8,11-16] For example, using clinical motor assessments of grip strength and hand dexterity, Harris and Eng[13] showed that in chronic stroke survivors who are pre-morbidly right-hand dominant, contralesional motor impairments were less severe in individuals who suffered damage in the dominant (i.e. left) hemisphere (LHD) compared to those who suffered damage in the non-dominant (right) hemisphere (RHD).[13,16] In contrast, considering ipsilesional motor deficits, the evidence is mixed concerning hemisphere-specific effects. For instance, some

studies reported that individuals with LHD exhibited more severe ipsilesional arm and hand deficits compared to those with RHD [4,16-18] while others have reported no difference in ipsilesional capacity between LHD and RHD.[2] In acute stroke survivors, Kust et al demonstrated that deficits in grip force of the ipsilesional hand were significantly associated with clinical measures of function of the contralesional hand *only* in LHD.[14] Contrary to this, de Paiva Silva et al found that the ipsilesional hand was significantly slower and less smooth in individuals with RHD who exhibited moderate-to-severe motor impairments (UEFM < 34) in the contralesional upper extremity compared to controls, LHD, and those with mild motor impairment.[8]

Taken together, there is converging evidence regarding the relationship between motor deficits of the contralesional and ipsilesional upper extremity, such that ipsilesional deficits are worse when contralesional impairment is greater (Figure 1A); however, it is uncertain whether this relationship between the two limbs depends on which hemisphere is damaged. In particular, motor deficits of the two limbs are most prominent for tasks that require dexterous motor control (e. e., grip force, tapping, tracking). For predominantly right-handed cohorts (as is the case in most studies), contralesional deficits appear to be more severe in those with RHD, in whom the contralesional limb is non-dominant; whereas ipsilesional deficits were more severe in those with LHD. An exception to this observation for those with RHD seems to be in the case when contralesional impairment is most severe (i.e., UEFM < 34).[78] Thus, one might predict that as contralesional impairment worsens, individuals with LHD would have proportionally worse ipsilesional deficits, but individuals with RHD (especially if say UEFM > 34) would not; see (Figure 1 B & C) for two alternative hypotheses. This prediction arising from an interaction between severity of contralesional deficits and the hemisphere affected by the stroke has not before been explicitly tested.

One reason that this prediction remains untested might be methodological in that in at least three of the studies discussed earlier, participants were categorically classified based on the degree of contralesional motor impairment (e.g., mild, moderate, severe).[7,8,10] While grouping in this manner may be somewhat useful for stratification and randomization purposes with large samples, categorization (or worse, dichotomization) of a continuous variable around arbitrarily set cut-off points presents several concerns. Of concern is a loss of measurement resolution, an assumption of discontinuity in the underlying construct (in this case motor impairment), unequal subgroup sizes (or biased sampling), and large unexplained residuals in regression models, to name a few.[19-21] Overall, if the objective is to understand the nature and extent of critical response-predictor relationships, then a categorical approach is particularly problematic.

Thus, the primary objective is to determine if the severity of deficits in the ipsilesional hand varies directly with that of the contralesional hand (using a continuous measure). Further, and more importantly, we seek to determine if this relationship differs based on the side of stroke lesion (i.e., an interaction effect). We predict that motor capacity of the ipsilesional hand will vary directly with the severity of the contralesional motor impairment and dexterous motor capacity, only in individuals with LHD, but not in individuals with RHD; see (Figure 1).

Figure 1. Hypothesized effects represented in schematic figure. A. The null hypothesis, wherein the relationship between contralesional (CL) impairment and ipsilesional (IL) motor capacity is not modified by the side of stroke lesion. B. Alternative hypothesis 1, wherein ipsilesional deficits are related to contralesional impairment but only in LHD (blue) and not in RHD (red). C. Alternate hypothesis 2, wherein ipsilesional deficits are related to contralesional impairment but only in LHD and in RHD with severe impairment (represented in the shaded dark-grey area).

Methods

Participants

Forty-two chronic stroke survivors (n = 21 left-hemisphere damage, LHD) provided informed consent to participate and in accordance with the 1964 Declaration of Helsinki and the guidelines of the Institutional Review Board for the Health Science Campus of the University of Southern California. These participants were enrolled as part of a larger phase-IIb clinical trial (Dose Optimization for Stroke Evaluation, ClinicalTrials.gov ID: NCT01749358).

Participants were included if they were: 1) between 21-75 years of age, 2) pre-morbidly right-handed, 3) ≥ 150 days post stroke (chronic phase), 4) had mild to moderate residual motor impairment (Upper Extremity Fugl-Meyer, UEFM score ≥ 19) with mostly resolved upper extremity paresis. Participants were excluded if they had: 1) severe sensory disturbances (no

response to light touch or complete loss of proprioception as indicated by the UEFM), 2) current major depressive disorder (score > 3 on PHQ2, depression screening survey) 3) a history of recent surgeries, significant orthopedic injuries, or pain affecting the upper extremity that would restrict shoulder and elbow movement, 4) severe cognitive deficits such as aphasia, apraxia or neglect that would preclude participants from comprehending test instructions or questionnaires.

Outcome Measures

Motor Component of the Upper Extremity Fugl-Meyer (UEFM)

The UEFM [22] is an assessment of motor impairment of the contralesional arm and hand after stroke and includes tests of strength and independent joint control. Item-wise scoring of the UEFM ranges from 0 (unable to perform) to 2 (able to perform completely) while total score ranges from 0 to 66, with a higher score indicating lesser impairment.

Wolf Motor Function Test (WMFT)

The WMFT is designed to assess upper extremity motor capacity through timed functional task performance (e.g., lifting a can, pencil, or paper clip). Originally designed for patients with moderate to severe upper extremity motor deficits, the test was later modified by Morris, Crago and Taub to accommodate individuals with mild motor impairments [23]. In a series of 15 tasks, the test administrator asks the participant to perform frontal or midsagittal plane motions with the shoulder and elbow, and dexterous tasks with the hand. Item-wise scoring entails a continuous capacity time-score. Traditionally, the test has been used to assess motor capacity of both the contralesional (CL) as well as the ipsilesional (IL) arm and hand, with the latter used as a reference for comparison within an individual.

We used the WMFT time-score to assess upper extremity motor capacity through timed functional task performance. A principle component analysis of WMFT scores revealed two clusters: one consisting of the proximal (#1-8, except 6, i.e., lifting weight to box) and other the distal (#9-17, except 14, i.e., grip strength),[24] with the latter serving as the primary measure of hand motor capacity. The distal battery (dWMFT) consists of the following 8 tasks: lift can, lift pencil, lift paper clip, stack checkers, flip cards, turn a key in a lock, fold towel, and lift basket. Hand motor capacity was assessed in both limbs.

Statistical Analyses

To test the hypothesis that the inter-limb relationship of motor capacity is modified by the side of stroke lesion, we used unadjusted coefficient of determination (R^2) with Thorndike type 2 correction for range restriction and multiple linear regression of the following form:

$$\textbf{Model 1}: y = \beta_0 + \beta_1(UEFM) + \beta_2(Side\ of\ Lesion) + \boldsymbol{\beta_3}(\textbf{CL\ UEFM} *$$
$$\boldsymbol{Side\ of\ Stroke}) + \epsilon$$

$$\textbf{Model 2}: y = \beta_0 + \beta_1(UEFM) + \beta_2(Side\ of\ Lesion) + \boldsymbol{\beta_3}(\textbf{CL\ }\boldsymbol{dWMFT} *$$
$$\boldsymbol{Side\ of\ Stroke}) + \epsilon$$

In both models, y is the average time score on the distal WMFT (dWMFT) of the ipsilesional (IL) hand. Using this multiple model, our hypotheses were that $\beta_1 \neq 0$ and $\beta_3 \neq 0$ (see Figure 1).

All continuous variables were assessed for normality using Lilliefors test (modified Kolmogorov-Smirnov test). Distributions for chronicity and average time-score for the distal WMFT were negatively skewed and were therefore log-transformed. All continuous variables

were standardized to unit variance (z-scored) for linear regression analysis. Welch's t-tests were used to compare age, chronicity, and Upper Extremity Fugl-Meyer scores between LHD and RHD, whereas Chi-square test was used to compare the proportion of females and males between the two groups. Significance level (α) was set at $p = 0.05$.

In order to select the predictor variables that best explain the response, we used a backward selection approach, in which we began by adding all predictor variables in each of the two above models to explain the response variable y. This included our hypothesized predictors, CL UEFM (or CL dWMFT) as well as the side of stroke lesion (LHD or RHD), and, potential confounders (age, chronicity, and sex). In a combined full model, those predictors that met a liberal cut-off of $p = 0.2$ were preserved in the final reduced model. Based on this selection process, we found sex to be a significant confounder ($p = 0.08$) in Model 1, and therefore included it as a predictor in the reduced Model 1. For Model 2, none of the confounders met the cut-off p-value, except our hypothesized predictors. Additional information on model selection and model diagnostics is included as supplementary materials. Standard errors and 95% CI of the estimates of regression coefficients were confirmed by performing 1000 bootstrap replicates and are reported in the next section.

Results

Descriptive statistics for all participants are provided in Table 1. On average, the 42 stroke survivors had moderate arm impairment (UEFM = 41.6), were approximately 60 years of age, 5.75 years post-stroke, and were predominantly male (74%). There were no significant differences between LHD and RHD in the level of impairment, chronicity or the number of

males. Individuals with RHD were younger compared to LHD (median age difference 8.7 years) but not statistically different.

Table 1. Descriptive Statistics for the full sample (N = 42), and for the two groups of interest, left hemisphere damage, LHD (n = 21) and right hemisphere damage, RHD (n = 21).

Variable	Overall (N = 42) Mean * (± SD)	LHD (n = 21) Mean (± SD)	RHD (n = 21) Mean (± SD)
Sex (Male) *	31 (73.8)	15 (71.4)	16 (76.2)
Age (in years)	59.16 (12.3)	60.71 (10.8)	57.62 (13.8)
Chronicity (in months)	69.37 (36.9)	63.32 (24.4)	75.41 (46.1)
UEFM Motor (/66)	41.57 (10.5)	40.67 (12.6)	42.48 (8.2)

* Count (percent) for categorical variables

Model 1: Side of lesion modifies the relationship between CL UEFM and IL motor capacity

Contralesional UEFM explains 42% of the variance in ipsilesional hand motor capacity in LHD ($p < 0.001$), but less than 1% in RHD (corrected for range restriction with Thorndike case 2 correction) ($p > 0.05$).

After adjusting for main effects and significant confounders using multiple regression, the final reduced form of Model 1 was statistically different from a null model ($F = 4.71$, $p = 0.003$, adjusted $R^2 = 0.26$). Based on estimates from Model 1, CL impairment (UEFM) was significantly associated with IL hand motor capacity, i.e., dWMFT, ($\beta_1 = -0.69 \pm 0.20$, $p < 0.001$) (Figure 2A). There was no significant effect of the side of lesion ($\beta_2 = 0.12 \pm 0.26$, $p = 0.66$). However, as predicted, there was a significant interaction between the side of lesion and CL impairment ($\beta_3 = 0.64 \pm 0.31$, $p = 0.038$). Figure 2 B and C illustrates the interaction.

Figure 2. Scatterplots show the relationship between contralesional motor impairment (CL UEFM) and ipsilesional distal motor performance (IL dWMFT) for the full sample (A), LHD (B), and RHD (C). Solid lines represent the linear prediction and shaded areas represent the 95% confidence interval.

Model 2: Side of lesion modifies the relationship between CL dWMFT and IL motor capacity

Contralesional dWMFT explains 65% of the variance in ipsilesional hand motor capacity in LHD ($p < 0.001$), but only 3.6% in RHD (corrected for range restriction with Thorndike case 2 correction) ($p > 0.05$).

After adjusting for main effects and confounders using multiple regression, the final reduced form of Model 2 was statistically different from a null model ($F = 11.04, p < 0.001$, adjusted $R^2 = 0.42$). Based on estimates from Model 2, CL hand motor capacity (dWMFT) was significantly associated with IL hand motor capacity ($\beta_1 = 0.83 \pm 0.17, p < 0.001$) (Figure 3A). There was no significant effect of the side of lesion ($\beta_2 = 0.28 \pm 0.24, p = 0.25$). As predicted, there was an interaction between the side of lesion and CL hand motor capacity ($\beta_3 = -0.52 \pm 0.35, p = 0.05$), however, this effect only trended towards significance. Figure 3 B and C illustrates the interaction. Not surprisingly, when the non-significant main effect of the side of

lesion (β_2) was removed from the model, the interaction effect achieved significance (β_3 = -0.54 ± 0.30, p = 0.04).

Figure 3. Scatterplots show relationship between contralesional distal motor performance (CL dWMFT) and ipsilesional distal motor performance (IL dWMFT) for the full sample (A), LHD (B), and RHD (C). Solid lines represent the linear prediction and shaded areas represent the 95% confidence interval.

Discussion

For the first time, we explicitly tested the hypothesis that motor capacity of the ipsilesional hand is influenced by an interaction between the severity of contralesional deficits and the side of stroke lesion. We found that ipsilesional motor capacity co-varies with the degree of impairment of the contralesional extremity, but only in individuals with LHD. Hints of this interaction were implicit in a few previous studies;[2,3,15] however, categorical reporting of the upper extremity Fugl-Meyer (UEFM) masked this interesting effect. In the following sections, we provide an analysis of the interaction effect, explore the insights obtained from the type of task, and discuss the role of the left hemisphere in organizing bilateral motor outputs. Finally, we identify the limitations of this study and suggest future research questions.

Analysis of the interaction effect

By preserving the continuity of the UEFM and utilizing continuous standardized z-scores, our statistical approach allowed a direct comparison of our regression parameters (i.e., βs), thus reflecting effect sizes of each of the candidate predictors—contralesional impairment, side of lesion and their interaction. While contralesional impairment alone bore the largest effect on ipsilesional motor capacity (β_1), the second largest effect was through the interaction between contralesional UEFM and the side of lesion (β_3), and not the side of lesion alone (β_2). In fact, unlike previous findings, we did not observe a significant effect of the side of lesion on ipsilesional motor capacity [17,18], nor on contralesional UEFM or dWMFT.[13]

One reason for this might be that the effect of the side of lesion observed in those previous studies may have arisen from its interaction with contralesional impairment. However, because an interaction effect was not explicitly tested and because contralesional impairment was either collapsed across the groups [23,15] or catagorical[78], variance in the ipsilesional capacity may have been conflated with the effect of the side of lesion, or, remained unexplained, especially for UEFM scores that fell at the boundaries of the pre-defined categories. To illustrate this point further, let us revisit Figure 1B. Here, if a cut-off for the x value (say, UEFM) was to be set at any point to the left of where the two lines intersect, one might conclude that ipsilesional deficits are greater in LHD compared to RHD, whereas a cut-off to the right of the intersection might suggest that ipsilesional deficits are greater in RHD. Both observations would indicate an effect of the side of lesion, but not the change in the direction of this effect, evident as an interaction.

Naturally, this brings to light another important issue and that is the choice of cut-off scores for categorization, as this would be crucial in determining the direction of the effect of the

side of lesion. Previously, various cut-off scores for the UEFM have been used to define

impairment categories.[7,8,10] Of these, our data suggest that a UEFM score of 42, which occurs at

the intersection of the linear fits for LHD and RHD, would best reflect the change in the

direction of effect, i.e., the interaction between side of lesion and contralesional impairment on

ipsilesional capacity. Interestingly, *at* this score of 42, there would appear to be no differences in

motor capacity of either hand between LHD and RHD, which might explain why a number of

large clinical trials (e.g. EXCITE,[25] ICARE[26]), designed for moderately impaired stroke survivors

(mean UEFM scores in these studies were 42.5 and 41.6 respectively) may not have observed, on

average, any differences in motor capacity based on the side of stroke.

Insights from the type of task

A common link between our study and past reports is that ipsilesional deficits were found to be

most pronounced for distal (dexterous) tasks. These tasks—lift can, lift pencil, lift paper clip,

stack checkers, flip cards, turn a key in a lock, fold towel, and lift basket—nearly always

involved object manipulation and inherently require dextrous motor control of the hands.

Sunderland and colleagues[4] demonstrated that early on after a stroke, spatial accuracy in dexterity

tasks performed with the ipsilesional hand correlated with cognitive deficits, such as apraxia, in

individuals with LHD. While individuals included in this study did not exhibit severe apraxia

and were approximately 5 years post-stroke, it is possible that mild cognitive deficits, including

apraxia, may have impacted dexterous task performance in those with LHD, especially in the

more severe ranges of UEFM. Furthermore, we note that our evaluation of dexterous task

performance was through timed tests, and not quality of movement or accuracy. It has been

suggested that the left hemisphere plays an important role in regulating the timing and speed of

movements[15], and thus, injury to the left hemisphere, particularly to premotor and fronto-parietal networks (e.g. IPC)[27,28] may impair planning and sequencing required for smooth and rapid performance of dexterous motor tasks.

That deficits are apparent for distal rather than proximal motor performance is also relevant to our understanding of the putative neural substrates responsible for ipsilesional deficits. In this regard, candidate substrates include direct ipsilateral corticofugal pathways or indirect inhibitory circuits in the cortex, sub-cortex (via the corpus callosum and other commissures), or spinal cord (via spinal inter-neuronal circuits). As for the indirect cortical inhibitory circuits, the classic view contends that, based purely on somatotopy, the control of distal musculature (such as of the hands) is represented further away laterally on the motor cortex and lacks transcallosal fibers. Based on this view, control of the ipsilesional hand or the lack thereof might not be directly attributable to interhemispheric interactions, but rather intra-hemispheric pathways. This view has been challenged based on empirical neurophysiologic evidence in animals and humans,[29,30] however, the presence of ipsilateral activations in acallosal patients[31] provides a further reason to suspect that intra-hemispheric pathways, or perhaps inhibitory circuits in the lower centers of the neuraxis, are likely responsible for the control of distal musculature of the ipsilateral hand. For a comprehensive review of this topic, please see Carson (2005).[32] Nevertheless, whether this type of ipsilateral control is normal or maladaptive is a controversial topic[33-35] and warrants further study.

The role of the left hemisphere in the bilateral control of hands

Our main observation that deficits in ipsilesional hand motor capacity scale with contralesional impairment only in LHD is qualitatively similar to previous clinico-behavioral[e.g. 17,18,36,37] and

phenomenological evidence.[e.g. 38–40] These findings are consistent with a rather simplified

organizational model of the nervous system in which certain aspects of motor and/or cognitive

control are lateralized to the left (or dominant) hemisphere, such that damage to the left

hemisphere results in deficits in skilled motor actions of both upper extremities. For example,

using EMG recordings of homologous muscles in the arm, Cernacek (1961) demonstrated that

the frequency of motor irradiations, i.e., unintended motor output in the ipsilateral hand, were

significantly higher from the dominant to the non-dominant extremity.[38] Similarly, Wyke (1968)

reported that while individuals with left-sided cerebral lesions exhibited bilateral motor deficits

in speed and limb postural control, deficits in those with right cerebral lesions were restricted to

the contralateral limb.[18] Lastly, in one of the earliest experiments using functional MRI, Kim and

colleagues (1993) showed that the task-evoked activation of the left hemisphere was

substantially greater for ipsilateral movements compared to the right hemisphere.[41] In later years,

a number of neuroimaging [42,43] and neurophysiologic[44–46] studies have provided confirmatory

evidence for the role of the dominant hemisphere in organizing bilateral motor outputs. Our

results of co-varying deficits between the contralesional and ipsilesional hand in LHD provides

further empirical support for the role of the left hemisphere (in our pre-morbidly right-handed

group) in the control of both hands.

Limitations and future considerations

Some of the most significant methodological shortcomings of this study are that: First, the design

was observational, and the statistical analyses were inferential. A prospective study or

independent validation in a separate cohort would be ideal, if larger samples were available.

Second, UEFM scores for the RHD group were restricted towards the more severe range, with

the most severely impaired individual's score being 28. This restriction, however, was not

present in LHD (min. UEFM = 19). Although estimates from our univariate analyses were

adjusted for range restriction with Thorndike type 2 corrections and were still non-significant, we

are cautious in generalizing our observations regarding the interaction effect to more severe

ranges of motor impairment in RHD. Indeed, it is possible that for this severe range in RHD,

there exists a linear relationship between contralesional and ipsilesional motor deficits as

illustrated in Figure 1C. Thus, while we can, with some confidence, reject the null hypothesis

(Figure 1A), our data are insufficient to differentiate between the two alternate hypotheses, and

warrant a follow-up study.

It must be emphasized that the absence of a relationship with contralesional impairment

in RHD should not be taken to mean that ipsilesional deficits are absent in this group. In fact,

there is substantial evidence to the contrary. Comparison with an appropriate control group

would be necessary to demonstrate the presence of ipsilesional deficits in RHD and the

functional implications of these deficits. As alluded to earlier, measuring the speed of

performance, as in the case of timed functional tasks assessed in this study, does not provide any

insight into the perceptual errors, spatial accuracy or visuomotor deficits, which, based on

previous evidence,[47,48] might be a more important index of motor performance in RHD.

In summary, our results suggest that ipsilesional motor deficits co-vary with the degree of

impairment in LHD but are less pronounced in RHD. This observation further underscores the

extensive motor experiences of the pre-morbidly dominant ipsilesional limb and the importance

of the left hemisphere for the bilateral control of timed tasks. Lastly, we propose that the

hypothetical model of bilateral deficits in LHD is readily testable using a bimanual experimental

paradigm with sensitive kinematic measures and would offer important insights into the role and

organization of each hemisphere for the control of unilateral and bilateral movements.

Funding

This research study is supported by the NICHD National Center for Medical Rehabilitation

Research of the National Institutes of Health under award numbers: F31HD098796 to Rini

Varghese and R01HD065438 to Nicolas Schweighofer and Carolee J. Winstein.

REFERENCES

1.   Winstein CJ, Pohl PS. Effects of unilateral brain damage on the control of goal-directed hand movements. *Exp Brain Res*. 1995;105(1):163-174. doi:10.1007/BF00242191

2.   Wetter S, Poole JL, Haaland KY. Functional implications of ipsilesional motor deficits after unilateral stroke. *Arch Phys Med Rehabil*. 2005;86(4):776-781. doi:10.1016/J.APMR.2004.08.009

3.   Chestnut C, Haaland KY. Functional Significance of Ipsilesional Motor Deficits After Unilateral Stroke. *Arch Phys Med Rehabil*. 2008;89(1):62-68. doi:10.1016/J.APMR.2007.08.125

4.   Sunderland A, Bowers MP, Sluman SM, Wilcock DJ, Ardron ME. Impaired dexterity of the ipsilateral hand after stroke and the relationship to cognitive deficit. *Stroke*. 1999. doi:10.1161/01.STR.30.5.949

5.   Boyd LA, Quaney BM, Pohl PS, Winstein CJ. Learning implicitly: Effects of task and severity after stroke. *Neurorehabil Neural Repair*. 2007;21(5):444-454. doi:10.1177/1545968307300438

6.   Rinne P, Hassan M, Fernandes C, et al. Motor dexterity and strength depend upon integrity of the attention-control system. *Proc Natl Acad Sci*. 2017;115(3):E536-E545. doi:10.1073/pnas.1715617115

7.   Maenza C, Good DC, Winstein CJ, et al. *Ipsilesional Arm Motor Deficits in Stroke Vary with Hemisphere and Contralesional Arm Impairment*.; 2019. (in review)

8.   de Paiva Silva FP, Freitas SMSF, Banjai RM, Alouche SR. Ipsilesional Arm Aiming Movements After Stroke: Influence of the Degree of Contralesional Impairment. *J Mot Behav*. 2018. doi:10.1080/00222895.2017.1306479

9.   Noskin O, Krakauer JW, Lazar RM, et al. Ipsilateral motor dysfunction from unilateral stroke : implications for the functional neuroanatomy of hemiparesis. 2008:401-406. doi:10.1136/jnnp.2007.118463

10.   Bustrén EL, Sunnerhagen KS, Alt Murphy M. Movement Kinematics of the Ipsilesional Upper Extremity in Persons with Moderate or Mild Stroke. *Neurorehabil Neural Repair*. 2017;31(4):376-386. doi:10.1177/1545968316688798

11.   Mani S, Mutha PK, Przybyla A, Haaland KY, Good DC, Sainburg RL. Contralesional motor deficits after unilateral stroke reflect hemisphere-specific control mechanisms. *Brain*. 2013;136(4):1288-1303. doi:10.1093/brain/aws283

12.   Schaefer SY, Haaland KY, Sainburg RL. Ipsilesional motor deficits following stroke reflect hemispheric specializations for movement control. *Brain*. 2007;130(8):2146-2158. doi:10.1093/brain/awm145

13.   Harris JE, Eng JJ. Individuals with the dominant hand affected following stroke demonstrate less impairment than those with the nondominant hand affected. *Neurorehabil Neural Repair*. 2006;20(3):380-389. doi:10.1177/1545968305284528

14.   Küst J, Nowak DA, Karbe H, Dafotakis M, Fink GR, Grefkes C. Dexterity is impaired at both hands following unilateral subcortical middle cerebral artery stroke. *Eur J Neurosci*. 2007;25(10):3173-3184. doi:10.1111/j.1460-9568.2007.05551.x

15.   Silva FPDP, De Freitas SMSF, Silva PV, Banjai RM, Alouche SR. Ipsilesional arm motor sequence performance after right and left hemisphere damage. *J Mot Behav*. 2014;46(6):407-414. doi:10.1080/00222895.2014.924473

16.   Hanna-Pladdy B, Mendoza JE, Apostolos GT, Heilman KM. Lateralised motor control: Hemispheric damage and the loss of deftness. *J Neurol Neurosurg Psychiatry*.

2002;73(5):574-577. doi:10.1136/jnnp.73.5.574

17.   Heap M, Wyke M. Learning of a Unimanual Motor Skill by Patients with Brain Lesions: An Experimental Study. *Cortex*. 1972;8(1):1-18. doi:10.1016/S0010-9452(72)80023-6

18.   Wyke M. The effect of brain lesions in the performance of an arm-hand precision task. *Neuropsychologia*. 1968;6(2):125-134. doi:10.1016/0028-3932(68)90054-7

19.   Collins GS, Ogundimu EO, Cook JA, Manach Y Le, Altman DG. Quantifying the impact of different approaches for handling continuous predictors on the performance of a prognostic model. *Stat Med*. 2016. doi:10.1002/sim.6986

20.   Royston P, Altman DG, Sauerbrei W. Dichotomizing continuous predictors in multiple regression: A bad idea. *Stat Med*. 2006. doi:10.1002/sim.2331

21.   Naggara O, Raymond J, Guilbert F, Roy D, Weill A, Altman DG. Analysis by categorizing or dichotomizing continuous variables is inadvisable: An example from the natural history of unruptured aneurysms. *Am J Neuroradiol*. 2011. doi:10.3174/ajnr.A2425

22.   Fugl-Meyer AR, Jääskö L, Leyman I, Olsson S, Steglind S. The post-stroke hemiplegic patient. 1. a method for evaluation of physical performance. *Scand J Rehabil Med*. 1975. doi:10.1038/35081184

23.   Wolf SL, Catlin PA, Ellis M, Archer AL, Morgan B, Piacentino A. Assessing Wolf Motor Function Test as outcome measure for research in patients after stroke. *Stroke*. 2001;32(7):1635-1639. doi:10.1161/01.STR.32.7.1635

24.   Woodbury M, Nichols-Larsen DS, Wolf SL, et al. Measurement Structure of the Wolf Motor Function Test: Implications for Motor Control Theory. *Neurorehabil Neural Repair*. 2010;24(9):791-801. doi:10.1177/1545968310370749

25.   Wolf SL, Winstein CJ, Miller JP, et al. Effect of Constraint-Induced Movement Therapy on Upper Extremity Function 3 to 9 Months After Stroke. *Jama*. 2006;296(17):2095. doi:10.1001/jama.296.17.2095

26.   Winstein CJ, Wolf SL, Dromerick AW, et al. Effect of a task-oriented rehabilitation program on upper extremity recovery following motor stroke the ICARE randomized clinical trial. *JAMA - J Am Med Assoc*. 2016;315(6):571-581. doi:10.1001/jama.2016.0276

27.   Harrington DL, Haaland KY. Hemispheric specialization for motor sequencing: Abnormalities in levels of programming. *Neuropsychologia*. 1991. doi:10.1016/0028-3932(91)90017-3

28.   Harrington DL, Haaland KY, Knight RT. Cortical networks underlying mechanisms of time perception. *J Neurosci*. 1998.

29.   Aizawa H, Mushiake H, Inase M, Tanji J. An output zone of the monkey primary motor cortex specialized for bilateral hand movement. *Exp Brain Res*. 1990. doi:10.1007/BF00230856

30.   Donchin O, Cardoso de Oliveira S, Vaadia E. Who tells one hand what the other is doing: The neurophysiology of bimanual movements. *Neuron*. 1999;23(1):15-18. doi:10.1016/S0896-6273(00)80748-5

31.   Meyer BU, Röricht S, Von Einsiedel HG, Kruggel F, Weindl A. Inhibitory and excitatory interhemispheric transfers between motor cortical areas in normal humans and patients with abnormalities of the corpus callosum. *Brain*. 1995. doi:10.1093/brain/118.2.429

32.   Carson RG. Neural pathways mediating bilateral interactions between the upper limbs. *Brain Res Rev*. 2005;49(3):641-662. doi:10.1016/j.brainresrev.2005.03.005

33.   Johansen-Berg H, Rushworth MFS, Bogdanovic MD, Kischka U, Wimalaratna S,

Matthews PM. The role of ipsilateral premotor cortex in hand movement after stroke.

Proc Natl Acad Sci. 2002;99(22):14518-14523. doi:10.1073/pnas.222536799

34.   Jankowska E, Edgley SA. How Can Corticospinal Tract Neurons Contribute to Ipsilateral

Movements? A Question With Implications for Recovery of Motor Functions. *Neurosci*.

2006;12(1):67-79. doi:10.1177/1073858405283392

35.   Jones TA. Motor compensation and its effects on neural reorganization after stroke. *Nat*

Publ Gr. 2017:1-14. doi:10.1038/nrn.2017.26

36.   Wyke M. Postural arm drift associated with brain lesions in man. *Arch Neurol*.

1966;15(3):329-334. doi:10.1001/archneur.1966.00470150107016

37.   Wyke M. The Effects of Brain Lesions on the Learning Performance of a Bimanual Co-

Ordination Task. *Cortex*. 1971;7(1):59-72. doi:10.1016/S0010-9452(71)80022-9

38.   Cernacek J. Contralateral Motor Irradiation-Cerebral Dominance: Its Changes in

Hemiparesis. *Arch Neurol*. 1961. doi:10.1001/archneur.1961.00450080047005

39.   Giuliani CA, Purser JL, Light KE, Genova PA. Impairments in arm control in subjects

with left and right hemisphere stroke. *NeuroRehabilitation*. 1997;9(1):71-87.

doi:10.1016/S1053-8135(97)00015-2

40.   Freitas SMSF, Gera G, Scholz JP. Timing variability of reach trajectories in left versus

right hemisphere stroke. *Brain Res*. 2011;1419:19-33. doi:10.1016/j.brainres.2011.08.039

41.   Kim SG, Ashe J, Hendrich K, et al. Functional magnetic resonance imaging of motor

cortex: Hemispheric asymmetry and handedness. *Science (80-)*. 1993;261(July):615-616.

doi:10.1126/science.8342027

42.   Serrien DJ, Cassidy MJ, Brown P. The importance of the dominant hemisphere in the

organization of bimanual movements. *Hum Brain Mapp*. 2003;18(4):296-305. doi:10.1002/hbm.10086

43. Bundy DT, Szrama N, Pahwa M, Leuthardt EC. Unilateral, 3D Arm Movement Kinematics Are Encoded in Ipsilateral Human Cortex. *J Neurosci*. 2018. doi:10.1523/jneurosci.0015-18.2018

44. Hess CW, Mills KR, Murray NMF. Magnetic stimulation of the human brain: Facilitation of motor responses by voluntary contraction of ipsilateral and contralateral muscles with additional observations on an amputee. *Neurosci Lett*. 1986. doi:10.1016/0304-3940(86)90565-3

45. Sohn YH, Jung HY, Kaelin-Lang A, Hallett M. Excitability of the ipsilateral motor cortex during phasic voluntary hand movement. *Exp Brain Res*. 2003. doi:10.1007/s00221-002-1292-5

46. Ganguly K, Secundo L, Ranade G, et al. Cortical Representation of Ipsilateral Arm Movements in Monkey and Man. *J Neurosci*. 2009. doi:10.1523/JNEUROSCI.2471-09.2009

47. Boll TJ. Right and left cerebral hemisphere damage and tactile perception: Performance of the ipsilateral and contralateral sides of the body. *Neuropsychologia*. 1974;12(2):235-238. doi:10.1016/0028-3932(74)90008-6

48. Bracewell RM, Husain M, Stein JF. Specialization of the right hemisphere for visuomotor control. *Neuropsychologia*. 1990. doi:10.1016/0028-3932(90)90001-5

SUPPLEMENTARY MATERIAL

The complete data table and codebook for statistical analysis, including additional resources, is available with this manuscript as well as through the first author's OSF Repository:

https://osf.io/jkfe6