Geometric coupling of helicoidal ramps and curvature-inducing proteins in organelle membranes

Morgan Chabanon[1] & Padmini Rangamani[1*]

[1] Department of Mechanical and Aerospace Engineering, University of California San Diego, La Jolla, CA, USA.
* Corresponding author. E-mail: prangamani@eng.ucsd.edu

Abstract

Cellular membranes display an incredibly diverse range of shapes, both in the plasma membrane and at membrane bound organelles. These morphologies are intricately related to cellular functions, enabling and regulating fundamental membrane processes. However, the biophysical mechanisms at the origin of these complex geometries are not fully understood from the standpoint of membrane-protein coupling. In this work, we focused on a minimal model of helicoidal ramps representative of specialized endoplasmic reticulum compartments. Given a helicoidal membrane geometry, we asked what is the distribution of spontaneous curvature required to maintain this shape at mechanical equilibrium? Based on the Helfrich energy of elastic membranes with spontaneous curvature, we derived the shape equation for minimal surfaces, and applied it to helicoids. We showed the existence of switches in the sign of the spontaneous curvature associated with geometric variations of the membrane structures. Furthermore, for a prescribed gradient of spontaneous curvature along the exterior boundaries, we identified configurations of the helicoidal ramps that are confined between two infinitely large energy barriers. Overall our results suggest possible mechanisms for geometric control of helicoidal ramps in membrane organelles based on curvature-inducing proteins.

Keywords: endoplasmic reticulum, helicoid, spontaneous curvature, lipid bilayer, minimal surface, Gaussian curvature

Introduction

Recent advances in microscopy, segmentation, and reconstruction methods have enabled us to visualize the plasma membrane and intracellular organelle membranes with a previously unforeseen level of spatial resolution [1–4]. These observations demonstrate the stunning complexity of membrane architecture within cells. Remarkably, despite this diversity, some common geometrical features are found across organelles and functions [5]. For instance budding/necking processes occur in the plasma membrane through a variety of endocytic and exocytic mechanisms (e.g., clathrin mediated [6–8], ESCRT [9–11], antimicrobial peptides [12, 13]), but also in the endoplasmic reticulum (ER) and Golgi in anterograde and retrograde trafficking [14, 15]. Similarly membrane tubes are present in the ER membrane network [1, 16–18], mitochondria [19–21], and in the form of cellular nanotubes [22–24]. Common to these seemingly disparate membrane structures is the catenoid-like geometry appearing at the neck of buds and the base of tubes [25]. Another striking example is the recently identified helicoidal structure connecting membrane sheets in the peripheral ER of neural and secretory cells [1, 26] as well as in the spine apparatus present in dendritic spines [4] (see Fig. 1). The molecular and biophysical mechanisms by which these helicoidal membrane geometries are produced and maintained remain under active investigation. However, the conserved features of membrane shapes across different cellular locations and biological functions suggest the possible existence of common design principles relating lipid membrane geometry, composition, and function [5].

We previously proposed a minimal model of membrane-protein interactions in catenoid-like necks [25], where our main finding was that at least two distinct curvature-inducing mechanisms are required to constrain a catenoid membrane neck below a critical radius. This has broad implications in ESCRT-mediated budding [9–11], yeast

Figure 1: Helicoidal membrane structures in the endoplasmic reticulum (ER). (a) Electron microscopy cross section of the ER in saliva secretory cell, (adapted from [26]). (b) Three-dimensional reconstruction of the black square in (a) showing a helicoidal ramp structure. (c) Spine apparatus in dendritic spine (adapted from [4]). (d) Geometry of a helicoidal ramp surface.

mitochondrial fission by dynamin [27], as well as yeast budding by septin filaments [28]. The fact that helicoids and catenoids belong to the same family of minimal surfaces – mathematical objects that minimize their surface energy by having a zero mean curvature and a negative Gaussian curvature everywhere – suggests that some of the insights gained from catenoids can be extended to helicoids. Following the idea that simple geometric arguments can provide insight into fundamental mechanisms underlying cellular organization [29], we sought to understand the relationship between curvature-inducing proteins and the geometry of intracellular lipid structures, in particular on helicoids representative of ER ramps (Fig. 1).

In a recent study that revealed the architecture of ER ramps to have helicoidal shapes, Terasaki *et al.* [26] proposed a physical model of the helicoidal membrane structures. They considered the mid-plane surface between the two ER bilayers – assumed to be at a constant distance from each other – bordered by an edge line representing the curved axis of the hemicylinder joining the two bilayers. They minimized the bending energy and the effective line energy of such a system. For an edge line having a negative spontaneous curvature [26, 30], they found that the structures satisfying the minimal energy are hollow helicoids such as the one shown in Fig. 1d. Extending this model to account for a mixture of membrane-proteins inducing either constant or negative edge curvatures, Shemesh *et al.* [30] numerically recapitulated various membranes structures present in the ER. In particular, they found that helicoid geometries are favored when the total amount of proteins is low with predominantly negative curvature-inducing proteins at the edges. While these studies lay the foundation of ER structure mechanics, they have focused on the influence of curvature-inducing proteins at the edges only [31], neglecting any potential coupling between curvature-inducing proteins at the sheet surface and the helicoid geometry.

Here we sought to understand the coupling between the distribution of curvature-inducing proteins and the geometry of a helicoidal membrane. We used *spontaneous curvature* as the mathematical catch-all for the curvature induced in the membrane by various molecular mechanisms. Using helicoids as geometric inputs, we computed the distribution of spontaneous curvature that satisfies the mechanical equilibrium of membrane bending [32]. This allowed us to map a geometric state-space for helicoidal membranes in which the bending energy is constrained by protein-induced curvature distribution.

Figure 2: (a-c) Helicoids are continuous and isometric transformations of catenoids. (a) Schematic of a full helicoid and definition of its parameters. (b) Isometric transformation from a helicoid to a catenoid (see Eq. S13). Note that every surface along the transformation is also a minimal surface. (c) Schematic of a catenoid whose boundaries have been transformed from the helicoid shown in (a). The distribution of Gaussian curvature is controlled by the helicoid pitch or equivalently by catenoid neck, through the relation $P = 2\pi r_n$. (d-g) The spontaneous curvature to maintain a helicoid (or a catenoid) shows a switch in sign at a critical pitch (or neck radius). All results here are for boundary conditions set to $C_0 = C_1$. (d) The computed distribution of spontaneous curvature (normalized by C_0) on full helicoids of varying pitches P. (e) Distribution of normalized spontaneous curvature along the radii of the helicoids shown in (d). The spontaneous curvature at the center ($r = 0$) switches sign at a critical pitch between $P/(2\pi L) = 0.3$ and 0.4. (f) Normalized spontaneous curvature at the helicoid center ($r = 0$) as a function of helicoid pitch. Color symbols correspond to the configurations in (e). (g) Corresponding dimensionless bending energy. An energy barrier occurs at a critical helicoid pitch. All results presented for helicoids hold for catenoids providing that the boundary conditions follow the same transformation.

Model development

Model assumptions

The main assumptions underlying the model are:

a) The lipid membrane is treated as a continuous thin elastic film of negligible thickness [32]. This assumption relies on the length-scale separation between the membrane thickness ($\sim 10^{-9}$ m) and typical radius of curvature of interest ($\sim 10^{-8} - 10^{-6}$ m).

b) Based on the same length-scale separation, the net effect of compositional asymmetry across the lipid bilayer

is accounted for by a "preferred" membrane curvature called *spontaneous curvature* (C). This can be induced by several molecular mechanisms, including asymmetric protein insertion in the bilayer, protein scaffolding, clustering, steric pressure, or lipid composition asymmetry [5, 33]. The spontaneous curvature is defined as a penalty for the mean curvature in the Helfrich energy [32] (see Eq. 1).

c) The system is at mechanical equilibrium. Any membrane deformation is much slower than the rate at which the membrane energy is dissipated. Given the coupling between spontaneous curvature and membrane geometry, this also implies no dynamic rearrangement of the membrane composition.

d) The lipid membrane is incompressible and inextensible. This assumption is supported by the orders of magnitude difference between the bending modulus ($\simeq 10^{-20}$ N m) and the stretching modulus ($\simeq 0.2$ N/m) of lipid membranes [34], allowing us to neglect area variations. This constraint is implemented through a Lagrangian multiplier on the surface area that has the physical interpretation of membrane tension [35, 36].

e) The lipid membrane is inviscid. Because of the fluid properties of lipid bilayers, we neglect any shear forces in the membrane. This simplification is in accordance with assumption (c) where dissipative processes are assumed to equilibrate instantly. Models accounting for in-plane lipid flow and diffusion can be found for instance in [37–39].

f) The lipid membrane is treated as an elastic manifold whose energy functional is the sum of the Helfrich bending energy [32] and a protein entropic contribution [36, 39–41].

g) The bending and Gaussian moduli are assumed to be homogeneous along the membrane surface.

Elastic lipid membrane with heterogeneous distribution of spontaneous curvature

The most common model of lipid membranes is based on the Helfrich energy [32], which can be extended to account for the entropic contribution of membrane-bound proteins [40] to the area free-energy functional such as

$$W(\sigma, H, K) = A(\sigma) + k[H - C(\sigma)]^2 + k_G K. \tag{1}$$

Here H is the membrane mean curvature, K is the Gaussian curvature, $A(\sigma)$ is the contribution of the membrane-bound proteins to the free energy, and σ is the surface density of proteins. Furthermore, k and k_G are the bending and Gaussian moduli respectively. $C(\sigma)$ is the spontaneous (mean) curvature, which is determined by the local membrane composition, say the surface density of a curvature-inducing protein σ. While it is certainly possible to propose explicit functions for $A(\sigma)$ and $C(\sigma)$ of the protein density (see [36, 40] for discussion on $A(\sigma)$, and [39, 41] for specific examples) we chose to retain their general form to obtain maximal insight. In the Supplementary Material Section S1, we briefly summarize the derivation of the governing equations of an elastic membrane at mechanical equilibrium to obtain the governing equations (Eqs. S11 and S12 (see [35, 38, 42–45] for detailed derivations).

The shape equation for a lipid membrane with protein-inducing spontaneous curvature is obtained by introducing the free energy density 1 into the normal force balance equation for elastic membranes Eq. S11, resulting in the following partial differential equation

$$\Delta \left[k(H - C) \right] + 2k(H - C)(H^2 + C - K) = p + 2\lambda H. \tag{2}$$

Here p is the pressure differential across the membrane and λ is a Lagrange multiplier that enforces the inextensibility constraint of the membrane, which is commonly interpreted as the membrane tension [35, 36]. Additionally, $\Delta(\cdot)$ is the surface Laplacian. The incompressibility equation for lipid membranes is obtained by introducing Eq. 1 into the tangential force balance equation Eq. S12, resulting in

$$\nabla \lambda = -W_\sigma \nabla \sigma , \tag{3}$$

where $(\cdot)_\sigma = \partial(\cdot)/\partial\sigma$ is the partial derivative with respect to σ and $\nabla(\cdot)$ is the surface gradient. W_σ is the chemical potential of the membrane protein, and given Eq. 1, we have $W_\sigma = A_\sigma - 2k(H - C)C_\sigma$.

Eqs. 2 and 3 describe the equilibrium configuration of lipid membrane subject to heterogeneous spontaneous curvature induced by proteins. An additional constraint for imposing the area incompressibility of the membrane requires the lipid velocity field, given by ($\mathbf{u} = u^{\alpha}\mathbf{a}_{\alpha} + w\mathbf{n}$), to satisfy $u^{\alpha}_{;\alpha} = 2Hw$ [39]. Provided a lipid velocity field satisfying the above incompressibility condition and suitable boundary conditions exists, the system given by the coupled Eqs. 2 and 3 fully describes the equilibrium configuration of a lipid membrane subject to a static distribution of curvature-inducing proteins. Although models for lipid flow within biological membranes have been proposed [35, 36, 38, 46], such an investigation is out of the scope of this study.

Distribution of curvature-inducing proteins on minimal surfaces

The general shape and incompressibility conditions expressed by Eqs. 2 and 3 relate the membrane geometry to its local composition. In this section, we specialize the governing equations to minimal surfaces, a mathematical family of surfaces that include helicoids, catenoids, and triply periodic minimal surfaces. We later apply this model to helicoids as minimal models of ER ramps. Using the fact that minimal surfaces have vanishing mean curvature ($H = 0$ everywhere on the membrane), in the absence of transmembrane pressure the shape equation (Eq. 2) reduces to a variable-coefficient Helmhotz equation for the spontaneous curvature

$$\Delta C(\sigma) - 2KC(\sigma) = 0. \tag{4}$$

This equation summarizes a key relationship between $C(\sigma)$ and K in our model. The spontaneous curvature induced by proteins $C(\sigma)$ reads the Gaussian curvature of the surface K, subject to boundary conditions. Both the energetic contribution of the proteins $A(\sigma)$ and the local Lagrange multiplier λ are now decoupled from the shape equation for minimal surfaces (Eq. 4), enabling us to solve for those quantities after solving the shape equation. However, any solution of Eq. 4 is restricted to the condition that Eq. 3 is satisfied. For minimal surfaces, this latter equation reduces to

$$\nabla\lambda = -\left[A(\sigma)_{\sigma} + 2kC(\sigma)C(\sigma)_{\sigma}\right]\nabla\sigma. \tag{5}$$

Using the identities $\nabla A(\sigma) = A(\sigma)_{\sigma}\nabla\sigma$ and $\nabla\left[C(\sigma)^2\right] = 2C(\sigma)C(\sigma)_{\sigma}\nabla\sigma$ and integrating, we get λ as a function of $A(\sigma)$ and $C(\sigma)$ up to a constant λ_0, such that

$$\lambda = -\left[A(\sigma) + kC(\sigma)^2\right] + \lambda_0, \tag{6}$$

which is the admissibility condition for the Lagrange multiplier field λ.

Now that the shape equation is uncoupled from the incompressibility equation, we can solve Eq. 4 for the distribution of spontaneous curvature C for a given minimal surface with Gaussian curvature K. We complete the system by specifying Dirichlet boundary conditions on the spontaneous curvature $C = C_0$ at the helicoid boundary ramp as indicated in each case [25], while applying a no-flux Neumann boundary condition on other boundaries. The Dirichlet boundary conditions in spontaneous curvature can be understood as the presence of curvature-inducing proteins at the edges of ER ramps and cisternae [30, 47]. On a minimal surface, Eq. 6 not only implies that imposing a Dirichlet condition on the spontaneous curvature is equivalent to imposing a membrane tension at the boundary, but also provides a direct relationship between spontaneous curvature and membrane tension everywhere on the surface.

Helicoid parametrization

A helicoid surface of diameter L and pitch P such as the one shown in Fig. 2a can be parametrized as [48, 49]

$$\begin{cases} x = r\cos(\phi) \\ y = r\sin(\phi) \\ z = P/(2\pi)\phi \end{cases} \quad \text{with} \quad \begin{cases} \phi \in [-\pi; \pi[\\ r \in [-L/2; L/2] \end{cases}. \tag{7}$$

The Gaussian curvature along the radius of such a helicoid is

$$K = -\left(\frac{P/(2\pi)}{[P/(2\pi)]^2 + r^2}\right)^2. \tag{8}$$

It should be noted that helicoids and catenoids belong to the same associated family of surfaces [49]. So by definition, there exists an isometric transformation from one surface to the other, as illustrated in Fig. 2b (see Supplementary Material Section S2 for details). In particular, the pitch (P) of a helicoid is related to the neck radius (r_n) of its corresponding catenoid by $r_n = P/(2\pi)$. This highlights how, in terms of Gaussian curvature, varying the pitch of a helicoid is equivalent to varying the neck radius of a catenoid. Since the Gaussian curvature enters in the source term of the shape equation for minimal surfaces (Eq.4), the distribution of spontaneous curvature on a helicoid is intrinsically related to its pitch, as a catenoid's distribution of spontaneous curvature is related to its neck radius [25].

Dimensionless system

The system has two length scales, a natural length scale L, which describes the system size, and an induced length scale, C_0, which arises due to the protein-induced spontaneous curvature. We therefore scale the geometric variables as $\bar{\theta}^\alpha = \theta^\alpha/L$ and $\bar{K} = KL^2$, and the spontaneous curvature as $\bar{C} = C/C_0$. Accordingly, the system can be written in its dimensionless form as

$$\Delta\bar{C} - 2\bar{K}\bar{C} = 0. \tag{9}$$

The total membrane bending energy, defined as the integral of the contribution from the curvature to the energy density over the surface (Ω), is given by

$$W_B = \int_\Omega \left(kC^2 + k_G K\right) d\omega. \tag{10}$$

In dimensionless form, the total bending energy is

$$\frac{W_B}{kC_0^2 L^2} = \int_\Omega \left(\bar{C}^2 + \frac{k_G}{k}\frac{1}{C_0^2 L^2}\bar{K}\right) d\bar{\omega}. \tag{11}$$

Dividing by the dimensionless area of the surface $\bar{\Omega} = \Omega/L^2$, the dimensionless energy per area is $W_B/(kC_0^2\Omega)$. Based on Hu *et al.* [50], we choose the value of the ratio of Gaussian to bending moduli as $k_G/k = -0.9$ for the reminder of our analysis.

Numerical implementation

We solved Eq. 9 subject to either Dirichlet or Neumann boundary conditions using the finite-element solver software COMSOL Multiphysics® v5.3a (COMSOL AB, Stockholm, Sweden) with the Surface Reaction Module. Parametric studies and quantitative data analysis were conducted in MATLAB R2018b (MathWorks, Inc., Natick, MA, USA) and the COMSOL With MATLAB interfacing module.

Results

Distribution of spontaneous curvature on a full helicoid is controlled by the pitch

We first consider helicoids with constant boundary conditions C_0 on both sides of the ramp (Fig. 2a with $C(\pm L/2) = C_0$), for various pitches. This configuration is formally equivalent to solving for the spontaneous curvature on a catenoid for various neck radius as was done in [25], and therefore serves to validate our model (see text associated with Eq. 8 and Supplementary Material Section S2). The resulting distribution of spontaneous curvature is

Figure 3: Hollow helicoidal ramps show a switch in the sign of spontaneous curvature that corresponds to an energy barrier at the inner radius when subject to constant exterior spontaneous curvature. (a) Geometry of a hollow helicoid with constant spontaneous curvature C_0 at the outer boundary. (b) Distribution of spontaneous curvature on the surface of hollow helicoids with various inner radii show a sign switch of spontaneous curvature at the center between $r_0/L = 0.04$ and 0.06. (c) Distribution of spontaneous curvature along the helicoid radius for various inner ramp radii. (d) Spontaneous curvature at the inner boundary as a function of the inner ramp radius shows a discontinuity at $r_0/L \simeq 0.045$. (e) Normalized bending energy of the helicoid shows an energy barrier at the critical radius of the inner hole. Color symbols in (d) and (e) match plots in (c). All computations are for helicoids with a pitch of $P/(2\pi L) = 0.05$.

presented as color maps in Fig. 2d and along the helicoid arclength in Fig. 2e. We find that although the maximum intensity of spontaneous curvature is always at the center of the helicoid, its value depends on the pitch. Furthermore, the spontaneous curvature exhibits switches in sign at a critical pitch between $P/(2\pi L) = 0.3$ and 0.4. For $P/(2\pi L) < 0.3$, the normalized spontaneous curvature C/C_0 is positive on the outer edge of the helicoid and negative towards the center. For $P/(2\pi L) > 0.4$, C/C_0 at the center switches sign and becomes positive. This switch is better seen in Fig. 2f, where the normalized spontaneous curvature at the center of the helicoid $C(r = 0)/C_0$ diverges at a critical pitch of about $P/(2\pi L) = 0.32$. The switch in sign of spontaneous curvature corresponds to an energy barrier in bending energy of the membrane (Fig. 2g).

These results, consistent with our previous results obtained for catenoids of varying neck radius [25], suggest that at least two distinct curvature-inducing mechanisms are required to admit helicoids below the critical pitch as energy minimizers and solutions to Eq. 9. Additionally, the existence of an energy barrier at a specific pitch can be interpreted as a possible biophysical mechanism preventing the collapse of a helicoidal membrane below a critical pitch, and therefore providing a geometrical control of the membrane structure through the Gaussian curvature.

Ratio between inner and outer ramp radius constrains the distribution of spontaneous curvature in hollow helicoids

Helicoidal ramps observed in organelles are often hollow with a single arm [1, 4, 26] (see Fig. 1a-c). Such a geometry can be constructed by varying $r \in [r_0, L/2]$ in Eq. 7, resulting in a hollow helicoid as depicted in Fig. 3a. We next investigated the relationship between the distribution of spontaneous curvature and the geometry of a hollow helicoid. The presence of the inner ramp radius at $r = r_0$ provides a new geometrical parameter relevant

to ER ramp shapes. A natural question that therefore arises is, how does the distance between the inner and outer ramp influences spontaneous curvature in hollow helicoidal membranes? To answer this question, we first impose uniform boundary conditions in spontaneous curvature ($C = C_0$) at the outer edge of the helicoid, while imposing no-flux conditions at the other boundaries. We solve Eq. 9 for the spontaneous curvature on the helicoid surface for fixed pitch ($P/2\pi L = 0.05$), and varying ratio between inner ramp radius and helicoid diameter (r_0/L).

The resulting distribution of spontaneous curvature is shown as a color map on the helicoid surfaces in Fig. 3b, and quantitatively along the helicoid radius in Fig. 3c. We find that for holes with large radii, the spontaneous curvature required to maintain the hollow helicoid is higher at the center, and remains of the same sign as C_0. However for small hole radii, C/C_0 is negative at the center, indicating the presence of a switch in sign at a critical inner radius. This switch, is better visualized when plotting C/C_0 at the inner boundary as a function of the inner radius (Fig. 3d), corresponds to a barrier in bending energy in the membrane (Fig. 3e). A local minimum in bending energy is obtained for values of the ratio between the inner and outer helicoid ramps smaller than for the barrier. Note that, although qualitatively similar to the results obtained for the full helicoid (Fig. 2), the position of the switch and the energy profile have different values.

Next, we explored how the pitch of the helicoid modulates the switch found in hollow helicoids and the associated energy barrier (see Figs. S1 and 4 respectively). As seen from Fig. 4, decreasing the helicoid pitch lowers the value of the radius ratio at which the energy barrier and local minimum occur. Additionally, the local minimum is deeper and narrower. For completeness, we investigate how imposing a constant spontaneous curvature at the inner boundary – instead of the outer – influences our findings. The resulting radial distribution and total bending energy are shown in Fig. S2 and S3 respectively for various hole radii and pitches. No switch-like behavior is observed for spontaneous curvature imposed at the inner boundary. We find that the intensity of spontaneous curvature and bending energy increases with decreasing hole radii and increasing pitch. It should be noted that, by correspondence with catenoids, for any configuration where $r_0 > 0$, no switch in the sign of the spontaneous curvature occurs if spontaneous curvature is imposed at both inner and outer boundary [25].

Figure 4: The pitch of a hollow helicoid controls the value of the critical inner radius and the width of the corresponding energy barrier. Helicoids with small pitch exhibit a deeper local energy minimum. (a) Helicoids with three different pitches. (b) Normalized bending energy of helicoids with pitches corresponding to (a), as a function of the inner radius.

Figure 5: Helicoid with a linear gradient of spontaneous curvature along the external boundary show two distinct sign switches in C at the center. (a) Hollow helicoid with an imposed spatial gradient of spontaneous curvature from C_0 to C_1 at the outer boundary. (b) Computed surface distribution of spontaneous curvature on hollow helicoids with $C_1 = 5C_0$ at various inner radii. Two distinct switches in spontaneous curvature are found at two critical inner ramp radius. All computations are made for helicoids of pitch $P/(2\pi L) = 0.05$. (c) Computed spontaneous curvature at the inner top (upward blue triangles) and inner bottom (downward red triangles) boundary for $C_1 = 5C_0$. Top and bottom points exhibit two switches in sign of C with varying inner radius, however their sign is opposite at the first switch (anti-aligned switch), but equal at the second switch (aligned switch). (d) Each switch corresponds to an energy barrier. The local energy minimum between the two energy barrier represents a possible mechanism for regulating the inner radius of an ER helicoidal ramp through spontaneous curvature.

Axial gradient of spontaneous curvature at the boundary induces two switches resulting in a locally favorable helicoidal geometry

Next, we investigate how a differential in curvature-inducing proteins along the helicoid axis affects the system. Such configuration is representative of possible inhomogeneous regions of protein source or consumption within the intracellular volume. We thus impose a linear gradient of spontaneous curvature ranging from C_0 to C_1 at the exterior boundary (Fig. 5a). Surprisingly, we find that two distinct switches in spontaneous curvature sign exist at different critical inner radii. These switches correspond to two different limits of C at the top and bottom points of the inner ramp and they behave differently for the two critical radii (Fig. 5b,c). At the largest critical radius, the spontaneous curvatures at the top and bottom of the inner ramp both tend to positive infinity for values of the inner ramp radius approaching the critical value from above, and tend to negative infinity when approaching from below. For simplicity, we call this type of switch in spontaneous curvature sign an "aligned switch". At the smallest critical radius, however, the spontaneous curvatures at the top and bottom ramps tend to opposite limits when approaching the critical value. We therefore call this behavior an "anti-aligned switch". Both switches correspond

Figure 6: The two switches induced by a gradient in spontaneous curvature reverse their sign at distinct values of the gradient. (a-g) Spontaneous curvature at the upper and lower inner boundary as a function of the inner radius for various gradients at the exterior boundary. (f) For $C_0 = C_1$, the distribution of C must be independent of z, requiring the suppression of the anti-aligned switch. (b) For $C_1 = C_0$, symmetry requirements induce the aligned switch to be suppressed.

to an energy barrier, therefore creating a new local minimum bound by virtually infinite energy barriers (Fig. 5d). As a consequence, if the exterior boundaries are subject to a gradient of spontaneous curvature such as one studied here, then a helicoidal membrane whose inner radius is constrained in-between the two energy barriers will tend to be energetically stable with respect to variations in the inner ramp radius.

The existence of two energy barriers corresponding to different switch mechanisms in spontaneous curvature leads to new questions: How is the two-switch behavior controlled by the boundary gradient? How can a membrane structure access the state of local energy minimum between two energy barriers? How does the system transition from a single energy barrier for homogeneous boundaries (Figs. 3 and 4) to two barriers when a gradient in spontaneous curvature at the boundaries is imposed? To answer these questions, we next investigated how the aligned and anti-aligned switches are controlled by the spontaneous curvature gradient at the exterior boundary. To do so, we compute the spontaneous curvature at the top and bottom of the inner ramp for various C_1/C_0 ratios and fixed helicoid pitch ($P/(2\pi L) = 0.05$). As shown in Fig. 6, the aligned and anti-aligned switches vanish at a distinct value of the boundary gradient in spontaneous curvature. This mechanism can be understood using simple symmetry arguments: for $C_1 = C_0$ no variation in spontaneous curvature in the z-direction can exist, implying that the anti-aligned switch must vanish (Fig. 5 and Fig. 6f). On the contrary, for $C_1 = -C_0$, the boundary condition imposes the spontaneous curvature along the z-axis to be symmetric with respect to the mid-plane of the helicoid. This therefore requires the aligned switch to vanish (Fig. 6b). Every time the inner ramp radius passes a critical value, the limits in spontaneous curvature associated with the vanishing switch take opposite signs.

The bending energy for each of the configurations examined in Fig. 6 are shown in Fig. S4, where it can be verified that the energy barrier associated with the switches indeed vanish at the corresponding spontaneous curvature gradients. Finally, we examine the effect of the helicoid pitch on the system. We find that, similarly to the case

where the boundary condition is homogeneous (Fig. 4), larger helicoid pitches lead to larger critical values in inner ramp radius, and higher energy values (Fig. S5).

Discussion

In this study, we proposed a minimal model for computing the spatial distribution of spontaneous curvature on helicoidal structures representative of ER ramps [1, 4, 26]. We identified conditions where continuous variations of the helicoid pitch and inner ramp radius are associated with discontinuous switches in the sign of spontaneous curvature, demonstrating a unique coupling between Gaussian curvature and protein-induced curvature. Membrane shapes associated with these switches correspond to energetically unfavorable configurations, resulting in energy barriers in the corresponding geometric parameter space. Surprisingly, we find that when a gradient of spontaneous curvature is imposed on the external ramp of hollow helicoids, the variation of the inner ramp radius is associated with two distinct switches in curvature sign and energy barriers. Our results suggest a geometric state-space for helicoidal membranes in which a well-constrained energetically favorable state can be accessed by turning-off one or the other energy barrier independently through curvature distribution.

In the context of biological membrane structures, the presence of a switch in sign of spontaneous curvature suggests the requirement for at least two distinct curvature-generating mechanisms to modulate the geometry. These can be thought of as either two different kind of membrane-bound proteins inducing spontaneous curvature of opposite sign, or as the same curvature-inducing protein but inserted on the opposite sides of the surface [33, 40]. Contrary to catenoid-shaped necks where such two-step molecular mechanisms have been identified in mammalian cells (e.g. ESCRT mediated budding [10, 51]) and yeast (e.g. mitochondrial fusion by synergetic action of Dnm1 and Fis1 [27]), the particular mechanisms that relate quantitative structural analysis to protein distribution and activity are not yet available for helicoidal ER geometries. However, a number of curvature-inducing proteins have been suggested to play a role in certain specific ER ramps, including reticulons and DP1/REEP5/Yop1 family proteins in highly curved membrane regions such as tube and cisternae edges [47, 52, 53].

The exquisite coupling between membrane geometry and distribution of curvature-inducing proteins in helicoids has several potential implications in the regulation of ER ramp shapes. The energy barrier associated with pitch and inner ramp radius might help prevent the collapse of ramps below critical geometrical parameter values. Additionally, in the case where a gradient of spontaneous curvature is imposed at the boundaries (Figs. 5c and S4), the helicoid geometry bound by the two energy barriers is energetically stable with respect to variations in the inner ramp radius. Therefore, inducing a gradient of curvature-inducing proteins along the vertical axis of a helicoidal ramp might serve as a possible molecular mechanism to regulate its geometry. Finally, from the reverse perspective, geometric control of the helicoid could present a mechanism to regulate the distribution of proteins on the ER surface. For instance, several proteins are found on the surface of ER cisternae, including Climp-63, kinectin, and p180 [47]. These proteins have transmembrane domains linked to coiled-coil structures on either side of the membrane, and interact with microtubules, ribosomes, or kinesins, and may induce curvature [47, 53]. The relationship between ER shape and distribution of spontaneous curvature identified in this study might underly spatial regulation mechanisms of surface proteins on helicoidal sheets.

Our ongoing work on spontaneous curvature distribution in minimal surfaces has implications for other membrane surfaces as well. In particular, triply periodic minimal surfaces (TPMS) have been observed in various membrane-bound organelles in stressed, diseased, and viral infected cells [54–58], as well as in reconstituted lipid membrane systems [12, 59–63]. In the context of the ER, cubic TPMS have been found in response to elevated levels of specific membrane resident proteins such as cytochrome b(5) [56]. When applied to P-Schwartz and D-Schwartz TPMS with prescribed surface average of spontaneous curvature per unit cell [64, 65], simulation results from our model for minimal surfaces showed that the dimensionless spontaneous curvature qualitatively follows the Gaussian curvature (Fig. 7, see Supplementary Material Section S3 for details [66]). Indeed, C/C_0 is smaller on "flatter" surfaces and larger on the more pronounced saddle-like surfaces.

Figure 7: Distribution of spontaneous curvature maintaining a triply periodic minimal surface (TPMS) structure for an imposed surface average of spontaneous curvature of C_0. (a) Cubic P-Schwartz, and (b) diamond D-Schwartz, shown in 3 by 3 arrays and in close up unit cells. Because P- and D-Schwartz surfaces belong to the same associated family of surfaces, the distribution of spontaneous curvature also follows an isometric transformation between the two structures.

In this study we considered helicoidal surfaces as a simplified model for ER ramps observed in the peripheral ER of neural and secretory cells [1, 26]. This simplification results in certain limitations. First, ER sheet structures are made of two lipid bilayer typically 40 nm apart [26, 52]. In the current model, the double bilayer is represented as a unique surface of zero thickness lying in between the two bilayer (see Model assumption (a)), in line with previous modeling studies [26, 30]. This approximation has two consequences on the interpretation of our results in the context of ER structures. First, the effective bending and Gaussian moduli have twice the value of a single bilayer, provided that both bilayers have the same spontaneous curvature [67]. This does not affect the results presented here because on minimal surfaces both moduli vanish from the shape equation and incompressibility condition, and only their ratio appears in the definition of the dimensionless bending energy (Eq. 11). Second, the edges of helicoidal ER ramps are bent surfaces connecting the two bilayers. To account for this geometry, previous modeling studies proposed effective boundary conditions that assume toroidal surface elements at the ramps' edges [26, 30]. The resulting edge energy was then expressed as analogously to the bending energy for the geodesic curvature, where edge scaffolding proteins induce a spontaneous edge curvature [30]. In the absence of available data supporting the detailed structures of ER sheet edges, we assumed for simplicity either Dirichlet or no-flux Neumann boundary conditions for the spontaneous curvature. The derivation of rigorous effective boundary conditions relating protein distribution and spontaneous curvature at open double-bilayer edges will likely require relaxing the Model assumption (f) to account for area difference elasticity between the leaflets as well as between the two bilayers, and possibly allow lipid tilt at the highly curved edges [40].

While in this paper we mostly discuss the role of protein-induced spontaneous curvature, it is worth noting that asymmetric lipid composition between the inner and outer leaflets can also lead to spontaneous curvature effects

(see Model assumption (b)). Depending on the ratio between the size of the lipid headgroup and the surface area occupied by their hydrocarbon chain, lipid shape can be divided between cylindrical, conical, and inverted conical. The two predominantly found lipids in mammalian ER membranes are the cylindrical phosphatidylcholine (PC), and the conical phosphatidylethanolamine (PE) [68]. As a consequence, an unbalanced PC/PE composition between these two lipids across the ER bilayer can generate spontaneous curvature, while an imposed membrane curvature can lead to lipid sorting to accommodate the constrained geometry [69, 70]. In addition to lipid packing effects, membrane electrostatics can also contribute to curvature generation. Indeed, in early endosomal membranes such as in the ER, phosphatidylserine (PS) – a negatively charged lipid primary responsible for membrane electrostatics in cells – is found almost exclusively in the luminal leaflet[71, 72]. It should be noted however that in the present model the surface represents two bilayers. Therefore, lipid composition asymmetry between the luminal and cytosolic sides of the ER are likely to be compensated by the symmetry of the two bilayers with respect to the mid-lumen plan.

Another key assumption of our model is the homogeneous values of bending and Gaussian moduli (Model assumption (g)). One can expect that a heterogeneous membrane composition – either in proteins or lipids – will induce spatial variations of the membrane mechanical properties [73]. The shape equation for heterogeneous membranes only has extra terms proportional to the covariant and surface derivatives of the Gaussian modulus [38, 42]. This means that in the case where only the bending modulus is dependent on the local composition ($k(\sigma)$ and k_G is a constant), Eq. (4) remains valid at the condition that spontaneous curvature is reinterpreted as the quantity $C^*(\sigma) = k(\sigma)C(\sigma)$. Additionally, it can be shown that in this case Eq. (6) remains unchanged, and therefore our results hold with $C = C^*$. Given the difficulties to measure and compute the value of the Gaussian modulus, such a scenario is a reasonable assumptions. In the case where the Gaussian modulus is also heterogeneous, extra coupling terms appear in the shape and incompressibility equations for the membrane, and a more general model would be required.

Overall this work provides a modeling framework to gain information on the distribution of curvature-inducing proteins on cellular membrane based on the knowledge of their shape. Despite its simplicity, we anticipate that the proposed approach has the potential to support super-resolution microscopy techniques by relating observed membrane structure to their biophysical properties.

Author Contributions

All authors contributed to the study design, model development, result analysis, and writing of the manuscript. MC carried the numerical implementation and data analysis.

Acknowledgments

The authors are grateful to Prof. Andreas Carlson, Jennifer Fromm, and Allen Leung for their critical comments.

Data Accessibility

The main Comsol file, Matlab code, and a sample data file, are available at the author's Github depository:
`github.com/mchabanon/helicoid_spontaneous_curvature`

Funding Statement

This work was supported by the ONR N00014-17-1- 2628 and NSF PHY 1505017 awards to PR.

References

[1] Nixon-Abell J, Obara CJ, Weigel AV, Li D, Legant WR, Xu CS, et al. Increased Spatiotemporal Resolution Reveals Highly Dynamic Dense Tubular Matrices in the Peripheral ER. Science. 2016 Oct;354(6311):aaf3928. doi:10.1126/science.aaf3928.

[2] Valm AM, Cohen S, Legant WR, Melunis J, Hershberg U, Wait E, et al. Applying Systems-Level Spectral Imaging and Analysis to Reveal the Organelle Interactome. Nature. 2017 Jun;546(7656):162–167. doi:10.1038/nature22369.

[3] Schroeder LK, Barentine AES, Merta H, Schweighofer S, Zhang Y, Baddeley D, et al. Dynamic Nanoscale Morphology of the ER Surveyed by STED Microscopy. J Cell Biol. 2019 Jan;218(1):83–96. doi:10.1083/jcb.201809107.

[4] Wu Y, Whiteus C, Xu CS, Hayworth KJ, Weinberg RJ, Hess HF, et al. Contacts between the Endoplasmic Reticulum and Other Membranes in Neurons. Proceedings of the National Academy of Sciences. 2017 Jun;114(24):E4859–E4867. doi:10.1073/pnas.1701078114.

[5] Voeltz GK, Prinz WA. Sheets, Ribbons and Tubules — How Organelles Get Their Shape. Nature Reviews Molecular Cell Biology. 2007 Mar;8(3):258–264. doi:10.1038/nrm2119.

[6] Kukulski W, Schorb M, Kaksonen M, Briggs JAG. Plasma Membrane Reshaping during Endocytosis Is Revealed by Time-Resolved Electron Tomography. Cell. 2012 Aug;150(3):508–520. doi:10.1016/j.cell.2012.05.046.

[7] Cocucci E, Aguet F, Boulant S, Kirchhausen T. The First Five Seconds in the Life of a Clathrin-Coated Pit. Cell. 2012 Aug;150(3):495–507. doi:10.1016/j.cell.2012.05.047.

[8] Hassinger JE, Oster G, Drubin DG, Rangamani P. Design Principles for Robust Vesiculation in Clathrin-Mediated Endocytosis. Proceedings of the National Academy of Sciences. 2017 Jan;p. 201617705. doi:10.1073/pnas.1617705114.

[9] Wollert T, Hurley JH. Molecular Mechanism of Multivesicular Body Biogenesis by ESCRT Complexes. Nature. 2010 Apr;464(7290):864–869. doi:10.1038/nature08849.

[10] Hurley JH, Hanson PI. Membrane Budding and Scission by the ESCRT Machinery: It's All in the Neck. Nature Reviews Molecular Cell Biology. 2010 Aug;11(8):556–566. doi:10.1038/nrm2937.

[11] Guizetti J, Gerlich DW. ESCRT-III Polymers in Membrane Neck Constriction. Trends in Cell Biology. 2012 Mar;22(3):133–140. doi:10.1016/j.tcb.2011.11.007.

[12] Schmidt NW, Wong GCL. Antimicrobial Peptides and Induced Membrane Curvature: Geometry, Coordination Chemistry, and Molecular Engineering. Current Opinion in Solid State and Materials Science. 2013 Aug;17(4):151–163. doi:10.1016/j.cossms.2013.09.004.

[13] Saikia K, Sravani YD, Ramakrishnan V, Chaudhary N. Highly Potent Antimicrobial Peptides from N-Terminal Membrane-Binding Region of E. Coli MreB. Scientific Reports. 2017 Feb;7:42994. doi:10.1038/srep42994.

[14] Lee C, Goldberg J. Structure of Coatomer Cage Proteins and the Relationship among COPI, COPII, and Clathrin Vesicle Coats. Cell. 2010 Jul;142(1):123–132. doi:10.1016/j.cell.2010.05.030.

[15] Campelo F, van Galen J, Turacchio G, Parashuraman S, Kozlov MM, García-Parajo MF, et al. Sphingomyelin Metabolism Controls the Shape and Function of the Golgi Cisternae. eLife. 2017 May;6:e24603. doi:10.7554/eLife.24603.

[16] Goyal U, Blackstone C. Untangling the Web: Mechanisms Underlying ER Network Formation. Biochimica et Biophysica Acta (BBA)-Molecular Cell Research. 2013;1833(11):2492–2498. doi:10.1016/j.bbamcr.2013.04.009.

[17] Puhka M, Joensuu M, Vihinen H, Belevich I, Jokitalo E. Progressive Sheet-to-Tubule Transformation Is a General Mechanism for Endoplasmic Reticulum Partitioning in Dividing Mammalian Cells. Molecular Biology of the Cell. 2012 Jan;23(13):2424–2432. doi:10.1091/mbc.E10-12-0950.

[18] Georgiades P, Allan VJ, Wright GD, Woodman PG, Udommai P, Chung MA, et al. The Flexibility and Dynamics of the Tubules in the Endoplasmic Reticulum. Scientific Reports. 2017 Nov;7(1):16474. doi:10.1038/s41598-017-16570-4.

[19] Rambold AS, Kostelecky B, Elia N, Lippincott-Schwartz J. Tubular Network Formation Protects Mitochondria from Autophagosomal Degradation during Nutrient Starvation. Proceedings of the National Academy of Sciences. 2011 Jun;108(25):10190–10195. doi:10.1073/pnas.1107402108.

[20] Rafelski SM. Mitochondrial Network Morphology: Building an Integrative, Geometrical View. BMC Biology. 2013 Jun;11(1):71. doi:10.1186/1741-7007-11-71.

[21] Wang C, Du W, Su QP, Zhu M, Feng P, Li Y, et al. Dynamic Tubulation of Mitochondria Drives Mitochondrial Network Formation. Cell Research. 2015 Oct;25(10):1108–1120. doi:10.1038/cr.2015.89.

[22] Davis DM, Sowinski S. Membrane Nanotubes: Dynamic Long-Distance Connections between Animal Cells. Nature Reviews Molecular Cell Biology. 2008 Jun;9(6):431–436. doi:10.1038/nrm2399.

[23] Gerdes HH, Rustom A, Wang X. Tunneling Nanotubes, an Emerging Intercellular Communication Route in Development. Mechanisms of Development. 2013 Jun;130(6):381–387. doi:10.1016/j.mod.2012.11.006.

[24] Marzo L, Gousset K, Zurzolo C. Multifaceted Roles of Tunneling Nanotubes in Intercellular Communication. Frontiers in Physiology. 2012;3. doi:10.3389/fphys.2012.00072.

[25] Chabanon M, Rangamani P. Gaussian Curvature Directs the Distribution of Spontaneous Curvature on Bilayer Membrane Necks. Soft Matter. 2018;14(12):2281–2294. doi:10.1039/C8SM00035B.

[26] Terasaki M, Shemesh T, Kasthuri N, Klemm RW, Schalek R, Hayworth KJ, et al. Stacked Endoplasmic Reticulum Sheets Are Connected by Helicoidal Membrane Motifs. Cell. 2013 Jul;154(2):285–296. doi:10.1016/j.cell.2013.06.031.

[27] Lee MW, Lee EY, Lai GH, Kennedy NW, Posey AE, Xian W, et al. Molecular Motor Dnm1 Synergistically Induces Membrane Curvature To Facilitate Mitochondrial Fission. ACS Central Science. 2017 Nov;3(11):1156–1167. doi:10.1021/acscentsci.7b00338.

[28] Beber A, Taveneau C, Nania M, Tsai FC, Cicco AD, Bassereau P, et al. Membrane Reshaping by Micrometric Curvature Sensitive Septin Filaments. Nature Communications. 2019 Jan;10(1):420. doi:10.1038/s41467-019-08344-5.

[29] Marshall WF. Differential Geometry Meets the Cell. Cell. 2013 Jul;154(2):265–266. doi:10.1016/j.cell.2013.06.032.

[30] Shemesh T, Klemm RW, Romano FB, Wang S, Vaughan J, Zhuang X, et al. A Model for the Generation and Interconversion of ER Morphologies. Proceedings of the National Academy of Sciences. 2014 Dec;111(49):E5243–E5251. doi:10.1073/pnas.1419997111.

[31] Guven J, Huber G, Valencia DM. Terasaki Spiral Ramps in the Rough Endoplasmic Reticulum. Physical Review Letters. 2014 Oct;113(18):188101. doi:10.1103/PhysRevLett.113.188101.

[32] Helfrich W. Elastic Properties of Lipid Bilayers: Theory and Possible Experiments. Zeitschrift fur Naturforschung Teil C. 1973;28(11):693–703. doi:10.1515/znc-1973-11-1209.

[33] Chabanon M, Stachowiak JC, Rangamani P. Systems Biology of Cellular Membranes: A Convergence with Biophysics. WIREs Systems Biology and Medicine. 2017 Apr;p. e01386. doi:10.1002/wsbm.1386.

[34] Evans E, Rawicz W. Entropy-Driven Tension and Bending Elasticity in Condensed-Fluid Membranes. Physical Review Letters. 1990 Apr;64(17):2094–2097. doi:10.1103/PhysRevLett.64.2094.

[35] Rangamani P, Mandadap KK, Oster G. Protein-Induced Membrane Curvature Alters Local Membrane Tension. Biophysical Journal. 2014 Aug;107(3):751–762. doi:10.1016/j.bpj.2014.06.010.

[36] Steigmann DJ. Mechanics and Physics of Lipid Bilayers. In: The Role of Mechanics in the Study of Lipid Bilayers. CISM International Centre for Mechanical Sciences. Springer, Cham; 2018. p. 1–61. 10.1007/978-3-319-56348-0-1.

[37] Rahimi M, Arroyo M. Shape Dynamics, Lipid Hydrodynamics, and the Complex Viscoelasticity of Bilayer Membranes. Physical Review E. 2012 Jul;86(1):011932. doi:10.1103/PhysRevE.86.011932.

[38] Rangamani P, Agrawal A, Mandadapu KK, Oster G, Steigmann DJ. Interaction between Surface Shape and Intra-Surface Viscous Flow on Lipid Membranes. Biomechanics and Modeling in Mechanobiology. 2013 Aug;12(4):833–845. doi:10.1007/s10237-012-0447-y.

[39] Agrawal A, Steigmann DJ. A Model for Surface Diffusion of Trans-Membrane Proteins on Lipid Bilayers. Zeitschrift für angewandte Mathematik und Physik. 2011 May;62(3):549. doi:10.1007/s00033-011-0132-5.

[40] Alimohamadi H, Rangamani P. Modeling Membrane Curvature Generation Due to Membrane–Protein Interactions. Biomolecules. 2018 Dec;8(4):120. doi:10.3390/biom8040120.

[41] Belay T, Kim CI, Schiavone P. Bud Formation of Lipid Membranes in Response to the Surface Diffusion of Transmembrane Proteins and Line Tension. Mathematics and Mechanics of Solids. 2017 Nov;22(11):2091–2107. doi:10.1177/1081286516657684.

[42] Steigmann DJ. Fluid Films with Curvature Elasticity. Archive for Rational Mechanics and Analysis. 1999 Dec;150(2):127–152. doi:10.1007/s002050050183.

[43] Agrawal A, Steigmann DJ. Boundary-Value Problems in the Theory of Lipid Membranes. Continuum Mechanics and Thermodynamics. 2009 Jun;21(1):57–82. doi:10.1007/s00161-009-0102-8.

[44] Agrawal A, Steigmann DJ. Modeling Protein-Mediated Morphology in Biomembranes. Biomechanics and modeling in mechanobiology. 2009;8(5):371–379. doi:10.1007/s10237-008-0143-0.

[45] Sokolnikoff IS. Tensor Analysis: Theory and Applications to Geometry and Mechanics of Continua. Wiley; 1964.

[46] Bahmani F, Christenson J, Rangamani P. Analysis of Lipid Flow on Minimal Surfaces. Continuum Mechanics and Thermodynamics. 2015 Jul;28(1-2):503–513. doi:10.1007/s00161-015-0458-x.

[47] Shibata Y, Shemesh T, Prinz WA, Palazzo AF, Kozlov MM, Rapoport TA. Mechanisms Determining the Morphology of the Peripheral ER. Cell. 2010 Nov;143(5):774–788. doi:10.1016/j.cell.2010.11.007.

[48] Struik DJ. Lectures on Classical Differential Geometry: Second Edition. Second edition ed. New York: Dover Publications; 1988.

[49] Kreyszig E. Differential Geometry. 1st ed. New York: Dover Publications; 1991.

[50] Hu M, Briguglio JJ, Deserno M. Determining the Gaussian Curvature Modulus of Lipid Membranes in Simulations. Biophysical Journal. 2012 Mar;102(6):1403–1410. doi:10.1016/j.bpj.2012.02.013.

[51] Lee IH, Kai H, Carlson LA, Groves JT, Hurley JH. Negative Membrane Curvature Catalyzes Nucleation of Endosomal Sorting Complex Required for Transport (ESCRT)-III Assembly. Proceedings of the National Academy of Sciences. 2015 Dec;112(52):15892–15897. doi:10.1073/pnas.1518765113.

[52] Westrate LM, Lee JE, Prinz WA, Voeltz GK. Form Follows Function: The Importance of Endoplasmic Reticulum Shape. Annual Review of Biochemistry. 2015;84(1):791–811. doi:10.1146/annurev-biochem-072711-163501.

[53] Zhang H, Hu J. Shaping the Endoplasmic Reticulum into a Social Network. Trends in Cell Biology. 2016 Dec;26(12):934–943. doi:10.1016/j.tcb.2016.06.002.

[54] Gunning BES. The Greening Process in Plastids. Protoplasma. 1965 Mar;60(1):111–130. doi:10.1007/BF01248133.

[55] Deng Y, Mieczkowski M. Three-Dimensional Periodic Cubic Membrane Structure in the Mitochondria of Amoebae Chaos Carolinensis. Protoplasma. 1998;203(1-2):16–25. doi:10.1007/BF01280583.

[56] Snapp EL, Hegde RS, Francolini M, Lombardo F, Colombo S, Pedrazzini E, et al. Formation of Stacked ER Cisternae by Low Affinity Protein Interactions. Journal of cell biology. 2003 Oct;163(2):257–269. doi:10.1083/jcb.200306020.

[57] Goldsmith CS, Tatti KM, Ksiazek TG, Rollin PE, Comer JA, Lee WW, et al. Ultrastructural Characterization of SARS Coronavirus. Emerging infectious diseases. 2004;10(2):320. doi:10.3201/eid1002.030913.

[58] Almsherqi ZA, Kohlwein SD, Deng Y. Cubic Membranes: A Legend beyond the Flatland* of Cell Membrane Organization. The Journal of Cell Biology. 2006 Jun;173(6):839–844. doi:10.1083/jcb.200603055.

[59] Schwarz US, Gompper G. Stability of Inverse Bicontinuous Cubic Phases in Lipid-Water Mixtures. Physical Review Letters. 2000 Aug;85(7):1472–1475. doi:10.1103/PhysRevLett.85.1472.

[60] Shearman GC, Ces O, Templer RH, Seddon JM. Inverse Lyotropic Phases of Lipids and Membrane Curvature. Journal of Physics: Condensed Matter. 2006;18(28):S1105. doi:10.1088/0953-8984/18/28/S01.

[61] Demurtas D, Guichard P, Martiel I, Mezzenga R, Hébert C, Sagalowicz L. Direct Visualization of Dispersed Lipid Bicontinuous Cubic Phases by Cryo-Electron Tomography. Nature Communications. 2015 Nov;6:8915. doi:10.1038/ncomms9915.

[62] Tran N, Mulet X, Hawley AM, Conn CE, Zhai J, Waddington LJ, et al. First Direct Observation of Stable Internally Ordered Janus Nanoparticles Created by Lipid Self-Assembly. Nano Letters. 2015 Jun;15(6):4229–4233. doi:10.1021/acs.nanolett.5b01751.

[63] Sankhagowit S, Lee EY, Wong GCL, Malmstadt N. Oxidation of Membrane Curvature-Regulating Phosphatidylethanolamine Lipid Results in Formation of Bilayer and Cubic Structures. Langmuir. 2016 Mar;32(10):2450–2457. doi:10.1021/acs.langmuir.5b04332.

[64] Chabanon M, David B, Goyeau B. Averaged Model for Momentum and Dispersion in Hierarchical Porous Media. Physical Review E. 2015 Aug;92(2):023201. doi:10.1103/PhysRevE.92.023201.

[65] Chabanon M, Valdés-Parada FJ, Ochoa-Tapia JA, Goyeau B. Large-Scale Model of Flow in Heterogeneous and Hierarchical Porous Media. Advances in Water Resources. 2017 Nov;109:41–57. doi:10.1016/j.advwatres.2017.08.018.

[66] Brakke KA. The Surface Evolver. Experimental Mathematics. 1992 Jan;1(2):141–165. doi:10.1080/10586458.1992.10504253.

[67] Siegel DP, Kozlov MM. The Gaussian Curvature Elastic Modulus of N-Monomethylated Dioleoylphosphatidylethanolamine: Relevance to Membrane Fusion and Lipid Phase Behavior. Biophysical Journal. 2004 Jul;87(1):366–374. doi:10.1529/biophysj.104.040782.

[68] van Meer G, de Kroon AIPM. Lipid Map of the Mammalian Cell. Journal of Cell Science. 2011 Jan;124(1):5–8. doi:10.1242/jcs.071233.

[69] Roux A, Cuvelier D, Nassoy P, Prost J, Bassereau P, Goud B. Role of Curvature and Phase Transition in Lipid Sorting and Fission of Membrane Tubules. The EMBO Journal. 2005 Apr;24(8):1537–1545. doi:10.1038/sj.emboj.7600631.

[70] Callan-Jones A, Sorre B, Bassereau P. Curvature-Driven Lipid Sorting in Biomembranes. Cold Spring Harbor Perspectives in Biology. 2011 Jan;3(2):a004648. doi:10.1101/cshperspect.a004648.

[71] Fairn GD, Schieber NL, Ariotti N, Murphy S, Kuerschner L, Webb RI, et al. High-Resolution Mapping Reveals Topologically Distinct Cellular Pools of Phosphatidylserine. The Journal of Cell Biology. 2011 Jul;194(2):257–275. doi:10.1083/jcb.201012028.

[72] Kay JG, Koivusalo M, Ma X, Wohland T, Grinstein S. Phosphatidylserine Dynamics in Cellular Membranes. Molecular Biology of the Cell. 2012 Apr;23(11):2198–2212. doi:10.1091/mbc.e11-11-0936.

[73] Dimova R. Recent Developments in the Field of Bending Rigidity Measurements on Membranes. Advances in Colloid and Interface Science. 2014 Jun;208:225–234. doi:10.1016/j.cis.2014.03.003.

Supplementary Material for:

Geometric coupling of helicoidal ramps and curvature-inducing proteins in organelle membranes

Morgan Chabanon[1] & Padmini Rangamani[1*]

[1] Department of Mechanical and Aerospace Engineering, University of California San Diego, La Jolla, CA, USA.

* Corresponding author. E-mail: prangamani@eng.ucsd.edu

Contents

S1   Local force balance of elastic surfaces at mechanical equilibrium

The equation of mechanical equilibrium of an elastic surface ω subject to a lateral pressure p can be written in the compact form

$$\boldsymbol{\Sigma}^\alpha_{;\alpha} + p\mathbf{n} = \mathbf{0} \,, \tag{S1}$$

where $\boldsymbol{\Sigma}^\alpha$ are the stress vectors and $\mathbf{n}$ is the unit normal to the local surface. Greek indices range over $1, 2$, and if repeated, are summed over this range. Semicolon identifies covariant differentiation with respect to the surface metric $a_{\alpha\beta} = \mathbf{a}_\alpha \cdot \mathbf{a}_\beta$ where $\mathbf{a}_\alpha = \mathbf{r}_{,\alpha}$ are the tangent vectors and $\mathbf{r}(\theta^\alpha)$ is the parametrization of the position field. The commas refer to partial derivatives with respect to the surface coordinates θ^α. With these definitions, the normal vector is given by $\mathbf{n} = (\mathbf{a}_1 \times \mathbf{a}_2)/ \mid \mathbf{a}_1 \times \mathbf{a}_2 \mid$. In Eq. S1, the differential operation represents the surface divergence defined as $\boldsymbol{\Sigma}^\alpha_{;\alpha} = (\sqrt{a})^{-1}(\sqrt{a}\boldsymbol{\Sigma}^\alpha)_{,\alpha}$ where $a = \det(a_{\alpha\beta})$. In surface theory, a manifold is described by the metric $a_{\alpha\beta}$ defined above, and the curvature tensor given by $b_{\alpha\beta} = \mathbf{n} \cdot \mathbf{r}_{,\alpha\beta}$.

For an elastic membrane whose energy surface density per unit mass depends on the metric and curvature only $F(a_{\alpha\beta}, b_{\alpha\beta}; \theta^\alpha)$, the stress vectors involved in the local force balance (Eq. S1) can be written as [42]

$$\boldsymbol{\Sigma}^\alpha = \mathbf{T}^\alpha + S^\alpha \mathbf{n} \,. \tag{S2}$$

Here the tangential stress vectors are

$$\mathbf{T}^\alpha = T^{\beta\alpha}\mathbf{a}_\beta \quad \text{with} \quad T^{\beta\alpha} = \Sigma^{\beta\alpha} + b^\beta_\mu M^{\mu\alpha} \,, \tag{S3}$$

and the components of the normal stress vectors are

$$S^\alpha = -M^{\alpha\beta}_{;\beta} \,, \tag{S4}$$

where $b^\beta_\alpha = a^{\beta\lambda}b_{\lambda\alpha}$. The components of the stress vectors depends on the energy density as [42]

$$\Sigma^{\alpha\beta} = \rho\left(\frac{\partial F}{\partial a_{\alpha\beta}} + \frac{\partial F}{\partial a_{\beta\alpha}}\right) \quad \text{and} \quad M^{\alpha\beta} = \frac{\rho}{2}\left(\frac{\partial F}{\partial b_{\alpha\beta}} + \frac{\partial F}{\partial b_{\beta\alpha}}\right) \,, \tag{S5}$$

where ρ is the surface mass density of the membrane. The tangential and normal local force balances can now be obtained by introducing Eqs. S2, S3, and S4 into Eq. S1, resulting in

$$T^{\beta\alpha}_{;\alpha} - S^\alpha b^\beta_\alpha = 0 \quad \text{and} \quad S^\alpha_{;\alpha} + T^{\beta\alpha}b_{\beta\alpha} + p = 0 \,, \tag{S6}$$

where we made use of the Gauss and Weingarten equations [45] $\mathbf{a}_{\alpha;\beta} = b_{\alpha\beta}\mathbf{n}$ and $\mathbf{n}_{,\alpha} = -b_\alpha^\beta \mathbf{a}_\beta$ respectively.

The free energy density can be written as a function of the mean curvature H and Gaussian curvature K. These are related to the metric and curvature by

$$H = \frac{1}{2}a^{\alpha\beta}b_{\alpha\beta} \quad \text{and} \quad K = \frac{1}{2}\varepsilon^{\alpha\beta}\varepsilon^{\lambda\mu}b_{\alpha\lambda}b_{\beta\mu} , \tag{S7}$$

where $a^{\alpha\beta} = (a_{\alpha\beta})^{-1}$ is the dual metric, and $\varepsilon^{\alpha\beta}$ is the permutation tensor defined by $\varepsilon^{12} = -\varepsilon^{21} = 1/\sqrt{a}$, $\varepsilon^{11} = \varepsilon^{22} = 0$. According to this definition (Eq. S7), the free energy density per unit mass can be re-written in terms of the mean and Gaussian curvatures $F(H, K; \theta^\alpha)$. Furthermore, lipid membranes are essentially incompressible (see assumption (d) in the Model Development Section of the main text). This is imposed using a Lagrange multiplier $\gamma(\theta^\alpha)$ to ensure that the local area dilatation $J = 1$, or equivalently, to constraint the constant surface density ρ of the membrane. Consequently we can define the surface energy density of the membrane as follows

$$F(\rho, H, K; \theta^\alpha) = \bar{F}(H, K; \theta^\alpha) - \frac{\gamma(\theta^\alpha)}{\rho} , \tag{S8}$$

and when introducing the surface energy per unit area $W(\rho, H, K; \theta^\alpha) = \rho\bar{F}(H, K; \theta^\alpha)$, the components of the stress vectors (Eqs. S5) can be written as [42]

$$\Sigma^{\alpha\beta} = (\lambda + W)a^{\alpha\beta} - (2HW_H + 2KW_K)\,a^{\alpha\beta} + W_H\tilde{b}^{\alpha\beta} \tag{S9}$$

$$M^{\alpha\beta} = \frac{1}{2}W_H a^{\alpha\beta} + W_K\tilde{b}^{\alpha\beta} \tag{S10}$$

where $\lambda(\theta^\alpha) = -[\gamma(\theta^\alpha) + W(H, K; \theta^\alpha)]$, and $\tilde{b}^{\alpha\beta} = 2Ha^{\alpha\beta} - b^{\alpha\beta}$ is the cofactor of the curvature. The subscripts H and K refer to the partial derivative of the energy with respect to the indicated variable. Note that the Lagrange multiplier γ can be interpreted as a surface pressure, and is not a material property of the surface [35, 38]. Consequently, λ can be interpreted as the surface tension based on comparisons with edge conditions on a flat surface [35].

Finally, introducing Eqs. S9 and S10 into Eqs. S3 and S4, we can rewrite the normal and tangential force balances (Eqs. S6) as

$$\Delta\left(\frac{1}{2}W_H\right) + (W_K)_{;\alpha\beta}\tilde{b}^{\alpha\beta} + W_H(2H^2 - K) + 2H(KW_K - W) = p + 2\lambda H , \tag{S11}$$

and

$$-(\gamma_{,\alpha} + W_K K_{,\alpha} + W_H H_{,\alpha})\,a^{\beta\alpha} = \left(\left.\frac{\partial W}{\partial\theta^\alpha}\right|_{\exp} + \lambda_{,\alpha}\right)a^{\beta\alpha} = 0 , \tag{S12}$$

where $\Delta(\cdot) = (\cdot)_{;\alpha\beta}a^{\alpha\beta}$ is the surface Laplacian (or Beltrami operator), and $\partial(\cdot)/\partial\theta^\alpha\,|_{\exp}$ is the explicit derivative with respect to θ^α.

Eqs. S11 and S12 are the general shape equation and incompressibility condition for an elastic surface with free energy per unit area $W(\rho, H, K; \theta^\alpha)$.

S2 Helicoid to catenoid transformation

According to the Gauss' *Theorema Egregium*, the distribution of Gaussian curvature on a minimal surface follows any isometric mapping of such surface [49]. As illustrated in Fig. 2b, such transformation exists between helicoids and catenoids.

Elementary surface	Surface evolution sequence
P-Schwartz	g 5; r; g 5; r; g 5; r; g 10; u; r; g 10; u; r; g 10; u; u; refine edge where on_constraint 2; refine edge where on_constraint 4; hessian; hessian; u ;
D-Schwartz	g 5; r; g 5; r; g 5; r; g 10; u; r; g 10; u; r; g 10; u; u; hessian; hessian; r ; g 100; hessian; hessian ; g 100; u; g 500 ; hessian; g 100;

Table S1: Command sequence to refine the surface meshes of the elementary P- and D-Schwartz elementary surfaces. Description of the commands can be found in the Surface Evolver documentation.

Helicoids and catenoids belong to the same associated family of surfaces. The (continuous) isometric transformation from one surface to the other can be written as

$$\begin{cases} x = \sin(\alpha)r_n \cosh(u/r_n)\sin(\phi) - \cos(\alpha)r_n \sinh(u/r_n)\cos(\phi) \\ y = -\sin(\alpha)r_n \cosh(u/r_n)\cos(\phi) - \cos(\alpha)r_n \sinh(u/r_n)\sin(\phi) \\ z = \sin(\alpha)u + \cos(\alpha)r_n\phi \end{cases} \quad \text{with} \quad \begin{cases} \phi \in [-\pi;\pi[\\ u \in [-H_c/2; H_c/2] \\ \alpha \in [0;\pi/2] \end{cases}, \tag{S13}$$

where P is the helicoid pitch, related to the catenoid neck radius by $r_n = P/(2\pi)$, and L is the helicoid diameter, related to the catenoid heigh by $H_c = 2r_n \sinh^{-1}(L/(2r_n))$.

For $\alpha = \pi/2$, this system describes a catenoid such as the one shown in Fig. 2c, while for $\alpha = 0$ it describes a helicoid as in Fig. 2a. The equivalence between Eqs. S13 and 7 when $\alpha = 0$ can be verified with the change of variable $r = r_n \sinh(u/r_n)$. Any intermediate values of α gives rise to a minimal surfaces belonging to the associated family of helicoids and catenoids, as illustrated in Fig. 2b.

The Gaussian curvature of these surfaces is given by

$$K = -\left(\frac{P/(2\pi)}{[P/(2\pi)]^2 + r^2}\right)^2 = -\left(\frac{r_n}{r_n^2 + [r_n \sinh(u/r_n)]^2}\right)^2. \tag{S14}$$

S3   Method for computing the distribution of spontaneous curvature on TPMS

Elementary surfaces P-Schwartz or D-Schwartz were downloaded from [1] and imported in Surface Evolver [66]. The surface energy was minimized through a succession of minimization and mesh refinement operations (see Table S1). The refined elementary surfaces were then duplicated and flipped to produce the periodic cubic unit cells, before to be exported in .stl format [2].

The cubic unit cell meshes were imported in Comosol Multiphysics, and Eq. 9 was solved using the "Surface reaction" module. Because of the periodic nature of the TPMS, instead of imposing Dirichlet boundary conditions at the boundaries, we chose periodic flux boundary conditions at the opposite edges, and imposed a surface average of C_0 on the unit cells. This choice is inspired from approaches to solve closure problems in transport in porous media, where the elementary unit volume is assumed to be periodic [64, 65].

---

[1]`facstaff.susqu.edu/brakke/evolver/examples/periodic/periodic.html` (consulted on January 29th, 2019)

[2]`github.com/kashif/evolver/blob/master/fe/stl.cmd` (consulted on January 29th, 2019)

S4 Supplementary Figures

Figure S1: Helicoid pitch controls the position of the energy the spontaneous curvature switch in sign at the inner boundary.

Figure S2: Distribution of spontaneous cutvature along the helicoid radius for C_0 imposed at the inner boundary of the catenoid for various pitch and inner radii. No switch is observed.

Figure S3: Normalized bending energy of helicoids with boundary conditions imposed on the inner bondary. No energy barrier is observed.

Figure S4: Effect of the imposed gradient in C at the external boundary on the bending energy of the helicoid.

Figure S5: Effect of the pitch on the switch and energy barrier of helicoids with imposed gradient of boundary conditions at the exterior boundary.