# Operating Characteristics of the Rank-Based Inverse Normal Transformation for Quantitative Trait Analysis in Genome-Wide Association Studies

**Zachary R. McCaw[1,*], Jacqueline M. Lane[2], Richa Saxena[2], Susan Redline[3], Xihong Lin[1,4,**]**

[1]Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA

[2]Center for Human Genetic Research, Massachusetts General Hospital, Boston, MA

[3]Division of Sleep and Circadian Disorders, Brigham and Women's Hospital, Boston, MA

[4]Department of Statistics, Harvard University, Cambridge, MA

*email: zrmacc@gmail.com

**email: xlin@hsph.harvard.edu

SUMMARY: Quantitative traits analyzed in Genome-Wide Association Studies (GWAS) are often non-normally distributed. For such traits, association tests based on standard linear regression have reduced power and inflated type I error. Applying the rank-based Inverse Normal Transformation (INT) to non-normally distributed traits has become common practice in GWAS. However, the different variations on INT-based association testing have not been formally defined, and guidance is lacking on when to use which approach. In this paper, we formally define and systematically compare the direct (D-INT) and indirect (I-INT) INT-based association tests. We discuss their assumptions, underlying generative models, and connections. We demonstrate that the relative powers of D-INT and I-INT depend on the underlying data generating process. Since neither approach is uniformly most powerful, we combine them into an adaptive, INT-based omnibus test (O-INT). O-INT is robust to model misspecification, protects the type I error, and is well powered against a wide range of non-normally distributed traits. Extensive simulations were conducted to examine the finite sample operating characteristics of these tests. Our results demonstrate that, for non-normally distributed traits, INT-based tests outperform the standard untransformed association test (UAT), both in terms of power and type I error rate control. We apply the proposed methods to GWAS of spirometry traits in the UK Biobank.

KEY WORDS: Direct and indirect rank-based inverse normal transformation; Non-normality; Omnibus test; Quantitative Traits; Transformation; Type I error rate.

This paper has been submitted for consideration for publication in *Biometrics*

## 1. Introduction

Linear regression is the standard method for analyzing quantitative (continuous) traits in Genome-wide Association Studies (GWAS). Few assumptions are placed on the form of the residual distribution: in samples of sufficient size, only mean zero and finite variance are required. However, the operating characteristics of linear regression are sensitive to the underlying residual distribution. Many complex, quantitative traits have markedly non-normal residuals. Examples include spirometry traits, such as peak expiratory flow (PEF), whose distribution is skewed and asymmetric even after log transformation. When the residual distribution is non-normal, linear regression is susceptible to inflated type I error at the genome-wide significance threshold ($\alpha = 5 \times 10^{-8}$). Even in samples of sufficient size to protect the type I error (often requiring $N > 10^5$), linear regression lacks power for detecting true associations when the residual distribution is non-normal.

The rank-based inverse normal transformation (INT) is commonly applied during GWAS of non-normally distributed traits. INT is a non-parametric mapping that replaces the sample quantiles by quantiles from the standard normal distribution. Regardless of its initial distribution, the limiting distribution of a continuous trait after INT is standard normal. INT of the outcome improves the performance of linear regression by symmetrizing and concentrating the residual distribution around zero. This in turn allows the sampling distribution of the association test statistic to approach its asymptotic limit more quickly.

We classify INT-based tests into direct and indirect methods. In the direct method (D-INT), INT is applied directly to the phenotype, and the INT-transformed phenotype is regressed on genotype and covariates. Covariates may include age, sex, and adjustments for population structure, such as genetic principal components (PCs). D-INT has been applied to GWAS of BMI (Scuteri et al., 2007), circulating lipids (Barber et al., 2010), polysomnography signals (Cade et al., 2016), and many quantitative traits in the UK Biobank (Abbott et al.,

2017). In the indirect method (I-INT), the phenotype is first regressed on covariates to obtain residuals, then the INT-transformed phenotypic residuals are regressed on genotype, with or without secondary adjustment for population structure. I-INT has been applied to GWAS of gene expression (Emilsson et al., 2008; Consortium et al., 2017), serum metabolites (Kettunen et al., 2012), and spirometry measurements (Repapi et al., 2010). However, the relative performance of D-INT versus I-INT has not been studied in detail. For the non-normal quantitative traits encountered in practice, it is unclear which of the these methods will more robustly control the type I error, and which will provide better power.

As discussed by Beasley and colleagues (Beasley et al., 2009), the question of whether INT-based methods have desirable operating characteristics in the GWAS context has not been critically evaluated. INT of the outcome in a regression model does not guarantee correct model specification. This is because standard linear regression, considered parametrically, requires normality of the residual distribution, not of the marginal distribution of the outcome. Here we systematically study the direct and indirect INT-based association tests, and provide recommendations on how to apply the INT in practice. We begin by formally defining D-INT and I-INT, studying their underlying assumptions and connections. We demonstrate that if the observed trait is generated by a non-linear, rank-preserving transformation of a latent normal trait, then INT provides an approximate inverse of the generative transformation, under the null and supposing the covariate effects are small. Moreover, if the mean of the observed trait is linear in covariates but the residual distribution is non-normal, then I-INT is asymptotically exact. Our derivation of I-INT agrees with recent work recommending double adjustment for covariates during INT-based association testing (Sofar et al., 2019).

Through extensive simulations covering a range of non-normal phenotypes, we compare D-INT and I-INT with the standard untransformed association test (UAT). Our results show that the INT-based tests robustly control the type I error and dominate the UAT in terms

of power. However, the relative power D-INT versus I-INT depends on the underlying data generating process, and neither approach was uniformly most powerful. D-INT performed best against non-linear transformations of a latent normal trait. I-INT performed best against traits with additive but non-normal residuals. Since the data generating mechanism is seldom known in practice, we next propose an adaptive, INT-based, omnibus test (O-INT) that synthesizes D-INT and I-INT. O-INT robustly controls the type I error, and for any given trait is nearly as powerful as the more effective of the component methods. We have implemented all candidate INT-based tests (D-INT, I-INT, and O-INT) in the R package `RNOmni`.

We applied the UAT and the INT-based association tests to GWAS of spirometry traits in the UK Biobank (Sudlow et al., 2015). To demonstrate the power advantage provided by O-INT, we compare the results from the overall analysis ($N = 292K$) with those from a subgroup analysis ($N = 29K$) among asthmatics. All associations identified by O-INT in the asthmatic subgroup were declared significant by UAT in the overall analysis. Hence UAT and O-INT tests agree as to the importance of these loci. However, the more-powerful O-INT test was able to detect them using only a fraction (9.7%) of the sample. In both the asthmatic subgroup and the overall analysis, O-INT realized empirical efficiency and discovery gains over the UAT.

The remainder of this paper is structured as follows. In the methods section, we present the UAT, formally define the INT, theoretically study the direct and indirect INT-based tests, and describe the omnibus test. In the simulation studies, we present evidence that INT-based association tests robustly control the type I error, whereas the UAT often does not, and demonstrate that INT-based association tests dominate the UAT in terms of power. In the data application, we perform GWAS of spirometry traits from the UK Biobank. We conclude with a discussion of the implications of our findings for quantitative trait GWAS.

## 2. Statistical Methods

### 2.1 *Setting*

For each of $N$ independent subjects, the following data are observed: a continuous (quantitative) phenotype $Y_i$, genotype $g_i$ at the locus of interest, and a $p \times 1$ vector $\boldsymbol{x}_i = (x_{i,1}, \cdots, x_{i,p})$ of covariates. In our data application, the phenotype $Y_i$ is a spirometry measurement, while the covariates include an intercept, age, sex, and genetic principal components (PCs). Let $\boldsymbol{y} = (Y_1, \cdots, Y_n)$ denote the $n \times 1$ sample phenotype vector, $\boldsymbol{g}$ the $n \times 1$ genotype vector, and $\boldsymbol{X}$ the $n \times p$ covariate design matrix.

### 2.2 *Untransformed Association Test*

The untransformed association test (UAT) is derived from the normal linear model:

$$\boldsymbol{y} = \boldsymbol{X}\boldsymbol{\beta}_x + \boldsymbol{g}\beta_g + \boldsymbol{\epsilon}, \tag{1}$$

where $\boldsymbol{\epsilon} \sim N\left(\boldsymbol{0}, \sigma^2 \boldsymbol{I}\right)$ is an $n \times 1$ residual vector, $\beta_g$ is the genetic effect, and $\boldsymbol{\beta}_x$ is the covariate effect. Define the *error projection matrix*, $\boldsymbol{P}_x = \boldsymbol{I} - \boldsymbol{X}(\boldsymbol{X}'\boldsymbol{X})^{-1}\boldsymbol{X}'$, and the *phenotypic residual* $\boldsymbol{e}_y = \boldsymbol{P}_x\boldsymbol{y}$, which is the residual after regressing $\boldsymbol{y}$ on $\boldsymbol{X}$, or the projection of $\boldsymbol{y}$ onto the orthogonal complement of the column space of $\boldsymbol{X}$. The efficient score for $\beta_g$ is $\mathcal{U}_{\beta_g} = \sigma^{-2}\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{y}$, and the score statistic assessing $H_0 : \beta_g = 0$ is:

$$T_U = \boldsymbol{y}'\boldsymbol{P}_x\boldsymbol{g}(\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{g})^{-1}\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{y}/\sigma^2, \tag{2}$$

which follows a $\chi_1^2$ distribution. Under $H_0$, an unbiased estimate of $\sigma^2$ is given by $\hat{\sigma}^2 = (n - p)^{-1}\boldsymbol{y}'\boldsymbol{P}_x\boldsymbol{y}$. The Wald statistic for assessing $H_0 : \beta_g = 0$ takes the same form as (2), save the residual variance is estimated as $\tilde{\sigma}^2 = (n - p - 1)^{-1}\boldsymbol{y}'\tilde{\boldsymbol{P}}_x\boldsymbol{y}$, where $\tilde{\boldsymbol{X}} = (\boldsymbol{X}, \boldsymbol{g})$ and $\tilde{\boldsymbol{P}}_x = \boldsymbol{I} - \tilde{\boldsymbol{X}}(\tilde{\boldsymbol{X}}'\tilde{\boldsymbol{X}})^{-1}\tilde{\boldsymbol{X}}' = \boldsymbol{P}_x - \boldsymbol{P}_x\boldsymbol{g}(\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{g})^{-1}\boldsymbol{g}'\boldsymbol{P}_x$.

If the normal residual assumption is relaxed to allow for an arbitrary distribution with mean zero and finite variance, then $T_U$ still follows an asymptotic $\chi_1^2$ distribution. Although (2) is eventually valid for any continuous trait with constant residual variance, when the residual

distribution exhibits excess skew or kurtosis, the sample size required for valid inference at $\alpha = 5 \times 10^{-8}$ may become impractically large. Moreover, as we will show, even in samples of sufficient size for valid inference, the UAT is generally less powerful than INT-based tests.

### 2.3 *Rank-Based Inverse Normal Transformation*

INT is a non-parametric mapping applicable to observations from any absolutely continuous distribution. The process may be decomposed into two steps. In the first, the observations are replaced by their fractional ranks. This is equivalent to transforming the observations by their empirical cumulative distribution function (ECDF) $\mathbb{F}_n$. If $W$ is any continuous random variable with CDF $F_W$, then the transformed random variable $U = F_W(W)$ is uniformly distributed (Casella and Berger, 2002). Since the empirical process $\mathbb{F}_n(\cdot)$ converges uniformly to the CDF $F_W$, in samples of sufficient size $\hat{U} = \mathbb{F}_n(W)$ is uniformly distributed (van der Vaart, 1998). After transformation by $\mathbb{F}_n$, the observations reside on the probability scale. In the next step, these probabilities are mapped to Z-scores using the *probit* function $\Phi^{-1}$. If $U$ is uniformly distributed, then $Z = \Phi^{-1}(U)$ follows the standard normal distribution. Consequently, in large samples, $\text{INT}(W) = \Phi^{-1}\{\mathbb{F}_n(W)\} \dot\sim N(0, 1)$, regardless of the initial distribution $F_W$.

In practice, an offset is introduced to ensures that all fractional ranks are strictly between zero and one, which in turn guarantees that all Z-scores are finite. Suppose that $W_i$ is observed for each of $N$ independent subjects. The modified INT is:

$$\text{INT}(W_i) = \Phi^{-1}\left\{\frac{\text{rank}(W_i) - c}{n + 1 - 2c}\right\}, \ c \in [0, 1/2]. \tag{3}$$

Hereafter we adopt the conventional *Blom* offset of $c = 3/8$ (Beasley et al., 2009). Since other choices for $c$ lead to Z-scores that are nearly linear transformations of one another, the choice of offset is considered immaterial.

2.4 *Direct Inverse Normal Transformation (D-INT)*

In direct INT (D-INT), the INT-transformed phenotype is regressed on genotype and co-variates according to the association model:

$$\boldsymbol{z} = \boldsymbol{X}\boldsymbol{\beta}_x + \boldsymbol{g}\beta_g + \boldsymbol{\epsilon}_D, \tag{4}$$

where $\boldsymbol{z} = \text{INT}(\boldsymbol{y})$ is the INT-transformed phenotype, and $\boldsymbol{\epsilon_D} \sim N(\boldsymbol{0}, \sigma_D^2 \boldsymbol{I})$. Model (4) is immediately comparable with model (1), the only difference being replacement of $\boldsymbol{y}$ by $\boldsymbol{z}$. Thus, the D-INT score statistic for assessing $H_0 : \beta_g = 0$ is:

$$T_D = \boldsymbol{z}'\boldsymbol{P}_x\boldsymbol{g}(\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{g})^{-1}\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{z}/\sigma_D^2. \tag{5}$$

A *p*-value is assigned with reference to the $\chi_1^2$ distribution. The score statistic estimates the residual variance $\sigma_D^2$ as $\hat{\sigma}_D^2 = (n-p)^{-1}\boldsymbol{z}'\boldsymbol{P}_x\boldsymbol{z}$, while the Wald statistic estimates the residual variance as $\tilde{\sigma}_D^2 = (n-p-1)^{-1}\boldsymbol{z}'\tilde{\boldsymbol{P}}_x\boldsymbol{z}$.

D-INT is adapted for data generating processes (DGPs) of the form:

$$\boldsymbol{y} = h(\boldsymbol{X}\boldsymbol{\beta}_x + \boldsymbol{g}\beta_g + \boldsymbol{\epsilon}_D^*), \tag{6}$$

where $\boldsymbol{\epsilon}_D^* \sim N(\boldsymbol{0}, \sigma_D^2 \boldsymbol{I})$, and $h(\cdot)$ is a rank-preserving transformation. An example is the log-normal phenotype, for which $h(t) = \exp(t)$. When $h(\cdot)$ is non-linear, the regression function $E(Y_i|\boldsymbol{x}_i, g_i)$ is non-linear in its parameters, and the residuals $\boldsymbol{\epsilon}_D^*$ have non-additive effects. However, there exists a transformed scale on which the mean model is linear and has additive normal residuals, namely: $h^{-1}(\boldsymbol{y}) = \boldsymbol{X}\boldsymbol{\beta}_x + \boldsymbol{g}\beta_g + \boldsymbol{\epsilon}_D$. Thus, under $H_0 : \beta_g = 0$, the efficient score for $\beta_g$ is $\mathcal{U}_{\beta_g} = \sigma_D^{-2}\boldsymbol{g}'\boldsymbol{P}_x h^{-1}(\boldsymbol{y})$. Since $h^{-1}(\cdot)$ is seldom known, D-INT makes the approximation $\text{INT}(y_i) = \Phi^{-1}\{\mathbb{F}_{y,n}(y_i)\} \approx \sigma_D^{-1} h^{-1}(y_i)$, where $\mathbb{F}_{y,n}$ is the *marginal* ECDF of the phenotype $Y_i$.

To justify this approximation, observe that under model (6), the *conditional* distribution of the transformed-scale residual $\epsilon_{D,i}^* = h^{-1}(y) - \boldsymbol{x}_i\boldsymbol{\beta}_x - g_i\beta_g$, given $\boldsymbol{x}_i$ and $g_i$, is $F_\epsilon(\epsilon_{D,i}^*|\boldsymbol{x}_i, g_i) =$

$\Phi(\epsilon_{D,i}^*/\sigma_D)$. The marginal and conditional CDFs of $Y_i$ are related via:

$$F_y(y_i) = \int \Phi\Big[\sigma_D^{-1}\{h^{-1}(y) - \boldsymbol{x}_i\boldsymbol{\beta}_x - g_i\beta_g\}\Big] dF(\boldsymbol{x}_i, g_i), \tag{7}$$

where $F(\boldsymbol{x}_i, g_i)$ is the joint CDF of $\boldsymbol{x}_i$ and $g_i$. The empirical counterpart to (7) is:

$$\mathbb{F}_{y,n}(y_i) = \frac{1}{n}\sum_{i=1}^{n} \Phi\big[\sigma_D^{-1}\{h^{-1}(y_i) - \boldsymbol{x}_i'\boldsymbol{\beta}_x - g_i\beta_g\}\big].$$

Under the strong $H_0 : (\boldsymbol{\beta}_x = 0)$ *and* $(\beta_g = 0)$, $\mathbb{F}_{y,n}(y_i)$ converges to $\Phi\{\sigma_D^{-1}h^{-1}(y_i)\}$, such that the D-INT approximation $\text{INT}(y_i) = \Phi^{-1}[\mathbb{F}_{y,n}(y_i)] \approx \sigma_D^{-1}h^{-1}(y_i)$ is asymptotically exact. Under the standard $H_0 : \beta_g = 0$, the approximation is accurate when $\boldsymbol{\beta}_x \approx \boldsymbol{0}$.

### 2.5 *Indirect Inverse Normal Transformation (I-INT)*

In indirect INT (I-INT), the phenotype is first regressed on covariates to obtain residuals, then the INT-transformed phenotypic residuals are regressed on genotype. Specifically, I-INT is based on the association model:

$$\tilde{\boldsymbol{z}} = \boldsymbol{e}_g\beta_g + \boldsymbol{\epsilon}_I, \tag{8}$$

where $\tilde{\boldsymbol{z}} = \text{INT}(\boldsymbol{e}_y)$ is the INT-transformed phenotypic residual (i.e. $\boldsymbol{e}_y = \boldsymbol{P}_x\boldsymbol{y}$), $\boldsymbol{e}_g = \boldsymbol{P}_x\boldsymbol{g}$ is the *genotypic residual*, which is the residual after regressing $\boldsymbol{g}$ on $\boldsymbol{X}$, and $\boldsymbol{\epsilon}_I \sim N(\boldsymbol{0}, \sigma_I^2\boldsymbol{I})$. The I-INT score statistic for assessing $H_0 : \beta_g = 0$ takes the form:

$$T_I = \tilde{\boldsymbol{z}}'\boldsymbol{P}_x\boldsymbol{g}(\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{g})^{-1}\boldsymbol{g}'\boldsymbol{P}_x\tilde{\boldsymbol{z}}/\sigma_I^2. \tag{9}$$

A $p$-value is assigned with reference to the $\chi_1^2$ distribution. The score statistic estimates the residual variance as $\hat{\sigma}_I^2 = n^{-1}\tilde{\boldsymbol{z}}'\tilde{\boldsymbol{z}} = 1$, while the Wald statistic estimates the residual variance as $\tilde{\sigma}_I^2 = (n-1)^{-1}\tilde{\boldsymbol{z}}'\boldsymbol{P}_{e_g}\tilde{\boldsymbol{z}}$, where $\boldsymbol{P}_{e_g} = \boldsymbol{I} - \boldsymbol{e}_g(\boldsymbol{e}_g'\boldsymbol{e}_g)^{-1}\boldsymbol{e}_g' = \boldsymbol{I} - \boldsymbol{P}_x\boldsymbol{g}(\boldsymbol{g}'\boldsymbol{P}_x\boldsymbol{g})^{-1}\boldsymbol{g}'\boldsymbol{P}_x$.

Since INT is a non-linear transformation, the INT transformed phenotypic residuals $\tilde{\boldsymbol{z}}$ are no longer orthogonal to the covariates. That is, the correlation between $\tilde{\boldsymbol{z}}$ and the columns of $\boldsymbol{X}$ is non-zero. Consequently, secondary adjustment for covariates has been recommended Sofar et al. (2019), as in the association model:

$$\tilde{\boldsymbol{z}} = \boldsymbol{X}\beta_x + \boldsymbol{g}\beta_g + \boldsymbol{\epsilon}_I. \tag{10}$$

We demonstrate that the score statistic from model (10), which adjusts twice for covariates, is equivalent to the score statistic from model (8), which instead adjusts for genotypic residuals.

I-INT is adapted for a DGP of the form:

$$\boldsymbol{y} = \boldsymbol{X}\boldsymbol{\beta}_x + \boldsymbol{g}\beta_g + \boldsymbol{\varepsilon}, \tag{11}$$

where the residuals $\boldsymbol{\varepsilon} \sim f_\varepsilon(\cdot)$ follow an arbitrary continuous distribution with mean zero and constant finite variance. Under (11), $F(y_i|\boldsymbol{x}_i, g_i) = F_\varepsilon(y_i - \boldsymbol{x}_i'\boldsymbol{\beta}_x - g_i\beta_g)$, such that under the strong null $H_0 : (\boldsymbol{\beta}_x = 0)$ *and* $(\beta_g = 0)$, the DGPs in (11) and (6) are equivalent.

To motivate I-INT, we begin by showing that the efficient score for $\beta_g$ from model (11) is consistently estimated by the score for $\beta_g$ from the model $\boldsymbol{e}_y = \boldsymbol{e}_g\beta_g + \boldsymbol{\varepsilon}$. Observe that, for any $f_\varepsilon$, the ordinary least squares estimator $\tilde{\beta}_x = (\boldsymbol{X}'\boldsymbol{X})^{-1}\boldsymbol{X}'(\boldsymbol{y} - \boldsymbol{g}\beta_g)$ remains consistent for $\boldsymbol{\beta}_x$. Thus, the profile log likelihood from (11) is consistently estimated by:

$$\tilde{\ell}_p(\beta_g) = \ln f_\varepsilon(\boldsymbol{y} - \boldsymbol{X}\tilde{\boldsymbol{\beta}}_x - \boldsymbol{g}\beta_g) = \ln f_\varepsilon\{\boldsymbol{P}_x(\boldsymbol{y} - \boldsymbol{g}\beta_g)\} \tag{12}$$

Letting $\mathcal{U}_\varepsilon(\cdot) = \partial \ln\{f_\varepsilon(\varepsilon)\}/\partial\varepsilon$, the efficient score for $\beta_g$ from (11) is consistently estimated by the gradient of (12), which is $\tilde{\mathcal{U}}_{\beta_g} = -\boldsymbol{g}'\boldsymbol{P}_x\mathcal{U}_\varepsilon\{\boldsymbol{P}_x(\boldsymbol{y} - \boldsymbol{g}\beta_g)\}$.

Now consider the following model for the phenotypic residual:

$$\boldsymbol{e}_y = \boldsymbol{X}\boldsymbol{\alpha}_x + \boldsymbol{g}\beta_g + \boldsymbol{\varepsilon}, \tag{13}$$

where $\boldsymbol{\varepsilon}$ is distributed as before. The profile log likelihood for $\beta_g$ in (13), with $\boldsymbol{\alpha}_x$ evaluated at the consistent estimator $\tilde{\boldsymbol{\alpha}}_x = (\boldsymbol{X}'\boldsymbol{X})^{-1}\boldsymbol{X}'(\boldsymbol{e}_y - \boldsymbol{g}\beta_g)$, coincides with the profile log likelihood in (12). Moreover, the log likelihood for $\beta_g$ from the following model, which relates $\boldsymbol{e}_y$ to the genotypic residual $\boldsymbol{e}_g$, is also identical to (12):

$$\boldsymbol{e}_y = \boldsymbol{e}_g\beta_g + \varepsilon. \tag{14}$$

Thus, the score for $\beta_g$ from (14) is consistent for the efficient score for $\beta_g$ from (11).

Now, under $H_0 : \beta_g = 0$, model (14) and model (13) with $\boldsymbol{\alpha}_x$ evaluated at its least squares estimate $\tilde{\boldsymbol{\alpha}}_x = \boldsymbol{0}$, both reduce to $\boldsymbol{e}_y = \boldsymbol{\varepsilon}$. Model (14), together with the observation that

$\tilde{z} = \text{INT}(e_y) = \text{INT}(\varepsilon) \stackrel{\cdot}{\sim} N(\mathbf{0}, \boldsymbol{I})$, motivate the I-INT association model in (8). Moreover, under $H_0 : \beta_g = 0$, the distributional assumption in (8) is asymptotically exact, with $\sigma_I^2 = 1$.

### 2.6 *Omnibus Inverse Normal Transformation Test (O-INT)*

As shown in the simulation studies, both D-INT and I-INT robustly controlled the type I error. However, neither D-INT nor I-INT was uniformly most powerful. We therefore propose combining the two approaches into a robust and broadly well powered omnibus test. The omnibus statistic is constructed using the method of Cauchy aggregation, in which the p-values from related hypothesis tests are converted to standard Cauchy random deviates then combined (Liu and Xie, 2019; Liu et al., 2019).

Let $p_D$ and $p_I$ denote the p-values from D-INT and I-INT. Define the O-INT statistic as:

$$T_O = -\frac{1}{2}\left\{F_C^{-1}(p_D) + F_C^{-1}(p_I)\right\} = \frac{1}{2}\tan\left\{\pi(0.5 - p_D)\right\} + \frac{1}{2}\tan\left\{\pi(0.5 - p_I)\right\}, \qquad (15)$$

where $F_C^{-1}(u) = \tan\left\{\pi(u - 0.5)\right\}$ is the inverse CDF of the standard Cauchy distribution. Under $H_0 : \beta_g = 0$, $p_D$ and $p_I$ are each uniformly distributed, such that $F_C^{-1}(p_D)$ and $F_C^{-1}(p_I)$ are standard Cauchy. Since the Cauchy distribution is symmetric and closed with respect to convolution, the omnibus statistic $T_O = -\{F_C^{-1}(p_D) + F_C^{-1}(p_I)\}/2$ is again standard Cauchy in the tail (Liu and Xie, 2019; Liu et al., 2019), even though $p_D$ and $p_I$ are in general positively correlated. Based on this fact, the p-value of the O-INT statistic (15) is:

$$p_O = P[T_O > t_O] = 1 - F_C(t_O) = \frac{1}{2} - \frac{1}{\pi}\arctan\left(t_O\right),$$

where $t_O$ is the observed value of $T_O$.

## 3. Simulation Studies

### 3.1 *Simulation Methods*

Extensive simulations were conducted to evaluate the size (i.e. type I error) and power of the UAT and INT-based association tests (D-INT, I-INT, O-INT). Genotypes exhibiting

linkage disequilibrium were randomly sampled from unrelated subjects in the UK Biobank. Simulated covariates included age and sex. Age was drawn from a gamma distribution with mean 50 and variance 10. Sex was drawn from a Bernoulli distribution with proportion $1/2$. To emulate population structure, the top 3 PCs of the empirical genetic relatedness matrix were included among covariates.

For size simulations, a subject-specific linear predictor $\eta_i$ was generated as $\eta_i = \boldsymbol{x}_i'\boldsymbol{\beta}_x$, where $\boldsymbol{x}_i$ included an intercept, age, sex, and 3 genetic PCs. Population regression coefficients were drawn independently on each simulation replicate from normal distributions with mean zero. The variance was tuned such that the proportion of phenotypic variation explained (PVE) by age and sex was 20%, and the PVE by PCs was 5%. For power simulations, the linear predictor included an additional contribution from genotype, and the PVE by genotype (i.e. the heritability) was 0.5%.

Phenotypes were generated either from models with additive residuals, as in $y_i = \eta_i + \epsilon_i$, or from non-linear transformations of such models, as in $y_i = h(\eta_i + \epsilon_i)$. Here, we report on four representative traits: three with additive residuals, and one with multiplicative residuals. The additive models were: (1) a reference trait, with $N(0,1)$ residuals; (2) a skewed trait, with $\chi_1^2$ residuals; and (3) a kurtotic trait, with $t_3$ residuals. In all cases, the residual distribution was centered and scaled to have mean zero and unit variance. For the multiplicative model, a log-normal phenotype was generated by exponentiating a latent normal trait: $y_i = \exp(\eta_i + \epsilon_i)$, where $\epsilon_i \sim N(0,1)$.

### 3.2 *Size Simulations*

A total of $R = 10^8$ simulation replicates were performed under $H_0 : \beta_g = 0$ at samples sizes of $N \in \{10^3, 10^4, 10^5\}$. On each simulation replicate, the four phenotypes (normal, skewed, kurtotic, log-normal) were generated independently and tested for association with genotype by each of the four association methods (UAT, D-INT, I-INT, O-INT).

[Figure 1 about here.]

The uniform QQ plots in Figure 1 summarize the distribution of association p-values at sample size $N = 10^3$ for each combination of phenotype (row) and association test (column). All association tests performed well against the normal phenotype (row 1), providing uniformly distributed p-values. The same was observed for increasing sample size $N \in \{10^4, 10^5\}$ (Web Figures S1-2). The UAT (column 1) exhibited inflated type I error against the non-normal phenotypes. Inflation was most severe at sample size $N = 10^3$, and attenuated with increasing sample size. In contrast, the INT-based tests provided valid p-values when applied to the non-normal phenotypes, even at sample size $N = 10^3$. D-INT maintained the type I error across all scenarios, while I-INT exhibited slight deflation in the case of the log-normal phenotype. Deflation was ameliorated with increasing sample size (Web Figures S1-2).

UAT incurred the most inflation against the log-normal (column 1, row 4). This is likely because the standard linear model is misspecified for a phenotype in which the residuals have multiplicative effects. Among the INT-based tests, I-INT exhibited slight deflation against the log normal phenotype. However, this does not invalidate inference. The p-values from D-INT and O-INT were uniformly distributed.

For the skewed $\chi_1^2$ phenotype (row 2), the UAT exhibited inflation at sample size $N = 10^3$, while the INT-based tests effectively maintained the type I error. By sample size $N = 10^4$, all methods performed well. For kurtotic $t_3$ phenotype (row 3), the UAT again exhibited inflation at sample size $N = 10^3$, and the inflation was still evident at sample size $N = 10^4$, notwithstanding the fact that the UAT is correctly specified for this phenotype. The INT-based methods performed well against the $t_3$ phenotype at all sample sizes.

[Table 1 about here.]

Type I error estimates at $\alpha = 10^{-6}$ and sample sizes $N \in \{10^3, 10^4, 10^5\}$ are presented in Table 1. For all non-normal phenotypes, the UAT had substantially inflated type I error

at sample size $N = 10^3$. This includes the skewed $\chi_1^2$ and kurtotic $t_3$ phenotypes, for which the UAT should provide asymptotically valid inference. Although the type I error rate approached its nominal level with increasing sample size, for the kurtotic and log-normal phenotypes the UAT still exhibited excess type I error at $N = 10^5$. For the non-normal phenotypes, D-INT generally provided nearly the nominal type I error, while I-INT was slightly conservative. However, this does not imply I-INT is less powerful for these phenotypes (see below). For all phenotypes and sample sizes considered, the omnibus test provided nominal control of the type I error.

### 3.3 *Power Simulations*

A total of $R = 10^6$ simulation replicates were performed under $H_A : \beta_g \neq 0$ at samples sizes of $N \in \{10^3, 10^4\}$. On each replicate, a randomly selected locus served as the causal locus. As before, the four phenotypes were generated independently and tested for association with genotype by each of the four association methods. The PVE by genotype at the causal locus was 0.5%. Power is considered even for the UAT, which did not consistently control the type I error, because this approach is still often applied in practice.

[Figure 2 about here.]

Figure 2 presents power at $\alpha = 10^{-6}$ for $N = 10^4$, and the relative efficiency (RE) of the INT-based tests to the UAT. RE was calculated as the ratio of the $\chi_1^2$ non-centrality parameters. This performance metric has the advantage of not depending on either the particular sample size or $\alpha$ level examined. For the normal phenotype, the UAT is theoretically most powerful. However, the INT-based tests were fully efficient, achieving a relative efficiency of one. Despite having inflated type I error under the null hypothesis, the UAT was consistently least powerful for detecting true associations with the non-normal phenotypes. Since the relative efficiencies of the INT-based tests always exceeded one for the non-normal phenotypes, this conclusion is expected to extend across sample sizes and

significance thresholds. For the log normal phenotype, D-INT was most powerful, achieving twice the efficiency of the UAT. For this phenotype, the log transform is theoretically optimal. Since the log transform maps the log normal phenotype to a normal phenotype, the power of the log transform against the log normal phenotype is identical to the power of the UAT against the normal phenotype. Comparing the power of D-INT against the log normal phenotype with that of UAT against the normal phenotype, we observe that D-INT attains optimal power.

For the skewed $\chi_1^2$ phenotype, I-INT was most powerful, achieving over 6 times the efficiency of the UAT, and over twice the efficiency of D-INT. For the kurtotic $t_3$ phenotype, the efficiency gains provided by the INT-based tests were more modest yet still noteworthy, at 53% for the I-INT and 38% for D-INT. By synthesizing D-INT and I-INT, the omnibus approach (O-INT) aims to provide a test that is well powered regardless of the phenotype's generative model. The power and relative efficiency of O-INT were intermediate between those of D-INT and I-INT. Since O-INT is a compromise between two complementary methods, it cannot be expected to attain optimal efficiency. However, for all phenotypes, the efficiency of O-INT was within 2% of the most efficient test considered. Thus, O-INT achieves robustness to the underlying data generating mechanism with little to no loss of efficiency.

## 4. Application to UK Biobank

### 4.1 *Application Methods*

We conducted GWAS of spirometry phenotypes within the UK Biobank (UKB). To mitigate confounding due to population structure, our study population was restricted to unrelated subjects of white, British ancestry. The phenotypes were forced expiratory volume in 1 second (FEV1), forced vital capacity (FVC), the FEV1 to FVC ratio (FEV1/FVC), and

the logarithm of peak expiratory flow (lnPEF). Our analyses focused on 360,761 directly genotyped, as opposed to imputed, autosomal SNPs, with sample minor allele frequencies exceeding 5%, and a per locus missingness rates of less than 10%. Covariates included an intercept, age, sex, BMI, two orthogonal polynomials in height, genotyping array, and 20 genetic PCs. Each locus was tested individually for association with the four spirometry phenotypes (FEV1, FVC, FEV1/FVC, lnPEF). The results were greedily 'clumped' in `PLINK` (Purcell et al., 2007) using a 1000 kb radius and an $r^2$ threshold of 0.2. The overall analysis consisted of $N = 292K$ subjects that met all inclusion criteria. A subgroup analyses was conducted among subjects with physician diagnosed asthma ($N = 29K$).

### 4.2 *Empirical Type I Error*

LD score regression (LDSC) was performed to assess inflation of the association test statistics due to confounding bias (Bulik-Sullivan et al., 2015). Briefly, in LDSC the test statistic for each locus is regressed on a local measure of linkage disequilibrium. An intercept exceeding one suggests inflation, whereas an intercept falling below one suggests deflation. The results from applying LDSC in the overall sample and in the asthmatic subgroup are presented in Web Tables S1-S2. Overall, there was no evidence of residual confounding due to population structure. Therefore, under the null hypothesis of no genetic effects, the marginal distribution of each spirometry trait is expected to be independent of genotype.

The empirical type I error of the association tests was assessed via a permutation analysis. Genotypes were first regressed on covariates to obtain residuals, then the genotypic residuals were permuted. Regressing genotypes on covariates accounts for potential confounding of the genotype-phenotype relationship, while permuting the genotypic residuals breaks the association between genotype and the spirometry traits, imposing the null hypothesis of no genetic effect. Since each subject retains their original covariates, this analysis preserves the association between covariates and the spirometry traits. Uniform QQ plots for the

association p-values after permutation are presented in Web Figure S5. For all association methods (columns) and spirometry traits (rows), the p-values were uniformly distributions, suggesting nominal control of the type I error for the observed residual distributions.

### 4.3 *Empirical Discovery and Efficiency Gains*

[Table 2 about here.]

Table 2 presents the average $\chi_1^2$ statistics across all loci that reached genome-wide significance ($\alpha = 5 \times 10^{-8}$) in the overall sample according to at least one of the association methods. Average $\chi_1^2$ for the asthmatic subgroup are presented in Web Table S3. The empirical *efficiency gain* (O-INT vs. UAT) was defined as the ratio of the non-centrality parameters minus one, where the non-centrality parameters were estimated using loci that reached significance according to at least one association method:

$$\text{Efficiency Gain} = \frac{\left(\bar{\chi}_{1,\text{O-INT}}^2 - 1\right)}{\left(\bar{\chi}_{1,\text{UAT}}^2 - 1\right)} - 1.$$

In all cases the average $\chi_1^2$ statistics of the INT-based tests exceeded those of UAT, both in the overall analysis and in the asthmatic subgroup. Table 2 also presents the counts of genome-wide significant associations after LD 'clumping' to reduce redundant signals. Counts for the asthmatic subgroup are presented in Web Table S3. The empirical *discovery gain* (O-INT vs. UAT) was defined as:

$$\text{Discovery Gain} = \frac{n_{\text{OINT}} - n_{\text{UAT}}}{n_{\text{OINT} \cup \text{UAT}}},$$

where $n_{\text{OINT}}$ is the number of associations identified by O-INT only, $n_{\text{UAT}}$ is the number of associations identified by UAT only (if any), and $n_{\text{OINT} \cup \text{UAT}}$ is the number of associations identified by either O-INT or UAT. In all cases, the INT-based tests discovered more independent (at $r^2 = 0.2$) associations with the target phenotype than the UAT. All associations that reach genome-wide significance in the asthmatic subgroup, according to either UAT or the INT-based tests, reached significance according to O-INT in the overall sample.

The efficiency and discovery gains were more dramatic for those traits whose residuals were less normally distributed. However, the INT-based tests were consistently more powerful than the UAT, even when the normal residual assumption was not unreasonable. Consistent with the simulations, the power of O-INT was always intermediate between that of D-INT and I-INT. For a given trait, the number of discoveries by O-INT was generally closer to the number of discoveries by the more powerful of D-INT and I-INT.

## 5. Discussion

In this paper, we have systematically investigated the utility of different INT-based association tests for GWAS of quantitative traits with non-normally distributed residuals. We formally defined the Direct (D-INT) and Indirect (I-INT) INT-based tests, demonstrating that these approaches are adapted to different underlying data generating processes. Our simulations confirmed that D-INT was most effective when the observed trait could have arisen from a rank-preserving transformation of a latent normal trait. I-INT was most effective when the observed trait is linear in covariates, but the residual distribution is skewed or kurtotic. Under the null hypothesis of no genetic effect, the two approaches are approximately equivalent when the covariate effects are small.

For non-normally distributed quantitative traits, both D-INT and I-INT provide better control of the type I error and higher power than the untransformed association test (UAT). We proposed an INT-based omnibus test (O-INT) as a robust and powerful approach to GWAS of quantitative traits with non-normally distributed residuals. O-INT is applicable to any trait with continuous residuals, and is nearly as powerful as deploying D-INT vs. I-INT based on prior knowledge of the underlying data generating mechanism. Moreover, O-INT was more powerful than the UAT even when the normal residual assumption was plausible. All INT-based tests (D-INT, I-INT, O-INT) have been implemented in the R package `RNOmni`,

which is available on CRAN. We further demonstrated the utility of INT-based association tests in GWAS of spirometry traits from the UK Biobank.

In D-INT, the INT is applied directly to the phenotype, and the transformed phenotype is regressed on genotype and covariates. Under the strong null hypothesis of no genetic or covariate effects, D-INT is asymptotically exact, and when the covariate effects are small, D-INT holds approximately. I-INT is a two-stage procedure. Different variants of I-INT have been considered in the literature. In all approaches, the phenotype is first regressed on covariates to obtain residuals. In the second stage, the INT-transformed residuals are regressed on genotype, with or without a secondary adjustment for genetic PCs. To provide guidance on which approach to use in practice, we formally derived I-INT, starting from the assumption that the observed phenotype follows a linear regression model with a non-normally distributed residual. Our derivations indicate that, during the second stage of I-INT, the transformed phenotypic residuals should be regressed on genotypic residuals, which are the residuals obtained by regressing genotype on covariates. This second stage is equivalent to regressing the INT-transformed phenotypic residuals on genotype while performing a secondary adjustment for covariates. Therefore, all covariates, including genetic PCs, should be included in both the first and second stage regressions. Under the standard null hypothesis of no genetic effects, I-INT is asymptotically exact, and under the strong null hypothesis of neither genetic nor covariate effects, D-INT and I-INT are asymptotically equivalent.

A limitation of INT-based tests is the restriction to absolutely continuous phenotypes. The INT cannot ensure asymptotic normality of a distribution with discrete probability masses. Moreover, this paper has focused on GWAS of independent subjects. However, INT-based tests can be extended to the correlated data setting using linear mixed models (LMMs). We plan to develop INT-based tests for LMMs in which that correlation across related subjects is modeled via a random effect whose covariance pattern depends on the genetic relatedness

matrix (Kang et al., 2010; Loh et al., 2015; Chen et al., 2016). A similar modeling strategy can accommodate longitudinal phenotypes arising from repeated measurements on the same subjects across time (Chen et al., 2019).

## Acknowledgments

This work was supported by F31 HL140822 (to Z. M.); by R35 HL135818 (to S. R.); and by R35 CA197449, P01-CA134294, U01-HG009088, U19-CA203654, and R01-HL113338 (to X. L.).

## Software

The association tests described in this paper (D-INT, I-INT, O-INT) are available in the R package on Github: `https://github.com/zrmacc/RNOmni` and on CRAN: `https://cran.r-project.org/web/packages/RNOmni/index.html`. Links to this and related software may also be found at `https://content.sph.harvard.edu/xlin/software.html`.

## Supporting Information

Web Appendix A, referenced in Sections 3 and 4, is available with this paper at the Biometrics website on Wiley Online Library.

## References

Abbott, L., Bryant, S., Churchhouse, C., Ganna, A., Howrigan, D., et al. (2017). http://www.nealelab.is/uk-biobank/.

Barber, M. J., Mangravite, L. M., Hyde, C. L., Chasman, D. I., Smith, J. D., et al. (2010). Genome-wide association of lipid-lowering response to statins in combined study populations. *PLOS One* **5,** e9763.

Beasley, T. M., Erickson, S., and Allison, D. B. (2009). Rank-based inverse normal transformations are increasingly used, but are they merited? *Behavioral Genetics* **39,** 580–95.

Bulik-Sullivan, B. K., Loh, P. R., Finucane, H. K., Ripke, S., Yang, J., et al. (2015). Ld score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nature Genetics* **47,** 291–295.

Cade, B. E., Chen, H., Stilp, A. M., Gleason, K. J., Sofer, T., et al. (2016). Genetic associations with obstructive sleep apnea traits in hispanic/latino americans. *American Journal of Respiratory and Critical Care Medicine* **194,** 886–897.

Casella, B. and Berger, R. (2002). *Statistical Inference.* Duxbury/Thomson Learning, Pacific Grove, CA, 2 edition.

Chen, H., Huffman, J., Brody, J. A., et al. (2019). Efficient variant set mixed model association tests for continuous and binary traits in large-scale whole-genome sequencing studies. *American Journal of Human Genetics* **104,** 260–274.

Chen, H., Wang, C., Conomos, M. P., et al. (2016). Control for population structure and relatedness for binary traits in genetic association studies via logistic mixed models. *American Journal of Human Genetics* **98,** 653–666.

Consortium, G. et al. (2017). Genetic effects on gene expression across human tissues. *Nature* **550,** 204–213.

Emilsson, V., Thorleifsson, G., Zhang, B., Leonardson, A. S., Zink, F., et al. (2008). Genetics of gene expression and its effect on disease. *Nature* **452,** 423–428.

Kang, H. M., Sul, J. H., Service, S. K., et al. (2010). Variance component model to account for sample structure in genome-wide association studies. *Nature Genetics* **42,** 348–354.

Kettunen, J., Tukiainen, T., Sarin, A. P., Ortega-Alonso, A., Tikkanen, E., et al. (2012). Genome-wide association study identifies multiple loci influencing human serum metabo-

lite levels. *Nature Genetics* **44,** 269–76.

Liu, Y., Chen, S., Li, Z., Morrison, A. C., Boerwinkle, E., and Lin, X. (2019). Acat: A fast and powerful p value combination method for rare-variant analysis in sequencing studies. *The American Journal of Human Genetics* **104,** 410–421.

Liu, Y. and Xie, J. (2019). Cauchy combination test: a powerful test with analytic p-value calculation under arbitrary dependency structures. *Journal of the American Statistical Association* .

Loh, P. R., Tucker, G., Bulik-Sullivan, B. K., et al. (2015). Efficient bayesian mixed-model analysis increases association power in large cohorts. *Nature Genetics* **47,** 284–290.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P., De Bakker, P. I. W., Daly, M. J., et al. (2007). Plink: a tool set for whole-genome association and population-based linkage analyses. *The American Journal of Human Genetics* **81,** 559–575.

Repapi, E., Sayers, I., Wain, L. V., Burton, P. R., Johnson, T., et al. (2010). Genome-wide association study identifies five loci associated with lung function. *Nature Genetics* **42,** 36–44.

Scuteri, A., Sanna, S., Chen, W. M., Uda, M., Albai, G., et al. (2007). Genome-wide association scan shows genetic variants in the fto gene are associated with obesity-related traits. *PLOS Genetics* **3,** e115.

Sofar, T., Zheng, X., Gogarten, S., Laurie, C., Grinde, K., et al. (2019). A fully adjusted two-stage procedure for rank-normalization in genetic association studies. *Genetic Epipemiology* pages 1–13.

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., et al. (2015). Uk biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLOS Medicine* **12,** e1001779.

van der Vaart, A. W. (1998). *Asymptotic Statistics*. Cambridge University Press, 1st edition.

**Figure 1. Simulation Studies of the Type I Error Rate at Sample Size $N = 10^3$ across $R = 10^8$ Simulation Replicates.** Rows correspond to different phenotype distributions. The first is normal, the second skewed $(\chi_1^2)$, the third kurtotic $(t_3)$, and the fourth log-normal. Columns correspond to different association tests. The first is the untransformed association test (UAT), the second is the direct inverse normal transformation (D-INT), the third is indirect INT (I-INT), and the fourth column is omnibus INT (O-INT).

**Figure 2. Simulation Studies of Power and Relative Efficiency at Sample Size $N = 10^4$ across $R = 10^6$ Simulation Replicates.** The proportion of phenotypic variation explained by genotype was 0.5%. The upper panel is power, as the proportion of null hypotheses correctly rejected at $\alpha = 10^{-6}$. The lower panel is relative efficiency, as the ratio of $\chi_1^2$ non-centrality parameters, which is independent of sample size and $\alpha$ level. Gray is the UAT, blue the D-INT, yellow the I-INT, red the O-INT. Within figures, from left to right, the first phenotype has normal residuals; the second has $\chi_1^2$ residuals; the third phenotype has $t_3$ residuals; and the log of the fourth phenotype has normal residuals. Note that this figure appears in color in the electronic version of this article.

**Table 1**
***Simulation Studies of the Empirical Type I Error Rate ($\times \mathbf{10^6}$) at $\boldsymbol{\alpha = 10^{-6}}$ across $\mathbf{R = 10^8}$ Simulation Replicates.*** *Size simulations were conducted under the $H_0 : \beta_g = 0$ at sample sizes ranging from $N = 10^3$ to $N = 10^5$. The following association tests were evaluated: the untransformed association test (UAT), direct inverse normal transformation (D-INT), indirect INT (I-INT), and omnibus INT (O-INT). Each test was applied to a normal phenotype, a skewed phenotype with $\chi_1^2$ residuals, a kurtotic phenotype with $t_3$ residuals, and a phenotype whose logarithm was normal.*

| Phenotype | Test | Sample Size | | |
| | | $N = 10^3$ | $N = 10^4$ | $N = 10^5$ |
|---|---|---|---|---|
| Normal | UAT | 1.04 | 0.93 | 1.03 |
| Normal | D-INT | 0.84 | 0.87 | 1.02 |
| Normal | I-INT | 0.97 | 0.93 | 1.00 |
| Normal | O-INT | 0.91 | 0.93 | 0.99 |
| Skewed | UAT | 8.03 | 1.87 | 1.43 |
| Skewed | D-INT | 1.20 | 1.10 | 1.05 |
| Skewed | I-INT | 0.67 | 0.84 | 0.89 |
| Skewed | O-INT | 1.10 | 1.01 | 0.98 |
| Kurtotic | UAT | 15.89 | 5.54 | 3.12 |
| Kurtotic | D-INT | 0.94 | 0.91 | 0.95 |
| Kurtotic | I-INT | 1.00 | 0.88 | 1.00 |
| Kurtotic | O-INT | 0.96 | 0.90 | 0.97 |
| Log-Normal | UAT | 59.34 | 11.01 | 7.52 |
| Log-Normal | D-INT | 0.74 | 1.02 | 1.02 |
| Log-Normal | I-INT | 0.40 | 0.40 | 0.43 |
| Log-Normal | O-INT | 0.55 | 0.74 | 0.76 |

**Table 2**
***Empirical Efficiency and Discovery Gains for Lung Function GWAS in the UK Biobank (N = 292K).***
*Genome-wide significance was declared at $\alpha = 5 \times 10^{-8}$. The average $\chi_1^2$ statistics are reported across all loci detected by at least one of the association tests. The empirical efficiency gain, comparing O-INT with UAT, is the ratio of the estimated $\chi_1^2$ non-centrality parameters minus 1. The counts of significant associations are reported after LD clumping within 1000 kb radii at $r^2 = 0.2$ to remove redundant signals. The discovery gain, comparing O-INT with UAT, is the ratio of the number of associations uniquely identified by O-INT to the total number of associations detected.*

## Average $\chi_1^2$

| Trait | UAT | D-INT | I-INT | O-INT | Efficiency Gain (%) |
|-------|-----|-------|-------|-------|---------------------|
| FEV1 | 56.77 | 57.55 | 64.69 | 63.59 | 12 |
| FVC | 37.14 | 46.91 | 51.47 | 50.59 | 37 |
| FEV1/FVC | 63.21 | 83.76 | 83.00 | 83.63 | 33 |
| lnPEF | 17.72 | 52.44 | 65.40 | 64.11 | 278 |

## Significant Associations

| Trait | UAT | D-INT | I-INT | O-INT | Discovery Gain (%) |
|-------|-----|-------|-------|-------|--------------------|
| FEV1 | 331 | 352 | 422 | 398 | 15 |
| FVC | 213 | 323 | 375 | 364 | 38 |
| FEV1/FVC | 450 | 653 | 649 | 652 | 28 |
| lnPEF | 39 | 202 | 270 | 251 | 79 |