Occurrence of marine sponge *Chelonaplysilla delicata* Pulitzer-Finali & Pronzato, 1999 (Porifera:

Demospongiae: Darwinellidae) from the Andaman Islands and the Indian Ocean with a baseline

information of epifauna on a mesophotic shipwreck

Rocktim Ramen Das[1,3]*, Titus Immanuel[1], Raj Kiran Lakra[1], Karan Baath[1] and Ganesh

Thiruchitrambalam[1]

[1]Department of Ocean Studies and Marine Biology, Pondicherry University, Brookshabad, Andaman

Islands, India; [2]Infinity Scuba Andaman's, Chidiyatapu, Andaman Islands, India; [3]Graduate School of

Engineering and Science, University of the Ryukyus, Nishihara, Okinawa, Japan.

Corresponding author: Rocktim Ramen Das; Email: k188604@cs.u-ryukyu.ac.jp

Titus Immanuel: titusimmanuel@gmail.com

Raj Kiran Lakra: rajkiranlakra@gmail.com

Karan Baath: karan123divewithme@gmail.com

T. Ganesh: tganesh.es@gmail.com

Occurrence of marine sponge *Chelonaplysilla delicata* Pulitzer-Finali & Pronzato, 1999 (Porifera: Demospongiae: Darwinellidae) from the Andaman Islands and the Indian Ocean with a baseline information of epifauna on a mesophotic shipwreck

Rocktim Ramen Das[1,3]*, Titus Immanuel[1], Raj Kiran Lakra[1], Karan Baath[1] and Ganesh Thiruchitrambalam[1]

[1]Department of Ocean Studies and Marine Biology, Pondicherry University, Brookshabad, Andaman Islands, India; [2]Infinity Scuba Andaman's, Chidiyatapu, Andaman Islands, India; [3]Graduate School of Engineering and Science, University of the Ryukyus, Nishihara, Okinawa, Japan.

Corresponding author: Rocktim Ramen Das; Email: k188604@cs.u-ryukyu.ac.jp

Abstract

During a biodiversity assessment on an upper mesophotic artificial reef of Andaman and Nicobar Islands (Shipwreck: HMIS Sophie Marie/HMIS SM), a single specimen of sponge *Chelonaplysilla delicata* was recorded. Our finding confirms the species taxonomy and highlights the current observation as a first report from the Andaman and Nicobar Islands and the Indian Ocean. The baseline diversity of epifaunal assemblage is further reported in this study.

Keywords: *Chelonaplysilla delicata*, Indian Ocean, Biodiversity, Porifera, Epifauna, Andaman and Nicobar Islands

Introduction

The Andaman and Nicobar Islands (ANI) (Figure 1) is an archipelago that lies in the eastern side of Bay of Bengal, bounded by the Andaman Sea on the west[1]. These islands support a diverse

coral reef ecosystem that harbours a plethora of marine organisms[2]. Most faunal groups that have been

studied extensively in these waters are large-sized, conspicuous and are of economic importance[3]

however, areas in the mesophotic depths remain poorly explored. Further, the study of sponge

distribution in the islands, though have existed since the early 19[th] century[4-11], recent works have

revealed many undocumented and new species[12-19] indicating its high diversity. Consisting of several

shipwrecks both in the shallow and mesophotic zones. These sunken structures act as an artificial reef

providing space for growth and establishment of various sessile marine communities creating a habitat

intricacy[20-22]. Being a dominant group in shipwreck driven ecosystem, poriferans are capable of

colonizing in a relatively short period[20, 23-27] along with which the substratum allows the growth of

other invasive and non-native fauna[28]. It is Interesting to note that these habitats remain less studied in

the mesophotic zones (F. Sinniger pers comm) thus a baseline assessment remains necessary, however

recent attempts to explore shallow wrecks in the area has revealed interesting results[22, 29-32].

Our findings document the presence of the marine sponge *Chelonaplysilla delicata* from the

Andaman and Nicobar Islands and the Indian Ocean along with providing baseline information relating

to the epifaunal assemblage on a mesophotic shipwreck.

Materials and Methods

The sponge *C.delicata* (Figure 2), collected from the shipwreck HMIS SM (Figure 2, 3)

during a survey conducted for documenting epifaunal diversity from February to March 2014. The

shipwreck is a 70m long Royal Indian navy minesweeper that sank in the year of 1942 (Figure 3). At a

depth of ≈33 meters, the wreck lies at the edge of the Macpherson strait near Chidiyatapu

(11º28'38.02"N 92º42'12.20"E), the southernmost tip of South Andaman Island (Fig. 1). Water

transparency and temperature were recorded with Secchi disc and dive calculator. Within 2 hours after

collection, the specimen was preserved in 100% ethanol. A surface peel of the easily separable cortex

of the specimen was removed and placed in xylene for 24 hours after which a permanent slide of the peel was mounted with DPX. A single fibre with its base and branches intact was removed from the sponge for species-level identification under a stereo microscope. (Figure 2B - D). The specimen was identified following Finali and Pronzato[33]. The preserved specimen is deposited in the National Zoological Collections (NZC) of the Andaman and Nicobar Regional Centre (ANRC), Zoological Survey of India (ZSI), Port Blair. Study maps were created using QGis.

Results

Taxonomy and Description

Phylum: Porifera

    Class: Demospongiae

        Subclass: Keratosa

            Order: Dendroceratida

                Family: Darwinellidae

                    Genus: *Chelonaplysilla*

                        Species: *C. delicata* Pulitzer-Finali & Pronzato, 1999

Materials Examined: 1 ex (Paratype)., ZSI/ANRC – 14321, India: Andaman Island: South Andaman: Chidiyatapu (11º28'38.02"N 92º42'12.20"E). Coll. RRD, 2014

Description: *C.delicata* predominantly thickly encrusting (< 10 mm) but has erect lobes that are about 4 - 5 cm high. The sponge surface is conulose, and the acute conules separated from each other

by 2 - 5 mm. Oscules 1 - 3 mm in diameter, flush with the surface and unevenly distributed all over on

sponge surface. The texture is soft collapsible and feeble. The fresh specimen was dark violet or purple

in colour and retained its colour even in the preserved condition. Sponge surface covered by structured

regular reticulation of sand and spicule detritus, which forms regular roundish or oval meshes of 90 -

155 µm. This reticulation is typical of the genus. Regular rounded fibrous pores, inhalant in nature are

enclosed within these rounded meshes (Figure 2D). The skeleton is dendritic, made up of pigmented

fibres fragile in nature with repeated branching that originate from a basal spongin plate (Fig. 4B and

4C) and extends towards the boundary. The primary fibre measured to be around 0.4 mm at its thickest.

Spicules are absent.

Distribution: India: ANI (South Andman, Present study). Elsewhere: Bismark Sea (Papua New

Guinea)[33], Indonesia (Sulawesi)[34], Palau[35], French Polynesia[36].

Similar Species: *C.delicata* is very similar to *C.erecta*[37]; however, the latter has fibres

anastomosing in nature whereas the thickness of fibres in *C.delicata* fades in diameter. Our specimen

was initially identified as *C.erecta*[30].

Remarks: The specimen mentioned in Finali and Pronzato[33] is gray whereas our specimen in

dark maroon in live condition.

Comments: The family Darwinellidae possesses sponging fibres with proper skeleton and

fibrous spicules[38,39]. It consists of four recognized genus and forty-seven accepted species (one under

"nomen nudum" status). *Chelonaplysilla* is the only genus, which is devoid of spicules but consists of a

fibrous dendritic skeleton that possesses distinct laminated bark surrounding a central pith region. A

structured and separable cortex that is reinforced by a delicate reticulation of sand grains[38]

distinguishes this genus.

Wreck Biodiversity: The surface of the wreck surveyed consisted of various sessile

communities[32], revised identification of which revealed the presence of encrusting Sponges, Tunicates,

Bryozoans, and Hydroids. Ahermatypic corals of family Dendrophylliidae (*Tubastrea*) were common

and easily visible (Figure 3). Solitary corals (*Heterocyathus*) were scattered in few vertical areas. Reef-

building corals were relatively less in abundance. The observed corals include *Favia*, *Symphyllia*,

Podabacia crustacea, and *Leptoseris*. A Gastropod (*Chicoreus* sp.) and few Crinoids were also present.

Poriferan families which were identified during the survey include Irciniidae (*Ircinia*), Chalinidae

(*Haliclona* (*Reniera*)); Thorectidae (*Hyrtios*), Iotrochotidae (*Iotrochota baculifera*), Thorectidae

(*Dactylospongia*). Tunicates comprised of Didemnidae (*Didemnum*), Perophoridae (*Perophora*) and

other unidentified sp. Macroalgae were absent however areas covered with a mix of turf algae and

encrusting crustose algae (ECA). The mean temperature in the area was 26.5º C.

Discussion

The Faunal organisms that thrives in Artificial Reefs (Shipwrecks) is an important part of the

marine community[27,40]. With increasing anthropogenic impacts on natural coral reef habitats, artificial

reefs are regarded as a successful alternative[25]. As a result, it becomes important to understand the

biological communities growing on these habitats[26]. Since a stable structural feature can lead a

centennial shipwreck to mimic a natural coral reef ecosystem[39,40], the necessity to investigate pre-

existing shipwrecks in a mesophotic zone can provide insights on these complex ecosystems. The

faunal assemblages in shipwrecks vary with horizontal and vertical orientation[39]. Such assemblages

enhance food resources for fishes, increasing its abundance thus improving diversity[20]. Sponges which

naturally occupy shipwrecks are one of the dominant organisms in such habitat, as evidenced by many

studies[20,21,26,27,41,42]. Their presence in large numbers also signifies advance stages of community

succession[20,43]. Sponges are known to play an important role in ecological recycling, habitat formation

for other organisms, acting as functional connecters between the benthic community and ocean

productivity and are also known to replace corals[44,45].

Environmental parameters like depth and light penetration play an important role in the

distribution and presence of communities in a shipwreck[20,39]. Lower light intensity in the study area can

contribute towards the abundance of poriferans and ahermatypic corals (*Tubastrea* sp.) (Figure 1), by

limiting macroalgal growth[32,40]. Yogesh-Kumar et al.[31] reported high sponge cover when studying

other wrecks in the region; however, the live coral coverage remains contrasting, indicating the role of

regional stressors and geolocation. Sponges that inhabit mesophotic areas mainly rely on plankton

feeding rather than photosymbionts and have higher growth rates due to the limited light intensity[46,47].

In the Caribbean's, sponge density seems to be directly proportional to depth[45]. Moreover, the

difference in current velocities affects the growth and development of filter feeders like sponges

present in the shipwrecks[40]. Higher current velocities through the Macpherson strait (Figure 1) can

effectively determine the faunal assemblage of HMIS SM but a long-term monitoring is necessary in

this regard.

The occurrence of a sponge from the deepest (Upper mesophotic region) and oldest wreck

(WWII) studied in the area sheds light on the vast and rich biodiversity that thrives on such habitats

and awaits much-needed attention from the scientific community. Technical gaps have always been a

hurdle to study mesophotic ecosystems but with the advancement of time, technologies like automated

underwater vehicle (AUV), remotely operated underwater vehicle (ROV), submersible's etc. can be

well utilized for the exploration of shipwrecks in such depths. It should be noted that the recent use of

ROV in ANI (see. Ramesh et al.[48]) and the documentation of Mesophotic reefs off Puducherry coast[49]

is an important step that will further lead towards the exploration of deeper ecosystems in the region.

Further, as only certain areas of the wreck were assessed, long-term repeated sampling, taking

structural heterogeny, hydrodynamics, and other environmental parameters into account will possibly give a detailed picture of the faunal assemblages.

Acknowledgements

The authors thank Dr. P.M Mohan Former HOD, DOSMB, Pondicherry University, Mr. G. Sakthivel for providing information on the wreck, Dr. S. Venu (DOSMB, Pondicherry University) for preliminary advice, Dr. E. Hirose (University of the Ryukyus, Japan) for identifying few Ascidians and to Dr. F. Sinniger (University of the Ryukyus, Japan) for providing information on the mesophotic ecosystem. RRD further acknowledges critical comments provided by Dr. K. Wangkulangkul (Prince of Shongkla University, Thailand) and to Dr. Yambem Tenjing Singh (National Centre for Sustainable Coastal Management, India) for valuable inputs and English correction on the early version of this manuscript.

References

1. Ubare, V.U. and Mohan, P.M., New Records and Range Extensions of Some Marine Sponges (Porifera: Demospongiae and Homoscleromorpha) from the Andaman Islands, India; Part of the Indo-Burma Biodiversity Hotspot. *Zool. Stud.,* 2018, DOI:10.6620/ZS.2018.57-03

2. Venkataraman K, *et al.,* 2012. Scleractinia of Andaman and Nicobar Islands. Kolkata: Zoological Survey of India.

3. Venkataraman, K. and Wafar, M., Coastal and Marine Biodiversity of India. *Indian J. Geomarine Sci.,* 2005, **34**(1): 57-75.

4. Schulze FE. 1902. An account of the Indian Triaxonia collected from the Royal Indian marine survey ship "Investigator" translated into English by R. von Lendenfeld. Trustees of the Indian museum, Calcutta, pp. 1-113

5. Schulze FE. 1904. Hexactinellida. Vol. 4 of the Wissenschaftliche Ergebnisse der Deutschen

Tiefsee-Expedition auf dem Dampfer" Valdivia" 1898-1899.

6. Dendy A. 1905. Report on the sponges collected by Professor Herdman at Ceylon in 1902.

Ceylon Pearl Oyster Fisheries, Supplementary reports Royal Society London **XVIII**: 57-246.

7. Annandale, N., Indian Boring Sponges of the Family Clionidae. *Rec. Indian Museum*, 1915, **11**:

1-24.

8. Dendy A, Burton M. 1926. Report on Some Deep-Sea Sponges from the Indian Museum

Collected by the R.I.M.S. 'Investigator'. Part I. Hexactinellida and Tetractinellida (PARS.). *Rec

Indian Museum* **28**(4): 225-248.

9. Burton, M. and Rao, H.S., Report on the Shallow-Water Marine Sponges in the Collection of

the Indian Museum. Part I. *Rec. Indian Museum,* 1932, **XXXXIV**: 299-358.

10. Burton M. 1937. Supplement to The Littoral Fauna of Krusadai Island in the Gulf of Manaar:

Porifera. 2 vols. Calcutta: Government Press.

11. Rao H. S. 1941. Indian and Ceylon Sponges in the Naturalistoriska Riksmuseet, Stockholm,

collected by K. Fistedt. *Rec Indian Museum* **XLIII**: 417–496.

12. Pattanayak JG. 2006. Marine Sponges of Andaman and Nicobar Islands, India. *Rec Zool

Surv India* **255**: 1–152.

13. Vinod K, *et al.,* 2007. An Account of Hexactinellid Sponge, *Hyalonema* (*Cyliconema*) *apertum

apertum* Collected from Andaman Waters. *JMBAI* **49**(2): 219-225.

14. Vinod K, *et al.,* 2012. *Semperella megaloxea* sp. nov. (Family: Pheronematidae): A new

hexactinellid sponge from Andaman waters, India. *Indian J Fish* **59**: 33-36

15. Immanuel T, Raghunathan C. 2011. First Record of the Sponge *Damiria toxifera*

(Poecilosclerida: Acarnidae) from the Andaman Sea, Andaman and Nicobar Islands. *Mar

Biodivers Rec* **4**: e32.

16. Immanuel T, *et al.,* 2014. An Updated Report on the Diversity of Marine Sponges of the

Andaman and Nicobar Islands. *In*: Venkataraman K, Sivaperuman C. (eds.) Marine Faunal

Diversity in India: Taxonomy, Ecology and Conservation, Elseiver, pp. 3-13.

17. Immanuel T, *et al.,* 2016. New Records of *Chondrilla australiensis* and *Chondrilla*

grandistellata (Demospongiae: Chondrosida: Chondrillidae) from Havelock Island, South

Andaman, India. *J Threat Taxa* **7**(14): 8167-8172.

18. Ubare VU, Mohan PM. 2016. A New Species of Genus *Plakortis* Schulze 1880 (Porifera:

Homoscleromorpha) from Badabalu, Andaman and Nicobar Islands, India. *Zool Stud* **55**(2):

doi:10.6620/ZS.2016.55-02

19. Periera P, Raghunathan C. 2018. Diversity of sponges in Marine protected areas of North

Andaman, India. *Thalassas*. https://doi.org/10.1007/s41208-018-0073-2

20. Walker SJ, *et al.,* 2007. Spatial Heterogeneity of Epibenthos on Artificial Reefs: Fouling

Communities in the Early Stages of Colonization on an East Australian Shipwreck. *Marine*

Ecology **28**: 435-445.

21. Lira SMA, *et al.,* 2010. Sessile and Sedentary Macrofauna from the Pirapama Shipwreck,

Pernambuco, Brazil. *Biota Neotrop **10**(9): 155-165.*

22. Mondal T, Raghunathan C. 2017. Shipwrecks in Andaman and Nicobar Islands: An Artificial

Habitat for Corals. *JMBAI* **59**(2): 92-101.

23. Aseltine-Neilson DA, *et al.*, 1999. Comparisons of Turf Communities from Pendleton Artificial

Reef, Torrey Pines Artificial Reef, and a Natural Reef using Multivariate Techniques. *Bull Mar*

Sci **65**(1): 37-57.

24. Clark S, Edwards AJ. 1999. An Evaluation of Artificial Reef Structures as Tools for Marine

Habitat Rehabilitation in the Maldives. *Aquat Conserv* **9**(1): 5-21.

25. Perkol-Finkel S, Benayahu Y. 2005. Recruitment of Benthic Organisms onto a Planned

Artificial Reef: Shifts in Community Structure One Decade Post Deployment. *Mar Environ*

Res **59**(2): 79–99.

26. Thanner SE, *et al.,* 2006. Development of Benthic and Fish Assemblages on Artificial Reef

Materials Compared to Adjacent Natural Reef Assemblages in Miami-Dade County,

Florida. *Bull Mar Sci* **78**(1): 57-70.

27. Amaral FMD, *et al.,* . 2009/2010. Benthic Macrofauna Inventory of Two Shipwrecks from

Pernambuco Coast, Northeastern of Brazil. *Revista Nordestina de Zoologia-Recife* **4**(1): 24-41.

28. Patro S, *et al.,* 2015. Snowflake coral, *Carijoa riisei* from Grand Island, Goa: a case of invasion

of an alien species or re-establishment of a native species.? *Curr Sci* 109(6): 1028-1030.

29. Hema M. 2013. Fish Assemblage on Artificial Reef and Natural Reef (A Comparative Study).

MSc Dissertation, Department of Ocean Studies and Marine Biology, Pondicherry University.

30. Das RR. 2014. An Inventory of Benthic Macrofauna from Two Shipwrecks along the Coast of

South Andaman. MSc Dissertation, Department of Ocean Studies and Marine Biology,

Pondicherry University.

31. Yogesh Kumar JS, *et al.,* 2015. An Assessment of Faunal Diversity and its conservation of

shipwrecks in Indian seas. *In*: Venkataraman K, Sivaperuman C (eds.) Marine Faunal Diversity

in India: Taxonomy, Ecology and Conservation, 441-450 USA: Elseiver, pp. 441-450.

32. Das RR, *et al.,* 2016. A preliminary macrobenthic study of world war two era wrecks along the

coast of South Andaman. *In*: Bijukumar A, Pradeep NS, Ajit Kumar KG, Rajendran PG. (eds.)

Perspectives on Biodiversity of India, Centre for Innovation in Science & Social Action

(CISSA), pp. 83-92.

33. Pulitzer-Finali G, Pronzato R. 1999. Horny Sponges from the North-Eastern Coast of Papua
New Guinea, Bismark Sea. *J Mar Biol Assoc U.K.* **79**(4): 593-607.

34. GBIF 2000

35. Ridley CP, *et al.,* 2005. Speciation and Biosynthetic variation in four dictyoceratid sponges and
their cyanobacterial symbiont, *Oscillatoria spongeliae*. *Chemistry and Biology* **12**: 397-406.

36. *Hall KA, et al., 2017. Sponges of Polynesia. Papeete (PYF): IRD. pp. 827.*

37. Tsurnamal M. 1967. *Chelonaplysilla erecta* n. sp. (Demospongiae, Keratosa) from
Mediterranean Coast of Israel. *Isr J Zool* **16**(2): 96-100.

38. Van Soest RWM. 1978. Marine Sponges from Curaçao and Other Caribbean localities Part I.
Keratosa. Studies on the Fauna of Curaçao and Other Caribbean Islands **56**(1): 1-94.

39. Bergquist PR, Cook SC. 2002. Family Darwinellidae Merejkowsky, 1879. In: Hooper JNA,
Van-Soest RWM. (eds.) Systema Porifera: A guide to the classification of Sponges, New York:
Springer, pp. 1068-1071.

40. Zintzen V, *et al.,* 2006. Epifaunal inventory of two shipwrecks from the Belgian Continental
Shelf. *Hydrobiologia* **555**(1): 207-219.

41. Perkol-Finkel S, *et al.,* 2006. Can Artificial Reefs Mimic Natural Reef Communities? The
Roles of Structural Features and Age. *Mar Environ Res* **61**(2): 121-135.

42. Diaz MC, Rützler K. 2001. Sponges: An Essential Component of Caribbean Coral Reefs. *Bull*
Mar Sci **69**(2): 535-546.

43. Cedro VR, *et al.,* 2007. Demospongiae (Porifera) of the shallow coral reefs of Maceió, Alagoas
State, Brazil. *In*: Custódio MR, Lôbo- Hajdu G, Muricy G. (eds.) Porifera research biodiversity,
innovation and sustainability, Rio de Janeiro: Museu Nacional, pp. 233-237.

44. Roberts DE, *et al.,* 2006. Structure and Dynamics of Sponge-Dominated Assemblages on

Exposed and Sheltered Temperate Reefs. *Mar Ecol Prog Ser* **321**: 19–30.

45. Lesser MP, Slattery M. 2013. Ecology of Caribbean Sponges: are Top-Down or Bottom-Up

Processes more Important?. *PloS one* **8**(11): e79799.

46. Sinniger F, *et al.,* 2016. Biodiversity of mesophotic coral ecosystems. *In*: Baker EK, Puglise

KA, Harris PT. (eds.) Mesophotic coral ecosystems A lifeboat for coral reefs.?, Arendal and

Nairobi: UNEP and GRID, pp. 50-58.

47. Baker E, *et al.,* 2016. What are mesophotic coral ecosystems.? *In*: Baker EK, Puglise KA,

Harris PT (eds.) Mesophotic coral ecosystems A lifeboat for coral reefs.?*,* Arendal and Nairobi:

UNEP and GRID, pp. 11-19.

48. Ramesh S, *et al.,* 2017. Application of Indigenously developed Remotely Operated Vehicle for

the study of Driving Parameters of Coral Reef Habitat of South Andaman Islands, India. *Curr*

Sci doi: 10.18520/cs/v113/i12/2353-2359.

49. Laxmilatha P, Jasmine S, Sreeram MP, Rengaiyan P. 2019. Benthic communities of mesophotic

coral ecosystem off Puducherry east coat of India. *Curr Sci* doi: 10.18520/cs/v116/i6/982-987.

Figure 1. Location of the study area (HMIS SM). (**A**) ANI, (**B**) South Andaman

Figure 2. *Chelonaplysilla delicata* [ZSI/ANRC-14321]: (A) Freshly collected specimen, (B) Branching fibres and basal sponging plate, (C) Closer view of pigmented, branching, dendritic spongin fibre, (D) Inhalant pores surrounded by rounded meshes reinforced by sand grains . Scale (A) 5mm (B) 2 mm, (C) 2 mm, (D) 155 μm.

288

Figure 3. A part of the wreck HMIS SM. (Arrow: high abundance of invasive *Tubastrea* cf.

micranthus)

292   **Figure 4.** Global distribution of *C.delicata* Pulitzer-Finali & Pronzato, 1999

 Table 1. Shipwrecks assessed in ANI

Wreck Name	Co-ordinates	Location	Date of Sinking	Depth (m)	Current Activities	Reference
SS Inchkeith	12°00'23.69"N 92°46'08.34"E	Kyd Island (South Andaman)	1955	14	Diving*	28,29,31
HMIS Sophie Marie	**11º28'38.02"N 92º42'12.20"E**	**Chidiyatapu (South Andaman)**	**1942**	**30 - 33**	**Diving***	29,31, **Current Study**
MV Mars	11°55'54.98"N 92°57'24.12"E	Havelock (Ritchie's Archipelago)	2006	10 - 16	Diving*	31
North Bay Wreck	11°43'00.56"N 92°45'60.60"E	Port Blair (South Andaman)	30 – 40 (yrs)	10	Diving* and Fishing	22,30
Peel Wreck	12°03'84.20"N 92°57'81.10"E	Havelock (Ritchie's Archipelago)	8 - 10	9 - 12.	Diving*	22,30

| Japan Wreck | 09°10'88.30"N 92°50'12.30"E | Car Nicobar (Nicobar Islands) | 40 - 50 | 28 | Fishing ground | 22,30 |
| Sinclair Bay Shipwreck | 11°39'873"N 92°45'488"E | Near Ross Island (South Andaman) | - | 8 | | 22 |

296 *Recreational/Water Tourism